



HAL
open science

Decreasing fallow duration in tropical slash-and-burn agriculture alters soil macro-invertebrate diversity: a case study in southern French Guiana

Jean-Pierre Rossi, Léonide Celini, Philippe Mora, Jerome Mathieu, Emmanuel Lapied, Johanne Nahmani, Jean-François Ponge, Patrick Lavelle

► **To cite this version:**

Jean-Pierre Rossi, Léonide Celini, Philippe Mora, Jerome Mathieu, Emmanuel Lapied, et al.. Decreasing fallow duration in tropical slash-and-burn agriculture alters soil macro-invertebrate diversity: a case study in southern French Guiana. *Agriculture, Ecosystems & Environment*, 2010, 135 (1-2), pp.148-154. <10.1016/j.agee.2009.08.012>. <hal-00493992>

HAL Id: hal-00493992

<https://hal.science/hal-00493992v1>

Submitted on 21 Jun 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons CC0 1.0 - Universal - International License

1
2
3
4
5
6
7
8
9 Decreasing fallow duration in tropical slash-and-burn
10 agriculture alters soil macro-invertebrate diversity: A
11 case study in southern French Guiana
12
13
14

15 J.P. Rossi^{*,a}, L. Celini^b, P. Mora^b, J. Mathieu^c, E. Lapied^d, J. Nahmani^e,
16 J.-F. Ponge^f, P. Lavelle^c
17
18

19 ^aINRA, UMR1202 BIOGECO, F-33612 Cestas, France

20 ^bUniversité Paris-XII, UMR 137 Biosol, F-94010 Créteil cedex, France

21 ^cUniversité Paris-VI, UMR 137 BioSol, F-93143 Bondy Cedex, France

22 ^dNorwegian University of Life Sciences, Department of Plant and Environmental
23 Sciences, Ås, Norway

24 ^eCNRS, UMR 7146 LIEBE, Université de Metz, F-57070 Metz, France

25 ^fMNHN, UMR 5176, F-91800 Brunoy, France
26
27
28
29

30 **Abstract**
31

32
33 In the humid tropics, slash-and-burn cultivation causes changes in the compo-
34 sition of soil biota communities. We investigated the soil macro-invertebrates
35 (body length ≥ 2 mm) in five sites, two at Maripasoula, an Aluku village
36 along the Maroni river (French Guiana), with short fallow (≈ 8 years), and
37 the other three at Elahe, a Wayana village along the same river, with long
38 fallow (≈ 25 years). We report observed species richness, the correspond-
39 ing estimates by bootstrap and its associated standard deviation. At both
40 sites the cultivation led to impoverished communities. The overall observed
41 species richness i.e. γ diversity was ca. twice as larger in Elahe than in
42 Maripasoula. The landscape at Maripasoula was dominated by highly dis-
43 turbed areas with the direct consequence that local species richness relied on
44
45
46
47
48
49
50
51
52
53

54
55 *Corresponding author

56 *Email address:* Jean-Pierre.Rossi@pierroton.inra.fr (J.P. Rossi)
57
58

59 *Preprint submitted to Agriculture, Ecosystems & Environment*

August 20, 2009
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9 colonization from an impoverished regional species pool. On the contrary,
10 in Elahe, crops formed small patches scattered across a landscape essentially
11 constituted of rich undisturbed or slightly disturbed forests hence higher γ
12 diversity. The proportion of rare species ranged from 44% to 54%. We
13 found 6 indicator species amongst which 5 were associated to the old sec-
14 ondary forest in Elahe and one, the earthworm *Pontoscolex corethrurus* was
15 associated to crop fields in Maripasoula (short fallow system). Results are
16 discussed in a landscape context in terms of conservation and management
17 of soil macrofaunal diversity in agro-ecosystems.

18
19 *Key words:* Soil macrofauna, species richness, slash-and-burn agriculture,
20 agriculture intensification, landscape, biodiversity.
21
22
23
24
25

26 27 28 29 30 31 **1. Introduction**

32
33
34 2 Soil invertebrates are key mediators of soil functions in agro-ecosystems.
35
36 3 They substantially affect many important processes that take place below-
37
38 4 ground like comminution and incorporation of litter into the soil, building and
39
40 5 maintenance of structural porosity and aggregation in soils through burrow-
41
42 6 ing, casting and nesting activities and control of microbial activities (Lavelle
43
44 7 et al., 2006, and references therein). Invertebrates therefore contribute to
45
46 8 the ecosystem services provided by soils and for this reason, they are in-
47
48 9 creasingly considered as a resource to be managed and protected. Amongst
49
50 10 soil biota, macrofauna (animals with body length ≥ 2 mm (Anderson and
51
52 11 Ingram, 1993)) are dramatically affected by cultural practices and various
53
54 12 authors have discussed the utility of managing their populations to improve
55
56 13 the sustainability of soil fertility especially in countries or regions where farm-

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

14 ers have limited access to mineral fertilizers (Matson et al., 1997; Brussaard
15 et al., 2007; Rossi and Blanchart, 2005).

16 In the tropics, the traditional slash-and-burn system (shifting cultivation)
17 consists of cutting the forest, burning the trees and settling familial agricul-
18 ture for several years. Long fallow periods follow the cropping period and
19 the regeneration of the vegetation combined with the recovery of soil fauna
20 contribute to restore soil organic content and structure which in turn affect
21 soil water and nutrient dynamics. These processes require a long fallow pe-
22 riod (Grandisson, 1997). Unfortunately, the changes from a traditional to
23 a permanent agriculture that accompany a population demographic growth
24 generally lead to a decrease in the fallow duration (Fleury, 1998). Ultimately,
25 the cropping period is followed by the establishment of permanent pastures
26 used for cattle ranching instead of fallows. This is the case for huge surfaces
27 of land in Brazil where these practices have a strong detrimental impact upon
28 soil physical and chemical properties of the soil as well as diversity and ac-
29 tivity of soil biota with dramatic impacts on the sustainability of agriculture
30 (Mathieu et al., 2005).

31 In French Guiana, the demographic pressure threatens the long-lasting
32 equilibrium between slash-and-burn agriculture and nature conservation, due
33 to the progressive disappearance of shifting cultivation. In southern Guiana
34 near the Suriname border, the duration of fallow in the slash-and-burn system
35 has decreased from 15 to 7-8 years in the last 30 years (Topoliantz et al.,
36 2006). The traditional shifting cultivation is still practised in that region by
37 Amerindian communities and the duration of the fallow ranges from 15 to
38 more than 100 years (Fleury, 1998). The aim of this study was to assess the

1
2
3
4
5
6
7
8
9
39 impact of slash-and-burn cultivation upon the diversity of soil macrofauna in
10 two agricultural systems that differed by the duration of the fallow period.
11
12
13 41 We investigated a traditional shifting cultivation system in a small Wayana
14
15 42 Amerindian village and an accelerated rotation cycle in a larger Aluku village
16
17 43 where the demographic pressure is strong. Crops are mostly manioc (*Manihot*
18
19 44 *esculenta* Cranz) i.e. cassava in both villages and the studied systems are
20
21 45 good examples of traditional shifting cultivation (Wayana Amerindians) and
22
23 46 change to permanent agriculture due to demographic pressure (Aluku village)
24
25 47 (Grandisson, 1997).
26
27

48 **2. Materials and methods**

49 *2.1. Sites*

50 The present survey was carried out in southern French Guiana near
51 the border between France and Suriname along the Maroni river. We in-
52
53 52 vestigated soil macrofauna diversity in agricultural fields of two communi-
54
55 53 ties, Wayanas (indians) and Alukus (maroons, of ancient African lineage).
56
57 54 Wayana Amerindians are still using the traditional slash-and-burn system
58
59 55 where short cropping periods (ca. 2–3 yr) alternate with long fallow periods
60
61 56 (≈ 25 yr). The fields (thereafter referred to as “abattis”) are settled by cut-
62
63 57 ting and burning forest plots and are planted with manioc which constitutes
64
65 58 the basic food. Soils are not tilled and manioc cuttings (from previous crops)
66
67 59 are planted after resprouting. Neither Wayanas nor Alukus use pesticides,
68
69 60 herbicides or fertilizers. Alukus are using a similar system with the difference
70
71 61 that cultivation does not exceed 1 year and fallows are shorter (≈ 8 yr on
72
73 62 average). The length of the cultivation period depends on the soil fertility
74
75 63
76
77
78
79
80
81
82
83
84
85

1
2
3
4
5
6
7
8
9
63 and the spontaneous regrowth of vegetation (Topoliantz et al., 2006).
10
11
12 64 *Ameridian village of Elahe (long fallow)*. We sampled soil macrofauna in the
13
14 65 Wayana (Amerindian) village of Elahe. This small village is situated on the
15
16 66 Tampock river which is a subsidiary of Maroni (3 ° 26'N, 53 ° 59'W). Three
17
18 67 contrasted situations were investigated. A field that had been cut, burnt and
19
20 68 cultivated by an Indian family 3 yrs before this study (EA). This field was
21
22 69 located next to a secondary forest and was itself a secondary forest before
23
24 70 its cultivation. It was sampled in July 1999. We sampled an old secondary
25
26 71 forest (EF) located nearby EA (ca. 100 m). The old secondary forest showed
27
28 72 woody species typical of mature forests (e.g. *Astrocaryum sciophilum* (Miq.)
29
30 73 Pulle, and *Dicorynia guianensis* Amsh. (Poncy et al., 2001) which indicated
31
32 74 that it had been left untouched for at least 100 years. The plot located in
33
34 75 the secondary forest was resampled in May 2000 after it had been cut and
35
36 76 burnt in December 1999 for cultivation (EB).

37
38 77 *Aluku village of Maripasoula (short fallow)*. The second site is located along
39
40 78 the Maroni river (3 ° 39'N, 54 ° 2'W) near the village of Maripasoula, ca. 25
41
42 79 km downstream of the first study site. Maripasoula is a large village (pop
43
44 80 1200 in 1999) mostly inhabited by Aluku people. The increase in population
45
46 81 density during the last 3 decades led to a decrease in the surface of cultivable
47
48 82 land and the subsequent decrease of fallow duration (Fleury, 1998). We
49
50 83 sampled a 1-yr old abattis (MA) at the end of the crop period. It had been
51
52 84 opened by an Aluku family by cutting and burning an 8-yr-old woody fallow
53
54 85 referred to as MF. MF was characterized by pioneer woody species such as
55
56 86 *Cecropia latiloba* Miq. and *Inga capitata* Desv. Both MA and MF plots were
57
58 87 sampled in July 1999. The MF plot was intended to be burnt in December

1
2
3
4
5
6
7
8
9
88 1999 and we planned to resampled in May 2000 but unfortunately the Aluku
10
11 89 family did not burn it as expected.

12
13 90 The mean annual temperature is 26 ° C and the mean annual rainfall is
14
15 91 2000 mm. There is a main dry season from September to December and a
16
17 92 shorter one between March and April. Sampled soils are sandy Oxisols at pH
18
19 93 of 5 and 4.7 on average in Maripasoula and Elahe, respectively (Topoliantz
20
21 94 et al., 2006). The average total C content was 24.6, 22.5, 25.8, 19.1 and 18.6
22
23 95 g kg⁻¹ while the total N content was 1.65, 1.48, 1.78, 1.39 and 1.35 g kg⁻¹
24
25 96 the in plots MA, MF, EA, EF and EB respectively (data from Table 1 in
26
27 97 Topoliantz et al., 2006). Other physico-chemical features of the soils at the
28
29 98 study sites are available in Topoliantz et al. (2006).

30 31 99 *2.2. Sampling*

32
33 100 We used the Tropical Soil Biology and Fertility (TSBF) procedure (An-
34
35 101 derson and Ingram, 1993). Sampling units consisted of 25 cm × 25 cm by
36
37 102 30 cm deep soil monoliths. As recommended in the TSBF procedure, we
38
39 103 used 10 monoliths per transect and carried out 3 transects per plot (i.e. 30
40
41 104 samples per plots). The distance separating monoliths was 5 m and transects
42
43 105 were 20 m distant from each other. The litter was collected at each sampling
44
45 106 point and a trench was then dug to a depth of 30 cm around the 25 × 25
46
47 107 cm² area to get a soil monolith. Macroinvertebrates from soil and litter were
48
49 108 hand-sorted and preserved in 4% formalin solution. Invertebrates were later
50
51 109 counted and identified in the laboratory. We grouped specimens in morphos-
52
53 110 pecies and identified most of them with the help of different taxonomists.
54
55 111 We excluded larvae from the statistical analyses because they were partially
56
57 112 redundant with adults found in the same samples. Overall, our estimation

1
2
3
4
5
6
7
8
9 of species richness is therefore underestimated.

10
11
12 114 *2.3. Data analysis*

13
14 115 *2.3.1. Community structure*

15
16 116 *Principal Coordinate Analysis (PCoA)*. The first step of data analysis con-
17
18 117 sisted of a Principal Coordinate Analysis (PCoA) of the raw data set (site-
19
20 118 species abundances). This multivariate analysis is fully described in Legendre
21
22 119 and Legendre (1998). It was first proposed by Gower (1966) and consists of
23
24 120 the Euclidean representation of a set of objects described by any similarity
25
26 121 or distance coefficients. We used PCoA in order to produce a general rep-
27
28 122 resentation of our soil fauna samples and to examine to which extent they
29
30 123 differed according to sites and land-use types. One advantage of this analysis
31
32 124 is that the user can select the most appropriate distance or similarity index
33
34 125 given the data at hand, which is not the case in more classical multivariate
35
36 126 analyses e.g. PCA or CoA. We used the Bray–Curtis index (Legendre and
37
38 127 Legendre, 1998, p. 287). The coefficient was computed using the R software
39
40 128 (R Development Core Team, 2008) and the `labdsv` package (Roberts, 2007).
41
42 129 The PCoA was done using the `ade4` package (Chessel et al., 2004). The
43
44 130 significance of PCoA axes were assessed by means of a bootstrap procedure
45
46 131 based on 1000 randomizations (Pillar, 1999).

47
48 132 *Species indicator value: IndVal*. Because the PCoA is based on the diagonal-
49
50 133 ization of a distance/similarity matrix there is no direct link between factorial
51
52 134 axes and original descriptors (i.e. macrofauna species). We thus examined
53
54 135 the presence of species associated to one or more samples or sites by means
55
56 136 of a specific method, the *IndVal* value (Dufrêne and Legendre, 1997). Indi-

1
2
3
4
5
6
7
8
9
10 137 cator species are species mostly present in one of the groups to be compared,
11 138 while being mostly absent in other groups. Dufrêne and Legendre (1997)
12
13 139 proposed to compute the *IndVal* index by combining two terms reflecting
14
15 140 the specificity and the fidelity of a species for the samples corresponding to
16
17 141 a certain land-use type. The specificity term is the mean abundance of a
18
19 142 species i in the samples of the land-use j compared to all land-uses.

$$A_{ij} = N_{individuals_{ij}}/N_{individuals_{i\bullet}}$$

20
21
22
23
24
25 The fidelity term is the relative frequency of occurrence of species i in samples
26
27 of land-use j .

$$B_{ij} = N_{sites_{ij}}/N_{sites_{\bullet j}}$$

28
29
30
31 The indicator value of a given species is the product of specificity and fidelity
32
33 terms in percent:

$$IndVal_{ij} = A_{ij} \times B_{ij} \times 100$$

34
35
36
37 143 The indicator value of a species i for an array of sites is the largest value of
38
39 144 $IndVal_{ij}$ observed over all sites j . In this paper we adopted the threshold
40
41 145 level of 25% for the index as suggested by Dufrêne and Legendre (1997),
42
43 146 i.e. species i is present in at least 50% of samples of land-use j and its
44
45 147 relative abundance in land-use j is at least 50%. We assessed the statistical
46
47 148 significance of observed *IndVal* values by means of a permutation test (1000
48
49 149 randomizations) as proposed by Dufrêne and Legendre (1997).

50
51
52 150 *Multi Response Permutation Procedure (MRPP)*. We tested the significance
53
54 151 of community dissimilarities among land-uses by means of the Multi Re-
55
56 152 sponse Permutation Procedure (MRPP) (Quinn and Keough, 2002). MRPP

1
2
3
4
5
6
7
8
9 153 tests whether there is a significant difference between two or more groups of
10 154 sampling units. The MRPP statistic δ is the overall weighted mean of within-
11 155 group means of pairwise dissimilarities among sampling units. The observed
12 156 value is statistically tested by mean of a permutation test where sampling
13 157 units and their associated pairwise distances are permuted N times and δ is
14 158 recalculated. The significance test is based on the proportion of permuted
15 159 δ s that are less than the observed δ . We used the Bray–Curtis distance to
16 160 quantify the dissimilarities between land-uses. MRPP was computed using
17 161 the `vegan` package (Oksanen et al., 2008).

162 2.3.2. *Species richness*

163 Samples were pooled to compute the observed bulk species richness for
164 each land-use type. We used bootstrap procedure to determine a possible bias
165 in the species richness estimator and to remove it from the observed value.
166 For a given land-use, a randomized sample of $n = 30$ sampling units was
167 constituted by randomly sampling with replacement amongst the 30 original
168 units and the observed cumulated species richness was computed. This was
169 repeated N times and constituted the bootstrap sample. The bias was defined
170 as the observed mean species richness minus the average of the N observed
171 species richnesses (Manly, 1997). A corrected estimate of species richness
172 was obtained by subtracting the bias to the observed species richness. We
173 approximated the standard error of the estimated species richness by the
174 standard deviation of bootstrap estimates following Manly (1997, p. 36).
175 Species rarity was measured as absolute and relative frequencies of singletons,
176 i.e. species with at most 1 individual per sample. We also report another
177 measure of rarity based on frequency of species: the number of unique species,

1
2
3
4
5
6
7
8
9
10 178 i.e. species that occurred in only one sample. The observed species richnesses
11 179 in the different land-uses were compared by mean of a one-factor analysis
12
13 180 of variance. Because our data did not satisfy most of the assumptions of
14
15 181 ANOVA we used the randomization test described by Manly (1997, p.117).
16
17 182 The observed F statistic was tested by comparison with corresponding values
18
19 183 in N ANOVAS after randomization of raw data.
20
21

22 184 **3. Results**

23 24 25 185 *3.1. General statistics*

26
27 186 Soil macrofauna communities differed markedly amongst land-uses and
28
29 187 between sites. The average macrofaunal density was higher in the secondary
30
31 188 forest (EF) in Elahe followed by the abbatis in Maripasoula (MA) and it was
32
33 189 lowest in the recently burnt secondary forest in Elahe (EB)(Table 1). The
34
35 190 observed species richness varied accordingly with a total of 121 species in the
36
37 191 secondary forest in Elahe (EF) and values ranging from 22 to 54 species in the
38
39 192 other plots. Corresponding bias-corrected values estimated from bootstrap
40
41 193 were somewhat higher (Table 1) but between-site ranking did not change at
42
43 194 all (there is a linear relationship between original and corrected values). The
44
45 195 number of singletons and unique species respectively ranged from 44% to 54
46
47 196 % and 55% to 72%, respectively, and was fairly homogeneous amongst sites
48
49 197 (Table 1). A grand total of $S = 186$ species was recorded (all sites pooled)
50
51 198 amongst which 42 (22.6%) were common to Elahe and Maripasoula (shared
52
53 199 species). One hundred and twelve (60.2%) species were encountered in the
54
55 200 site of Elahe whereas 32 (17.2%) species were only recorded in Maripasoula.
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
201 *3.2. Community structure*

202 First and second eigenvalues of PCoA significantly differed from those
203 stemming from $N = 1000$ randomizations and accounted for 20.7% and 16%
204 of the total inertia, respectively. Axis 1 clearly reflected changes in macro-
205 fauna community structure according to the type of agriculture: Maripasoula
206 plots (slash-and-burn agriculture with short fallow periods) are opposed to
207 Elahe plots (traditional shifting agriculture) along Axis 1 (Fig. 1). EA, the
208 abbatis from the Elahe village was intermediate. Interestingly, plots EA, EF
209 and EB were clearly different and much more heterogeneous than MA and
210 MF. Within-plot heterogeneity was graphically represented by the scatter-
211 ing of samples around each centre of inertia. Axis 2 mainly reflected within
212 site variability and did not discriminate land-uses. MRPP (1000 random-
213 izations) showed that community dissimilarity among the five land-uses was
214 highly significant ($p < 0.001$).

215 *3.3. Species richness within and between sites*

216 A total of 145 species were found in Elahe amongst which 121 were
217 recorded in the secondary forest (EF) i.e. 83% of the total (Table 2). The
218 abattis (EA) hosted 45 species among which 17% were shared with EF. The
219 recently burnt forest (EB) harboured a total of 22 species, ca. 15% of the
220 total richness (Table 2). The number of exclusive species i.e. those species
221 that were found in only one land-use was 93, 20 and 12 for EF, EA and EB,
222 respectively. The proportion of shared species between EB and both EF and
223 EA was low and ranged from 6.7% to 9.8% (Table 2). In the short fallow
224 system (Maripasoula) the total species richness (all sites pooled) was 77. The
225 woody fallow that served as reference hosted 54 species (70% of the total)

1
2
3
4
5
6
7
8
9 226 while the abattis hosted 41 species (53% of the total) (Table 3). The propor-
10 tion of shared species was 28% (Table 3) which was higher than Elahe (Table
11 227 tion of shared species was 28% (Table 3) which was higher than Elahe (Table
12 228 2). The observed species richness significantly differed among land-uses as
13 229 indicated by the ANOVA test based on 1000 randomizations ($p < 0.001$).
14
15
16
17

18 230 3.4. *Indicator species*

19
20 231 Six morphospecies showed *IndVal* values $\geq 25\%$ and significant associ-
21 ated probabilities at $\alpha = 0.05$: two earthworms, two centipedes, a termite
22 232 and a bristletail (Table 4). Five of these were associated to the old secondary
23 233 forest (traditional slash-and-burn shifting agriculture) while one earthworm
24 234 species, *Pontoscolex corethrurus*, was indicator of the abattis in the short
25 235 fallow system in the Aluku village of Maripasoula.
26
27
28
29
30
31
32

33 237 4. Discussion

34 238 4.1. *Forest logging and fire*

35
36 239 Forest clearing and burning had a strong impact upon soil macrofauna
37 species richness in Elahe. Soil macrofauna communities were dramatically
38 240 impoverished soon after fire (plot AB) and remained strongly affected three
39 241 years after crop establishment (plot EA). Tree harvesting is known to affect
40 242 the composition of soil faunal communities as well as food web structure in
41 243 various ways (Bengtsson et al., 1997). One direct and strong impact is the
42 244 change in soil climatic conditions (temperature, moisture) that is associated
43 245 to the direct exposure of soil to solar radiation (Matlack, 1993). These effects
44 246 are magnified by fire associated with forest logging. During fire, the temper-
45 247 ature of the surface soil can exceed 400°C (Gimeno-García et al., 2004) with
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9 249 important consequences upon soil dwelling organisms and most notably non-
10 250 mobile species that inhabit litter layers and/or upper soil strata (Mathieu
11 251 et al., 2005; Abbott et al., 2003). Fire also indirectly affects soil fauna by
12 252 destroying epigeic microhabitats like decaying wood, fine twigs, dead plant
13 253 stems and leaves that serve as trophic resources or habitat for numerous
14 254 species (Mathieu et al., 2005).

20 255 Not surprisingly, the species pool observed in the recently burnt plot EB
21 256 was constituted either by species that withstand forest logging and burning
22 257 or by colonisers originating from adjacent areas.

26 258 After a few years of cropping, the species richness had increased but re-
27 259 mained low (plot EA) and ca. half the species richness of the abattis was
28 260 constituted by species also present in the forest (Table 2). One may hy-
29 261 pothesize that these species are eurytopic (species with large niche breadth)
30 262 that can withstand the environmental conditions that prevail in crop plots.
31 263 These results are fully in accordance with data collected in eastern Amazonia
32 264 (Brazil) (Mathieu et al., 2005).

39 265 Interestingly, forest clearing and the set up of manioc crop has a lower
40 266 impact on the observed richness in the region of Maripasoula where crop
41 267 rotation is more rapid. The abattis (MA) and the woody fallow (MF) share
42 268 more species than corresponding plots in Elahe (EA and EF): one reason for
43 269 this is that the plots that are logged and burnt in Maripasoula are woody
44 270 fallows ca. 7–8 yr-old and by no means old growth forests as in Elahe. The
45 271 species richness of such plots is low and roughly comparable to what is ob-
46 272 served in the abattis in the traditional slash-and-burn system in the native
47 273 Amerindian area of Elahe. These results indicate that agricultural intensi-

1
2
3
4
5
6
7
8
9 274 fication leads to impoverishment of species richness and homogenization of
10
11 275 soil macrofaunal communities.

12
13 276 All but one species that were found to be indicators were associated to the
14
15 277 old secondary forest in Elahe (EF). Given huge environmental differences that
16
17 278 exist between the forest and other land-uses, forest specialists could be ex-
18
19 279 pected to have high *IndVal* values. This index encapsulates two contrasted
20
21 280 and complementary information, specificity and fidelity. Soil macroorgan-
22
23 281 isms generally display a highly aggregated spatial distribution (Ettema and
24
25 282 Wardle, 2002) and this is even more marked in the case of social insects
26
27 283 like termites or ants. This leads to high variance of abundance data and
28
29 284 may cause low values of the fidelity term B_{ij} and subsequently low and/or
30
31 285 not-significant *IndVal* index (Nahmani et al., 2006). Apart from specialist
32
33 286 species from the secondary forest, only one species was associated to crops
34
35 287 with short fallow periods, the pantropical endogeic earthworm *P. corethrurus*
36
37 288 (*Oligochaeta*, *Glossoscolecidae*). This peregrine species has been dispersed
38
39 289 worldwide by man and is probably indigenous to the Guianas plateau in
40
41 290 South America (Righi, 1984). This species exhibits a very good aptitude to
42
43 291 withstand soil disturbance and scarcity of organic matter and is abundant
44
45 292 in open areas over a wide range of tropical regions (Lavelle et al., 1987).
46
47 293 *P. corethrurus* has been shown to have the capacity to invade Amazonian
48
49 294 pastures and to cause soil degradation through physical compaction of the
50
51 295 soil surface (Chauvel et al., 1999). However, other studies showed that it
52
53 296 can incorporate charcoal to the topsoil thereby increasing its humus content
54
55 297 of the topsoil by adding finely powdered black carbon (Ponge et al., 2006;
56
57 298 Topoliantz and Ponge, 2003, 2005).

1
2
3
4
5
6
7
8
9 299 *4.2. Landscape context and species diversity*

10
11 300 Our sampled soils differed in terms of physico-chemical parameters (e.g.
12
13 301 total C content) and these differences can be explained by land management
14
15 302 practices among which fallow duration. Land management can have direct
16
17 303 effects upon soil biodiversity as well as indirect effects through changes in
18
19 304 soil parameters like or C or N content. Both direct and indirect effects
20
21 305 convey land-use impacts on soil biodiversity. In that context, our plots can
22
23 306 be compared although our data do not allow unravelling such direct and
24
25 307 indirect effects.

26
27 308 Comparing Elahe and Maripasoula sites allows us to discuss the effects of
28
29 309 the reduction of fallow duration upon γ , i.e. regional diversity. The impact of
30
31 310 reduced fallow on the pooled richness was huge (145 versus 77 species in Elahe
32
33 311 and Maripasoula, respectively). The main difference between these species
34
35 312 pools is due to the low richness of the woody fallow (MF) in Maripasoula as
36
37 313 compared to the old secondary forest (EF) in Elahe. The woody fallow is the
38
39 314 typical habitat that is cut and burnt before crop settlement in the reduced
40
41 315 fallow system. There is no secondary forest nor older fallow left in the region
42
43 316 of Maripasoula where all areas accessible by foot or by canoe have been and
44
45 317 are still used for cultivation (Topoliantz et al., 2006). On the other hand,
46
47 318 abattis are similar in Elahe and Maripasoula in terms of richness (roughly
48
49 319 40 to 45 species) although the proportion of shared species is low ($\approx 21\%$).

50 320 Various mechanisms contribute to link α diversity to neighbouring land-
51
52 321 scape configuration (Dunning et al., 1992; Rossi and van Halder, in press).
53
54 322 Species richness in the abattis could be affected by the neighbouring species-
55
56 323 rich secondary forest or by woody fallows through the so-called “vicinism” or

1
2
3
4
5
6
7
8
9 324 “mass” effect (Zonneveld, 1995) whereby a flow of individuals originates from
10 325 species rich habitat and allows the presence of species in neighbouring habi-
11 326 tats where they are not self-maintaining (Shmida and Wilson, 1985). This
12
13 327 corresponds to the “spillover effect” by which species can be maintained in
14
15 328 unsuitable habitat patches in the framework of source-sink dynamics (Holt,
16
17 329 1997). In the context of accelerated crop rotation, the amount of land covered
18
19 330 by pristine habitats is very low if not zero. In the site of Elahe, the situation
20
21 331 is inverse, crops constituting a habitat distributed as small patches scattered
22
23 332 across a landscape essentially constituted of rich undisturbed or slightly dis-
24
25 333 turbed forests. In highly anthropised landscapes, the overall proportion of
26
27 334 species with high capacity to colonize abattis is larger and the landscape con-
28
29 335 text tends to favour a rapid colonization of newly settled fields. This is not the
30
31 336 case in the Amerindian region where abattis are hardly spatially connected.
32
33
34

337 4.3. *The importance of rare species*

338 We defined rare species as species collected as single individuals, single-
339 tons. All studied sites exhibited roughly 50% of singletons and an even larger
340 proportion of unique species. Such high values were reported e.g. for insect
341 communities associated to tropical trees (Novotný and Basset, 2000) and soil
342 macrofauna in Amazonian pastures (Rossi et al., 2006). Rarity can be a prob-
343 lem during data analysis and various indices have been proposed to correct
344 estimates of species richness and shared species (Colwell and Coddington,
345 1994; Chao et al., 2005). In this study we used a bootstrap correction of
346 species richness estimates which led to corrected estimates that were linearly
347 correlated to observed species richnesses. This indicated that our perception
348 of between-site similarity in terms of species richness were correct even when

1
2
3
4
5
6
7
8
9 349 solely based on raw data.

10
11 350 Statistics aside, rare species raise the question why are they so numerous.
12
13 351 First, rare species could be species that were inadequately sampled and this
14
15 352 must be considered with caution. Indeed, soils are highly variable habitats
16
17 353 with various nested sources of spatial heterogeneity that affected species dis-
18
19 354 tribution and hence short-scale diversity patterns (Mathieu et al., 2009). The
20
21 355 problem of sampling deficiency in estimating biodiversity in soils is discussed
22
23 356 elsewhere (Rossi et al., 2006) and could possibly be responsible for a certain
24
25 357 proportion of our rare species. It must be also noted that the TSBF sampling
26
27 358 protocol may not be optimal for certain type of organisms and notably litter
28
29 359 inhabiting species. Other sampling methods like Winkler bags may also be
30
31 360 used (Smith et al., 2008) while specific strategies have been developped for
32
33 361 social insects (termites) (Jones and Eggleton, 2000). On the other hand, the
34
35 362 “spillover effect” could lead to the presence of rare species simply because
36
37 363 they are transient species i.e. species that cannot maintain in one habitat
38
39 364 but originates from a neighbouring habitat that acts as a permanent source.
40
41 365 Distinguishing transient species is a very difficult task (Novotný and Basset,
42
43 366 2000) that would require a huge amount of additional field data (and direc-
44
45 367 tional activity traps) in our case. However, this is the only way to refine our
46
47 368 estimates of species richness while controlling for close neighbourhood effects
48
49 369 and therefore assessing landscape effects upon crop field species richness.

50
51 370 *4.4. Conservation and management of soil macrofaunal diversity in agro-*
52
53 371 *ecosystems*

54
55 372 It is broadly acknowledged that the conservation of soil macrofauna in
56
57 373 agroecosystems is an important aspect of sustainable agriculture (Brussaard

1
2
3
4
5
6
7
8
9 374 et al., 2007) but the spatial facet of that question have not yet received the
10 375 attention it requires (but see Tschardt et al., 2005). The landscape at Mari-
11 376 pasoula is dominated by highly disturbed areas and the direct consequence
12 377 is an erosion of γ diversity. Local species richness thus rely on colonization
13 378 from impoverished regional species pool. Swift et al. (2004) argue that the
14 379 management of biodiversity in agricultural context is more effective at the
15 380 landscape scale than at the local plot scale where strictly utilitarian options
16 381 prevail. Following this line of reasoning, and given that most forest species
17 382 are unlikely to maintain themselves within abattis, the management of γ
18 383 diversity may be based on the maintenance of fallow plots of a large range
19 384 of age and not only of pristine ecosystems. It is additionally important to
20 385 consider the spatial arrangement of these plots across the landscape. They
21 386 act as source of colonisers for newly abandoned field crops and as such their
22 387 spatial location may be of prime importance. The dispersal abilities of most
23 388 soil invertebrates are not very well documented but are certainly limited for
24 389 many species and this reinforces the importance of the distribution of the
25 390 source of colonisers within agricultural landscapes.

391 5. Acknowledgements

392 We are greatly indebted Gabriel Fossé (Maripasoula) who guided us
393 through the forest and the rivers, tremendously contributed to field work
394 and had the role of interpreter when we were at the village of Elahe. We also
395 want to thank everybody at Elahe and Maripasoula for their warm welcome
396 and giving access to their abbatis. We thank Céline Houdin, Périne Lavelle,
397 Jocelyne Roman, Corinne Rouland and Stéphanie Topoliantz who helped in

1
2
3
4
5
6
7
8
9 398 the fieldwork. We are indebted to Guy Pallaud (ONF, Maripasoula) for his
10 399 help and interest in this project when we stayed in Maripasoula in 2000. This
11
12 400 work was conducted in the framework of the SOFT programme “Effets des
13 401 pratiques culturelles traditionnelles sur la fertilité des sols et la régénération
14 402 de la forêt du sud de la Guyane” coordinated by J. M. Betsch et S. Bahuchet
15 403 and funded by the French Ministry of the Environment. We thank two
16 404 anonymous reviewers for their useful comments on an earlier version of the
17 405 manuscript
18
19
20
21
22
23
24
25

26 406 **References**

- 27
28 407 Abbott, I., Burbidge, T., Strehlow, K., Mellican, A., Wills, A., 2003. Log-
29 408 ging and burning impacts on cockroaches, crickets and grasshoppers, and
30 409 spiders in Jarrah forest, Western Australia. *Forest. Ecol. Manage.* 174,
31 410 383–399.
- 32
33 411 Anderson, J., Ingram, J., 1993. *Tropical Soil Biology and Fertility: a Hand-*
34 412 *book of Methods.* CAB International, Oxford.
- 35
36 413 Bengtsson, J., Persson, T., Lundkvist, H., 1997. Long-term effects of logging
37 414 residue addition and removal on macroarthropods and enchytraeids. *J.*
38 415 *Appl. Ecol.* 34, 1014–1022.
- 39
40 416 Brussaard, L., de Ruiter, P. C., Brown, G. G., 2007. Soil biodiversity for
41 417 agricultural sustainability. *Agric. Ecosyst. Environ.* 121, 233–244.
- 42
43 418 Chao, A., Chazdon, R. L., Colwell, R. K., Shen, T.-J., 2005. A new statistical
44 419 approach for assessing similarity of species composition with incidence and
45 420 abundance data. *Ecol. Lett.* 8, 148–159.
- 46
47
48
49
50
51
52
53
54
55
56
57
58

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

421 Chauvel, A., Grimaldi, M., Barros, E., Blanchart, E., Desjardins, T., Sar-
422 razin, M., Lavelle, P., 1999. Pasture damage by an Amazonian earthworm.
423 Nature 398, 32–33.

424 Chessel, D., Dufour, A.-B., Thioulouse, J., 2004. The ade4 package-I: One-
425 table methods. R News 4, 5–10.

426 Colwell, R. K., Coddington, J. A., 1994. Estimating terrestrial biodiversity
427 through extrapolation. Philos. Trans. R. Soc. Lond. B 345, 101–118.

428 Dufrière, M., Legendre, P., 1997. Species assemblages and indicator species:
429 the need for a flexible asymmetrical approach. Ecol. Monogr. 67, 345–366.

430 Dunning, J. B., Danielson, B. J., Pulliam, H. R., 1992. Ecological processes
431 that affect populations in complex landscapes. Oikos 65, 169–175.

432 Ettema, C., Wardle, D., 2002. Spatial soil ecology. Trend. Ecol. Evol. 17,
433 177–183.

434 Fleury, M., 1998. Les populations du Haut-Maroni et le projet du Parc Na-
435 tional de la Guyane. JATBA 40, 577–610.

436 Gimeno-García, E., Andreu, V., Rubio, J., 2004. Spatial patterns of soil
437 temperatures during experimental fires. Geoderma 118, 17–38.

438 Gower, J., 1966. Some distance properties of latent root and vector methods
439 used in multivariate analysis. Biometrika 53, 325–338.

440 Grandisson, M., 1997. Diagnostic de la reproductibilité de la fertilité dans les
441 abattis de la région ouest-Guyane: étude des stratégies de gestion et anal-
442 yse des trajectoires d'évolution des composantes de la fertilité en fonction

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

443 du milieu et des modes de mise en valeur. Ph. thesis, Université Antilles-
444 Guyane, Cayenne, France.

445 Holt, R. D., 1997. From metapopulation dynamics to community structure
446 - some consequences of spatial heterogeneity In: Hanski, I. A., Gilpin,
447 M. E. (Eds.), Metapopulation Biology - Ecology, Genetics, and Evolution.
448 Academic Press, San Diego, pp. 149–164.

449 Jones, D. T., Eggleton, P., 2000. Sampling termite assemblages in tropical
450 forests: testing a rapid biodiversity assessment protocol. J. Appl. Ecol. 37,
451 191–203.

452 Lavelle, P., Barois, I., Cruz, C., Hernandez, A., Pineda, A., Rangel, P.,
453 1987. Adaptative strategies of *Pontoscolex corethrurus* (Glossoscolecidae,
454 Oligochæta), a peregrine geophagous earthworm of the humid tropics. Biol.
455 Fert. Soils 5, 188–194.

456 Lavelle, P., Decaëns, T., Aubert, M., Barot, S., Blouin, M., Bureau, F.,
457 Margerie, P., Mora, P., Rossi, J.-P., 2006. Soil invertebrates and ecosystem
458 services. Eur. J. Soil Biol. 42, S3–S15.

459 Legendre, P., Legendre, L., 1998. Numerical Ecology, Second English Edition.
460 Elsevier, Amsterdam.

461 Manly, B., 1997. Randomization and Monte Carlo methods in Biology. Chap-
462 man & Hall, London.

463 Mathieu, J., Grimaldi, M., Rouland, C., Lavelle, P., Desjardin, T., Rossi, J.-

- 1
2
3
4
5
6
7
8
9
10 464 P., 2009. Variations in soil macrofauna biodiversity with spatial patterns
11 465 of grasses in Amazonian pastures. *Soil Biol. Biochem.* 41, 586–593.
12
13
14 466 Mathieu, J., Rossi, J.-P., Grimaldi, M., P., M., Lavelle, P., Rouland, C.,
15
16 467 2004. A multi-scale study of soil macrofauna biodiversity in Amazonian
17
18 468 pastures. *Biol. Fert. Soils* 40, 300–305.
19
20 469 Mathieu, J., Rossi, J.-P., Mora, P., Lavelle, P., Martins, P. F. D. S., Rouland,
21
22 470 C., Grimaldi, M., 2005. Recovery of soil macrofauna communities after
23
24 471 forest clearance in Eastern Amazonia, Brazil. *Conserv. Biol.* 19, 1598–
25
26 472 1605.
27
28
29 473 Matlack, G. R., 1993. Microenvironment variation within and among forest
30
31 474 edge sites in the eastern United States. *Biol. Conserv.* 66, 185–194.
32
33
34 475 Matson, P. A., Parton, W. J., Power, A. G., Swift, M. J., 1997. Agricultural
35
36 476 intensification and ecosystem properties. *Science* 277, 504–509.
37
38
39 477 Nahmani, J., Lavelle, P., Rossi, J.-P., 2006. Does changing the taxonomical
40
41 478 resolution alter the value of soil macroinvertebrate as bioindicators of metal
42
43 479 pollution? *Soil Biol. Biochem.* 38, 385–396.
44
45 480 Novotný, V., Basset, Y., 2000. Rare species in communities of tropical insect
46
47 481 herbivores: pondering the mystery of singletons. *Oikos* 89, 564–572.
48
49
50 482 Oksanen, J., Kindt, R., Legendre, P., O’Hara, B., Simpson, G., Stevens, M.,
51
52 483 Wagner, H., 2008. *vegan: Community Ecology Package*. R package version
53
54 484 1.13-1. <http://vegan.r-forge.r-project.org/>
55
56
57
58
59
60
61
62
63
64
65

- 1
2
3
4
5
6
7
8
9
485 Pillar, V. D., 1999. The bootstrapped ordination re-examined. *J. Veg. Sci.*
10
486 10, 895–902.
11
12
13
487 Poncy, O., Sabatier, D., Prévost, M.F., Hardy, I., 2001. The lowland high
14
488 rainforest: structure and tree species diversity. In: Bongers, F., Charles-
15
489 Dominique, P., Forget, P.M., Théry, M. (Eds.), *Nouragues: Dynamics and*
16
490 *Plant-Animal Interactions in a Neotropical Rainforest*. Academic Press,
17
491 San Diego, pp. 149–164.
18
19
20
21
22
23
492 Ponge, J., Topoliantz, S., Ballof, S., Rossi, J.-P., Lavelle, P., Betsch, J.,
24
493 Gaucher, P., 2006. Ingestion of charcoal by the amazonian earthworm
25
494 *Pontoscolex corethrurus*: a potential for tropical soil fertility. *Soil Biol.*
26
495 *Biochem.* 38, 2008–2009.
27
28
29
30
31
32
496 Quinn, G., Keough, M., 2002. *Experimental Design and Data Analysis for*
33
497 *Biologists*. Cambridge University Press, Cambridge.
34
35
36
37
498 R Development Core Team, 2008. *R: A Language and Environment for Sta-*
38
499 *tistical Computing*. R Foundation for Statistical Computing, Vienna, Aus-
39
40
41
500 tria. ISBN 3-900051-07-0, URL <http://www.R-project.org>.
42
43
501 Righi, G., 1984. *Pontoscolex* (Oligochæta, Glossoscolecidae): a new evalua-
44
45
46
502 tion. *Stud. Neotrop. Fauna Environ.* 19, 159–177.
47
48
503 Roberts, D., 2007. labdsv: Ordination and Multivariate Analysis for Ecology.
49
50
504 R package version 1.3-1. <http://ecology.msu.montana.edu/labdsv/R>
51
52
53
505 Rossi, J.-P., Blanchart, E., 2005. Seasonal and land-use induced variations
54
55
56
57
58

- 1
2
3
4
5
6
7
8
9 506 of soil macrofauna composition in the Western Ghats (South India). *Soil*
10 *Biol. Biochem.* 37, 1093–1104.
11
12
13
14 508 Rossi, J.-P., van Halder, I., in press. Towards indicators of butterfly bio-
15 diversity based on a multiscale landscape description. *Ecol. Indic.*, DOI:
16 10.1016/j.ecolind.2009.07.016.
17
18
19
20 511 Rossi, J.-P., Mathieu, J., Cooper, M., Grimaldi, M., 2006. Soil macrofaunal
21 biodiversity in Amazonian pastures: matching sampling with patterns. *Soil*
22 *Biol. Biochem.* 38, 2178–2187.
23
24
25
26
27 514 Shmida, A., Wilson, M. V., 1985. Biological determinants of species diversity.
28 *J. Biogeogr.* 12, 1–20.
29
30
31
32 516 Smith, J., Potts, S., Eggleton, P., 2008. Evaluating the efficiency of sampling
33 methods in assessing soil macrofauna communities in arable systems. *Eur.*
34 *J. Soil Biol.* 44, 271–276.
35
36
37
38 519 Swift, M. J., Izac, A. M. N., van Noordwijk, M., 2004. Biodiversity and
39 ecosystem services in agricultural landscapes—are we asking the right ques-
40 tions? *Agric. Ecosyst. Environ.* 104, 113–134.
41
42
43
44 522 Topoliantz, S., Ponge, J. F., 2003. Burrowing activity of the geophagous
45 earthworm *Pontoscolex corethrurus* (Oligochaeta : Glossoscolecidae) in
46 the presence of charcoal. *Appl. Soil Ecol.* 23, 267–271.
47
48
49
50
51 525 Topoliantz, S., Ponge, J. F., 2005. Charcoal consumption and casting activity
52 by *Pontoscolex corethrurus* (Glossoscolecidae). *Appl. Soil Ecol.* 28, 217–
53 224.
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

528 Topoliantz, S., Ponge, J. F., Lavelle, P., 2006. Humus components and bio-
529 genic structures under tropical slash-and-burn agriculture. *Eur. J. Soil Sci.*
530 57, 269–278.

531 Tschardtke, T., Klein, A. M., Kruess, A., Steffan-Dewenter, I., Thies, C.,
532 2005. Landscape perspectives on agricultural intensification and biodiver-
533 sity – ecosystem service management. *Ecol. Lett.* 8, 857–874.

534 Zonneveld, I. S., 1995. Vicinism and mass effect. *J. Veg. Sci.* 6, 441–444.

Table 1: Soil macrofaunal diversity in southern French Guiana. EF and EB correspond to the same plot sampled in 1999 and 2000 was burnt in between.

Sites	Acronym	n	date	mean density (ind.m ⁻²)	S_{obs}	S_{corr}	singleton	unique
Elahe (traditional slash-and-burn)								
Secondary forest (≥ 100 -yr)	EF	30	July 1999	1781.9 (289.8)	121 (8.3)	151.2	56 (46.3%)	76 (62.8%)
Abbatis (3-yr)	EA	30	July 1999	541.3 (92)	45 (4.2)	56.8	20 (44.4%)	29 (64.4%)
Recently burnt forest	EB	30	May 2000	224.5 (60.8)	22 (2.1)	27	11 (50%)	12 (54.5%)
Maripasoula								
Woody fallow (8-yr)	MF	30	May 2000	645.3 (76.5)	41 (3.7)	51.6	21 (51.2%)	26 (63.4%)
Abbatis (1-yr)	MA	30	July 1999	1109.9 (271.2)	54 (4.6)	69.4	29 (53.7%)	39 (72.2%)

S_{obs} : observed species richness, S_{corr} : bias-corrected species richness, singleton: species with at most one individual, unique: species encountered in only one sample. Percentages for singletons and uniques are computed with reference to the total species richness in the site. Standard errors for mean density and S_{obs} are indicated in parentheses.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Table 2: Observed species richness and shared species of soil macrofauna in different land-uses in a traditional Amerindian slash-and-burn system (region of Elahe, southern French Guiana).

$S_{\bullet} = 145$	EF	EA	EB
EF	121 (83.4%)	24 (142)	9 (134)
EA	17%	45 (31%)	6 (61)
EB	6.7%	9.8%	22 (15.2%)

539 EF: Secondary forest, EA: Abbatis, EB: recently burnt forest, S_{\bullet} : total
540 species richness (all sites pooled). On diagonal: observed species richness and
541 corresponding ratio to the total number of species (in parentheses). Above
542 diagonal: absolute number of species shared by site pairs and total number of
543 species collected in site pairs. Below diagonal: ratio of the number of shared
544 species to the total number of species in site pairs.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Table 3: Observed species richness and shared species of soil macrofauna in different land-uses in a short-fallow system (region of Maripasoula, southern French Guiana).

$S_{\bullet}=77$	MF	MA
MF	54 (70.1%)	21 (74)
MA	28.4%	41 (53.2%)

MF: Secondary forest, MA: Abbatis, S_{\bullet} : total species richness (all sites pooled). On diagonal: observed species richness and corresponding ratio to the total number of species (in parentheses). Above diagonal: absolute number of species shared by site pairs and total number of species collected in site pairs. Below diagonal: ratio of the number of shared species to the total number of species in site pairs.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Table 4: Soil macrofaunal indicator species in agricultural landscapes in southern French Guiana. IndVal values were tested using permutation tests ($n = 1000$). See text for definitions and Table 1 for land-use acronyms.

Species	group	Specificity	Fidelity	<i>IndVal</i>	<i>p</i>	Land-use
<i>Pontoscolex corethrurus</i>	Oligochaeta	0.38	0.97	36.9	0.001	MA
morphospecies 1	Diplura	0.92	0.33	30.4	0.001	EF
morphospecies 2	Isoptera	0.84	0.57	47.9	0.001	EF
morphospecies 3	Chilopoda	0.79	0.43	34	0.001	EF
morphospecies 4	Chilopoda	0.68	0.43	29.2	0.001	EF
morphospecies 1	Lumbricida	0.51	0.67	34.2	0.001	EF

1
2
3
4
5
6
7
8
9
551 **Figure captions**

10
11
12
13 553 **Figure 1**

14
15 554 Principal Coordinates Analysis (PCoA) of soil macrofaunal communities in
16
17 555 different land-uses in southern French Guiana. Projection of site scores in
18
19 556 the plane defined by Axes 1 (horizontal) and 2 (vertical). Labels are placed
20
21 557 at the centre of gravity of each site and are linked to each sample of the
22
23 558 corresponding site. See Table 1 for site codes.

Figure
[Click here to download Figure: f1.eps](#)

