



HAL
open science

Compact Kähler 3-manifolds without non-trivial subvarieties

Frédéric Campana, Jean-Pierre Demailly, Misha Verbitsky

► **To cite this version:**

Frédéric Campana, Jean-Pierre Demailly, Misha Verbitsky. Compact Kähler 3-manifolds without non-trivial subvarieties. 2013. hal-00819044v1

HAL Id: hal-00819044

<https://hal.science/hal-00819044v1>

Submitted on 29 Apr 2013 (v1), last revised 14 Jan 2014 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Compact Kähler 3-manifolds without non-trivial subvarieties

Frédéric Campana, Jean-Pierre Demailly, Misha Verbitsky¹

Abstract

We prove that any compact Kähler 3-dimensional manifold which has no non-trivial complex subvarieties is a torus. This is a very special case of a general conjecture on the structure of ‘simple manifolds’, central in the bimeromorphic classification of compact Kähler manifolds. The proof follows from the Brunella pseudo-effectivity theorem, combined with fundamental results of Siu and of the second author on the Lelong numbers of closed positive $(1, 1)$ -currents, and with a version of the hard Lefschetz theorem for pseudo-effective line bundles, due to Takegoshi and Demailly-Peternell-Schneider. In a similar vein, we show that a normal compact and Kähler 3-dimensional analytic space with terminal singularities and nef canonical bundle is a cyclic quotient of a simple non-projective torus if it carries no effective divisor. This is a crucial step to complete the bimeromorphic classification of compact Kähler 3-folds.

1 Introduction

We consider here connected compact Kähler manifolds X of (complex) dimension $n > 1$. An irreducible compact analytic subset Z of X will be said to be a ‘subvariety’ of X . It is said to be ‘non-trivial’ if its (complex) dimension is neither 0, nor n .

The bimeromorphic classification of compact Kähler manifolds can be reduced, by means of suitable functorial¹ fibrations², to the following two extreme particular cases: either X is projective, or X is ‘simple’, which means that its ‘general’³ point x of X is not contained in any non-trivial subvariety of X ⁴.

The present text is concerned with the bimeromorphic classification of such ‘simple’ X . Its difficulty, in contrast to the projective case, is not due to the abundance and complexity of the examples, but to their (expected) scarceness and ‘simple structure’, which makes essentially all usual invariants of the classification vanish.

¹Misha Verbitsky is partially supported by RFBR grants 12-01-00944-a, AG Laboratory NRI-HSE, RF government grant, ag. 11.G34.31.0023, and NRI-HSE Academic Fund Program in 2013-2014, research grant 12-01-0179.

²In the category of connected compact Kähler manifolds, morphisms being the dominant rational maps with connected fibres.

³Relative algebraic reductions, and relative Albanese maps, see [C1], [C4], [F].

⁴‘general’ means: in the intersection of countably many dense Zariski open subsets.

⁵If X is a non-projective torus, this new notion of ‘simplicity’ coincides with the classical one, defined by the absence of non-trivial subtori.

For example, if X is ‘simple’, its algebraic dimension⁵ $a(X)$ vanishes, which implies, by [U], Lemma 13.1 and Lemma 13.6, that its Albanese map is surjective and has connected fibres. Because the fibres must then have dimension either 0 or n , we get: either $q = 0$, or X is bimeromorphic to its Albanese torus. Thus only the case where $q = 0$ needs to be considered.

The known⁶ examples of ‘simple’ compact Kähler manifolds are (up to bimeromorphic equivalence) ‘general’ complex tori, $K3$ surfaces, or the ‘general’ member of the deformation families of hyperkähler manifolds⁷. ‘Simple’ surfaces are thus classified: either Tori ($q > 0$), or $K3$ ($q = 0$). But when $n \geq 3$, the classification is open. The situation is however expected to be similar to the surface case in higher dimensions (see Conjecture 1.1), where the only known examples are constructed out of surfaces which are either $K3$, or tori.

The following converse was essentially formulated in [C3], as question 1.4.

Conjecture 1.1: Let X be a ‘simple’ compact Kähler manifold. Then:

1. either X has a finite étale cover bimeromorphic to a complex torus, or $H^0(X, \Omega_X^2)$ is generated by some σ which is ‘generically symplectic’ (i.e. $n = 2m$ is even, and $\wedge^m \sigma \neq 0$).

We should thus have: $\kappa(X) = 0$. If $\dim(X)$ is odd, then X should be bimeromorphic to a complex torus, possibly after some finite étale cover. Such a cover will be needed, as shown by the Kummer quotient by the -1 involution.

2. When X does not contain any nontrivial subvariety, X should be either a complex torus, or an irreducible hyperkähler manifold⁸.

Thus K_X should be trivial in this case, the two cases being told by $q > 0$, or $q = 0$.

3. If X is ‘generically symplectic’, then $\pi_1(X)$ should be finite, of cardinality at most 2^m , where $2m = n$.

The assertion 3 follows from [C4], if $\chi(\mathcal{O}_X) \neq 0$ for X ‘simple’ and ‘generically symplectic’.

⁵Recall that $a(X) \in \{0, \dots, n\}$ is the transcendence degree of the field of global meromorphic functions on X , $a(X) = n$ if and only if X is projective.

⁶It is shown in [KV1] that a stable bundle which does not degenerate to a direct sum of stable bundles on the generic deformation of a Hilbert scheme of $K3$ has a compact moduli space; this may lead to new examples of hyperkähler manifolds. The ‘rigidity’ of the category of coherent sheaves on the general members of these families is established in [V02], [V03].

⁷Recall that X is ‘hyperkähler’ if it is Kähler and admits a holomorphic symplectic form. It is said to be ‘irreducible’ if, moreover, $\pi_1(X)$ is finite, and $h^{2,0}(X) = 1$.

⁸Notice that a general deformation of a Hilbert scheme of a $K3$ surface has no subvarieties [V98], while a general deformation of a generalized Kummer variety has some subvarieties, partially classified in [KV2].

In the sequel we prove the second assertion of the conjecture for $n = 3$.

This conjecture can be motivated by the conjectural existence of minimal models in the bimeromorphic category of connected compact Kähler manifolds. Indeed, if such a theory exists, and if X is simple, we have $a(X) = 0$, which brings $\kappa(X) \leq 0$. The possibility $\kappa(X) = -\infty$ is excluded, since X should then be uniruled. This implies $\kappa(X) = 0$.

In this case, a Kähler minimal model is a map $X \dashrightarrow X'$ such that X' has terminal singularities and $K_{X'} \equiv 0$. A (conjectural) Bogomolov-type decomposition ([Bo1], [Bea]) for manifolds with terminal singularities and $K_{X'} \equiv 0$ could be used to represent a finite cover of X' as a product. However X is simple, hence either X' is covered by a simple torus (with finite locus of ramification), or X' carries a symplectic holomorphic 2-form, establishing the conjecture above.

The above conjecture can thus be reformulated in terms of conjectural ‘minimal models’ for compact Kähler manifolds, defined as normal compact complex spaces bimeromorphic to compact Kähler manifolds, and having \mathbb{Q} -factorial terminal singularities. The ‘simple’ ones should then have a torsion canonical sheaf, and be either of the form T/G , quotients of simple non-projective tori by a finite group of automorphisms, or carry a multiplicatively unique holomorphic 2-form, symplectic on the regular locus.

Theorem 1.2: Let X be a compact Kähler 3-dimensional manifold. Assume that X has no non-trivial subvariety. Then X is a torus.

The proof will be given as Corollary 4.4. Although it is an easy combination of known results, it seems worth to present it, because it indicates that the above conjecture might be accessible in dimension 3 with the actually existing techniques. The proof also gives some information on the higher dimensional case (see Corollary 4.5).

We also show:

Theorem 1.3: Let X be a three-dimensional Kähler normal connected compact complex space with terminal singularities. Assume that K_X is nef, and that X does not contain any effective divisor⁹. Then $X = T/G$ is isomorphic to the quotient of a non-projective simple torus by a finite cyclic group of automorphisms.

We will use the following simple criterion:

Lemma 1.4: Let X be a ‘simple’ compact Kähler threefold. If $\chi(\mathcal{O}_X) = 0$, then X is bimeromorphic to its Albanese torus.

⁹The non-existence of effective divisors linearly equivalent to a positive multiple of K_X actually suffices for the conclusion.

Proof: By the above remarks, it is sufficient to show that $q > 0$. But

$$0 = \chi(\mathcal{O}_X) = 1 - q + h^{2,0} - h^{3,0}.$$

By Kodaira's theorem, $h^{2,0} > 0$ since X is not projective, and $h^{3,0} \leq 1$ since $a(X) = 0$. Thus $0 = 1 - q + h^{2,0} - h^{3,0} \geq 1 - q + 1 - 1 = 1 - q$, and $q > 0$. ■

The proof of Theorem 1.2 is thus reduced to showing that $\chi(\mathcal{O}_X) = 0$, which is the objective of the following sections.

Finally, let us mention that compact (connected) complex manifolds, not necessarily Kähler, without non-trivial subvarieties are also quite important in model theory, giving examples of so-called 'trivial' Zariski geometries ([Moo], [MP] [MMT]). Our arguments below show that compact three-dimensional complex manifolds without non-trivial subvarieties are tori, provided their canonical bundle are pseudo-effective. Brunella's theorem and the hard Lefschetz theorem appear to be the only steps of the proof requiring the Kähler hypothesis in depth.

2 Pseudo-effective and nef line bundles

We refer to [D3] for the basic notions and properties of currents on complex manifolds, and only recall briefly some of the facts needed here.

Let X be a compact, n -dimensional complex manifold, and L a holomorphic line bundle on X . Assume that L is equipped with a singular hermitian metric h with local weights $e^{-\varphi}$ which are locally integrable. Its curvature current is then $\Theta(L, h) := i\partial\bar{\partial}\varphi$, and its first Chern class $c_1(L)$ is represented by the cohomology class $\frac{1}{2\pi} [\Theta(L, h)]$ in the Bott-Chern cohomology group $H_{\text{BC}}^{1,1}(X)$, where

$$H_{\text{BC}}^{p,q}(X) = \{d\text{-closed } (p, q)\text{-currents}\} / \{\partial\bar{\partial}\text{-exact } (p, q)\text{-currents}\}.$$

In case X is Kähler, $H_{\text{BC}}^{1,1}(X)$ is isomorphic to the Dolbeault cohomology group $H^{1,1}(X)$ and can be viewed as a subspace of $H^2(X, \mathbb{C})$.

Definition 2.1: A $(1, 1)$ cohomology class $\{\alpha\} \in H_{\text{BC}}^{1,1}(X)$ is said to be **pseudo-effective** if it contains a closed positive $(1, 1)$ -current Θ . The class $\{\alpha\}$ is said to be **nef** if for a given smooth positive $(1, 1)$ -form $\omega > 0$ on X and every $\varepsilon > 0$, it contains a smooth closed $(1, 1)$ -form α_ε such that $\alpha_\varepsilon \geq -\varepsilon\omega$ [or alternatively, in the Kähler case, if $\{\alpha\}$ is a limit of Kähler classes $\{\alpha + \varepsilon\omega\}$, where ω is Kähler]. A class $\{\alpha\}$ is said to be **big** if it contains a Kähler current, namely a closed $(1, 1)$ -current Θ such that $\Theta \geq \varepsilon\omega$ for some smooth positive $(1, 1)$ -form ω and some $\varepsilon > 0$. A line bundle L on X is said to be nef (resp. pseudo-effective, resp. big) if its first Chern class $c_1(L) \in H_{\text{BC}}^{1,1}(X)$ has the same property.

It follows from the Bochner-Kodaira technique that a big line bundle $L \rightarrow X$ has maximal Kodaira dimension $\kappa(L) = \dim X$. Therefore, a compact complex manifold X carrying a big line bundle is Moishezon, i.e. bimeromorphic to a projective algebraic manifold; in particular, X is not simple. Now, we have:

Theorem 2.2: ([DP]) Let X be a compact Kähler manifold. Let $\{\alpha\} \in H^{1,1}(X)$ be a nef class. If $\int_X \alpha^n > 0$ then $\{\alpha\}$ is big. As a consequence, if X carries a nef line bundle L such that $c_1(L)^n > 0$, then L is big and X cannot be simple.

This special case where $\{\alpha\} = c_1(L)$ (the only one that we need here) can also be obtained as a consequence of holomorphic Morse inequalities [D1]. It can be seen as a strengthening of the Grauert-Riemenschneider conjecture proved by Siu [Si2] (although the latter is also valid in the non Kähler case).

Definition 2.3: The **Lelong number** $\nu_x(\Theta(L, h)) = \nu(\varphi, x)$ of the $(1, 1)$ -current $\Theta(L, h)$ at $x \in X$, is defined as $\liminf_{z \rightarrow x} \frac{\varphi(z)}{|z-x|}$, for a local metric on X near x .

Definition 2.4: For a positive real number $c > 0$, the **Lelong set** F_c of a $(1, 1)$ -current η is the set of points $x \in M$ with $\nu(\eta, x) \geq c$. By a well-known theorem of Siu ([Si1]) any Lelong set of a positive, closed current is a complex analytic subvariety of M (the proof of this difficult result has been considerably simplified, using regularization of currents and the Ohsawa-Takegoshi extension theorem, see [D2].)

Theorem 2.5: ([D2]) Let X be a compact complex manifold. Let L be a pseudo-effective holomorphic line bundle on X , with singular hermitian metric h with positive curvature current $\Theta(L, h)$. Assume that the Lelong sets of $\Theta(L, h)$ are all zero-dimensional. Then L is nef, and big as soon as there is at least one nonzero Lelong number.

Proof: The first assertion is ([D2], corollary 6.4). The second follows from ([D2], corollary 7.6). See also [V09], theorem 3.12. ■

Corollary 2.6: If X is a compact Kähler manifold without nontrivial subvarieties, and with pseudo-effective canonical bundle, then K_X is nef, and for any singular hermitian metric h on K_X , $m > 0$ with positive curvature current, the Lelong number vanish at any point. In particular, the associated multiplier ideal sheaves on $K_X^{\otimes m}$ are all trivial for any $m > 0$ (i.e. $e^{-m\varphi}$ is integrable for any $m > 0$).

3 Hard Lefschetz theorem for the cohomology of pseudo-effective line bundles

We recall the version of the Hard Lefschetz theorem which is going to be used here:

Theorem 3.1: ([T], [DPS]) Let (X, ω) be a compact Kähler manifold, of dimension n with Kähler form ω , let K_X be its canonical bundle, and L be a pseudo-effective holomorphic line bundle on X equipped with a singular Hermitian metric h . Assume that the curvature Θ of (L, h) is a positive current on X , and denote by $\mathcal{I}(h)$ the corresponding multiplier ideal sheaf. Then the wedge multiplication operator $\eta \rightarrow \omega^i \wedge \eta$ induces a surjective map

$$H^0(X, \Omega_X^{n-i} \otimes L \otimes \mathcal{I}(h)) \xrightarrow{\omega^i \wedge} H^i(X, K \otimes L \otimes \mathcal{I}(h)).$$

Here ω is considered as an element in $H^1(X, \Omega_X^1)$, and multiplication by ω maps $H^k(X, \Omega_X^{n-l} \otimes F)$ to $H^{k+1}(X, \Omega_X^{n-l+1} \otimes F)$. ■

This theorem was obtained under various hypothesis during the decade 1990: see [E], and [Mou]. The most general form given here is due to [DPS], Theorem 2.1.1. It was proved in [T] when L nef.

Corollary 3.2: Let X be a compact Kähler manifold of dimension $n > 1$ without any non-trivial subvariety. Assume that K_X is pseudo-effective. Then $h^i(X, m.K_X) \leq \binom{n}{i}$, for any $i \geq 0$, and the polynomial $P(m) := \chi(X, m.K_X)$ is constant, equal to $\chi(\mathcal{O}_X)$.

Proof: Since $n > 1$ and X has no non-trivial subvarieties, the algebraic dimension of X vanishes: $a(X) = 0$. Therefore $h^0(X, E) \leq \text{rank}(E)$, for any holomorphic vector bundle E on X . Combining with the Hard Lefschetz theorem (Theorem 3.1), this gives the first claim of Corollary 3.2. The second claim is clear because a polynomial function $P(m)$ which remains bounded when $m \rightarrow +\infty$ is necessarily constant. ■

Corollary 3.3: Let X be a compact Kähler manifold of dimension 3 without any non-trivial subvariety. Assume that K_X is pseudo-effective. The polynomial $P(m) := \chi(X, m.K_X)$ is constant, equal to $\chi(\mathcal{O}_X) = 0$.

Proof: The intersection number K_X^3 vanishes (either by Theorem 2.5, or because it is the leading term of $P(m)$, up to the factor 3!). The Riemann-Roch formula then gives: $P(m) := \frac{(1-12m)}{24} \cdot c_1(X) \cdot c_2(X)$. The boundedness of $P(m)$ then implies that $24 \cdot \chi(\mathcal{O}_X) = c_1(X) \cdot c_2(X) = 0$. ■

Remark 3.4: Arguments close to some of the ones presented here have been already used in the proof [DPS], theorem 2.7.3: if X is a compact Kähler manifold with pseudo-effective canonical bundle admitting a metric with weights

φ having analytic singularities and positive curvature current, then either $H^0(X, \Omega_X^i \otimes (m.K_X)) \neq 0$ for infinitely many $m > 0$ and some $i \geq 0$, or $\chi(X, \mathcal{O}_X) = 0$.

4 Brunella’s pseudo-effectivity criterion.

The rest of our arguments is based on the following strong (and very difficult) theorem by Brunella.

Theorem 4.1: ([Br]) Let X be a compact Kähler manifold with a 1-dimensional holomorphic foliation F given by a nonzero morphism of vector bundle $L \rightarrow T_X$, where L is a line bundle on X , and T_X is its holomorphic tangent bundle. If L^{-1} is not pseudo-effective, the closures of the leaves of F are rational curves (and X is thus uniruled).

The following corollary had already been observed in [HPR], proposition 4.2.

Corollary 4.2: If X is an n -dimensional compact Kähler manifold with $H^0(X, \Omega_X^{n-1}) \neq 0$. Then K_X is pseudo-effective.

Proof: Ω_X^{n-1} is canonically isomorphic to $K_X \otimes T_X$. Any nonzero section thus provides a nonzero map $K_X^{-1} \rightarrow T_X$, and an associated foliation ■

Corollary 4.3: If X is a 3-dimensional non-projective compact Kähler manifold. Then: either K_X is pseudo-effective, or X is uniruled. If X is ‘simple’, K_X is pseudo-effective.

Proof: Kodaira’s theorem implies that any compact Kähler manifold with $H^{2,0}(X) = 0$ is projective. Thus $H^{2,0}(X) \neq 0$, and the preceding corollary applies (since $2 = (n - 1)$, here) and gives the first claim. If X is uniruled, it is not simple, hence the second assertion. ■

By combining Corollary 4.3, Corollary 3.3, and Lemma 1.4, we get our main result:

Corollary 4.4: If X is a 3-dimensional compact Kähler manifold without non-trivial subvariety, then X is a complex torus.

In higher dimensions, we can replace the assumption that K_X was pseudo-effective from our preceding Corollary 3.2 by the existence of a holomorphic $(n - 1)$ -form:

Corollary 4.5: Let X be a compact Kähler manifold of dimension $n > 1$ without non-trivial subvariety. Assume that $H^0(X, \Omega_X^{n-1}) \neq 0$. Then $h^i(X, m.K_X) \leq$

$\binom{n}{i}$, for any $i \geq 0$, and the polynomial $P(m) := \chi(X, m.K_X)$ is constant, equal to $\chi(\mathcal{O}_X)^1$.

The following result (announced as Theorem 1.3) strengthens Theorem 1.2:

Theorem 4.6: Let X be a three-dimensional Kähler normal connected compact complex space with terminal² singularities. Assume that K_X is nef, and that X does not contain any effective divisor³. Then $X = T/G$ is isomorphic to the quotient of a non-projective simple torus by a finite cyclic group of automorphisms.

Proof: By [DPe], theorem 7.1, $\kappa(X) = 0$. Thus mK_X is a trivial line bundle for some $m > 0$. Let $X' \rightarrow X$ be the cyclic cover of X defined by a nonzero section of $m.K_X$ so that $m.K_X$ is Cartier. The singularities of X' are thus terminal as well ([KM], 5.20). They are also rational, since $K_{X'}$ is Cartier (by [KM], 5.24). We thus have $\chi(\mathcal{O}_{X'}) = \chi(\mathcal{O}_{X^n})$ for any smooth model X'' of X' . But $\chi(\mathcal{O}_{X'}) = \frac{-K_{X'} \cdot c_2(X')}{24} = 0$, since $K_{X'}$ is locally free trivial (see for example [Fl]). The conclusion now follows from Lemma 1.4 ■

Remark 4.7: In order to show the conjecture for ‘simple’ Kähler threefolds, it would be sufficient:

1. to show the following statement (which is a form of the abundance conjecture in this context): If X is a ‘simple’ (normal, terminal, \mathbb{Q} -factorial, Kähler) threefold, and D an effective divisor on X , then D is not nef. A possible approach might consist in adapting the proof in the projective case. A more direct approach were however desirable.

2. to show the existence of ‘minimal models’ such as the ones appearing in Theorem 1.3 for any compact Kähler threefold X .⁴

Acknowledgements: We are grateful to the Tata Institute of Fundamental research in Mumbai, India, where this paper was conceived, and especially to I. Biswas and A.J Parameswaran, organizers of our stay there.

References

[Bea] A. Beauville, *Variétés Kähleriennes dont la première classe de Chern est nulle*, J. Differential Geom. 18 (1983), no. 4, 755782.

¹The intersection numbers $K_X^j \cdot \text{Todd}_{n-j}(X)$ thus all vanish when $j > 0$, as expected since K_X should then be trivial.

²we refer to [KM] for this notion, which makes sense without change in the analytic category. In particular, the sheaf K_X is then \mathbb{Q} -Cartier.

³Divisors linearly equivalent to a positive multiple of K_X actually suffice. This second hypothesis should follow from the first if X is ‘simple’.

⁴This step has just been achieved in [HP], where a conjecture equivalent to the one above is formulated in dimension 3.

- [Bo1] F.A. Bogomolov, *On the decomposition of Kähler manifolds with trivial canonical class*, Mat. USSR-Sb. 22 (1974), no. 4, 580583.
- [Br] M. Brunella, *Uniformization of foliations by curves*, Holomorphic dynamical systems, 105-163, Lecture Notes in Math., 1998, Springer, Berlin, 2010. Also arXiv.math/0802.4432.
- [C1] F. Campana, *Réduction algébrique d'un morphisme faiblement Kählérien propre*, Math. Ann. 256 (1980), 157-189.
- [C2] F. Campana, *Réduction d'Albanese d'un morphisme propre et faiblement Kählérien et applications (I, II)*, Comp. Math. 54 (1985), 373-416.
- [C3] F. Campana, *Isotrivialité de certaines familles Kählériennes de variétés non projectives*, Math. Z. 252 (2006), 147-156 . Also arXiv.math/ 0408148.
- [C4] F. Campana, *Fundamental group and positivity of cotangent bundles of compact Kähler manifolds*, J. Algebraic Geom., 4(3):487-502, 1995.
- [D1] J.-P. Demailly, *Holomorphic Morse inequalities*, Several complex variables and complex geometry, Part 2 (Santa Cruz, CA, 1989), 93–114, Proc. Sympos. Pure Math., 52, Part 2, Amer. Math. Soc., Providence, RI, 1991.
- [D2] J.-P. Demailly, *Regularization of closed positive currents and Intersection Theory*, J. Alg. Geom. 1 (1992) 361-409
- [D3] J.-P. Demailly, *Analytic methods in algebraic geometry*, Lecture Notes, École d'été de Mathématiques de Grenoble "Géométrie des variétés projectives complexes : programme du modèle minimal" (June-July 2007)
- [DP] Demailly, J.-P., Paun, M., *Numerical characterization of the Kähler cone of a compact Kähler manifold*, math.AG/0105176, Annals of Mathematics, **159** (2004), 1247-1274.
- [DPe] J.-P. Demailly, Th. Peternell, *A Kawamata-Viehweg Vanishing Theorem on compact Kähler manifolds*, J. Differential Geom. **63** (2003), no. 2, 231277.
- [DPS] J.-P. Demailly, Th. Peternell, M. Schneider, *Pseudo-effective line bundles on compact Kähler manifolds*, International Journal of Math. **6** (2001), pp. 689-741.
- [E] I. Enoki, *Strong-Lefschetz-type theorem for semi-positive line bundles over compact Kähler manifolds*, Geometry and global analysis (Sendai, 1993), 211–212, Tohoku Univ., Sendai, 1993
- [Fl] A.R. Fletcher *Contributions to Riemann-Roch on projective threefolds with only canonical singularities* Proc. Symp. Pure. Math. 46 (1987), 221-231.
- [F] A. Fujiki. *On the structure of compact complex manifolds in \mathbb{C}* . Algebraic varieties and analytic varieties (Tokyo, 1981), 231-302, Adv. Stud. Pure Math., 1, North-Holland, Amsterdam, 1983.
- [HP] A. Höring, T. Peternell *Minimal Models for Kähler Threefolds*. arXiv/math.1304.4013.

- [HPR] A. Höring, T. Peternell, I. Radloff. *Uniformisation in dimension four: toward a conjecture of Iitaka*. arXiv/math. 1103.5392.
- [KV1] D. Kaledin, M. Verbitsky. *Hyperholomorphic sheaves and new examples of hyperkähler manifolds*. alg-geom 9712012, a chapter in a book “Hyperkähler manifolds” International Press, Boston, 2001.
- [KV2] D. Kaledin, M. Verbitsky. *Partial resolutions of Hilbert type, Dynkin diagrams, and generalized Kummer varieties*. 33 pages, arXiv:math/9812078
- [KM] J. Kollár, S. Mori. *Birational Geometry of Algebraic Varieties* Cambridge Tracts in Mathematics 134. CUP (1998).
- [Moo] R. Moosa, *The model theory of compact complex spaces*, Logic Colloquium 2001, volume 20 of Lect. Notes Log., pages 317-349, Assoc. Symbol. Logic, 2005.
- [MMT] R. Moosa, R. Moraru, M. Toma, *An essentially saturated surface not of Kaehler-type*, Bull. Lond. Math. Soc., volume 40 (2008), number 5, 845854.
- [MP] R. Moosa, A. Pillay, *Model theory and Kaehler geometry*, in: Model Theory with Applications to Algebra and Analysis Volume 1 (Eds. Z. Chatzidakis, D. Macpherson, A. Pillay, A. Wilkie), London Mathematical Society Lecture Note Series Nr 349, Cambridge University Press 2008, 167195.
- [Mou] Ch. Mourougane, *Théorèmes d’annulation génériques pour les fibrés vectoriels semi-négatifs*, Bull. Soc. Math. Fr. 127 (1999) 115–133.
- [P1] Th. Peternell, *Towards a Mori theory on compact Kähler 3-folds, II*. Math. Ann. 311 (1998), 729-764.
- [P2] Th. Peternell, *Towards a Mori theory on compact Kähler 3-folds, III*. Bull. Soc. Math. France. 129 (2001), 339-356.
- [Si1] Y.T. Siu., *Analyticity of sets associated to Lelong numbers and the extension of closed positive currents*, Invent. Math., 27 (1974), 53-156.
- [Si2] Y.T Siu, *A vanishing theorem for semi-positive line bundles over non-Kähler manifolds*. J. Diff. Geom. 19 (1984), 431-452.
- [U] K. Ueno, *Classification theory of compact analytic spaces and algebraic varieties*. LNM 439. Springer 1975.
- [T] K. Takegoshi, *On cohomology groups of nef line bundles tensorized with multiplier ideal sheaves on compact Kähler manifolds*, Osaka J. Math. 34 (1997) 783–802.
- [V98] M. Verbitsky, *Trianalytic subvarieties of the Hilbert scheme of points on a K3 surface*, Geom. Funct. Anal. 8 (1998), no. 4, 732782
- [V02] M. Verbitsky, *Coherent sheaves on general K3 surfaces and tori*, arXiv:math/0205210, Pure Appl. Math. Q. 4 (2008), no. 3, part 2, 651714.
- [V03] M. Verbitsky, *Coherent sheaves on generic compact tori*, math.AG/0310329, CRM Proc. and Lecture Notices vol. 38 (2004), 229-249.

[V09] M. Verbitsky, *Parabolic nef currents on hyperkaehler manifolds*,
arXiv:0907.4217, 22 pages.

FRÉDÉRIC CAMPANA INSTITUT ELIE CARTAN
UNIVERSITÉ HENRI POINCARÉ
BP 239
F-54506. VANDOEUVRE-LES-NANCY CÉDEX
ET: INSTITUT UNIVERSITAIRE DE FRANCE
frederic.campana@univ-lorraine.fr

JEAN-PIERRE DEMAILLY
ACADÉMIE DES SCIENCES, ET:
UNIVERSITÉ DE GRENOBLE I
INSTITUT FOURIER, LABORATOIRE DE MATHÉMATIQUES
UMR 5582 DU CNRS, BP 74
100 RUE DES MATHS, F-38402 SAINT-MARTIN D'HÈRES,
jean-pierre.demailly@ujf-grenoble.fr

MISHA VERBITSKY
LABORATORY OF ALGEBRAIC GEOMETRY, SU-HSE,
7 VAVILOVA STR. MOSCOW, RUSSIA, 117312
verbit@maths.gla.ac.uk, verbit@mcme.ru