



HAL
open science

Volume entropy of Hilbert Geometries

Gautier Berck, Andreas Bernig, Constantin Vernicos

► **To cite this version:**

Gautier Berck, Andreas Bernig, Constantin Vernicos. Volume entropy of Hilbert Geometries. Pacific Journal of Mathematics, 2010, 245, pp.201-225. 10.2140/pjm.2010.245.201 . hal-00819143

HAL Id: hal-00819143

<https://hal.science/hal-00819143v1>

Submitted on 5 Jun 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

VOLUME ENTROPY OF HILBERT GEOMETRIES

G. BERCK, A. BERNIG & C. VERNICOS

ABSTRACT. It is shown that among all plane Hilbert geometries, the hyperbolic plane has the maximal volume entropy. More precisely, it is shown that the volume entropy is bounded above by $\frac{2}{3-d} \leq 1$, where d is the Minkowski dimension of the extremal set of K . An explicit example of a plane Hilbert geometry with non-integer volume entropy is constructed. In arbitrary dimension, the hyperbolic space has maximal entropy among all Hilbert geometries satisfying some additional technical hypothesis. To achieve this result, a new projective invariant of convex bodies, similar to the centro-affine area, is constructed.

1. INTRODUCTION

In his famous 4-th problem, Hilbert asked to characterize metric geometries whose geodesics are straight lines. He constructed a special class of examples, nowadays called *Hilbert geometries* [20, 21]. These geometries have attracted a lot of interest, see for example the works of Y. Nasu [38], P. de la Harpe [16], A. Karlsson & G. Noskov [26], E. Socie-Methou [41], T. Foertsch & A. Karlsson [18], Y. Benoist [8], B. Colbois & C. Vernicos [13] and the two complementary surveys by Y. Benoist [6] and the last named author [44].

A Hilbert geometry is a particularly simple metric space on the interior of a compact convex set K (see definition below). This metric happens to be a complete Finsler metric whose set of geodesics contains the straight lines. Since the definition of the Hilbert geometry only uses cross-ratios, the Hilbert metric is a projective invariant. In the particular case where K is an ellipsoid, the Hilbert geometry is isometric to the usual hyperbolic space.

An important part of the above mentioned works, and of older ones, is to study how different or close to the hyperbolic geometry these geometries can be. For instance, if K is not an ellipsoid, then the metric is never Riemannian, see D.C. Kay [27, Corollary 1]. This last result is actually related to the fact that among all finite dimensional normed vector spaces, many notions of curvatures are only satisfied by the Euclidean spaces (see also P. Kelly & L. Paige [28], P. Kelly

MSC classification: 53C60, 53A20, 51F99

The first two authors were supported by the Schweizerischer Nationalfonds grants SNF PP002-114715/1 and 200020-121506 /1.

& E. Strauss [29, 30]). However, if ∂K is sufficiently smooth then the flag curvature, an analog of the sectional curvature, of the Hilbert metric is constant and equals -1 , see for example Z. Shen [40, Example 9.2.2]. Hence a question one can ask is whether or not these geometries behave like negatively curved Riemannian manifold. The example of the triangle geometry which is isometric to a two dimensional normed vector space (see P. De la Harpe [16]) shows that things are a little more involved (see also theorems cited below). The present work is partially inspired by the feeling that Hilbert geometries might be thought as geometries with Ricci curvature bounded from below, and focuses on the volume growth of balls.

Unlike the Riemannian case, where there is only one natural choice of volume, there are several good choices of volume on a Finsler manifold. We postpone this issue to section 2 and fix just one *volume* (like the n -dimensional Hausdorff measure) for the moment.

Let $B(o, r)$ be the metric ball of radius r centered at o . The volume entropy of K is defined by the following limit (provided it exists)

$$(1) \quad \text{Ent } K := \lim_{r \rightarrow \infty} \frac{\log \text{Vol } B(o, r)}{r}.$$

The entropy does not depend on the particular choice of the base point $o \in \text{int } K$ nor on the particular choice of the volume. If $h = \text{Ent } K$, then $\text{Vol } B(o, r)$ behaves roughly as e^{hr} .

It is well-known and easy to prove (see, e.g., S. Gallot, D. Hulin & J. Lafontaine [19, Section III.H]) that the volume of a ball of radius r in the n -dimensional hyperbolic space is given, with ω_n the volume of the Euclidean unit ball of dimension n , by

$$n\omega_n \int_0^r (\sinh s)^{n-1} ds = O(e^{(n-1)r}).$$

It follows that the entropy of an ellipsoid equals $n - 1$.

In general, it is not known whether the above limit exists. If the convex set K is divisible, which means that a discrete subgroup of the group of isometries of the Hilbert geometry acts cocompactly, then the entropy is known to exist, see Y. Benoist [7]. If the convex set is sufficiently smooth, e.g., C^2 with positive curvature suffices, then the entropy exists and equals $n - 1$ (see the theorem of B. Colbois & P. Verovic below). In general, one may define lower and upper entropies $\underline{\text{Ent}}$, $\overline{\text{Ent}}$ by replacing the limit in the definition (1) by \liminf or \limsup .

There is a well-known conjecture (whose origin seems difficult to locate) saying that the hyperbolic space has maximal entropy among all Hilbert geometries of the same dimension.

Conjecture. *For any n -dimensional Hilbert geometry,*

$$\overline{\text{Ent}} K \leq n - 1.$$

Notice that such a result is a consequence of Bishop's volume comparison theorem for a complete Riemannian manifold of Ricci curvature bounded by $-(n-1)$ (see [19, theorem 3.101, i]).

Several particular cases of the conjecture were treated in the literature. The following one shows that the volume entropy does not characterize the hyperbolic geometry among all Hilbert geometries.

Theorem. (B. Colbois & P. Verovic [15])

If K is C^2 -smooth with strictly positive curvature, then the Hilbert metric of K is bi-Lipschitz to the hyperbolic metric and therefore

$$\text{Ent } K = n - 1.$$

The case of convex polytopes is rather well understood.

Theorem. (A. Bernig [9], C. Vernicos [43])

The Hilbert metric associated to a convex body K is bi-Lipschitz to a normed space if and only if K is a polytope. In particular, the entropy of a polytope is 0.

The two-dimensional case of this theorem was earlier obtained by B. Colbois, C. Vernicos & P. Verovic in [14].

Instead of taking the volume of balls, another natural choice is to study the volume growth of the metric spheres $S(o, r)$. One may define a (spherical) entropy by

$$(2) \quad \text{Ent}^s K := \lim_{r \rightarrow \infty} \frac{\log \text{Vol } S(o, r)}{r},$$

provided the limit exists. In general, one may define upper and lower spherical entropies $\overline{\text{Ent}}^s K$ and $\underline{\text{Ent}}^s K$ by replacing the limits in the definition (2) by a lim sup or lim inf.

The following theorem is a spherical version of the theorem of B. Colbois & P. Verovic.

Theorem. (A.A. Borisenko & E.A. Olin [11])

If K is an n -dimensional convex body of class C^3 with positive Gauss curvature, then $\text{Ent}^s = n - 1$.

Our first main theorem treats the two-dimensional case. Recall that an extremal point of a convex body K is a point which is not a convex combination of two other points of K .

Theorem (First Main Theorem). *Let K be a two-dimensional convex body. Let d be the upper Minkowski dimension of the set of extremal points of K . Then the entropy of K is bounded by*

$$(3) \quad \overline{\text{Ent}} K \leq \frac{2}{3-d} \leq 1.$$

The inequality is sharp if K is smooth or contains some positively curved smooth part in the boundary. In this case the upper Minkowski dimension of $\text{ex} K$ and the entropy both are 1. On the other hand, for polygons the upper Minkowski dimension of the set of extremal points and the entropy both vanish (see the theorem of B. Colbois, C. Vernicos & P. Verovic above), and the inequality is not sharp in this case.

It should be noted that the entropy behaves in a rather subtle way (see also C. Vernicos [42] for a technical and complementary study, to this paper, of the entropy). As we have seen above, the entropy of a polygon vanishes. In contrast to this, we will construct a convex body with piecewise affine boundary whose entropy is between $\frac{1}{4}$ and $\frac{3}{4}$.

Our second main theorem applies in all dimensions. It weakens in a substantial way the assumptions in the theorem of B. Colbois & P. Verovic and strengthens its conclusions for not only does it give the precise value of the entropy but also the *entropy coefficient*. In order to state it, we introduce a projective invariant of convex bodies interesting in itself.

Let V be an n -dimensional vector space with origin o . Given a convex body K containing o in the interior, we define a positive function a on the boundary by the condition that for $p \in \partial K$ we have $-a(p)p \in \partial K$. The letter a stands for *antipodal*. If V is endowed with a Euclidean scalar product, we let $k(p)$ be the Gauss curvature and $n(p)$ be the outer normal vector at a boundary point p (whenever they are well-defined, which is almost everywhere the case following A.D. Alexandroff [1]).

Definition. *The centro-projective area of K is*

$$(4) \quad \mathcal{A}_p(K) := \int_{\partial K} \frac{\sqrt{k}}{\langle n, p \rangle^{\frac{n-1}{2}}} \left(\frac{2a}{1+a} \right)^{\frac{n-1}{2}} dA.$$

It is not quite obvious (but true, as we shall see) that this definition does not depend on the choice of the scalar product. In fact, the centro-projective area is invariant under *projective transformations* fixing the origin. The reader familiar with the theory of valuations may notice the similarity with the centro-affine surface area, whose definition is the same except that the second factor (containing the function a) does not appear. We refer to the books by Laugwitz [32] and Leichtweiss [34] for more information on affine and centro-affine differential geometry.

Theorem (Second Main Theorem). *If ∂K is $C^{1,1}$, then*

$$(5) \quad \lim_{r \rightarrow \infty} \frac{\text{Vol} B(o, r)}{\sinh^{n-1} r} = \frac{1}{n-1} \mathcal{A}_p(K) \neq 0$$

and $\text{Ent} K = n - 1$. Moreover, without any assumption on K , if $\mathcal{A}_p(K) \neq 0$ then $\underline{\text{Ent}} K \geq n - 1$.

Our next theorem, together with the previous ones, shows in particular that it suffices to assume K to be merely of class $C^{1,1}$ in the theorem of A.A. Borisenko & E.A. Olin.

Theorem. *For each convex body K ,*

$$\begin{aligned}\underline{\text{Ent}}^s K &= \underline{\text{Ent}} K, \\ \overline{\text{Ent}}^s K &= \overline{\text{Ent}} K.\end{aligned}$$

Plan of the paper. In the next section, we collect some well-known facts about convex bodies, Hilbert geometries and volumes on Finsler manifolds. A number of easy lemmas is proved which will be needed in the proof of our main theorem. Using some inequalities for volumes in normed spaces, we show that entropy and spherical entropy coincide for general convex bodies.

In section 3, we give the proofs of our main theorems. In the final section 4, we give an intrinsic definition of the centro-projective surface area and study some of its properties. In particular, we show that it is upper semi-continuous with respect to Hausdorff topology.

Acknowledgements. We wish to thank Bruno Colbois and Daniel Hug for interesting discussions and Franz Schuster for useful remarks on an earlier version of this paper.

2. PRELIMINARIES ON CONVEX BODIES AND HILBERT GEOMETRIES

2.1. Convex bodies. Let V be a finite-dimensional real vector space. By *convex body*, we mean a compact convex set $K \subset V$ with non-empty interior (note that this last condition is sometimes not required in the literature). Most of the time, the convex bodies will be assumed to contain the origin in their interiors. In such a case, we will call as usual *Minkowski functional* the positive, homogeneous of degree one function whose level set at height 1 is the boundary ∂K . It is a convex function and by Alexandroff's theorem, it admits a quadratic approximation almost everywhere (see e.g. A.D. Alexandroff [1] or L.C. Evans & R.F. Gariepy [17, p. 242]). In the following, boundary-points where Alexandroff's theorem applies will be called *smooth*. Assuming the vector space to be equipped with an inner product, the principal curvatures of the boundary and its Gauss curvature k are well defined at every smooth point.

We will be concerned with generalizations and variations of *Blaschke's rolling theorem*, a proof of which may be found in K. Leichtweiß [33].

Theorem 2.1 (W. Blaschke, [10]). *Let K be a convex body in \mathbb{R}^n whose boundary is C^2 with everywhere positive Gaussian curvature.*

Then there exist two positive radii R_1 and R_2 such that for every boundary point p , there exists a ball of radius R_1 (resp. R_2) containing p on its boundary and contained in K (resp. containing K).

We first remark that for the “inner part” of Blaschke’s result, the regularity of the boundary may be lowered. Recall that the boundary of a convex body is $C^{1,1}$ provided it is C^1 and the Gauss map is Lipschitz-continuous. Roughly speaking, the second condition says that the curvature of the boundary remains bounded, even if it is only almost everywhere defined. The following proposition then gives a geometrical characterization of such bodies, see L. Hörmander [22, proposition 2.4.3] or V. Bangert [4] and D. Hug [25].

Proposition 2.2. *The boundary of a convex body K is $C^{1,1}$ if and only if there exists some $R > 0$ such that K is the union of balls with radius R .*

Without assumption on the boundary, there is still an integral version of Blaschke’s rolling theorem.

Theorem 2.3 (C. Schütt & E. Werner, [39]). *For a convex body K containing the unit ball of a Euclidean space and $p \in \partial K$, let $R(p) \in [0, \infty)$ be the radius of the biggest ball contained in K and containing p . Then for all $0 < \alpha < 1$*

$$(6) \quad \int_{\partial K} R^{-\alpha} d\mathcal{H}^{n-1} < \infty.$$

We will need the following refinement of this theorem.

Proposition 2.4. *In the same situation as in Theorem 2.3, for each Borel subset $B \subset \partial K$ we have*

$$(7) \quad \int_B R^{-\alpha} d\mathcal{H}^{n-1} \leq 2(n-1)^\alpha \left(\frac{2^\alpha}{1-2^{\alpha-1}} \right)^\alpha (\mathcal{H}^{n-1}(B))^{1-\alpha} (\mathcal{H}^{n-1}(\partial K))^\alpha.$$

In particular for some constant C depending on K we have

$$(8) \quad \int_B R^{-\frac{1}{2}} d\mathcal{H}^{n-1} \leq C (\mathcal{H}^{n-1}(B))^{\frac{1}{2}}.$$

Proof. By ([39], Lemma 4), we have for $0 \leq t \leq 1$

$$(9) \quad \mathcal{H}^{n-1}(\{p \in \partial K | R(p) \leq t\}) \leq (n-1)t \mathcal{H}^{n-1}(\partial K),$$

from which we deduce that, for each $0 < \epsilon < 1$

$$\begin{aligned}
 (10) \quad \int_{\partial K \cap \{R < \epsilon\}} R^{-\alpha} d\mathcal{H}^{n-1} &= \sum_{i=0}^{\infty} \int_{\partial K \cap \{\epsilon 2^{-i-1} \leq R < 2^{-i}\epsilon\}} R^{-\alpha} d\mathcal{H}^{n-1} \\
 &\leq \sum_{i=0}^{\infty} (\epsilon 2^{-i-1})^{-\alpha} \mathcal{H}^{n-1}(\partial K \cap \{\epsilon 2^{-i-1} \leq R < 2^{-i}\epsilon\}) \\
 &\leq \sum_{i=0}^{\infty} (\epsilon 2^{-i-1})^{-\alpha} (n-1) 2^{-i} \epsilon \mathcal{H}^{n-1}(\partial K) \\
 &= \epsilon^{1-\alpha} (n-1) \frac{2^\alpha}{1-2^{\alpha-1}} \mathcal{H}^{n-1}(\partial K).
 \end{aligned}$$

It follows that

$$\begin{aligned}
 \int_B R^{-\alpha} d\mathcal{H}^{n-1} &= \int_{B \cap \{R < \epsilon\}} R^{-\alpha} d\mathcal{H}^{n-1} + \int_{B \cap \{R \geq \epsilon\}} R^{-\alpha} d\mathcal{H}^{n-1} \\
 &\leq \epsilon^{1-\alpha} (n-1) \frac{2^\alpha}{1-2^{\alpha-1}} \mathcal{H}^{n-1}(\partial K) + \epsilon^{-\alpha} \mathcal{H}^{n-1}(B).
 \end{aligned}$$

Choosing

$$\epsilon := \frac{1-2^{\alpha-1}}{2^\alpha(n-1)} \frac{\mathcal{H}^{n-1}(B)}{\mathcal{H}^{n-1}(\partial K)}$$

yields the inequality of the lemma. \square

2.2. Hilbert geometries. Given two distinct points $x, y \in \text{int } K$, the *Hilbert distance* between x and y is defined by

$$d(x, y) := \frac{1}{2} |\log[a, b, x, y]|,$$

where a and b are the intersections of the line passing through x and y with the boundary ∂K , and $[a, b, x, y]$ denotes the cross-ratio (with the convention of [12]).

This distance is invariant under projective transformations. If K is an ellipsoid, the Hilbert geometry on $\text{int } K$ is isometric to hyperbolic n -space.

Unbounded closed convex sets with non-empty interiors and not containing a straight line are projectively equivalent to convex bodies. Therefore, the definition of the distance naturally extends to the interiors of such convex sets. In particular the convex sets bounded by parabolas are also isometric to the hyperbolic space.

Let us assume the origin o lies inside the interior of K . We will write $B(r)$ for the *metric ball* of radius r and centered at o . Its boundary, the *metric sphere*, will be denoted by $S(r)$. Let $a: \partial K \rightarrow \mathbb{R}_+$ be defined by the equation

$$-a(p)p \in \partial K,$$

so the letter a refers to the antipodal point. It is an easy exercise to check that metric spheres are parameterized by the boundary ∂K as

$$S(r) = \{\phi(p, r) : p \in \partial K\},$$

where

$$(11) \quad \begin{aligned} \phi : \partial K \times \mathbb{R}_+ &\rightarrow \text{int } K \\ (p, r) &\mapsto a \frac{e^{2r} - 1}{ae^{2r} + 1} p. \end{aligned}$$

The Hilbert distance comes from a Finsler metric on the interior of K . Given $x \in \text{int } K$ and $v \in T_x V$, the Finsler norm of v is given by

$$(12) \quad \|v\|_x = \frac{1}{2} \left(\frac{1}{t_1} + \frac{1}{t_2} \right),$$

where $t_1, t_2 > 0$ are such that $x \pm t_i v \in \partial K$. Again, we do not exclude that one of the t_i 's is infinite. Equivalently, if F_x is the Minkowski functional of $K - x$, then

$$\|v\|_x = \frac{1}{2} (F_x(v) + F_x(-v)).$$

The Finsler metric makes it possible to measure the length of a differentiable curve $c : I \rightarrow \text{int } K$ by

$$l(c) := \int_I \|c'(t)\|_{c(t)} dt.$$

It is less trivial to measure the area (or volume) of higher dimensional subsets of $\text{int } K$. In fact, different notions of volume are being used. The most important ones are the Busemann definition (which equals the Hausdorff n -dimensional measure) and the Holmes-Thompson definition. In the following, only the axioms of a *volume* as defined in [3] will be used. We will make use of the following properties:

- Vol is a Borel measure on $\text{int } K$ which is absolutely continuous with respect to Lebesgue measure.
- If $A \subset K \subset L$, where K, L are compact convex sets, then the measure of A with respect to K is larger than the measure of A with respect to L .
- If K is an ellipsoid, then $\text{Vol}(A)$ is the hyperbolic volume of A .

The following projective invariants of convex bodies will be our main subjects of investigation.

Definition 2.5. The *upper (resp. lower) volume entropy* of K is

$$\begin{aligned} \overline{\text{Ent}}(K) &:= \limsup_{r \rightarrow \infty} \frac{\log(\text{Vol } B(r))}{r}, \\ \underline{\text{Ent}}(K) &:= \liminf_{r \rightarrow \infty} \frac{\log(\text{Vol } B(r))}{r}. \end{aligned}$$

If the upper and lower volume entropies of K coincide, their common value is called volume entropy of K and denoted by $\text{Ent } K$.

Note that these invariants are independent of the choice of the center and of the choice of the volume definition.

2.3. Busemann's density. For simplicity, we restrict ourselves to Busemann's volume, although all results remain true for every other choice of volume. The reason is that the proofs of the crucial propositions 2.7 and 2.8 below do not use any particular property of Busemann's volume, but only the axioms satisfied by every definition of volume.

The density of Busemann's volume (with respect to some Lebesgue measure \mathcal{L}) is given by

$$\sigma(x) = \frac{\omega_n}{\mathcal{L}(B_x)},$$

where B_x is the tangent unit ball of the Finsler metric at x and ω_n is the (Euclidean) volume of the unit ball in \mathbb{R}^n . The volume of a Borel subset $A \subset \text{int } K$ is thus given by

$$\text{Vol}(A) = \int_A \sigma \, d\mathcal{L}.$$

We now state and prove some propositions concerning upper bounds and asymptotic behaviors of Busemann's densities for points which are close to the boundary of particular convex sets. We will make use of an auxiliary inner product, calling \mathcal{L} and μ the corresponding Lebesgue measure and volume n -form. Busemann densities are defined with this particular choice of measure.

Proposition 2.6. *Let K, K' be closed convex sets not containing any straight line and $\sigma : \text{int } K \rightarrow \mathbb{R}$, $\sigma' : \text{int } K' \rightarrow \mathbb{R}$ their corresponding Busemann densities. Let $p \in \partial K$, E_0 a support hyperplane of K at p and E_1 a hyperplane parallel to E_0 intersecting K . Suppose that K and K' have the same intersection with the strip between E_0 and E_1 (in particular $p \in \partial K'$). Then*

$$\lim_{y \rightarrow p} \frac{\sigma(y)}{\sigma'(y)} = 1.$$

Proof. Let d be the distance between E_0 and E_1 and (y_i) a sequence of points of $\text{int } K$ converging to p . We may suppose that the distance d_i between y_i and E_0 is strictly less than d . For every fixed point y_i and non-zero tangent vector $v \in T_{y_i}K$, let $t_1, t_2 \in \mathbb{R}_+ \cup \{\infty\}$ be such that $y_i \pm t_{1,2}v \in \partial K$; let t'_1, t'_2 be the corresponding numbers for K' . Since at least one of $y_i + t_1v$ and $y_i - t_2v$ is inside the strip, say $y_i + t_1v$, we must have $t_1 = t'_1$.

Either $t_2 = t'_2$ and $\|v\|_i = \|v\|'_i$, or $t_2 \neq t'_2$, in which case

$$\frac{t_1}{t_2}, \frac{t'_1}{t'_2} \leq \frac{d_i}{d - d_i}.$$

Therefore,

$$\frac{d - d_i}{d} \leq \frac{\|v\|_i}{\|v\|'_i} = \frac{1 + \frac{t_1}{t_2}}{1 + \frac{t'_1}{t'_2}} \leq \frac{d}{d - d_i}$$

which shows that, as functions on $\mathbb{R}P^{n-1}$, $\|\cdot\|_i/\|\cdot\|'_i$ uniformly converge to 1. Hence, for every ϵ and every i large enough,

$$(1 - \epsilon)B_{y_i} \subset B'_{y_i} \subset (1 + \epsilon)B_{y_i},$$

which implies the convergence of σ/σ' to 1. \square

Proposition 2.7. *Let $V = \mathbb{R}^n$ with its usual scalar product. Let P be the convex set bounded by the parabola $y = \sum_{i=1}^{n-1} \frac{c_i}{2} x_i^2$, $c_1, \dots, c_{n-1} > 0$. Then*

$$(13) \quad \sigma(0, \dots, 0, 1 - \lambda) = \frac{\sqrt{c}}{(2(1 - \lambda))^{\frac{n+1}{2}}},$$

where $c = \prod_{i=1}^{n-1} c_i$.

Proof. By the invariance of the Hilbert metric under projective transformations, the tangent unit sphere at any point of $\text{int } P$ is an ellipse. At the point $(0, \dots, 0, 1 - \lambda)$, the symmetry implies that the principal axes of this ellipse are parallel to the coordinate axes. Hence

$$\sigma = \frac{1}{\prod_{i=1}^n l_i},$$

where the l_i 's, $i = 1, \dots, n$, are the Euclidean lengths of the principal half-axes.

Now $l_i = \sqrt{\frac{2(1-\lambda)}{c_i}}$, $i = 1, \dots, n - 1$ and $l_n = 2(1 - \lambda)$. \square

Proposition 2.8. *Assume the origin o is inside $\text{int } K$. For a smooth point p of ∂K , let $n(p)$ be the outward normal vector and let $k(p)$ be the Gauss curvature of ∂K at p . Then*

$$(14) \quad \lim_{\lambda \rightarrow 1} \sigma(\lambda p)(1 - \lambda)^{\frac{n+1}{2}} = \frac{\sqrt{k(p)}}{(2\langle p, n(p) \rangle)^{\frac{n+1}{2}}}.$$

Proof. Let us choose a frame $(p; v_1, \dots, v_{n-1}, v_n)$ where $v_1, \dots, v_{n-1} \in T_p \partial K$ are unit vectors tangent to the principal curvature directions of ∂K at p and $v_n = -p$. In these coordinates, the boundary of K is

locally the graph of a function: $y = \sum_{i=1}^{n-1} \frac{c_i}{2} x_i^2 + R(|x|)$ with $R(|x|) = o(|x|^2)$ and $c_1, \dots, c_{n-1} \geq 0$. We set

$$c := \prod_{i=1}^{n-1} c_i.$$

A small computation shows that

$$dx_1 \wedge \dots \wedge dx_{n-1} \wedge dy = \frac{1}{m} \mu,$$

where μ is the Euclidean n -form and $m := \mu(v_1, \dots, v_n) = \langle p, n(p) \rangle$. Also, the Gauss curvature at p is given by

$$k(p) = cm^{n-1}.$$

Let us fix $\epsilon > 0$. Locally, the parabola defined by

$$y = \sum_{i=1}^{n-1} \frac{c_i + \epsilon}{2} x_i^2$$

lies inside K . Cutting it with some horizontal hyperplane, we obtain a convex body K' inside K . In particular, the metric of K' is greater than or equal to the metric of K , hence $\sigma'(\lambda p) \geq \sigma(\lambda p)$ for λ near 1.

Then by propositions 2.6 and 2.7,

$$\begin{aligned} \limsup_{\lambda \rightarrow 1} \sigma(\lambda p)(1 - \lambda)^{\frac{n+1}{2}} &\leq \lim_{\lambda \rightarrow 1} \sigma'(\lambda p)(1 - \lambda)^{\frac{n+1}{2}} \\ (15) \qquad \qquad \qquad &= \frac{\sqrt{\prod_{i=1}^{n-1} (c_i + \epsilon)}}{2^{\frac{n+1}{2}} m}. \end{aligned}$$

Note that $\sigma > 0$, hence this already settles the case $k = c = 0$ since ϵ was arbitrary small.

If $c > 0$ and $0 < \epsilon < \min\{c_1, \dots, c_{n-1}\}$, the parabola P defined by

$$y = \sum_{i=1}^{n-1} \frac{c_i - \epsilon}{2} x_i^2$$

locally contains K . Cutting it with some horizontal hyperplane, we obtain a convex body K' inside P . By propositions 2.6 and 2.7 again,

$$\begin{aligned} \liminf_{\lambda \rightarrow 1} \sigma(\lambda p)(1 - \lambda)^{\frac{n+1}{2}} &\geq \liminf_{\lambda \rightarrow 1} \sigma'(\lambda p)(1 - \lambda)^{\frac{n+1}{2}} \\ (16) \qquad \qquad \qquad &= \frac{\sqrt{\prod_{i=1}^{n-1} (c_i - \epsilon)}}{2^{\frac{n+1}{2}} m}. \end{aligned}$$

From (15) and (16) (with $\epsilon \rightarrow 0$) we get

$$\lim_{\lambda \rightarrow 1} \sigma(\lambda p)(1 - \lambda)^{\frac{n+1}{2}} = \frac{\sqrt{c}}{2^{\frac{n+1}{2}} m}.$$

□

To state precisely our main theorem in section 3 we need to introduce the pseudo-Gauss curvature of the boundary of a convex set K in \mathbb{R}^n .

For a smooth point $p \in \partial K$, let $n(p)$ be the outward normal of ∂K at p . For each unit vector $e \in T_p \partial K$, let $H_e(p)$ be the affine plane containing p and directed by the vectors e and $n(p)$. We define R_e as the radius of the biggest disc containing p inside $K_e := K \cap H_e(p)$.

Definition 2.9. The *pseudo Gauss-curvature* $\bar{k}(p)$ of ∂K at p is the minimum of the numbers

$$\prod_{i=1}^{n-1} R_{e_i}(p)^{-1},$$

where e_1, \dots, e_{n-1} ranges over all orthonormal bases of $T_p \partial K$.

Proposition 2.10. *Let V be a Euclidean vector space of dimension n . Let K be a convex body containing the unit ball B . Then for $\frac{1}{2} \leq \lambda < 1$ and $p \in \partial K$*

$$(17) \quad \sigma(\lambda p) \leq \frac{\omega_n n!}{2^n (1-\lambda)^{\frac{n+1}{2}}} \bar{k}(p)^{1/2}.$$

Proof. We use the same notation as in the definition of \bar{k} . We may suppose that for all i , $R_i := R_{e_i}(p) > 0$, otherwise the statement is trivial. By definition of R_i , there is a 2-disc $B_i(p)$ of radius R_i inside K_{e_i} containing p . Let us denote by $B(e_i)$ the intersection of B with the affine plane $p + H_{e_i}$. Since $B(e_i), B_i(p) \subset K$, one has

$$\hat{C}_i := \text{conv}(B(e_i) \times \{0\} \cup B_i(p) \times \{1\}) \subset K_{e_i} \times [0, 1].$$

Note that \hat{C}_i is a truncated cone. Let E_i be the plane containing the line that is parallel to $T_p \partial K_{e_i}$ and that passes through the points $o \times \{0\}$ and $p \times \{1\}$. With $\pi : V \times [0, 1] \rightarrow V$ the projection on the first component, $C_i := \pi(E_i \cap \hat{C}_i) \subset K$ is bounded by a truncated conic.

In the non-orthogonal frame $(o; p, e_i)$, C_i is given by

$$(2R_i - 1)x^2 + 2(1 - R_i)x + y_1^2 \leq 1, \quad 0 \leq x \leq 1.$$

Now let C be the convex hull of the union of the C_i . Then the polytope P with vertices

$$\left(\lambda, 0, \dots, \pm \sqrt{(1-\lambda)(2\lambda R_i - \lambda + 1)}, 0, \dots, 0 \right), (1, \vec{0}), (2\lambda - 1, \vec{0})$$

lies inside C , with all but the last vertex being on the boundaries of the respective C_i 's.

Its volume is given by

$$\begin{aligned}
 \mathcal{L}(P) &= \frac{2^n \langle p, n(p) \rangle}{n!} (1 - \lambda)^{\frac{n+1}{2}} \prod_{i=1}^{n-1} (2\lambda R_i - \lambda + 1)^{\frac{1}{2}} \\
 &\geq \frac{2^n}{n!} (1 - \lambda)^{\frac{n+1}{2}} (R_1 \cdot R_2 \cdots R_{n-1})^{\frac{1}{2}} \\
 (18) \quad &= \frac{2^n}{n!} (1 - \lambda)^{\frac{n+1}{2}} \bar{k}^{-\frac{1}{2}}(p).
 \end{aligned}$$

The factor $\langle p, n(p) \rangle$ in the first line is due to the fact that our coordinate system is not orthonormal. Since the unit ball is contained in K , this factor is at least 1.

From $P \subset C \subset K$ and the fact that P is centered at λp , we deduce that

$$\sigma(\lambda p) \leq \frac{\omega_n}{\mathcal{L}(P)} \leq \frac{\omega_n n!}{2^n} (1 - \lambda)^{-\frac{n+1}{2}} \bar{k}^{-\frac{1}{2}}(p).$$

□

The next proposition will be needed in the construction of a convex body with entropy between 0 and 1.

Proposition 2.11. *Let $K = oab$ be a triangle with $1 \leq oa, ob \leq 2$ and such that the distance from o to the line passing through a and b is at least 1. Let p be a point in the interior of the side ab and suppose that $\min\{ap, bp\} \geq \epsilon > 0$. Then for $\lambda \geq \frac{1}{2}$ Busemann's density of K at λp is bounded above by*

$$\sigma(\lambda p) \leq 32\pi \max \left\{ \frac{1}{\epsilon(1 - \lambda)}, \frac{1}{\epsilon^2} \right\}.$$

Proof. The hypothesis on the triangle implies that $\sin(abo), \sin(bao) \geq \frac{1}{2}$.

Let a' be the intersection of the line passing through a and $z := \lambda p$ with ob and define b' similarly.

The unit tangent ball at z is a hexagon centered at z . The length of one of its half-diagonals is the harmonic mean of za and za' ; the length of the second half-diagonal is the harmonic mean of zb and zb' and the third half-diagonal has length $\frac{2op}{\frac{1}{\lambda} + \frac{1}{1-\lambda}} \geq 1 - \lambda$.

An easy geometric argument shows that $za', zb \geq \frac{1}{2}pb \sin(abo) \geq \frac{1}{4}\epsilon$ and $za, zb' \geq \frac{1}{2}pa \sin(bao) \geq \frac{1}{4}\epsilon$.

The area A of the hexagon is at least half of the minimal product of two of its half-diagonals, hence

$$A \geq \min \left\{ \frac{1}{8}\epsilon(1 - \lambda), \frac{1}{32}\epsilon^2 \right\}.$$

□

2.4. Volume entropy of spheres. By definition, the entropy controls the volume growth of metric balls in Hilbert geometries. We show in this section that it coincides with the growth of areas of metric spheres. Again, there are several definitions of area of hypersurfaces in Finsler geometry. For simplicity, we consider Busemann's definition which gives the Hausdorff $(n - 1)$ -measure of these hypersurfaces.

We will need the following two lemmas:

Lemma 2.12 (Rough monotonicity of area). *There exist a monotone function f and a constant $C_1 > 1$ such that for all $r > 0$*

$$(19) \quad C_1^{-1}f(r) \leq \text{Area}(S(r)) \leq C_1f(r).$$

Proof. Let $f(r)$ be the Holmes-Thompson area of $S(r)$. Since all area definitions agree up to some universal constant, inequality (19) is trivial. It remains to show that f is monotone.

If ∂K is C^2 with everywhere positive Gaussian curvature then the tangent unit spheres of the Finsler metric are quadratically convex. According to [2, theorem 1.1 and remark 2] there exists a Crofton formula for the Holmes-Thompson area, from which the monotonicity of f easily follows.

Such smooth convex bodies are dense in the set of all convex bodies for the Hausdorff topology (see e.g. [22, lemma 2.3.2]). By approximation, it follows that f is monotone for arbitrary K . \square

Lemma 2.13 (Co-area inequalities). *There exists a constant $C_2 > 1$ such that for all $r > 0$*

$$C_2^{-1}\text{Area}(S(r)) \leq \frac{\partial}{\partial r} \text{Vol}(B(r)) \leq C_2\text{Area}(S(r)).$$

Proof. Let $\mu := \sigma dx_1 \wedge \cdots \wedge dx_n$ be the volume form, and let α be the $n - 1$ -form on $S(r)$ whose integral equals the area.

Since

$$\text{Vol}(B(r)) = \int_0^r \int_{S(s)} i_{\partial_r} \mu \, ds,$$

where ∂_r at $\lambda p \in S(s)$ is the tangent vector multiple of $\vec{\sigma p}$ with unit Finsler norm, we have to compare $i_{\partial_r} \mu$ and α .

We will assume that $S(r)$ is differentiable at λp . The section of the unit tangent ball by the tangent space $T_{\lambda p}S(r)$ will be called γ . By definition of Busemann area, the area of γ measured with the form α is the constant

$$\alpha(\gamma) = \omega_{n-1}.$$

In the same way, calling Γ the half unit ball containing ∂_r and bounded by γ , one has

$$\mu(\Gamma) = \frac{1}{2}\omega_n.$$

Since Γ is convex it contains the cone with base γ and vertex ∂_r . Therefore,

$$(20) \quad \frac{1}{n} i_{\partial_r} \mu(\gamma) \leq \frac{1}{2} \omega_n.$$

By Brunn's theorem (see e.g. [31, theorem 2.3]), the sections of the tangent unit ball with hyperplanes parallel to γ have an area lesser than or equal to the area of γ . Also the tangent unit ball has a supporting hyperplane at ∂_r which is parallel to γ . Therefore, by Fubini's theorem, the cylinder $\gamma \times ([0, 1] \cdot \partial_r)$ has a volume greater than or equal to the volume of Γ (even if it generally does not contain Γ). Hence,

$$(21) \quad \frac{1}{2} \omega_n \leq i_{\partial_r} \mu(\gamma).$$

Inequalities (20) and (21) give

$$\frac{1}{2} \frac{\omega_n}{\omega_{n-1}} \alpha(\gamma) \leq i_{\partial_r} \mu(\gamma) \leq \frac{n}{2} \frac{\omega_n}{\omega_{n-1}} \alpha(\gamma),$$

from which the result easily follows. \square

Theorem 2.14. *The spherical entropy coincides with the entropy. More precisely,*

$$\begin{aligned} \limsup_{r \rightarrow \infty} \frac{\log \text{Area}(S(r))}{r} &= \overline{\text{Ent}} K, \\ \liminf_{r \rightarrow \infty} \frac{\log \text{Area}(S(r))}{r} &= \underline{\text{Ent}} K. \end{aligned}$$

Proof. For convenience, let

$$\begin{aligned} V(r) &:= \text{Vol } B(r), \\ A(r) &:= \text{Area } S(r). \end{aligned}$$

Using the previous two lemmas, one has for all $r > 0$

$$\begin{aligned} V(r) &= \int_0^r V'(s) ds \leq C_2 \int_0^r A(s) ds \leq C_1 C_2 \int_0^r f(s) ds \\ &\leq C_1 C_2 f(r) r \leq C_1^2 C_2 A(r) r. \end{aligned}$$

It follows that

$$\begin{aligned} \overline{\text{Ent}} K &= \limsup_{r \rightarrow \infty} \frac{\log V(r)}{r} \leq \limsup_{r \rightarrow \infty} \frac{\log C_1^2 C_2 A(r) r}{r} \\ &= \limsup_{r \rightarrow \infty} \frac{\log \text{Area}(S(r))}{r}. \end{aligned}$$

Similarly, for each $\epsilon > 0$

$$\begin{aligned} V(r(1+\epsilon)) &= \int_0^{r(1+\epsilon)} V'(s) ds \geq C_1^{-1} C_2^{-1} \int_0^{r(1+\epsilon)} f(s) ds \\ &\geq C_1^{-1} C_2^{-1} \int_r^{r(1+\epsilon)} f(s) ds \geq C_1^{-1} C_2^{-1} f(r) r \epsilon \geq C_1^{-2} C_2^{-1} A(r) r \epsilon \end{aligned}$$

and hence

$$\begin{aligned} (1+\epsilon) \overline{\text{Ent}} K &= (1+\epsilon) \limsup_{r \rightarrow \infty} \frac{\log V(r(1+\epsilon))}{r(1+\epsilon)} \geq \limsup_{r \rightarrow \infty} \frac{\log C_2^{-1} C_1^{-2} A(r) r \epsilon}{r} \\ &= \limsup_{r \rightarrow \infty} \frac{\log \text{Area}(S(r))}{r}. \end{aligned}$$

Letting $\epsilon \rightarrow 0$ gives the first equality. The second one follows in a similar way. \square

3. ENTROPY BOUNDS

3.1. Upper entropy bound in arbitrary dimension. We may now state and prove the second main theorem.

Theorem 3.1. *Let K be an n -dimensional convex body and $o \in \text{int } K$. For any point $p \in \partial K$ we denote by $\bar{k}(p)$ its pseudo-Gauss curvature as in definition 2.9. If*

$$(22) \quad \int_{\partial K} \bar{k}^{\frac{1}{2}}(p) dp < \infty,$$

then

$$(23) \quad \lim_{r \rightarrow \infty} \frac{\text{Vol } B(o, r)}{\sinh^{n-1} r} = \frac{1}{n-1} \mathcal{A}_p(K).$$

In particular,

$$\overline{\text{Ent}} K \leq n-1,$$

and if $\mathcal{A}_p(K) \neq 0$, then $\overline{\text{Ent}} K = n-1$.

Proof. Using the parameterization (11), the volume of metric balls is given by

$$\text{Vol}(B(r)) = \int_0^r \int_{\partial K} F(p, r) d\mathcal{H}^{n-1},$$

where

$$F(p, r) := \sigma(\phi(p, r)) \text{Jac } \phi(p, r).$$

The Jacobian may be explicitly computed:

$$\text{Jac } \phi(p, r) = \frac{(e^{2r} - 1)^{n-1} e^{2r}}{(ae^{2r} + 1)^{n+1}} 2a^n (1+a) \langle p, n(p) \rangle.$$

In particular,

$$(24) \quad \lim_{r \rightarrow \infty} e^{2r} \text{Jac } \phi(p, r) = \frac{2(1+a)\langle p, n(p) \rangle}{a}.$$

On the other hand, for each smooth boundary point p we have, by proposition 2.8,

$$(25) \quad \lim_{r \rightarrow \infty} \frac{\sigma(\phi(p, r))}{e^{(n+1)r}} = \frac{\sqrt{k(p)} a^{\frac{n+1}{2}}}{\left(2\langle p, n(p) \rangle\right)^{\frac{n+1}{2}} (1+a)^{\frac{n+1}{2}}}.$$

Then, by proposition 2.10 and the hypothesis (22),

$$(26) \quad \begin{aligned} \lim_{r \rightarrow \infty} \frac{1}{e^{(n-1)r}} \int_{\partial K} F(p, r) d\mathcal{H}^{n-1} &= \int_{\partial K} \lim_{r \rightarrow \infty} \frac{F(p, r)}{e^{(n-1)r}} d\mathcal{H}^{n-1} \\ (27) \quad &= \int_{\partial K} \lim_{r \rightarrow \infty} \frac{\sigma(\phi(p, r))}{e^{(n+1)r}} \lim_{r \rightarrow \infty} e^{2r} \text{Jac } \phi(p, r) d\mathcal{H}^{n-1} \\ &= \int_{\partial K} \frac{\sqrt{k(p)}}{\left(2\langle p, n(p) \rangle\right)^{\frac{n-1}{2}}} \left(\frac{a}{1+a}\right)^{\frac{n-1}{2}} d\mathcal{H}^{n-1} \\ &= \frac{1}{2^{n-1}} \mathcal{A}_p(K). \end{aligned}$$

By L'Hospital's rule we get

$$\lim_{r \rightarrow \infty} \frac{\text{Vol}(B(r))}{e^{(n-1)r}} = \lim_{r \rightarrow \infty} \frac{\int_0^r \int_{\partial K} F(p, s) d\mathcal{H}^{n-1} ds}{(n-1) \int_0^r e^{(n-1)s} ds} = \frac{1}{2^{n-1}(n-1)} \mathcal{A}_p(K).$$

□

Remark: The metric balls $B(r)$ are projective invariants of K . There is an affine version of the previous theorem using the affine balls $B_a(r) := \tanh(r)K$ (where multiplication is with respect to the center o). Under the same assumptions as in theorem 3.1, we obtain that

$$\lim_{r \rightarrow \infty} \frac{\text{Vol } B_a(r)}{e^{(n-1)r}} = \frac{1}{2^{n-1}(n-1)} \mathcal{A}_a(K)$$

where $\mathcal{A}_a(K)$ is the centro-affine area (see section 4). The proof goes as the previous one by replacing the function a by 1.

Corollary 3.2. *Suppose K is an n -dimensional convex body of class $C^{1,1}$. Then*

$$\text{Ent } K = n - 1.$$

Proof. For any $p \in \partial K$, $R(p)$ is the biggest radius of a ball in K containing p . By proposition 2.2, there exists a constant $R > 0$ such that $R(p) \geq R$ for all $p \in \partial K$. It follows that the hypothesis (22) is satisfied and therefore $\text{Ent } K \leq n - 1$.

The Gauss map $\mathcal{G}: \partial K \rightarrow S^{n-1}$ is well-defined and continuous. As a consequence of theorem 2.3 in Hug [24] and equation 2.7 in Hug [23], the standard measure on the unit sphere is the push-forward of $k \cdot d\mathcal{H}^{n-1}$, i.e.

$$\mathcal{G}_*(k \cdot d\mathcal{H}^{n-1}|_{\partial K}) = d\mathcal{H}^{n-1}|_{S^{n-1}},$$

hence the curvature has a positive integral. Therefore, $\mathcal{A}_p(K) > 0$, and equation (23) implies that $\text{Ent } K = n - 1$. \square

Corollary 3.3. *If K is an arbitrary n -dimensional convex body with $\mathcal{A}_p(K) \neq 0$, then $\overline{\text{Ent}} K \geq n - 1$.*

Proof. Arguing as in the proof of theorem 3.1 and using Fatou's lemma instead of the dominated convergence theorem gives the result. \square

3.2. The plane case. Let us now assume that $n = 2$. By theorem 2.3, the hypothesis (22) is satisfied for each convex body K . Therefore

$$(28) \quad \overline{\text{Ent}} K \leq 1$$

and

$$\lim_{r \rightarrow \infty} \frac{\text{Vol } B(o, r)}{\sinh r} = \mathcal{A}_p(K).$$

Next, we are going to prove a better bound for $\overline{\text{Ent}} K$. In order to state our main result, we need to recall some basic notions of measure theory in a Euclidean space and refer to P. Mattila [37] for details. For a non-empty bounded set A , let $N(A, \epsilon)$ be the minimal number of ϵ -balls needed to cover A . Then the upper Minkowski dimension of A is defined as

$$\overline{\text{dim}} A := \inf \left\{ s : \limsup_{\epsilon \rightarrow 0} N(A, \epsilon) \epsilon^s = 0 \right\}.$$

One should note that this dimension is invariant under bi-Lipschitz maps. In particular, it does not depend on a particular choice of inner product and moreover it is invariant under projective maps provided the considered subsets are bounded.

Recall that a point $p \in K$ is called *extremal* if it is not a convex combination of other points of K . The set of extremal points is a subset of ∂K , which we denote by $\text{ex } K$.

Theorem 3.4. *Let K be a plane convex body and d be the upper Minkowski dimension of $\text{ex } K$. Then the entropy of K is bounded by*

$$\overline{\text{Ent}} K \leq \frac{2}{3-d} \leq 1.$$

Proof. Since the entropy is independent of the choice of the center, we may suppose that the Euclidean unit ball around o is the maximum

volume ellipsoid inside K . Then K is contained in the ball of radius 2 (see [5]).

Set $\epsilon := e^{-\alpha r}$, where $\alpha \leq 1$ will be fixed later. Divide the boundary of K into two parts:

$$\partial K = \mathcal{B} \cup \mathcal{G},$$

where \mathcal{B} (the bad part) is the closed ϵ -neighborhood around the set of extremal points of K and \mathcal{G} (the good part) is its complement.

Using proposition 2.4 and equalities (24), (25), we get the following upper bound for large values of r ,

$$(29) \quad \int_{\frac{r}{2}}^r \int_{\mathcal{B}} \sigma(\phi(p, s)) \text{Jac } \phi(p, s) d\mathcal{H}^1 ds \leq O\left(e^r \sqrt{\mathcal{H}^1(\mathcal{B})}\right).$$

Next, let $p \in \mathcal{G}$. The endpoints of the maximal segment in ∂K containing p are extremal points of K and hence of distance at least ϵ from p . Therefore K contains a triangle as in proposition 2.11 and if $s \geq r/2$ and r is sufficiently large

$$\sigma(\phi(p, s)) = \sigma(\lambda \cdot p) \leq 32 \max\left\{\frac{1}{\epsilon(1-\lambda)}, \frac{1}{\epsilon^2}\right\} = \frac{32}{\epsilon(1-\lambda)}.$$

Integrating this from $r/2$ to r yields

$$(30) \quad \int_{\frac{r}{2}}^r \int_{\mathcal{G}} \sigma(\phi(p, s)) \text{Jac } \phi(p, s) d\mathcal{H}^1 ds = O(e^{\alpha r}).$$

Let d be the upper Minkowski dimension of the set of extremal points of K . Then, for each $\eta > 0$, $N(\text{ex } K, \epsilon) = o(\epsilon^{-d-\eta})$ as $\epsilon \rightarrow 0$. By definition of N , there is a covering of $\text{ex } K$ by $N(\text{ex } K, \epsilon)$ balls of radius ϵ . Hence there is a covering of \mathcal{B} by $N(\text{ex } K, \epsilon)$ balls of radius 2ϵ . The intersection of a 2ϵ -ball with ∂K has length less than $4\pi\epsilon$. It follows that

$$\mathcal{H}^1(\mathcal{B}) = o(\epsilon^{-d-\eta+1}).$$

Since the volume of $B(r/2)$ is bounded by $O(e^{r/2})$ (see (28)), the volume of $B(r)$ is bounded by

$$\begin{aligned} \text{Vol } B(r) &= \text{Vol } B(r/2) + \int_{\frac{r}{2}}^r \int_{\mathcal{B}} \sigma(\phi(p, s)) \text{Jac } \phi(p, s) d\mathcal{H}^1 ds \\ &\quad + \int_{\frac{r}{2}}^r \int_{\mathcal{G}} \sigma(\phi(p, s)) \text{Jac } \phi(p, s) d\mathcal{H}^1 ds \\ &= O(e^{\frac{r}{2}}) + O\left(e^{r(1-\frac{\alpha(1-d-\eta)}{2})}\right) + O(e^{\alpha r}). \end{aligned}$$

We fix α such that $1 - \alpha\frac{1-d-\eta}{2} = \alpha$, i.e. $\alpha := \frac{2}{3-d-\eta} > \frac{2}{3}$. Then

$$\text{Vol } B(r) = O(e^{\alpha r}),$$

which implies that the (upper) entropy of K is bounded by α . Since $\eta > 0$ was arbitrary, the result follows. \square

3.3. An example of non-integer entropy. We will construct an example of a plane convex body with piecewise affine boundary whose entropy is strictly between 0 and 1.

Let us choose a real number $s > 2$ and set $\alpha_i := \frac{C_s}{i^s}$ where $C_s > 0$ is sufficiently small such that

$$3 \sum_{i=1}^{\infty} \alpha_i < \pi.$$

Consider a centrally symmetric sequence E of points on S^1 such that the angles between consecutive points are $\alpha_1, \alpha_1, \alpha_1, \alpha_2, \alpha_2, \alpha_2, \dots$ (each angle appearing three times).

Theorem 3.5. *The entropy of $K = \text{conv}(E)$ is bounded by*

$$0 < \frac{1}{s} \leq \underline{\text{Ent}}K \leq \overline{\text{Ent}}K \leq \frac{2s-2}{3s-4} < 1.$$

Proof. Lower bound

The unit sphere of radius r in the Hilbert geometry K is $\tanh rK$ and consists of an infinite number of segments.

An easy geometric computation shows that the middle segment $S_i(r)$ corresponding to $\alpha := \alpha_i$ has for each $r \geq 0$ length bounded from below by

$$l(S_i(r)) \geq \log \left(\frac{2 \tanh r}{1 - \tanh r} \tan(\alpha/2) \sin(\alpha) + 1 \right).$$

Set

$$i_0(r) := \left\lfloor (2C_s)^{\frac{1}{s}} e^{\frac{r}{s}} \right\rfloor.$$

Then, for sufficiently large r ,

$$\frac{2 \tanh r}{1 - \tanh r} \tan(\alpha_i/2) \sin(\alpha_i) \leq 1 \quad \forall i \geq i_0(r).$$

By concavity of the log-function, we have $\log(1+x) \geq x \log 2 \geq \frac{x}{2}$ for $0 \leq x \leq 1$. Therefore

$$l(S(r)) \geq \sum_{i=i_0}^{\infty} \frac{\tanh r}{1 - \tanh r} \tan(\alpha_i/2) \sin(\alpha_i).$$

For sufficiently large r , the first factor is bounded from below by $\frac{e^{2r}}{4}$, while the second is bounded from below by α_i^2 . We thus get

$$l(S(r)) \geq \frac{e^{2r}}{4} \sum_{i=i_0}^{\infty} \alpha_i^2 = C_s^2 \frac{e^{2r}}{4} \sum_{i=i_0}^{\infty} \frac{1}{i^{2s}} \geq C_s^2 \frac{e^{2r}}{4} \int_{i_0}^{\infty} \frac{1}{x^{2s}} dx = C_s^2 \frac{e^{2r}}{4(2s-1)i_0^{2s-1}}.$$

Replacing our explicit value for i_0 gives

$$l(S(r)) \geq C e^{\frac{r}{s}}$$

for sufficiently large r and some constant C (again depending on s). Hence $\underline{\text{Ent}}K \geq \frac{1}{s}$.

Upper bound

For the upper bound in the statement, we apply our main theorem. For this, we have to find an upper bound on the Minkowski dimension of $\text{ex } K = E$.

Since the Minkowski dimension is invariant under bi-Lipschitz maps, we may replace distances on the unit circle by angular distances.

E has two accumulation points $\pm x_0$. For $\epsilon > 0$, let $N(\epsilon)$ be the number of ϵ -balls needed to cover E . We take one such ball around $\pm x_0$ and one further ball for each point in E not covered by these two balls.

The three points corresponding to the angle α_i are certainly in the ϵ -neighborhood of $\pm x_0$ provided

$$3 \sum_{j=i}^{\infty} \alpha_j \leq \epsilon.$$

Now we compute that

$$\sum_{j=i}^{\infty} \alpha_j = C_s \sum_{j=i}^{\infty} \frac{1}{j^s} \leq C_s \int_{i-1}^{\infty} \frac{1}{x^s} dx = \frac{C_s}{s-1} \frac{1}{(i-1)^{s-1}}.$$

It follows that all $i \geq i_0 := \left(\frac{3C_s}{s-1}\right)^{\frac{1}{s-1}} \epsilon^{\frac{1}{1-s}} + 1$ satisfy the inequality above and hence

$$N(\text{ex } K, \epsilon) \leq 6i_0 + 2 \leq C\epsilon^{-\frac{1}{s-1}}.$$

It follows that the upper Minkowski dimension is not larger than $\frac{1}{s-1}$. The upper bound of theorem 3.4 gives

$$\overline{\text{Ent}} K \leq \frac{2s-2}{3s-4}.$$

□

4. CENTRO-PROJECTIVE AND CENTRO-AFFINE AREAS

In this section, we will take a closer look at the centro-projective area which was introduced (in a non-intrinsic way) in definition 1.

4.1. Basic definitions and properties. Geometrically speaking, both centro-affine and centro-projective areas are Riemannian volumes of the boundary ∂K .

We first give intrinsic definitions of the centro-affine metric and area. Let K be a convex body with a distinguished interior point which we may suppose to be the origin o of V . The Minkowski functional of K is the unique positive function F that is homogeneous of degree one and whose level set at height 1 is the boundary ∂K . This function is convex and, according to Alexandroff's theorem, has almost everywhere a quadratic approximation.

Definition 4.1. Let v be a tangent vector to ∂K at a smooth point p . Then the *centro-affine semi-norm* of v is

$$\|v\|_a := \sqrt{\text{Hess}_p F(v, v)}.$$

The square of the centro-affine semi-norm is a quadratic function on the tangent, hence we may define as usual a volume form, say ω_a (which vanishes if $\|\cdot\|_a$ is not definite).

Definition 4.2. The *centro-affine area* of K is

$$\mathcal{A}_a(K) := \int_{\partial K} |\omega_a|.$$

It easily follows from the definitions that the centro-affine area is indeed an affine invariant of pointed convex bodies. Moreover, it is finite and vanishes on polytopes. The next proposition relates our definitions with the classical ones, its proof is a straightforward computation.

Proposition 4.3. *If the space is equipped with a Euclidean inner product, then the centro-affine area is given by*

$$\mathcal{A}_a(K) = \int_{\partial K} \frac{\sqrt{k}}{\langle n, p \rangle^{\frac{n-1}{2}}} dA,$$

where k is the Gaussian curvature of ∂K at p , n the unit vector normal to $T_p \partial K$ and dA the Euclidean area.

In order to introduce the centro-projective area, we will consider a compact convex subset of the (real) n -dimensional projective space. Here the word ‘‘convex’’ means that each intersection with a projective line is connected.

The definitions of the centro-projective semi-norm and area are merely the same as the centro-affine ones, but one has to replace the Minkowski functional by a projectively invariant function.

Definition 4.4. Let $K \subset \mathbb{P}^n$ be a convex body and $o \in \text{int } K$. The *projective gauge function* is

$$\begin{aligned} G_K : \mathbb{P}^n \setminus \{o\} &\rightarrow \mathbb{R} \cup \{\infty\}, \\ x &\mapsto 2[q_1, o, x, q_2] \end{aligned}$$

where q_1 and q_2 are the two intersections of ∂K with the line going through o and x .

Since the order of q_1 and q_2 is not fixed, this function is multi-valued (in fact 2-valued). Identifying $\mathbb{R} \cup \{\infty\}$ with \mathbb{P}^1 , this function is continuous.

If p belongs to the boundary of K , then the two values of $G_K(p)$ are different, one of them being 2, the other being ∞ . Hence there is some neighborhood U of p such that the restriction of G_K to U is

the union of two continuous (in fact smooth) functions G_K^+, G_K^- on U , where $G_K^+(p) = 2$ and $G_K^-(p) = \infty$.

Let v be a tangent vector to ∂K at a smooth point p . Since the restriction of G_K^+ to $\partial K \cap U$ is constant, the derivative of G_K^+ in the direction of v vanishes. Therefore, the Hessian of the restriction of G_K^+ to the tangent line is well-defined.

Definition 4.5. The *centro-projective semi-norm* of v is

$$\|v\|_p := \sqrt{\text{Hess}_p G_K^+(v, v)}.$$

Calling ω_p the induced volume form on ∂K , the *centro-projective area* of K is

$$\mathcal{A}_p(K) := \int_{\partial K} |\omega_p|.$$

As a consequence of the definition, one has

Proposition 4.6. *In a Euclidean space,*

$$\mathcal{A}_p(K) = \int_{\partial K} \frac{\sqrt{k}}{\langle n, p \rangle^{\frac{n-1}{2}}} \left(\frac{2a}{1+a} \right)^{\frac{n-1}{2}} dA.$$

In particular, the intrinsic definition of \mathcal{A}_p agrees with the definition given in the introduction.

Proof. An easy computation shows that

$$[q_1, o, x, q_2] = \frac{1 + a(q_2)}{F(x) + a(q_2)} F(x).$$

Then, if p is a smooth point of ∂K and $v \in T_p \partial K$,

$$\text{Hess}_p G_K(v, v) = \frac{2a(p)}{1 + a(p)} \text{Hess}_p F(v, v).$$

□

4.2. Properties of the centro-projective area. Both centro-affine and centro-projective areas vanish on polytopes, hence they are not continuous with respect to the Hausdorff topology on (pointed) bounded convex bodies. Nevertheless, the centro-affine area is upper-semicontinuous (see [36]). The same holds true for the centro-projective area as shown in the next theorem.

Theorem 4.7. *The centro-projective area is finite, invariant under projective transformations and upper-semicontinuous.*

Proof. From the above intrinsic definition, it follows that \mathcal{A}_p is invariant under projective transformations. Also, since the function a on the boundary is bounded and positive and since the centro-affine area is finite, it follows from proposition 4.6 that the centro-projective area is

also finite. It remains to show that it is upper-semicontinuous. Our proof is based on the fact that the centro-affine surface area \mathcal{A}_a is semicontinuous, see E. Lutwak [36].

Let K be a bounded convex body containing the origin in its interior and (K_i) a sequence of convex bodies with the same properties converging to K . Set

$$\tau(p) := \left(\frac{2a(p)}{1+a(p)} \right)^{\frac{n-1}{2}}, \quad p \in \partial K$$

which is a continuous function on ∂K .

For each i , if a_i is the function corresponding to K_i and p_i is the radial projection of p on ∂K_i , define $\tau_i \in C(\partial K)$ by

$$\tau_i(p) := \left(\frac{2a_i(p_i)}{1+a_i(p_i)} \right)^{\frac{n-1}{2}}.$$

Since $K_i \rightarrow K$, τ_i converges uniformly to τ . Therefore, for fixed $\epsilon > 0$ and all sufficiently large i ,

$$\|\tau_i - \tau\|_\infty < \epsilon$$

Take a triangulation of the sphere and let $\partial K = \cup_{j=1}^m \Delta_j$ (resp. $\partial K_i = \cup_{j=1}^m \Delta_{ij}$) be its radial projection.

Choosing this triangulation sufficiently thin, there exist $t_1, \dots, t_m \in \mathbb{R}_+$ with

$$|\tau(p) - t_j| < \epsilon$$

on Δ_j . By the triangle inequality, $|\tau_i(p) - t_j| < 2\epsilon$ on Δ_{ij} .

We define

$$\mathcal{A}_p(K_i, \Delta_{ij}) := \int_{\Delta_{ij}} \frac{\sqrt{k(x)}}{\langle n(x), x \rangle^{\frac{n-1}{2}}} \tau_i d\mathcal{H}^{n-1}(x).$$

Clearly, $\mathcal{A}_p(K_i) = \sum_{j=1}^m \mathcal{A}_p(K_i, \Delta_{ij})$. In a similar way, we define $\mathcal{A}_p(K, \Delta_j)$, $\mathcal{A}_a(K_i, \Delta_{ij})$ and $\mathcal{A}_a(K, \Delta_j)$.

Fix p_j in the interior of Δ_j and consider the convex hull $\widehat{\Delta}_i$ (resp. $\widehat{\Delta}_{ij}$) of Δ_j (resp. Δ_{ij}) and $-p_j$. The boundary of $\widehat{\Delta}_{ij}$ is a union of Δ_{ij} and line segments, hence $\mathcal{A}_a(K_i, \Delta_{ij}) = \mathcal{A}_a(\widehat{\Delta}_{ij})$. By the semicontinuity of \mathcal{A}_a , we obtain

$$\limsup_{i \rightarrow \infty} \mathcal{A}_a(K_i, \Delta_{ij}) = \limsup_{i \rightarrow \infty} \mathcal{A}_a(\widehat{\Delta}_{ij}) \leq \mathcal{A}_a(\widehat{\Delta}_j) = \mathcal{A}_a(K, \Delta_j).$$

It follows that

$$\begin{aligned} \limsup_{i \rightarrow \infty} \mathcal{A}_p(K_i) &= \limsup_{i \rightarrow \infty} \sum_{j=1}^m \mathcal{A}_p(K_i, \Delta_{ij}) \\ &\leq \limsup_{i \rightarrow \infty} \sum_{j=1}^m \mathcal{A}_a(K_i, \Delta_{ij})(t_j + 2\epsilon) \\ &\leq \sum_{j=1}^m \mathcal{A}_a(K, \Delta_j)(t_j + 2\epsilon) \end{aligned}$$

On the other hand,

$$\mathcal{A}_p(K) = \sum_{j=1}^m \mathcal{A}_p(K, \Delta_j) \geq \sum_{j=1}^m \mathcal{A}_a(K, \Delta_j)(t_j - \epsilon)$$

from which we deduce that

$$\limsup_{i \rightarrow \infty} \mathcal{A}_p(K_i) \leq \mathcal{A}_p(K) + 3\epsilon \mathcal{A}_a(K).$$

□

The centro-affine surface area has the following important properties:

- (1) \mathcal{A}_a is a valuation on the space of compact convex subsets of V containing o in the interior. This means that whenever $K, L, K \cup L$ are such bodies, then

$$\mathcal{A}_a(K \cup L) = \mathcal{A}_a(K) + \mathcal{A}_a(L) - \mathcal{A}_a(K \cap L).$$
- (2) \mathcal{A}_a is upper semi-continuous with respect to the Hausdorff topology.
- (3) \mathcal{A}_a is invariant under $GL(V)$.

A recent theorem by M. Ludwig & M. Reitzner [35] states that the vector space of functionals with these three properties is generated by the constant valuation and \mathcal{A}_a . The centro-projective surface area satisfies the last two conditions, but is not a valuation.

REFERENCES

- [1] A. D. Alexandroff. Almost everywhere existence of the second differential of a convex function and some properties of convex surfaces connected with it. *Leningrad State Univ. Annals [Uchenye Zapiski] Math. Ser.*, 6:3–35, 1939.
- [2] J.C. Álvarez Paiva and E. Fernandes. Crofton formulas in projective Finsler spaces. *Electron. Res. Announc. Am. Math. Soc.*, 4(13):91–100, 1998.
- [3] J.C. Álvarez Paiva and A.C. Thompson. Volumes on normed and Finsler spaces. Bao, David (ed.) et al., *A sampler of Riemann-Finsler geometry*. Cambridge: Cambridge University Press. Mathematical Sciences Research Institute Publications 50, 1-48, 2004.
- [4] V. Bangert. Convex hypersurfaces with bounded first mean curvature measure. *Calc. Var. Partial Differ. Equ.*, 8(3):259–278, 1999.

- [5] A. Barvinok. *A course in convexity*. Graduate Studies in Mathematics. 54. Providence, RI: American Mathematical Society (AMS), 2002.
- [6] Y. Benoist. A survey on divisible convex sets. Written for the Morningside center conference in Beijing 2006.
- [7] Y. Benoist. Convexes divisibles. I. In *Algebraic groups and arithmetic*, pages 339–374. Tata Inst. Fund. Res., Mumbai, 2004.
- [8] Y. Benoist. Convexes hyperboliques et quasiisométries. (Hyperbolic convexes and quasiisometries.). *Geom. Dedicata*, 122:109–134, 2006.
- [9] A. Bernig. Hilbert geometry of polytopes. arXiv:0812.1080.
- [10] W. Blaschke. *Kreis und Kugel. 2. durchgesehene und verbesserte Auflage*. Berlin: Walter de Gruyter & Co VIII, 167 S., 27 Fig., 1956.
- [11] A. A. Borisenko and E. A. Olin. Asymptotic Properties of Hilbert Geometry. arXiv:0711.0446.
- [12] M. Bridson and A. Haefliger. *Metric spaces of non-positive curvature*. Grundlehren der Mathematischen Wissenschaften. 319. Berlin: Springer. xxi, 643 p., 1999.
- [13] B. Colbois and C. Vernicos. Les géométries de Hilbert sont à géométrie locale bornée. (Hilbert geometries have bounded local geometry.). *Ann. Inst. Fourier*, 57(4):1359–1375, 2007.
- [14] B. Colbois, C. Vernicos, and P. Verovic. Hilbert geometry for convex polygonal domains. arXiv:0804.1620.
- [15] B. Colbois and P. Verovic. Hilbert geometry for strictly convex domains. *Geom. Dedicata*, 105:29–42, 2004.
- [16] P. de la Harpe. On Hilbert’s metric for simplices. Niblo, Graham A. (ed.) et al., Geometric group theory. Volume 1. Proceedings of the symposium held at the Sussex University, Brighton (UK), July 14-19, 1991. Cambridge: Cambridge University Press. Lond. Math. Soc. Lect. Note Ser. 181, 97-119 (1993), 1993.
- [17] L. C. Evans and R. F. Gariepy. *Measure theory and fine properties of functions*. Studies in Advanced Mathematics. Boca Raton: CRC Press. viii, 268 p. , 1992.
- [18] T. Foertsch and A. Karlsson. Hilbert metrics and Minkowski norms. *J. Geom.*, 83(1-2):22–31, 2005.
- [19] S. Gallot, D. Hulin, and J. Lafontaine. *Riemannian geometry. 3rd ed.* Universitext. Berlin: Springer. xv, 322 p. , 2004.
- [20] D. Hilbert. Ueber die gerade Linie als kürzeste Verbindung zweier Punkte. *Math. Ann.*, XLVI:91–96, 1895.
- [21] D. Hilbert. *Foundations of geometry*. Teubner-Archiv zur Mathematik. Stuttgart: B. G. Teubner. xxiii, 408 S., 1999.
- [22] L. Hörmander. *Notions of convexity. Reprint of the 1994 edition*. Modern Birkhäuser Classics. Basel: Birkhäuser. viii, 414 p., 2007.
- [23] D. Hug. Absolute continuity for curvature measures of convex sets. I. *Math. Nachr.*, 195:139–158, 1998.
- [24] D. Hug. Absolute continuity for curvature measures of convex sets. II. *Math. Z.*, 232(3):437–485, 1999.
- [25] D. Hug. *Measures, curvatures and currents in convex geometry*. Habilitationsschrift, Universität Freiburg, 1999.
- [26] A. Karlsson and G.A. Noskov. The Hilbert metric and Gromov hyperbolicity. *Enseign. Math.*, 48(1-2):73–98, 2002.
- [27] David C. Kay. The ptolemaic inequality in Hilbert geometries. *Pacific J. Math.*, 21:293–301, 1967.
- [28] P. Kelly and L. Paige. Symmetric perpendicularity in Hilbert geometries. *Pacific J. Math.*, 2:319–322, 1952.

- [29] P. Kelly and E. Straus. Curvature in Hilbert geometries. *Pacific J. Math.*, 8:119–125, 1958.
- [30] P. Kelly and E. Straus. Curvature in Hilbert geometries. II. *Pacific J. Math.*, 25:549–552, 1968.
- [31] A. Koldobsky. *Fourier analysis in convex geometry*. Mathematical Surveys and Monographs 116. Providence, RI: American Mathematical Society (AMS). vi, 170 p., 2005.
- [32] D. Laugwitz. *Differentialgeometrie in Vektorräumen unter besonderer Berücksichtigung der unendlichdimensionalen Räume*. Braunschweig: Friedr. Vieweg und Sohn; Berlin: VEB Deutscher Verlag der Wissenschaften. VI, 89 p., 1965.
- [33] K. Leichtweiß. Convexity and differential geometry. Gruber, P. M. (ed.) et al., Handbook of convex geometry. Volume B. Amsterdam: North-Holland. 1045-1080 (1993), 1993.
- [34] K. Leichtweiß. *Affine geometry of convex bodies*. Heidelberg: Johann Ambrosius Barth. x, 310 p., 1998.
- [35] M. Ludwig and M. Reitzner. A classification of $SL(n)$ invariant valuations. To appear in *Ann. Math.*
- [36] E. Lutwak. The Brunn-Minkowski-Firey theory. II: Affine and geominimal surface areas. *Adv. Math.*, 118(2):244–294, 1996.
- [37] P. Mattila. *Geometry of sets and measures in Euclidean spaces. Fractals and rectifiability. 1st paperback ed.* Cambridge Studies in Advanced Mathematics. 44. Cambridge: Cambridge University Press. xii, 343 p., 1999.
- [38] Y. Nasu. On Hilbert geometry. *Math. J. Okayama Univ.*, 10:101–112, 1961.
- [39] C. Schütt and E. Werner. The convex floating body. *Math. Scand.*, 66(2):275–290, 1990.
- [40] Z. Shen. *Lectures on Finsler geometry*. Singapore: World Scientific. xiv, 307 p., 2001.
- [41] E. Socié-Méthou. Behaviour of distance functions in Hilbert-Finsler geometry. *Differ. Geom. Appl.*, 20(1):1–10, 2004.
- [42] C. Vernicos. Entropie volumique des géométries de Hilbert. to appear in *Actes de Séminaire de Théorie Spectrale et Géométrie. Année 2007–2008*.
- [43] C. Vernicos. Lipschitz characterisation of polytopal Hilbert geometries. arXiv:0812.1032.
- [44] C. Vernicos. Introduction aux géométries de Hilbert. In *Actes de Séminaire de Théorie Spectrale et Géométrie. Vol. 23. Année 2004–2005*, volume 23 of *Sémin. Théor. Spectr. Géom.*, pages 145–168. Univ. Grenoble I, Saint, 2005.

DÉPARTEMENT DE MATHÉMATIQUES, CHEMIN DU MUSÉE 23, 1700 FRIBOURG, SWITZERLAND

E-mail address: gautier.berck@unifr.ch

E-mail address: andreas.bernig@unifr.ch

DEPARTMENT OF MATHEMATICS, LOGIC HOUSE, SOUTH CAMPUS, NUI MAYNOOTH, CO. KILDARE, IRELAND

E-mail address: Constantin.Vernicos@maths.nuim.ie