



**HAL**  
open science

# Universal Superfluid Transition and Transport Properties of Two-Dimensional Dirty Bosons

Giuseppe Carleo, Guilhem Boéris, Markus Holzmann, Laurent Sanchez-Palencia

► **To cite this version:**

Giuseppe Carleo, Guilhem Boéris, Markus Holzmann, Laurent Sanchez-Palencia. Universal Superfluid Transition and Transport Properties of Two-Dimensional Dirty Bosons. 2013. hal-00822056v1

**HAL Id: hal-00822056**

**<https://hal.science/hal-00822056v1>**

Preprint submitted on 13 May 2013 (v1), last revised 28 Aug 2013 (v2)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Universal Superfluid Transition and Transport Properties of Two-Dimensional Dirty Bosons

Giuseppe Carleo,<sup>1</sup> Guilhem Bo ris,<sup>1</sup> Markus Holzmann,<sup>2,3</sup> and Laurent Sanchez-Palencia<sup>1</sup>

<sup>1</sup>*Laboratoire Charles Fabry, Institut d'Optique, CNRS, Univ Paris Sud 11,  
2 avenue Augustin Fresnel, F-91127 Palaiseau cedex, France*

<sup>2</sup>*LPTMC, UMR 7600 of CNRS, Universit  Pierre et Marie Curie, 75005 Paris, France*

<sup>3</sup>*Universit  Grenoble 1/CNRS, LPMMC, UMR 5493, B.P. 166, 38042 Grenoble, France*

We study the phase diagram of two-dimensional, interacting bosons in the presence of a correlated disorder in continuous space, using large-scale finite temperature quantum Monte Carlo simulations. We show that the superfluid transition is strongly protected against disorder. It remains of the Berezinskii-Kosterlitz-Thouless type up to disorder strengths comparable to the chemical potential. Moreover, we study the transport properties in the strong disorder regime where a zero-temperature Bose-glass phase is expected. We show that the conductance exhibits a thermally activated behavior and strictly vanishes only at zero temperature. Our results do not show any evidence of a finite-temperature localization transition, and point towards the existence of Bose bad-metal phase as a precursor of the Bose-glass phase.

PACS numbers: 05.30.Jp, 02.70.Ss, 67.25.dj, 72.15.Rn

*Introduction* – Transport properties in quantum materials are governed by a complex interplay of disorder and interactions. While disorder tends to localize particles, interactions may strongly alter the over-idealized single-particle picture by either reinforcing or suppressing localization. Dramatic effects are expected in two dimensions (2D), such as metal-insulator transitions [1–3], suppression of superfluidity [4–8], and presumably high- $T_c$  superconductivity [9]. Possible phase transitions are however particularly elusive owing to absence of true long-range order even in extended phases [10, 11], and many questions remain open. For instance, resistance measurements in Si-MOFSETs suggest a metal-insulator transition [1], which may be attributed to quantum localization or classical percolation [2], but experiments on GaAs heterostructures point towards a crossover behavior [3]. Studies of the superfluid Berezinskii-Kosterlitz-Thouless (BKT) transition [12, 13] have also been reported for  $^4\text{He}$  films adsorbed on porous media [4, 5]. While the BKT transition is unaffected in the weak disorder limit [14], the question of its relevance for strong disorder is left open owing to the difficulty to identify a universal jump of the superfluid density [6–8]. Moreover, a question that is attracting much debate is whether many-body localization effects [15–19] can drive a finite-temperature metal-insulator transition in two dimensions.

Ultracold quantum gases in controlled disorder offer a unique tool to address these questions in a unified way [20]. In clean or disordered quasi-2D geometries, direct consequences of vortex pairing [21], superfluidity [22], quasi-long-range phase coherence [23, 24], and resistance measurements [25] have been reported. On the theoretical side, most knowledge rely on lattice models with uncorrelated disorder [26–30]. Conversely, little is known about the novel models of disorder introduced by ultracold atoms, which are continuous, correlated, and sustain classical percolation thresholds [31–35].

In this Letter, we report on an *ab-initio* path-integral quantum Monte Carlo study of the phase diagram of interacting

2D bosons in such a continuous model of disorder. We show that, although the density profile exhibits large spatial modulations, the superfluid BKT transition is strongly protected against disorder. This holds up to disorder strengths comparable to the chemical potential, where the zero-temperature Bose-glass transition is expected. The critical properties of the *dirty* superfluid transition can be understood in terms of a universal description including a simple renormalization of the critical parameters. In particular, we find that the superfluid transition occurs while the fluid percolates with a density significantly above the critical density of the clean system, which allows us to rule out the classical percolation scenario. Moreover, we study the conductance by means of a novel quantum Monte Carlo estimator. Deep in the strong disorder regime, we find direct evidence of an insulating behavior, characterized by a conductance that decreases with the temperature. Our data is consistent with a thermally-activated behavior of the Arrhenius type, indicating that the conductance vanishes only at zero temperature.

*System and methods* – We consider a two-dimensional quantum fluid of interacting bosons at temperature  $T$  characterized by short-range repulsive interactions of effective coupling parameter  $\tilde{g} = mg/\hbar^2$ , where  $m$  is the particle mass, and subjected to a disordered potential  $V(\mathbf{r})$ . The latter is a correlated, isotropic, and continuous speckle disorder, as realized by laser light diffusion through a ground-glass plate [23–25, 31]. Its average value coincides with its standard deviation  $V_r$ , and its two-point correlation function is a Gaussian of r.m.s. radius  $\sigma_r$ . In the following, we use the chemical potential  $\mu$  and the disorder correlation radius  $\sigma_r$  as units of energy and length, and the parameters  $\tilde{g} = 0.1$  and  $\sigma_r = \sqrt{5}\xi_0$ , where  $\xi_0 = \hbar/\sqrt{2m\mu}$  is the healing length in the absence of disorder. They are typical values in ultracold-atom experiments [23, 24]. For instance, for  $^{87}\text{Rb}$  atoms and the chemical potential  $\mu/k_B \simeq 56\text{ nK}$ , it corresponds to  $\sigma_r = 0.5\ \mu\text{m}$ .

Our study is based on a fully *ab-initio* path-integral quantum Monte Carlo (QMC) approach, which allows for unbiased calculations of equilibrium, finite-temperature properties of

interacting bosons [36] in terms of a discretized path-integral representation using  $M$  time slices. We use the worm algorithm [37] to sample the grand-canonical partition function at inverse temperature  $\beta = 1/k_B T$ . Whereas disorder is taken into account at the level of the primitive-approximation, the two-body interaction is treated in the pair-product approximation [36, 38]. The two-particle propagator is derived by solving the s-wave scattering problem at high temperatures,  $\varepsilon^{-1} = M k_B T$ , which yields

$$\langle \mathbf{r} | e^{-\varepsilon H_2} | \mathbf{r}' \rangle \simeq \frac{m}{4\pi\hbar^2\varepsilon} e^{-m|\mathbf{r}-\mathbf{r}'|^2/4\hbar^2\varepsilon} + \frac{\tilde{g}}{8\pi} \int_0^\infty dk k e^{-\varepsilon \frac{\hbar^2 k^2}{m}} [J_0(kr)Y_0(kr') + J_0(kr')Y_0(kr)], \quad (1)$$

where  $H_2$  is the Hamiltonian of the relative motion of two particles,  $\mathbf{r}$  and  $\mathbf{r}'$  are the relative coordinates at different times, whereas  $J_0$  and  $Y_0$  are Bessel functions of the first and second kind, respectively. Here, we report results from simulations of up to  $N = 10^5$  particles in a box of linear extension  $L$  with periodic boundary conditions, averaged over  $\approx 40$  disorder realizations.

*Superfluid transition* – We first study the superfluid to normal fluid transition in the presence of disorder. To elucidate the critical properties of this transition, we calculate the static superfluid susceptibility,  $\chi_s = \frac{1}{L^2} \int d\mathbf{r} d\mathbf{r}' \langle \Psi^\dagger(\mathbf{r}) \Psi(\mathbf{r}') \rangle$ , and the field correlation function,  $g_1(r) = \langle \Psi^\dagger(\mathbf{r}) \Psi(0) \rangle$ , where  $\Psi$  is the field operator, and  $\langle \dots \rangle$  denotes thermal and disorder average. In the superfluid phase of the clean system the decay of field correlations is algebraic,  $g_1(r) \sim r^{-\eta}$ , and the superfluid susceptibility diverges as  $\chi_s \sim L^{2-\eta}$ . The exponent  $\eta$  is directly related to the superfluid density  $n_s$  as  $\eta = \frac{mk_B T}{2\pi\hbar^2 n_s}$ . In the normal phase, field correlations are exponentially suppressed,  $g_1(r) \sim e^{-r/\gamma}$ , and the susceptibility remains finite in the thermodynamic limit.

Analysis of our QMC data allows us to study both qualitatively and quantitatively the superfluid to normal transition in the presence of disorder (Fig. 1). The inset of Fig. 1 shows the superfluid susceptibility,  $\chi_s$ , as a function of the system size for various temperatures at intermediate disorder amplitude,  $V_R/\mu = 0.3$ . For high temperatures (curves in the light-orange zone),  $\chi_s$  converges to a finite value, characteristic for the normal fluid phase. Above the critical temperature (curves in the blue zone),  $\chi_s$  shows an algebraic divergence. At the critical point, our QMC results are compatible with the scaling  $\chi_s \sim L^{7/4}$ , i.e.  $\eta \simeq 1/4$ , as expected for the BKT transition in the clean system [12, 13]. It is confirmed by the behavior of the rescaled superfluid susceptibility  $\chi_s/L^{7/4}$ , plotted in the main panel of Fig. 1 as a function of temperature for various values of  $L$ , where all curves cross at a single point. Further, our data for the superfluid density calculated from the winding number estimator [36] (not shown) is consistent with the universal jump at the transition temperature in the thermodynamic limit, expected for a BKT transition. We conclude that the superfluid transition of dirty bosons remains in the universality class of the clean BKT transition. To some extent,




Figure 1: Scaled superfluid susceptibility against temperature for system sizes ranging from  $L_R = L/\sigma_R = 5 - 40$  at disorder strength  $V_R/\mu = 0.3$  and interaction strength  $\tilde{g} = 0.1$ . Darker curves mark increasingly large size. In the inset, the susceptibility is shown as a function of  $L_R$  (temperature increases from top to bottom). The divergence predicted by the BKT transition for clean systems ( $\eta = 1/4$ ) is indicated by the dashed line.

this result is expected for weak disorder. According to Harris argument [14], local fluctuations of the disorder potential are smoothed out at the scale of the diverging correlation length at the Kosterlitz-Thouless transition, thus introducing only a renormalization of the effective parameters. Our calculations show that it holds also for strong disorder (up to  $V_R \simeq \mu$ ).

The intersection point of the rescaled superfluid susceptibility (curves as in Fig. 1) precisely determines the critical temperature at fixed disorder amplitude. In Fig. 2, we show the resulting critical temperatures versus disorder strength and draw the superfluid to normal fluid phase diagram of dirty 2D bosons. Within numerical accuracy, the critical line is described by  $T_c(V_R) \simeq T_c^0(\mu - V_R)$ , where  $T_c^0 \simeq \frac{\pi\mu\hbar^2}{m\tilde{g}k_B} / \log(13.2/\tilde{g})$  is the critical temperature for the disorder-free system [39]. The shift of the critical temperature can be understood in terms of the leading order renormalization of the chemical potential,  $\mu_c(V_R) - V_R = \mu_c(0)$ . Similar to  $T_c$ , the critical density also decreases approximately linear from the disorder-free value  $n_c\sigma_R^2 \simeq 21.1$  to  $n_c\sigma_R^2 \approx 9$  for  $V_R \simeq \mu$ . For lower densities, the system remains normal for any of our temperatures,  $k_B T \gtrsim \mu$ , consistent with a Bose-glass phase at zero temperature, as indicated in Fig. 2.

In our analysis we find that the BKT scaling regime is attained for *increasingly large* system sizes, upon increasing the disorder strength. For the lowest reported transition temperature, we find that systems of linear size  $L/\sigma_R \sim 40 - 80$  are needed to attain the scaling regime. This should be contrasted with the typically (much) smaller sizes of the inner critical regions probed in experiments [6–8, 23, 24] and in some numerical simulations [40].

Let us compare our results to the possibility of a percolation driven transition. In the superfluid phase, the Bose gas den-




Figure 2: Phase diagram of two-dimensional interacting bosons at fixed chemical potential and interaction strength  $\tilde{g} = 0.1$ . The dark dashed line is the critical temperature of the clean system with a renormalized chemical potential; the lighter dashed line is the prediction of a combination of LDA and percolation theory (see text). For strong disorder, the normal system goes to the Bose-glass phase in the zero- $T$  limit.

sity for each realization of the disorder may be approximated using the local density approximation (LDA),  $n_{\text{LDA}}(\mathbf{r}, V_R) = n_0[\mu - V(\mathbf{r})]$ , where  $n_0(\mu)$  is the density of the clean system at chemical potential  $\mu$ . The quantitative agreement between this approximation and our QMC results is reasonably good in the superfluid phase. We consider percolating clusters in which the local density stays everywhere higher than the critical density of the clean system, i.e.  $n_{\text{LDA}}(\mathbf{r}, V_R) \gtrsim \frac{mk_B T}{2\pi\hbar^2} \log(380/\tilde{g})$  [39]. The occurrence of such percolating superfluid clusters yields the orange line reported in Fig. 2, which shows that the percolation scenario strongly overestimates the critical disorder strength. Superfluidity disappears while there still exist large percolating clusters above the critical density of clean system. Notice that we have disregarded the weak penetration of the density into the disorder potential barriers due to the finite healing length in the superfluid phase [41]. However, this effect does not alter the above conclusions since it would further raise the percolation line in Fig. 2.

*Transport Properties* – We now address the strongly disordered regime, where superfluid coherence is lost and disorder substantially affects mass transport. A long-debated issue is to understand whether interacting systems preserve Anderson-like localization properties, as found in the single particle case [15–19]. At zero temperature, the conditions for a localized (Bose glass) phase to exist have been established in seminal works [42, 43]. At finite temperature, however, the existence of a perfectly insulating state, characterized by an exactly vanishing conductance ( $G_{\text{DC}} = 0$ ) is intrinsically more complex and theoretically plausible scenarios have been put forward only recently. The *many-body localization* transition scenario [15, 19] implies the existence of a critical tem-

perature,  $T_{\text{loc}}$ , separating the perfect insulator behavior at low energies where  $G_{\text{DC}}(T < T_{\text{loc}}) = 0$ , from a delocalized, diffusive phase with  $G_{\text{DC}}(T > T_{\text{loc}}) \neq 0$ . This transition, however, has not been directly observed in experiments and remains at the center of intense theoretical activity, for instance in out-of-equilibrium phenomena [44–46].

In order to identify possible signatures of many-body localization at finite temperature, we have computed transport properties in the strongly disordered region. Such transport properties are traditionally addressed in disordered materials subjected to an external electric bias. They were recently shown to be also accessible in a new generation of ultracold-atoms experiments [25].

To obtain the zero-frequency (DC) conductance within QMC, we compute the imaginary-time correlations of the density current operator along a given direction  $\alpha$ ,  $\Lambda_\alpha(\tau, q) = \langle J_\alpha(\tau, q) J_\alpha(0, -q) \rangle$ . It allows us to reconstruct the associated longitudinal conductance  $G_{\text{reg}}(\omega)$  [57] from numerical analytical continuation [47–49] of the Laplace transform [50]

$$\Lambda_\alpha(\tau, 0) = 2\hbar \int_{-\infty}^{+\infty} d\omega \frac{\exp(-\omega\tau) \omega}{1 - \exp(-\beta\hbar\omega)} G_{\text{reg}}(\omega). \quad (2)$$

Imaginary-time correlations of high statistical quality are essential for a reliable reconstruction of the dynamical response  $G_{\text{reg}}(\omega)$ . Direct estimators of current correlations successfully used for lattice simulations [51, 52] present a diverging statistical error in the small imaginary-time step limit in continuum systems, preventing its use for numerical analytical continuation for our purposes. To overcome this serious problem, we introduce a novel estimator that significantly reduces




Figure 3: Finite-temperature, zero-frequency longitudinal conductance for an intensity of the disorder  $V_R = 2.4 \mu$ . Error bars refer to averages over disorder and the dashed line is a fit to a simple exponential Arrhenius law  $G_{\text{DC}}(T) \propto e^{-\frac{AE}{k_B T}}$ . In the upper inset, the complete frequency dependence of the conductance is exemplified for  $k_B T = 2\mu$ . In the lower inset, the conductance at constant temperature ( $k_B T = \mu$ ) and varying disorder strength is also shown. In all cases darker curves mark increasingly large sizes, in the range  $L/\sigma_R = 20 - 40$ .

the statistical error making possible a reliable determination of the conductance. The key observation is that the imaginary-time derivatives of the correlations are well-behaved in the small time-step limit  $\varepsilon \rightarrow 0$ , whereas the imaginary-time integral of the correlation function is itself well-behaved and directly related to the superfluid density  $n_s$  [53]. We can therefore express the current-current correlations by means of the integral expression

$$\Lambda_\alpha(\tau, 0) = \frac{1}{\beta m} (n - n_s) + \int_0^\beta d\tau_1 \Lambda'_\alpha(\tau_1, 0) + \frac{1}{\beta} \int_0^\beta d\tau_1 \int_0^{\tau_1} d\tau_2 \Lambda'_\alpha(\tau_2, 0), \quad (3)$$

where the imaginary-time derivative of the correlation function reads

$$\varepsilon \Lambda'_\alpha(\tau, 0) = \frac{1}{mL^2} \sum_i \langle [\nabla_i^\alpha V(\tau)] \times D^\alpha(0) \rangle + \frac{1}{2mL^2} \sum_{i,j,\beta} \langle D_j^\beta(\tau) \times [\nabla_i^\alpha \nabla_j^\beta V(\tau)] \times D^\alpha(0) \rangle + \mathcal{O}(\varepsilon^{\frac{3}{2}}). \quad (4)$$

In the latter expression  $\nabla_i^\alpha$  is the derivative with respect to the position of the  $i$ -th particle,  $D^\alpha(\tau) = \sum_j^N D_j^\alpha(\tau) = \sum_j^N [R_j^\alpha(\tau) - R_j^\alpha(\tau - \varepsilon)]$  is proportional to the imaginary-time displacement of the center of mass and  $V(\tau)$  is a generic one-body potential which in our specific case is the disordered speckle potential.

We first study the behavior of the DC longitudinal conductance  $G_{\text{DC}} = G_{\text{reg}}(\omega = 0)$  versus the temperature deep in the strong disorder regime. The outcome is shown in the main panel of Fig. 3 for a fixed disorder strength,  $V_R = 2.4\mu$ . The conductance monotonously decreases with temperature, a behavior which is typically observed in related experiments with disordered materials. Extrapolating to zero temperature, the longitudinal conductance vanishes, as expected for the Bose-glass phase [43]. The overall behavior of  $G_{\text{DC}}$ , however, points towards a thermally activated transport (see fit in Fig. 3), at variance with a perfectly insulating, many-body localized phase. The energy broadening of the reconstructed dynamical conductances (see upper inset of Fig. 3) remains finite and does not show any indication for a finite-temperature mobility edge.

In the lower inset of Fig. 3 we further show the behavior of the longitudinal conductance at fixed temperature versus the disorder strength  $V_R$ , ranging from the edge of the superfluid transition  $V_R \sim \mu$  to the strong disorder regime, with  $V_R \sim 4\mu$ . Compatibly with the thermally activated scenario, we find that the transport is exponentially suppressed with increasing disorder, but never vanishes, in agreement with experimental results [25]. Size effects, though present, do not indicate any phase transition, at least for mesoscopic samples. Therefore, our *ab-initio* analysis hints to a delocalized *bad Bose-metal* phase at finite-temperature.

*Conclusions* – We have studied two-dimensional interacting bosons in the presence of a correlated disordered potential using *ab-initio* quantum Monte Carlo calculations. We have found that the disorder renormalizes the low-energy Hamiltonian in the neighborhood of the superfluid transition, but the critical line remains in the universality class of the Berezinskii-Kosterlitz-Thouless transition. It holds for arbitrary strong disorder up the zero-temperature Bose-glass transition. Moreover, we have developed a new estimator for the conductance, which does not suffer from diverging statistical errors. Deep in the strong disorder regime, the mass transport exhibits a thermally activated behavior and it is strictly suppressed only at zero temperature. It points towards the existence of a *Bose bad-metal* phase, as the finite-temperature precursor of the Bose-glass insulator. We have found no indication of a finite-temperature many-body localization transition. Our results provide new theoretical insights both in the general understanding of disordered interacting systems and in the interpretation of experiments with both ultracold atoms and disordered materials.

*Acknowledgments* – We acknowledge discussions with B. Altshuler, D. Basko, T. Bourdel, T. Giamarchi, T. Roscilde, S. Moroni, and T. Ziman. This research was supported by the European Research Council (FP7/2007-2013 Grant Agreement No. 256294), Ministère de l'Enseignement Supérieur et de la Recherche, and Agence Nationale de la Recherche (“DisorderTransitions”). It was performed using HPC resources from GENCI-CCRT/CINES (Grants t2012056853 and t2013056853). Use of the computing facility cluster GM-PCS of the LUMAT federation (FR LUMAT 2764) is also acknowledged. Our simulations make use of the ALPS [58] scheduler library and statistical analysis tools [54–56].

- 
- [1] E. Abrahams, S. V. Kravchenko, and M. P. Sarachik, *Rev. Mod. Phys.* **73**, 251 (2001).
  - [2] L. A. Tracy, E. H. Hwang, K. Eng, G. A. Ten Eyck, E. P. Nordberg, K. Childs, M. S. Carroll, M. P. Lilly, and S. Das Sarma, *Phys. Rev. B* **79**, 235307 (2009).
  - [3] G. Allison, E. A. Galaktionov, A. K. Savchenko, S. S. Safonov, M. M. Fogler, M. Y. Simmons, and D. A. Ritchie, *Phys. Rev. Lett.* **96**, 216407 (2006).
  - [4] P. A. Crowell, F. W. Van Keuls, and J. D. Reppy, *Phys. Rev. Lett.* **75**, 1106 (1995).
  - [5] P. A. Crowell, F. W. Van Keuls, and J. D. Reppy, *Phys. Rev. B* **55**, 12620 (1997).
  - [6] D. R. Luhman and R. B. Hallock, *Phys. Rev. Lett.* **93**, 086106 (2004).
  - [7] D. R. Luhman and R. B. Hallock, *Phys. Rev. B* **74**, 014510 (2006).
  - [8] R. J. Lazarowich and P. Taborek, *Phys. Rev. B* **74**, 024512 (2006).
  - [9] S. H. Pan, J. P. O’Neal, R. L. Badzey, C. Chamon, H. Ding, J. R. Engelbrecht, Z. Wang, H. Eisaki, S. Uchida, A. K. Gupta, et al., *Nature* **413**, 282 (2001).
  - [10] N. D. Mermin and H. Wagner, *Phys. Rev. Lett.* **17**, 1133 (1966).
  - [11] P. C. Hohenberg, *Phys. Rev.* **158**, 383 (1967).

- [12] V. L. Berezinskii, *Sov. Phys. JETP* **34**, 610 (1971).
- [13] J. M. Kosterlitz and D. J. Thouless, *J. Phys. C* **6**, 1181 (1973).
- [14] A. B. Harris, *J. Phys. C* **7**, 1671 (1974).
- [15] D. Basko, I. Aleiner, and B. Altshuler, *Ann. Phys.* **321**, 1126 (2006).
- [16] V. Oganesyan and D. A. Huse, *Phys. Rev. B* **75**, 155111 (2007).
- [17] P. Lugan, D. Clément, P. Bouyer, A. Aspect, and L. Sanchez-Palencia, *Phys. Rev. Lett.* **99**, 180402 (2007).
- [18] P. Lugan and L. Sanchez-Palencia, *Phys. Rev. A* **84**, 013612 (2011).
- [19] I. L. Aleiner, B. L. Altshuler, and G. V. Shlyapnikov, *Nat. Phys.* **6**, 900 (2010).
- [20] L. Sanchez-Palencia and M. Lewenstein, *Nat. Phys.* **6**, 87 (2010).
- [21] Z. Hadzibabic, P. Kruger, M. Cheneau, B. Battelier, and J. Dalibard, *Nature* **441**, 1118 (2006).
- [22] R. Desbuquois, L. Chomaz, T. Yefsah, J. Léonard, J. Beugnon, C. Weitenberg, and J. Dalibard, *Nat. Phys.* **8**, 645 (2012).
- [23] M. C. Beeler, M. E. W. Reed, T. Hong, and S. L. Rolston, *New J. Phys.* **14**, 073024 (2012).
- [24] B. Allard, T. Plisson, M. Holzmann, G. Salomon, A. Aspect, P. Bouyer, and T. Bourdel, *Phys. Rev. A* **85**, 033602 (2012).
- [25] S. Krinner, D. Stadler, J. Meineke, J.-P. Brantut, and T. Esslinger, *Phys. Rev. Lett.* **110**, 100601 (2013).
- [26] F. Alet and E. S. Sørensen, *Phys. Rev. E* **67**, 015701 (2003).
- [27] N. Prokof'ev and B. Svistunov, *Phys. Rev. Lett.* **92**, 015703 (2004).
- [28] G. M. Wysin, A. R. Pereira, I. A. Marques, S. A. Leonel, and P. Z. Coura, *Phys. Rev. B* **72**, 094418 (2005).
- [29] S. G. Söyler, M. Kiselev, N. V. Prokof'ev, and B. V. Svistunov, *Phys. Rev. Lett.* **107**, 185301 (2011).
- [30] F. Lin, E. S. Sørensen, and D. M. Ceperley, *Phys. Rev. B* **84**, 094507 (2011).
- [31] D. Clément, A. F. Varón, J. A. Retter, L. Sanchez-Palencia, A. Aspect, and P. Bouyer, *New J. Phys.* **8**, 165 (2006).
- [32] L. Pezzé, M. Robert-de-Saint-Vincent, T. Bourdel, J.-P. Brantut, B. Allard, T. Plisson, A. Aspect, P. Bouyer, and L. Sanchez-Palencia, *New J. Phys.* **13**, 095015 (2011).
- [33] S. Pilati, S. Giorgini, and N. Prokof'ev, *Phys. Rev. Lett.* **102**, 150402 (2009).
- [34] S. Pilati, S. Giorgini, M. Modugno, and N. Prokof'ev, *New J. Phys.* **12**, 073003 (2010).
- [35] T. Bourdel, *Phys. Rev. A* **86**, 063626 (2012).
- [36] D. M. Ceperley, *Rev. Mod. Phys.* **67**, 279 (1995).
- [37] M. Boninsegni, N. Prokof'ev, and B. Svistunov, *Phys. Rev. Lett.* **96**, 070601 (2006).
- [38] W. Krauth, *Phys. Rev. Lett.* **77**, 3695 (1996).
- [39] N. Prokof'ev, O. Ruebenacker, and B. Svistunov, *Phys. Rev. Lett.* **87**, 270402 (2001).
- [40] G. Astrakharchik, K. Krutitsky, and P. Navez, arXiv:1205.2967 (2012).
- [41] L. Sanchez-Palencia, *Phys. Rev. A* **74**, 053625 (2006).
- [42] T. Giamarchi and H. J. Schulz, *Phys. Rev. B* **37**, 325 (1988).
- [43] M. P. A. Fisher, P. B. Weichman, G. Grinstein, and D. S. Fisher, *Phys. Rev. B* **40**, 546 (1989).
- [44] E. Canovi, D. Rossini, R. Fazio, G. E. Santoro, and A. Silva, *Phys. Rev. B* **83**, 094431 (2011).
- [45] G. Carleo, F. Becca, M. Schiro, and M. Fabrizio, *Sci. Rep.* **2**, 243 (2012).
- [46] E. Canovi, D. Rossini, R. Fazio, G. E. Santoro, and A. Silva, *New J. Phys.* **14**, 095020 (2012).
- [47] A. W. Sandvik, *Phys. Rev. B* **57**, 10287 (1998).
- [48] J. E. Gubernatis, M. Jarrell, R. N. Silver, and D. S. Sivia, *Phys. Rev. B* **44**, 6011 (1991).
- [49] G. Carleo, S. Moroni, and S. Baroni, *Phys. Rev. B* **80**, 094301 (2009).
- [50] R. Kubo, *Rep. Prog. Phys.* **29**, 255 (1966).
- [51] D. J. Scalapino, S. R. White, and S. Zhang, *Phys. Rev. B* **47**, 7995 (1993).
- [52] N. Trivedi, R. T. Scalettar, and M. Randeria, *Phys. Rev. B* **54**, R3756 (1996).
- [53] E. L. Pollock and D. M. Ceperley, *Phys. Rev. B* **36**, 8343 (1987).
- [54] M. Troyer, B. Ammon, and E. Heeb, *Lect. Notes Comput. Sci.* **1505**, 191 (1998).
- [55] A. Albuquerque, F. Alet, P. Corboz, P. Dayal, A. Feiguin, S. Fuchs, L. Gamper, E. Gull, S. Guertler, A. Honecker, et al., *J. Magn. Magn. Mater.* **310**, 1187 (2007).
- [56] B. Bauer, L. D. Carr, H. G. Evertz, A. Feiguin, J. Freire, S. Fuchs, L. Gamper, J. Gukelberger, E. Gull, S. Guertler, et al., *J. Stat. Mech.* **05**, P05001 (2011).
- [57] Notice that we refer here to mass transport. The conventional electric conductance is  $\tilde{G}_{\text{reg}}(\omega) = q^2 G_{\text{reg}}(\omega)$  for particles of charge  $q$ .
- [58] <http://alps.comp-phys.org/>