



HAL
open science

The complexity of deciding whether a graph admits an orientation with fixed weak diameter

Julien Bensmail, Romaric Duvignau, Sergey Kirgizov

► **To cite this version:**

Julien Bensmail, Romaric Duvignau, Sergey Kirgizov. The complexity of deciding whether a graph admits an orientation with fixed weak diameter. 2014. hal-00824250v5

HAL Id: hal-00824250

<https://hal.science/hal-00824250v5>

Preprint submitted on 18 Sep 2014 (v5), last revised 16 Aug 2016 (v6)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The complexity of deciding whether a graph admits an orientation with fixed weak diameter

Julien Bensmail^{a,b,*}, Romaric Duvignau^{a,b}, Sergey Kirgizov^{c,d}

^aUniv. Bordeaux, LaBRI, UMR 5800, 33400 Talence, France

^bCNRS, UMR 5800, 33400 Talence, France

^cUniv. Pierre et Marie Curie, LIP6, UMR 7606, 75252 Paris, France

^dCNRS, UMR 7606, 75252 Paris, France

Abstract

An oriented graph \vec{G} is said weak (resp. strong) if, for every pair $\{u, v\}$ of vertices of \vec{G} , there are directed paths joining u and v in either direction (resp. both directions). In case, for every pair, some of these directed paths have length at most k , we call \vec{G} k -weak (resp. k -strong). We consider several problems asking whether an undirected graph G admits orientations satisfying some connectivity and distance properties. As a main result, we show that deciding whether G admits a k -weak orientation is NP-complete for every $k \geq 2$. This notably implies the NP-completeness of several problems asking whether G is an extremal graph for some vertex-colouring problems (in terms of needed colours).

Keywords: oriented graph, weak diameter, strong diameter, complexity

1. Introduction

Let G be a simple undirected graph with vertex set $V(G)$ and edge set $E(G)$. By *orienting* every edge uv of G , either from u to v or from v to u , one obtains an *orientation* \vec{G} of G . This oriented graph \vec{G} has the same vertex set as G , i.e. $V(\vec{G}) = V(G)$, and, for every edge $uv \in E(G)$, we have either $\vec{uv} \in E(\vec{G})$ or $\vec{vu} \in E(\vec{G})$ depending on the orientation assigned to uv .

The *distance* $\text{dist}(G, u, v)$ from u to v in G is the minimal length of a path joining u and v . We refer to the maximum distance between two vertices of G as its *diameter*, and denote it $\text{diam}(G)$. These definitions can be naturally adapted to the context of oriented graphs. A *dipath* of \vec{G} is a sequence (v_1, v_2, \dots, v_k)

*Corresponding author. Mail: julien.bensmail@labri.fr. Phone: +33-(0)5-40-00-35-47. Fax: +33-(0)5-40-00-66-69

Email addresses: julien.bensmail@labri.fr (Julien Bensmail), romaric.duvignau@labri.fr (Romaric Duvignau), sergey.kirgizov@lip6.fr (Sergey Kirgizov)

of distinct vertices such that $\overrightarrow{v_i v_{i+1}} \in E(\vec{G})$ for every $i \in \{1, 2, \dots, k-1\}$. Such a dipath has length $k-1$ and is written $\overrightarrow{v_1 v_2 \dots v_k}$. The *directed distance* $\text{dist}(\vec{G}, u, v)$ from u to v in \vec{G} is the minimal length of a dipath starting from u and ending at v . Note that, contrary to the undirected case, we may have $\text{dist}(\vec{G}, u, v) \neq \text{dist}(\vec{G}, v, u)$. Therefore, two definitions of the oriented diameter can be adopted. Let

$$\text{dist}_w(\vec{G}, u, v) = \min\{\text{dist}(\vec{G}, u, v), \text{dist}(\vec{G}, v, u)\}$$

and

$$\text{dist}_s(\vec{G}, u, v) = \max\{\text{dist}(\vec{G}, u, v), \text{dist}(\vec{G}, v, u)\}.$$

These two measures are called the *weak distance* and *strong distance*, respectively, from u to v in \vec{G} . The *weak diameter* of \vec{G} , denoted $\text{diam}_w(\vec{G})$, is the maximum weak distance from a vertex to another one. The *strong diameter* of \vec{G} , denoted $\text{diam}_s(\vec{G})$, is the maximum strong distance from a vertex to another one. The weak diameter can intuitively be seen as an optimistic measure of the directed distances in an oriented graph (basically two vertices u and v are considered close when, say, u can reach v with few moves, and this no matter how many moves needs v to reach u (if possible)). We call \vec{G} *k-weak* (resp. *k-strong*) if it has weak (resp. strong) diameter at most k . More generally, we say that \vec{G} is *weak* (resp. *strong*) if it is *k-weak* (resp. *k-strong*) for some value of k .

Several appealing and attractive problems in graph theory are about deducing graph orientations with particular properties. Such problems find natural applications in real-world problems (e.g. traffic problems). In this paper, we mainly focus on the existence of (either weak or strong) orientations of some undirected graph G in which the diameter is preserved, i.e. as close to $\text{diam}(G)$ as possible. Though the question of deciding whether G admits a weak or strong orientation can be answered easily by using several classic results of graph theory (see Sections 2.1 and 3), the hardness of deciding the same when a (weak or strong) diameter restriction is required was mostly unknown. Our main contribution is an indication of the complexity of answering this question. In particular, we show that deciding whether G admits a *k-weak* orientation is NP-complete for every $k \geq 2$, and suggest that the same should be true for *k-strong* orientations, completing a result of Chvátal and Thomassen.

This paper is organized as follows. In Section 2, we first consider the questions above for the weak notions of distance, orientation, and diameter. Then, we consider, in Section 3, the same questions but for the strong notions of distance, orientation, and diameter. Consequences of our results are then discussed in Section 4. In particular, as side results we get that deciding whether an undirected graph is extremal for some vertex-colouring problems (in terms of needed colours) is NP-complete.

2. Weak orientations

This section is devoted to the following related two decision problems.

WEAK ORIENTATION

Instance: A graph G .

Question: Does G admit a weak orientation?

k -WEAK ORIENTATION

Instance: A graph G .

Question: Does G admit a k -weak orientation?

Using two classic tools of graph theory, we prove, in Theorem 3 below, that WEAK ORIENTATION can be answered in linear time. Then, we prove that k -WEAK ORIENTATION is in P for $k = 1$, and NP-complete otherwise, i.e. whenever $k \geq 2$ (see Theorem 5). These two results confirm that imposing a (even constant) weak diameter condition is a strong restriction which makes the problem more difficult.

2.1. Complexity of WEAK ORIENTATION

We here show that WEAK ORIENTATION can be solved in linear time, and is hence in P. For this purpose, we need to recall the following two classic results. Recall that a *bridge* of a graph is an edge whose removal disconnects the graph.

Theorem 1 (Tarjan [10]). The bridges of a graph can be found in linear time.

Theorem 2 (Robbins [7]). A strong orientation of a bridgeless undirected graph can be computed in linear time.

We also need the notion of *B-contraction*. Given an undirected graph G , its B -contraction is the graph obtained as follows. Let first e_1, e_2, \dots, e_x denote the bridges of G , and B_1, B_2, \dots, B_y denote the (bridgeless) components of $G - \{e_1, e_2, \dots, e_x\}$. Then the B -contraction of G is obtained by associating a vertex v_{B_i} with each component B_i , and in which two vertices v_{B_i} and v_{B_j} are joined by an edge if and only if there is a bridge joining B_i and B_j in G . Clearly the B -contraction of any graph is a tree.

We are now ready to introduce the result of this section.

Theorem 3. An undirected graph admits a weak orientation if and only if its B -contraction is a path.

Proof. We start by proving the sufficiency. Assume G is a connected undirected graph whose B -contraction is a path with successive edges e_1, e_2, \dots, e_x , and denote B_1, B_2, \dots, B_y the components of $G - \{e_1, e_2, \dots, e_x\}$. We obtain a weak orientation \vec{G} of G as follows. First orient the edges e_1, e_2, \dots, e_x towards the same direction, i.e. following the natural first-last ordering of the B -contraction. Then orient the edges of every B_i to form a strong component. Such an orientation exists according to Theorem 2 since each B_i has no bridge. Clearly \vec{G}

is weak since every two vertices within a same B_i can reach each other and the orientation of the edges e_1, e_2, \dots, e_x form a dipath in the B -contraction.

We now prove the necessity by contradiction. Assume the B -contraction of G is not a path, but G admits a weak orientation \vec{G} . Clearly the orientation of \vec{G} , restricted to the B -contraction, should be weak. But this is impossible as the B -contraction of G has a node with degree at least 3, and every B -contraction is a tree. A contradiction. \square

Since the B -contraction of any graph can be computed in linear time (due notably to Theorem 1), we can answer in linear time to every instance of WEAK ORIENTATION. Actually, since the bridges of a graph and a strong orientation of every bridgeless undirected graph can be found in linear time (recall Theorems 1 and 2), the algorithm described in the sufficiency part of the proof of Theorem 3 can even be implemented to efficiently construct, i.e. in linear time, a weak orientation (if any) of every undirected graph.

Corollary 4. WEAK ORIENTATION is in P.

2.2. Complexity of k -WEAK ORIENTATION

Clearly the answer to every instance of 1-WEAK ORIENTATION is *yes* if and only if G is complete. So 1-WEAK ORIENTATION is in P. The complexity of every remaining problem k -WEAK ORIENTATION (i.e. with $k \geq 2$) was mentioned and asked in several references of the literature (notably in [5, 8, 9]) due to its relationship with other problems of graph theory (see concluding Section 4). We herein settle the complexity of these problems by showing them to be NP-complete in general.

Theorem 5. k -WEAK ORIENTATION is NP-complete for every $k \geq 2$.

Proof. The problem k -WEAK ORIENTATION is in NP for every fixed k since one can, given an orientation \vec{G} of G , check whether $\text{diam}_w(\vec{G}) \leq k$. For this purpose, one just has to check, for every possible pair $\{u, v\}$ of distinct vertices of G , whether either u and v are adjacent, or there exist $k' \leq k - 1$ vertices $w_1, w_2, \dots, w_{k'}$ such that $\overrightarrow{uw_1w_2\dots w_{k'}v}$ or $\overrightarrow{vw_1w_2\dots w_{k'}u}$ is a dipath of \vec{G} . This naive witness algorithm runs in polynomial time compared to the order of G .

Let $k \geq 2$ be fixed. We show that k -WEAK ORIENTATION is NP-hard by reduction from the following problem, which is shown to be NP-complete in [4].

2-VERTEX-COLOURING OF 3-UNIFORM HYPERGRAPHS

Instance: A 3-uniform hypergraph H .

Question: Is H 2-colourable, i.e. can we colour each vertex of H either blue or red so that each edge of H has at least one blue vertex and one red vertex?

Throughout this proof, we denote the vertices and edges of any hypergraph H with order n and size m by x_1, x_2, \dots, x_n and E_1, E_2, \dots, E_m , respectively. For every $i \in \{1, 2, \dots, n\}$, we further denote by $n_i \geq 1$ the number of distinct edges of H which contain the vertex x_i . From a 3-uniform hypergraph H , we produce




Figure 1: The core subgraph G_H^c of G_H obtained assuming H has two edges $E_1 = \{x_1, x_2, x_3\}$ and $E_2 = \{x_3, x_4, x_5\}$.

a graph G_H such that H is 2-colourable if and only if G_H admits a k -weak orientation \vec{G}_H . This reduction is achieved in polynomial time compared to the size of H .

We first describe the *core* G_H^c of G_H , i.e. the subgraph of G_H from which the equivalence with H follows. The subgraph G_H^c does not have diameter k , but G_H will be augmented later so that it has diameter k , and this without altering the equivalence. The core G_H^c has the following vertices (see Figure 1). With each vertex x_i of H , we associate $n_i + 2$ vertices u_i , u'_i , and $v_{i,j_1}, v_{i,j_2}, \dots, v_{i,j_{n_i}}$ in G_H^c , where j_1, j_2, \dots, j_{n_i} are the distinct indices of the edges of H which contain x_i . We now associate additional vertices in G_H^c with each edge E_j of H , where $j \in \{1, 2, \dots, m\}$. This association depends on the parity of k .

- If k is even, then add two vertices a_j and a'_j to G_H^c .
- Otherwise, if k is odd, then add two cycles $a_j b_j c_j a_j$ and $a'_j b'_j c'_j a'_j$ with length 3 to G_H^c .

We now link the vertices of G_H^c by means of several vertex-disjoint paths. By “joining a pair of vertices $\{u, v\}$ by a path”, we mean that we identify the endvertices of a new path with u and v , respectively. Since this operation is used at most once for joining any pair $\{u, v\}$ of G_H^c , we use the notation uPv to denote the resulting path (if any). First, join every pair $\{u_i, u'_i\}$ by a path with length $\lfloor \frac{k}{2} \rfloor$. Then also join every pair $\{u'_i, v_{i,j}\}$ by a path with length $\lceil \frac{k}{2} \rceil$. Now consider each edge $E_j = \{x_{i_1}, x_{i_2}, x_{i_3}\}$ of H , and add the following paths to G_H^c .

- If k is even, join every pair of $\{v_{i_1,j}, v_{i_2,j}, v_{i_3,j}\} \times \{a_j, a'_j\}$ by means of a path with length $\frac{k}{2}$.
- Otherwise, if k is odd, then join every pair of $\{v_{i_1,j}\} \times \{a_j, a'_j\}$, $\{v_{i_2,j}\} \times \{b_j, b'_j\}$, and $\{v_{i_3,j}\} \times \{c_j, c'_j\}$ by a path with length $\lfloor \frac{k}{2} \rfloor$.

Note that, by construction, exactly one pair $\{v_{i,j}, s(v_{i,j})\}$ (resp. $\{v_{i,j}, s'(v_{i,j})\}$) was joined by a path with length $\lfloor \frac{k}{2} \rfloor$, where $s(v_{i,j})$ (resp. $s'(v_{i,j})$) is a vertex of the form a_j, b_j or c_j (resp. a'_j, b'_j or c'_j). The notation $s(v_{i,j})$ and $s'(v_{i,j})$ are used throughout this section. In particular, observe that if k is even, then we have $s(v_{i_1,j}) = s(v_{i_2,j}) = s(v_{i_3,j}) = a_j$ for every edge $E_j = \{x_{i_1}, x_{i_2}, x_{i_3}\}$ of H . We analogously have $s'(v_{i_1,j}) = s'(v_{i_2,j}) = s'(v_{i_3,j}) = a'_j$.

A pair $\{u, v\}$ of distinct vertices of G_H^c is said *representative* whenever it matches one of the following forms.

1. $\{u_i, v_{i,j}\}$ where $i \in \{1, 2, \dots, n\}$, $j \in \{1, 2, \dots, m\}$, and $x_i \in E_j$.
2. $\{u'_i, s(v_{i,j})\}$ where $i \in \{1, 2, \dots, n\}$, $j \in \{1, 2, \dots, m\}$, and $x_i \in E_j$.
3. $\{v_{i_1,j}, v_{i_2,j}\}$ where $i_1, i_2 \in \{1, 2, \dots, n\}$, $j \in \{1, 2, \dots, m\}$, and $x_{i_1}, x_{i_2} \in E_j$.

An orientation of G_H^c is *good* if two vertices forming a representative pair are linked by a k -dipath in either direction. Note that, in this definition, there is no requirement on the oriented distance between two vertices which are at distance at least $k + 1$. A representative pair is a pair of vertices which are not adjacent in G_H , and for which there are at most two paths with length at most k joining it. All of these paths belong to G_H^c so that the existence of a k -weak orientation of G_H depends on the existence of a good orientation of G_H^c .

We prove below that we have an equivalence between finding a proper 2-vertex-colouring of H and a good orientation of G_H^c . The proof relies on the following claims.

Claim 1. Suppose the vertex x_i belongs to the edges $E_{j_1}, E_{j_2}, \dots, E_{j_{n_i}}$ of H . Then, in any good orientation $\overrightarrow{G_H^c}$ of G_H^c , $\overrightarrow{u_i P u'_i P v_{i,j} P s(v_{i,j})}$ is a dipath for every $j \in \{j_1, j_2, \dots, j_{n_i}\}$, or $\overrightarrow{s(v_{i,j}) P v_{i,j} P u'_i P u_i}$ is a dipath for every $j \in \{j_1, j_2, \dots, j_{n_i}\}$.

Proof. Note that because $u_i P u'_i P v_{i,j_1}$ is the only path with length at most k joining u_i and v_{i,j_1} in G_H^c , either $\overrightarrow{u_i P u'_i P v_{i,j_1}}$ or $\overrightarrow{v_{i,j_1} P u'_i P u_i}$ must be a dipath

of $\overrightarrow{G_H^c}$. Assume $\overrightarrow{u_i P u'_i P v_{i,j_1}}$ is a dipath of $\overrightarrow{G_H^c}$. Since $\overrightarrow{u_i P u'_i}$ is now a dipath of $\overrightarrow{G_H^c}$, $\overrightarrow{u'_i P v_{i,j}}$ must also be a dipath for every $j \in \{j_1, j_2, \dots, j_{n_i}\}$ since $\overrightarrow{u_i P u'_i P v_{i,j}}$ is the only path with length at most k joining u_i and $v_{i,j}$ in G_H^c .

Similarly, since, for every $j \in \{j_1, j_2, \dots, j_{n_i}\}$, the only path with length at most k joining u'_i and $s(v_{i,j})$ in G_H^c is $\overrightarrow{u'_i P v_{i,j} P s(v_{i,j})}$, and $\overrightarrow{u'_i P v_{i,j}}$ is a dipath of $\overrightarrow{G_H^c}$, then $\overrightarrow{v_{i,j} P s(v_{i,j})}$ has to be a dipath of $\overrightarrow{G_H^c}$. Thus, $\overrightarrow{u_i P u'_i P v_{i,j} P s(v_{i,j})}$ belongs to $\overrightarrow{G_H^c}$ for every $j \in \{j_1, j_2, \dots, j_{n_i}\}$ assuming that $\overrightarrow{u_i P u'_i P v_{i,j_1}}$ belongs to the orientation. The claim follows analogously from the assumption that $\overrightarrow{v_{i,j_1} P u'_i P u_i}$ is a dipath of $\overrightarrow{G_H^c}$. \square

Claim 2. Suppose k is even, and $E_j = \{x_{i_1}, x_{i_2}, x_{i_3}\}$ is an edge of H . Then, in any good orientation $\overrightarrow{G_H^c}$ of G_H^c , either $\overrightarrow{v_{i,j} P s(v_{i,j})}$ or $\overrightarrow{s(v_{i,j}) P v_{i,j}}$ is a dipath for every $i \in \{i_1, i_2, i_3\}$. Furthermore, these three dipaths cannot be all directed from or towards the $s(v_{i,j})$'s.

Proof. Recall that $s(v_{i_1,j}) = s(v_{i_2,j}) = s(v_{i_3,j}) = a_j$ and $s'(v_{i_1,j}) = s'(v_{i_2,j}) = s'(v_{i_3,j}) = a'_j$ when k is even. Note further that there are only two paths with length at most k joining any two of $v_{i_1,j}$, $v_{i_2,j}$, and $v_{i_3,j}$. These include a_j and a'_j , respectively. If the statement of the claim is not fulfilled, then there is no k -dipath of $\overrightarrow{G_H^c}$ joining any two of $v_{i_1,j}$, $v_{i_2,j}$, and $v_{i_3,j}$ including a_j . So there must be three k -dipaths joining these vertices including a'_j , but this is impossible. \square

Claim 3. Suppose k is odd, and $E_j = \{x_{i_1}, x_{i_2}, x_{i_3}\}$ is an edge of H . Then, in any good orientation $\overrightarrow{G_H^c}$ of G_H^c , either $\overrightarrow{v_{i,j} P s(v_{i,j})}$ or $\overrightarrow{s(v_{i,j}) P v_{i,j}}$ is a dipath for every $i \in \{i_1, i_2, i_3\}$. Besides these three dipaths cannot be all directed from or towards the $s(v_{i,j})$'s.

Proof. Similarly as for previous Claim 2, if the statement of the claim is not fulfilled by $\overrightarrow{G_H^c}$, then there is no dipath with length at most k joining any two of $v_{i_1,j}$, $v_{i_2,j}$, and $v_{i_3,j}$ including the $s(v_{i,j})$'s. Then there cannot be three k -dipaths, including the $s'(v_{i,j})$'s, joining every pair of these vertices, and this no matter how the paths $\overrightarrow{v_{i_1,j} s'(v_{i_1,j})}$, $\overrightarrow{v_{i_2,j} s'(v_{i_2,j})}$ and $\overrightarrow{v_{i_3,j} s'(v_{i_3,j})}$ are oriented, and how the edges of the cycles $a_j b_j c_j a_j$ and $a'_j b'_j c'_j a'_j$ are oriented. \square

Regarding previous Claims 2 and 3, remark that if two of the dipaths obtained by orienting the paths $\overrightarrow{v_{i_1,j} P s(v_{i_1,j})}$, $\overrightarrow{v_{i_2,j} P s(v_{i_2,j})}$ and $\overrightarrow{v_{i_3,j} P s(v_{i_3,j})}$ have the same direction, i.e. from or towards the $s(v_{i,j})$'s while the third one is oriented in the opposite direction, then we can obtain three k -dipaths joining any two of $v_{i_1,j}$, $v_{i_2,j}$, and $v_{i_3,j}$. Suppose e.g. that $\overrightarrow{v_{i_1,j} P s(v_{i_1,j})}$, $\overrightarrow{v_{i_2,j} P s(v_{i_2,j})}$ and $\overrightarrow{s(v_{i_3,j}) P v_{i_3,j}}$ are dipaths of $\overrightarrow{G_H^c}$. So far, note that there are two k -dipaths starting from $v_{i_1,j}$ and $v_{i_2,j}$, respectively, and ending at $v_{i_3,j}$ (when k is odd, these are obtained by adding $\overrightarrow{s(v_{i_1,j}) s(v_{i_3,j})}$ and $\overrightarrow{s(v_{i_2,j}) s(v_{i_3,j})}$ to $E(\overrightarrow{G_H^c})$). The last

k -dipath starting from $v_{i_1,j}$ and ending at $v_{i_2,j}$ can be obtained e.g. by orienting the edges of G_H^c in such a way that $\overrightarrow{v_{i_1,j}Ps'(v_{i_1,j})}$ and $\overrightarrow{s'(v_{i_2,j})Pv_{i_2,j}}$ are dipaths, and $\overrightarrow{s'(v_{i_1,j})s'(v_{i_2,j})}$ is an arc when k is odd.

According to Claims 1, 2 and 3, we have an equivalence between finding a proper 2-vertex-colouring of H and a good orientation of G_H^c . Indeed, assume that having the dipath $\overrightarrow{u_iPu'_i}$ (resp. $\overrightarrow{u'_iPu_i}$) in an orientation of G_H^c simulates that the vertex x_i of H is coloured blue (resp. red), and that having the dipath $\overrightarrow{v_{i,j}Ps(v_{i,j})}$ (resp. $\overrightarrow{s(v_{i,j})Pv_{i,j}}$) simulates the fact that the vertex x_i is counted as a blue (resp. red) vertex in E_j . Claim 1 reflects the fact that if x_i is coloured, say, blue by a proper 2-vertex-colouring of H , then x_i counts as a blue vertex in every edge which contains it. Claims 2 and 3 depict the fact that all vertices from a single edge of H cannot have the same colour. Thus, from a proper 2-vertex-colouring of H we can deduce a good orientation of G_H^c , and vice-versa.

We now augment G_H with additional vertices so that there is a path with length at most k joining every two non-adjacent vertices of G_H^c that do not form a representative pair. This is done in such a way that there is an orientation of the edges of $E(G_H) - E(G_H^c)$ so that every two vertices of G_H that do not form a representative pair are joined by a dipath with length at most k . In this way, the existence of a k -weak orientation of G_H only relies on the existence of a good orientation of G_H^c .

The augmentation consists in associating a gadget G_v with each vertex v of G_H^c , and then connecting all the resulting gadgets in such a way there is a path with length at most k between any two vertices from different gadgets G_u and G_v . In the case where $\{u, v\}$ is not a representative pair, we add a *shortcut* between G_u and G_v , i.e. an alternative shorter path for joining two vertices of G_u and G_v . This is done in such a way that every vertex u' of G_u is at distance at most k from any vertex v' of G_v , unless $u' = u$, $v' = v$ and $\{u, v\}$ is a representative pair. However, in the situation where $\{u, v\}$ is not representative, there is a path with length k joining u and v that uses the shortcut between G_u and G_v .

Set $x = \lfloor \frac{k}{2} \rfloor$. For every $i \in \{1, 2, \dots, x\}$, add two new vertices s_v^i and p_v^i to G_v . These two vertices form the i^{th} level of G_v , and are said to be i -vertices. Next, for every $i \in \{1, 2, \dots, x-1\}$, add all possible edges between the i - and $(i+1)$ -vertices of G_v so that two consecutive levels of G_v form a clique on 4 vertices. Finally, add an edge between v and every 1-vertex of G_v .

We finish the construction of G_H by adding some connection between the gadgets. We distinguish two cases depending on the parity of k .

- If k is even, then we turn the subgraph induced by all x -vertices of G_H into a clique. Next, for every pair $\{u, v\}$ of vertices of G_H^c which is not representative, add a shortcut vertex $e_{u,v}$ to the clique constructed just before. Finally, add every edge between $e_{u,v}$ and the vertices from the $(x-1)^{\text{th}}$ levels of G_u and G_v if $k \geq 4$, or the edges $ue_{u,v}$ and $e_{u,v}v$ when $k = 2$.




Figure 2: The gadgets G_u and G_v obtained for a pair $\{u, v\}$ which is not representative.

- Otherwise, if k is odd, then add a new vertex z to G_H , and add all possible edges between z and x -vertices. For every pair $\{u, v\}$ of G_H^c that is not representative, also add the shortcut edges $s_u^x p_v^x$ and $p_u^x s_v^x$ to G_H .

This construction is illustrated in Figure 2 for $k = 6$ and $k = 7$. Note that no new path with length at most k between two vertices composing a representative pair of G_H arose from the modifications. Therefore, the equivalence between finding a proper 2-vertex-colouring of H and a good orientation of G_H^c is preserved. We finally show that there is an orientation of the edges we just added so that every pair of vertices of G_H which is not representative is joined by a k -dipath in either direction.

Define an arbitrary ordering $\sigma = (v_1, v_2, \dots, v_{|V(G_H^c)|})$ over all vertices of G_H^c , and consider the following partial orientation. First, for every vertex v of G_H^c , let $\overrightarrow{vs_v^1}$ and $\overrightarrow{p_v^1 v}$ be arcs. Then, for every level $i \in \{1, 2, \dots, x\}$ of G_v , let $\overrightarrow{p_v^i s_v^{i+1}}$ be an arc. Next, for every $i \in \{1, 2, \dots, x-1\}$, add the arcs $\overrightarrow{s_v^i s_v^{i+1}}$, $\overrightarrow{p_v^{i+1} p_v^i}$, $\overrightarrow{p_v^i s_v^{i+1}}$ and $\overrightarrow{p_v^{i+1} s_v^i}$ to the partial orientation. The partial orientation is completed depending on the parity of k .

- If k is even, then, for every shortcut vertex e of G_H , add the arcs $\overrightarrow{s_v^x e}$ and $\overrightarrow{ep_v^x}$. Next, for every $i < j$ consider $\overrightarrow{s_{v_i}^x s_{v_j}^x}$, $\overrightarrow{s_{v_i}^x p_{v_j}^x}$, $\overrightarrow{p_{v_i}^x s_{v_j}^x}$ and $\overrightarrow{p_{v_i}^x p_{v_j}^x}$ as arcs of the partial orientation. Additionally, if $\{v_i, v_j\}$ is not a representative pair, then let $\overrightarrow{s_{v_j}^{x-1} e_{v_i, v_j}}$, $\overrightarrow{e_{v_i, v_j} p_{v_i}^{x-1}}$, $\overrightarrow{s_{v_i}^{x-1} e_{v_i, v_j}}$, and $\overrightarrow{e_{v_i, v_j} p_{v_j}^{x-1}}$ be arcs if $k \geq 4$, or $\overrightarrow{v_i e_{v_i, v_j}}$ and $\overrightarrow{e_{v_i, v_j} v_j}$ be arcs when $k = 2$.
- If k is odd, then let $\overrightarrow{s_{v_i}^x z}$ and $\overrightarrow{z p_{v_i}^x}$ be arcs. Finally, if $\{v_i, v_j\}$ is not representative, then let $\overrightarrow{s_{v_i}^x p_{v_j}^x}$ and $\overrightarrow{s_{v_j}^x p_{v_i}^x}$ be arcs.

Note that, under the partial orientation given above, any vertex u' from a gadget G_u can directly “access” the upper or lower level of G_u . Besides, there is a dipath with length at most k joining u' and any vertex v' from another gadget

G_v , unless $u' = u$, $v' = v$, and $\{u, v\}$ is a representative pair. Such a path typically goes up across G_u , then exits G_u to enter G_v (either directly from the x^{th} levels or via z), and finally goes down across G_v . Because the gadgets have $x = \lfloor \frac{k}{2} \rfloor$ levels, the length of such a path does not exceed k . Finally observe that if $\{u, v\}$ is representative, then there is no path with length at most k joining u and v going across the gadgets. On the contrary, if $\{u, v\}$ is not a representative pair, then there is a path with length exactly k joining u and v . This path necessarily includes the shortcut between G_u and G_v , i.e. the vertex $e_{u,v}$ if k is even or an edge linking the x^{th} levels of G_u and G_v otherwise.

Hence, G_H admits a k -weak orientation if and only if G_H^c admits a good orientation. Besides, G_H^c admits a good orientation if and only if H is 2-colourable. By transitivity, we get that G_H admits a k -weak orientation if and only if H is 2-colourable, and thus that k -WEAK ORIENTATION is NP-complete. \square

3. Strong orientations

We herein consider the following strong analogues of WEAK ORIENTATION and k -WEAK ORIENTATION.

STRONG ORIENTATION

Instance: A graph G .

Question: Does G admit a strong orientation?

k -STRONG ORIENTATION

Instance: A graph G .

Question: Does G admit a k -strong orientation?

Using Theorems 1 and 2, we can answer to every instance G of STRONG ORIENTATION in linear time via the following algorithm. Clearly, if G has bridges, then it admits no strong orientation. Otherwise, a strong orientation can be obtained according to Theorem 2. All these steps can be achieved in linear time, so the whole algorithm indeed runs in linear time.

Theorem 6. STRONG ORIENTATION is in P.

Clearly, no instance of 1-STRONG ORIENTATION is positive since we consider oriented graphs only (no symmetric arc is allowed). So 1-STRONG ORIENTATION is trivially in P. Besides, it was proved by Chvátal and Thomassen that 2-STRONG ORIENTATION is NP-complete in general (see [1]). For the other values of $k \geq 3$, we strongly believe that the NP-completeness of the remaining problems could be proved by slightly modifying the reduction scheme given in the proof of Theorem 5. Namely, consider e.g. the following modifications. First, the core graph G_H^c would be obtained in the same way. Then, when constructing the big clique subgraph (which would be intended to have the same purpose, namely to have a lot of pairs of vertices being joined by a lot of paths with length at most k), one would have to make sure that the following additional paths exist:

- at least two new paths with length k joining every non-representative pair;
- one new path with length k joining every representative pair.

Then note that if these modifications are performed, then, in order to get a strong orientation of G_H , for every representative pair $\{u, v\}$, we would need to have the path of length k joining u and v in G_H^c being oriented, say, from u to v , and the additional path (of the clique subgraph) oriented from v to u . The hard part would be to make sure that the clique subgraph can be always oriented correctly (as in the second part of the proof of Theorem 5), but this should not be a problem due to its large number of paths with length at most k (basically the clique subgraph could be less dense in the original construction, but its large size facilitates the proof process).

4. Discussion

In this paper, we have considered the complexity of orienting an undirected graph in such a way that the distances between its vertices are preserved. As a main result, we have proved the same result as Chvátal and Thomassen for weak orientations, hence proving that the weak and strong versions of all these problems are theoretically as hard as each other in essence.

It is worth mentioning that Theorem 5 has consequences on some special vertex-colouring problems. These consequences are related to the following context. Usually, a *proper vertex-colouring* of an undirected graph G is an assignment of colours to its vertices such that no two adjacent vertices receive the same colour. It is well-known that extremal graphs for the notion of proper vertex-colouring (i.e. the graphs which need the most colours compared to their order) are complete graphs. But for augmented kinds of graphs and vertex-colourings, the notion of extremal graph is not as obvious. Actually, the NP-completeness of every problem k -WEAK ORIENTATION (in particular for $k = 2$) implies that, in some contexts, an easy characterization of these extremal graphs in terms of underlying undirected graph cannot exist (unless $P=NP$).

4.1. Oriented vertex-colouring of oriented graphs

Let \vec{G} be an oriented graph. An *oriented vertex-colouring* of \vec{G} is a vertex-colouring c satisfying the following two properties:

- for every two adjacent vertices u and v , we have $c(u) \neq c(v)$,
- for every two arcs \vec{xy} and \overleftarrow{uv} , if $c(x) = c(v)$ then $c(y) \neq c(u)$.

We call \vec{G} an *oriented clique* if it does not admit an oriented vertex-colouring using strictly less than $|V(\vec{G})|$ colours. It is known that \vec{G} is an oriented clique if and only if every two vertices of \vec{G} are at weak distance at most 2, i.e. \vec{G} is 2-weak, see e.g. [2, 3]. From the undirected graphs point of view, the case $k = 2$ of Theorem 5 hence implies the following.

Corollary 7. It is NP-complete to decide whether an undirected graph G is the underlying graph of an oriented clique.

4.2. Proper vertex-colouring of signified graphs

A *signified graph* $G = (V, E^-, E^+)$ is basically an undirected graph whose each edge is either *negative* (i.e. in E^-) or *positive* (i.e. in E^+), refer e.g. to [6, 8] for more details. A *proper vertex-colouring* of G is then a vertex-colouring c such that:

- for every two adjacent vertices u and v , we have $c(u) \neq c(v)$,
- for every negative edge xy and positive edge uv , if $c(x) = c(u)$ then $c(y) \neq c(v)$.

Similarly as in the previous section, in case G cannot be properly vertex-coloured with strictly less than $|V(G)|$ colours, we call G a *signified clique*. According to the definition, G is a signified clique if and only if every two of its vertices are either adjacent, or joined by a path of length 2 whose one edge is positive and the other one is negative (see notably [8] for more details). Actually the reduction given in the proof of Theorem 5 can be modified to prove the following.

Corollary 8. It is NP-complete to decide whether an undirected graph G is the underlying graph of a signified clique.

The modifications are mainly the following. Instead of orienting the edges of G_H^c (and G_H), we now basically want to sign each of them either positive or negative. The reduced core graph G_H^c remains the same. A signing of the edges of G_H^c is *good* if, for every representative pair $\{u, v\}$ of vertices, the unique path with length 2 joining u to v has one positive edge and one negative edge. Then it can be easily checked that the cornerstone property of overlapping unique short paths remains applicable in this context. Namely, in a good signing, signing the edges of a unique path with length 2 in the core “forces” the signing of other unique paths overlapping it. The rest of the reduction, i.e. the construction of G_H from G_H^c , is similar.

Acknowledgements

We would like to thank Professor Éric Sopena for bringing the problem investigated in this paper to our attention, and for his valuable comments on the first drafts of this paper.

References

- [1] V. Chvátal and C. Thomassen. Distances in orientations of graphs. *J. Combin. Theory, Ser. B*, 24(1):61–75, 1978.
- [2] W. F. Klostermeyer and G. MacGillivray. Analogues of cliques for oriented coloring. *Discuss. Math. Graph Theory*, 24:373–387, 2004.

- [3] A. V. Kostochka, T. Łuczak, G. Simonyi, and É. Sopena. On the minimum number of edges giving maximum oriented chromatic number. *In Dimacs/Dimatia conference "Contemporary Trends in Discrete Mathematics", Stirin, Czech Rep., May 1997, Dimacs Series in Discrete Math. Theor. Comput. Sci.*, 49:179–182, 1999.
- [4] L. Lovász. Coverings and coloring of hypergraphs. *Proc. 4th South-Eastern Conference on Combinatorics, Graph Theory, and Computing, Boca Raton, Florida*, page 3–12, 1973.
- [5] A. Nandy, S. Sen, and É. Sopena. Outerplanar and planar oriented cliques. 2014. Preprint available online at <http://www.labri.fr/perso/sopena/pmwiki/uploads/Publications/nss14.pdf>.
- [6] R. Naserasr, E. Rollová, and É. Sopena. Homomorphisms of signed graphs. 2014. To appear in *J. Graph Theory*.
- [7] H.E. Robbins. A theorem on graphs, with an application to a problem in traffic control. *Amer. Math. Monthly*, 46:281–283, 1939.
- [8] S. Sen. *A contribution to the theory of graph homomorphisms and colorings*. PhD thesis, University of Bordeaux, 2014.
- [9] É. Sopena. Complete oriented colourings and the oriented achromatic number. *Discrete Appl. Math.*, 173:102–112, 2014.
- [10] R.E. Tarjan. A note on finding the bridges of a graph. *Inf. Proc. Lett.*, 2:160–161, 1974.