



HAL
open science

Poly[[μ -aqua-tetraaquabis(μ -2-hydroxy-4-oxocyclobut-1-ene-1,3-diolato)strontium] hemihydrate].

Amira Bouhali, Chahrazed Trifa, Sofiane Bouacida, Chaouki Boudaren,
Thierry Bataille

► To cite this version:

Amira Bouhali, Chahrazed Trifa, Sofiane Bouacida, Chaouki Boudaren, Thierry Bataille. Poly[[μ -aqua-tetraaquabis(μ -2-hydroxy-4-oxocyclobut-1-ene-1,3-diolato)strontium] hemihydrate].. Acta Crystallographica Section E: Structure Reports Online [2001-2014], 2011, 67 (Pt 8), pp.m1130-1. 10.1107/S1600536811028704 . hal-00825567

HAL Id: hal-00825567

<https://hal.science/hal-00825567>

Submitted on 24 May 2013

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Poly[[μ -aqua-tetraaquabis(μ -2-hydroxy-4-oxocyclobut-1-ene-1,3-diolato)-strontium] hemihydrate]

Amira Bouhali,^a Chahrazed Trifa,^a Sofiane Bouacida,^{a*} Chaouki Boudaren^a and Thierry Bataille^b

^aUnité de Recherche de Chimie de l'Environnement et Moléculaire Structurale, CHEMS, Université Mentouri–Constantine, 25000 Algeria, and ^bSciences Chimiques de Rennes, UMR 6226 CNRS - Université de Rennes 1, Avenue du Général Leclerc, 35042 Rennes cedex, France

Correspondence e-mail: Bouacida_Sofiane@yahoo.fr


Received 30 June 2011; accepted 17 July 2011

Key indicators: single-crystal X-ray study; $T = 150$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.028; wR factor = 0.068; data-to-parameter ratio = 12.6.

In the title coordination polymer, $[\text{Sr}(\text{C}_4\text{HO}_4)_2(\text{H}_2\text{O})_5] \cdot 0.5\text{H}_2\text{O}$, the Sr^{2+} ion is coordinated by three monodentate hydrogensquarate (hsq) anions and six aqua ligands in a distorted SrO_9 monocapped square-antiprismatic geometry. The hsq anions and water molecules bridge the metal ions into infinite sheets lying parallel to (100). The O atom of the uncoordinated water molecule lies on a crystallographic twofold axis. The packing is stabilized by numerous $\text{O} \cdots \text{H} \cdots \text{O}$ hydrogen bonds.

Related literature

For the isostructural mixed-metal Ba/Sr analogue of the title compound and background references, see: Trifa *et al.* (2011).



Experimental

Crystal data

$[\text{Sr}(\text{C}_4\text{HO}_4)_2(\text{H}_2\text{O})_5] \cdot 0.5\text{H}_2\text{O}$

$M_r = 412.81$

Monoclinic, $C2/c$

$a = 24.885$ (3) Å

$b = 8.8026$ (9) Å

$c = 13.8918$ (17) Å

$\beta = 119.609$ (4)°

$V = 2645.7$ (5) Å³

$Z = 8$

Mo $K\alpha$ radiation

$\mu = 4.15$ mm⁻¹

$T = 150$ K

$0.57 \times 0.27 \times 0.10$ mm

Data collection

Bruker APEXII diffractometer

Absorption correction: multi-scan

SADABS (Bruker, 2006)

$T_{\min} = 0.365$, $T_{\max} = 0.660$

9165 measured reflections

3006 independent reflections

2243 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.037$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.028$

$wR(F^2) = 0.068$

$S = 1.03$

3006 reflections

239 parameters

2 restraints

H atoms treated by a mixture of independent and constrained refinement

$\Delta\rho_{\text{max}} = 0.42$ e Å⁻³

$\Delta\rho_{\text{min}} = -0.51$ e Å⁻³

Table 1

Selected bond lengths (Å).

Sr1—O1	2.691 (2)	Sr1—O6	2.6179 (18)
Sr1—O2	2.642 (2)	Sr1—O12	2.6646 (16)
Sr1—O3	2.690 (2)	Sr1—O14 ⁱ	2.5906 (16)
Sr1—O4	2.641 (3)	Sr1—O3 ⁱⁱ	2.7154 (19)
Sr1—O5	2.572 (2)		

Symmetry codes: (i) $x, y + 1, z$; (ii) $-x + 2, -y + 2, -z + 1$.

Table 2

Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
O1—H1A \cdots O9 ⁱⁱⁱ	0.76 (4)	2.27 (4)	2.983 (3)	158 (4)
O1—H1B \cdots O7 ^{iv}	0.79 (4)	1.99 (4)	2.715 (2)	153 (4)
O1W—H1W \cdots O13 ⁱⁱ	0.76 (2)	2.39 (2)	2.801 (2)	115 (2)
O1W—H1W \cdots O8 ^v	0.76 (2)	2.51 (3)	3.118 (2)	138 (3)
O2—H2A \cdots O14 ^{vi}	0.92 (4)	1.91 (4)	2.794 (3)	161 (4)
O2—H2B \cdots O1W ^{vii}	0.70 (4)	2.52 (4)	3.165 (3)	154 (4)
O3—H3A \cdots O13 ⁱⁱ	0.93 (4)	1.79 (4)	2.712 (3)	174 (3)
O3—H3B \cdots O4 ^{viii}	0.76 (4)	2.59 (4)	3.172 (3)	136 (3)
O4—H4A \cdots O7 ^{ix}	0.78 (4)	2.01 (4)	2.785 (3)	177 (4)
O4—H4B \cdots O1W	0.87 (4)	2.03 (4)	2.871 (3)	163 (3)
O5—H5A \cdots O1 ^x	0.72 (4)	2.09 (4)	2.787 (3)	166 (4)
O5—H5B \cdots O8 ^{xi}	0.90 (4)	1.82 (4)	2.716 (3)	175 (4)
O9—H9 \cdots O12	0.82	1.74	2.548 (3)	169
O11—H11 \cdots O6 ^{vii}	0.82	1.77	2.580 (2)	172

Symmetry codes: (ii) $-x + 2, -y + 2, -z + 1$; (iii) $-x + \frac{3}{2}, y + \frac{1}{2}, -z + \frac{1}{2}$; (iv) $-x + \frac{3}{2}, y - \frac{1}{2}, -z + \frac{1}{2}$; (v) $x + \frac{1}{2}, -y + \frac{5}{2}, z + \frac{1}{2}$; (vi) $-x + 2, -y + 1, -z + 1$; (vii) $x, y - 1, z$; (viii) $-x + 2, y, -z + \frac{1}{2}$; (ix) $-x + \frac{3}{2}, -y + \frac{5}{2}, -z$; (x) $x, -y + 2, z - \frac{1}{2}$; (xi) $-x + \frac{3}{2}, -y + \frac{3}{2}, -z$.

Data collection: *APEX2* (Bruker, 2006); cell refinement: *SAINTE* (Bruker, 2006); data reduction: *SAINTE*; program(s) used to solve structure: *SIR2002* (Burla *et al.*, 2003); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997) and *DIAMOND* (Brandenburg & Berndt, 2001); software used to prepare material for publication: *WinGX* publication routines (Farrugia, 1999).

The authors thank Dr Thierry Roisnel, Centre de Diffraction X (CDIFX) de Rennes, Université de Rennes 1, France, for his technical assistance with the single-crystal X-ray data collection. This work was supported by the Unité de Recherche de Chimie de l'Environnement et Moléculaire Structurale, CHEMS, Université Mentouri–Constantine 25000 Algeria.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB5939).

References

- Brandenburg, K. & Berndt, M. (2001). *DIAMOND*. Crystal Impact, Bonn, Germany.
- Bruker (2006). *APEX2, SAINT and SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Burla, M. C., Camalli, M., Carrozzini, B., Cascarano, G. L., Giacovazzo, C., Polidori, G. & Spagna, R. (2003). *J. Appl. Cryst.* **36**, 1103.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Trifa, C., Bouhali, A., Bouacida, S., Boudaren, C. & Bataille, T. (2011). *Acta Cryst.* **E67**, m275–m276.