



HAL
open science

Quasilinear Elliptic Hamilton-Jacobi Equations on Complete Manifolds

Marie-Françoise Bidaut-Veron, Marta Garcia-Huidobro, Laurent Veron

► **To cite this version:**

Marie-Françoise Bidaut-Veron, Marta Garcia-Huidobro, Laurent Veron. Quasilinear Elliptic Hamilton-Jacobi Equations on Complete Manifolds. 2013. hal-00827016v2

HAL Id: hal-00827016

<https://hal.science/hal-00827016v2>

Preprint submitted on 1 Jun 2013 (v2), last revised 4 Jun 2013 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**QUASILINEAR ELLIPTIC HAMILTON-JACOBI
EQUATIONS ON COMPLETE MANIFOLDS**

Marie-Françoise Bidaut-Véron¹, Marta Garcia-Huidobro², Laurent Véron³

RÉSUMÉ. Let (M^n, g) be a n -dimensional complete, non-compact and connected Riemannian manifold with Ricci tensor $Ricc_g$ and sectional curvature Sec_g . Assume $Ricc_g \geq (1 - n)B^2$ and either $p > 2$ and $Sec_g(x) = o(dist^2(x, a))$ when $dist^2(x, a) \rightarrow \infty$ for $a \in M$, or $1 < p < 2$ and $Sec_g(x) \leq 0$. If $q > p - 1 > 0$ any C^1 solution of (E) $-\Delta_p u + |\nabla u|^q = 0$ on M satisfies $|\nabla u(x)| \leq c_{n,p,q} B^{\frac{1}{q+1-p}}$ for some constant $c_{n,p,q} > 0$. As a consequence there exists $c_{n,p} > 0$ such that any positive p -harmonic function v on M satisfies $v(a)e^{-c_{n,p}Bdist(x,a)} \leq v(x) \leq v(a)e^{c_{n,p}Bdist(x,a)}$ for any $x \in M$.

**Equations de Hamilton-Jacobi quasilineaires sur une variété
complète**

RÉSUMÉ. Soit (M^n, g) une variété riemannienne n -dimensionnelle complète, non compacte et connexe de courbures de Ricci $Ricc_g$ et sectionnelle Sec_g . On suppose $Ricc_g \geq (1 - n)B^2$ et $Sec_g(x) = o(dist^2(x, a))$ si $dist^2(x, a) \rightarrow \infty$ pour $a \in M$ si $p > 2$, ou $Sec_g(x) \leq 0$ si $1 < p < 2$. Si $q > p - 1 > 0$, toute solution de classe C^1 de (E) $-\Delta_p u + |\nabla u|^q = 0$ sur M satisfait à $|\nabla u(x)| \leq c_{n,p,q} B^{\frac{1}{q+1-p}}$ pour une constante $c_{n,p,q} > 0$. On en déduit qu'il existe $c_{n,p} > 0$ tel que toute fonction p -harmonique positive v sur M vérifie $v(a)e^{-c_{n,p}Bdist(x,a)} \leq v(x) \leq v(a)e^{c_{n,p}Bdist(x,a)}$ pour tout $x \in M$.

Version française abrégée. Soit (M^n, g) une variété riemannienne complète, connexe et non-compacte de courbure de Ricci $Ricc_g$ et courbure sectionnelle Sec_g . Si $p > 1$, on dénote par $u \mapsto \Delta_p u := div(|\nabla u|^{p-2} \nabla u)$ le p -Laplacien sur M pour la métrique g . Notre résultat principal est le suivant

Theorème 1. Soit $B \geq 0$ tel que $Ricc_g \geq (1 - n)B^2$ et $q > p - 1 > 0$. On suppose

$$(1) \quad \lim_{dist(x,a) \rightarrow \infty} \frac{Sec_g(x)}{(dist(x, a))^2} = 0$$

1. Laboratoire de Mathématiques et Physique Théorique, CNRS UMR 7350, Faculté des Sciences, 37200 Tours France. E-mail : veronmf@univ-tours.fr

2. Departamento de Matematicas, Pontifica Universidad Catolica de Chile Casilla 307, Correo 2, Santiago de Chile. E-mail : mgarcia@mat.puc.cl

3. Laboratoire de Mathématiques et Physique Théorique, CNRS UMR 7350, Faculté des Sciences, 37200 Tours France. E-mail : veronl@univ-tours.fr

pour $a \in M$ si $p > 2$ ou $Sec_g \leq 0$ si $1 < p < 2$. Il existe alors $c_{n,p,q} > 0$ telle que toute solution $u \in C^1(M)$ de

$$(2) \quad -\Delta_p u + |\nabla u|^q = 0 \quad \text{sur } M$$

vérifie

$$(3) \quad |\nabla u(x)| \leq c_{n,p,q} B^{\frac{1}{q+1-p}} \quad \forall x \in M.$$

Une des conséquences est un théorème de type Liouville.

Corollaire 2. *Supposons $Ricc_g \geq 0$, $q > p - 1 > 0$ et que les hypothèses du Théorème 1 portant sur la courbure scalaire sont vérifiées si $p \neq 2$. Alors toute solution $u \in C^1(M)$ de (2) est constante.*

Si v est une fonction p -harmonique positive sur M , la fonction $u := -(p-1) \ln v$ vérifie

$$(4) \quad -\Delta_p u + |\nabla u|^p = 0 \quad \text{sur } M.$$

En utilisant le résultat du théorème 1, on en déduit

Théorème 3. *Supposons $p > 1$ et que les hypothèses du Théorème 1 portant sur la courbure sont vérifiées. Il existe alors une constante $c_{n,p} > 0$ telle que toute fonction p -harmonique et positive v sur M vérifie*

$$(5) \quad v(a)e^{-c(n,p)B \text{dist}(x,a)} \leq v(x) \leq v(a)e^{c(n,p)B \text{dist}(x,a)} \quad \forall (a, x) \in M \times M.$$

Quand $p = 2$ Cheng et Yau [1] ont montré que toute fonction harmonique positive sur une variété riemannienne complète à courbure de Ricci positive est une constante. Dans le cas des fonctions p -harmoniques positives et sous l'hypothèse de minoration uniforme de la courbure sectionnelle, $Sec_g \geq -B^2$, Kotschwar et Ni [4] montrent l'estimation suivante que vérifie toute fonction p -harmonique positive v sur M .

$$(6) \quad \frac{|\nabla v|}{v} \leq (p-1)B.$$

Notons que leur hypothèse implique $Ricc_g \geq -(n-1)B^2$.

Let (M^n, g) be a complete, connected and non compact Riemannian manifold with Ricci curvature $Ricc_g$ and sectionnal curvature Sec_g . For $p > 1$ we denote by Δ_p the p -Laplacian defined in the metric g by

$$\Delta_p u := \text{div} \left(|\nabla u|^{p-2} \nabla u \right),$$

and thus Δ_2 is the Laplace-Beltrami operator on M . If $p = 2$ a classical result due to Cheng and Yau [1] asserts that if $Ricc_g$ is nonnegative, any nonnegative harmonic function v is a constant. In [4], Kotschwar et Ni obtained sharper results dealing with positive p -harmonic functions under the

assumption that $Sec_g \geq -B^2$. They proved that if v is such a function, it satisfies

$$(1) \quad \frac{|\nabla v|}{v} \leq (p-1)B.$$

Their assumption on Sec_g implies $Ricc_g \geq (1-n)B^2$. They also noticed that if $p = 2$ their estimate holds under the previous lower estimate on the Ricci curvature. In this note we give an extension of their result in imbedding it the more general class of quasilinear Hamilton-Jacobi type equations

$$(2) \quad -\Delta_p u + |\nabla u|^q = 0 \quad \text{on } M.$$

Our main result is the following

Theorem 1. *Let $B \geq 0$ such that $Ricc_g \geq (1-n)B^2$. We assume*

$$(3) \quad \lim_{\text{dist}(x,a) \rightarrow \infty} \frac{Sec_g(x)}{(\text{dist}(x,a))^2} = 0$$

for $a \in M$ if $p > 2$, or $Sec_g \leq 0$ if $1 < p < 2$. Then there exists $c_{n,p,q} > 0$ such that any solution $u \in C^1(M)$ of (2) satisfies

$$(4) \quad |\nabla u(x)| \leq c_{n,p,q} B^{\frac{1}{q+1-p}} \quad \forall x \in M.$$

A clear consequence of (3) is the following Liouville theorem

Corollary 2. *Assume $Ricc_g \geq 0$ and that the assumptions of Theorem 1 concerning Sec_g hold if $p \neq 2$. Then any solution $u \in C^1(M)$ of (2) is constant.*

If v is a positive p -harmonic function on M , then $u := -(p-1) \ln v$ satisfies

$$(5) \quad -\Delta_p u + |\nabla u|^p = 0 \quad \text{on } M.$$

Therefore estimate (4) yields to the following result

Theorem 3. *Assume $p > 1$ and that the curvature assumptions of Theorem 1 hold. There exists a constant $c_{n,p} > 0$ such that any positive p -harmonic function v on M satisfies*

$$(6) \quad v(a)e^{-c(n,p)B \text{dist}(x,a)} \leq v(x) \leq v(a)e^{c(n,p)B \text{dist}(x,a)} \quad \forall (a,x) \in M \times M.$$

Proof of Theorem 1. Let $M_+ := \{x \in M : |\nabla u(x)| > 0\}$. Then M_+ is open and $u \in C^3(M_+)$ since the equation is no longer degenerate. The proof is based upon the fact that $z = |\nabla u|^2$ is a subsolution of an elliptic differential inequality with a super linear absorption term (see [5] for other applications). We denote by TM the tangent bundle of M and by $\langle \cdot, \cdot \rangle$ the

scalar product induced by the metric g . We recall that any C^3 -function u verifies the Böchner-Weitzenböck formula; combined with Schwarz inequality it yields to

$$(7) \quad \begin{aligned} \frac{1}{2} \Delta_2 |\nabla u|^2 &= |D^2 u|^2 + \langle \nabla \Delta_2 u, \nabla u \rangle + Ricc_g(\nabla u, \nabla u) \\ &\geq \frac{1}{n} |\Delta_2 u|^2 + \langle \nabla \Delta_2 u, \nabla u \rangle + Ricc_g(\nabla u, \nabla u), \end{aligned}$$

where $D^2 u$ is the Hessian. If u is a C^1 solution of (2), then $z = |\nabla u|^2$ satisfies

$$(8) \quad -\Delta_2 z - \frac{p-2}{2} \frac{\langle \nabla z, \nabla u \rangle}{z} + z^{\frac{q+2-p}{2}} = 0$$

on M_+ . Replacing $\Delta_2 u$ in (7) it follows that, for any $a > 0$,

$$(9) \quad \begin{aligned} \Delta_2 z + (p-2) \frac{\langle D^2 z(\nabla u), \nabla u \rangle}{z} &\geq \frac{2a^2}{N} z^{q+2-p} - \frac{1}{Na^2} \frac{\langle \nabla z, \nabla u \rangle^2}{z^2} - \frac{(p-2)}{2} \frac{|\nabla z|^2}{z} \\ &\quad + (p-2) \frac{\langle \nabla z, \nabla u \rangle^2}{z^2} + (q+2-p) z^{\frac{q-p}{2}} \langle \nabla z, \nabla u \rangle - (N-1) B^2 z. \end{aligned}$$

Since $z^{\frac{q-p}{2}} |\langle \nabla z, \nabla u \rangle| \leq z^{\frac{q+1-p}{2}} \frac{|\nabla z|}{\sqrt{z}}$, we can take $a = a(p, q) > 0$ large enough so that the right-hand side of (9) is bounded from below by $Cz^{q+2-p} - D \frac{|\nabla z|^2}{z}$ for some $C, D > 0$ which depend only on p and q . We set

$$\mathcal{A}(v) := -\Delta_2 v - (p-2) \frac{\langle D^2 v(\nabla u), \nabla u \rangle}{|\nabla u|^2} = - \sum_{i,j=1}^N a_{ij} v_{x_i x_j}$$

where the a_{ij} depend on ∇u and satisfy

$$\theta |\xi|^2 := \min\{1, p-1\} |\xi|^2 \leq \sum_{i,j=1}^N a_{ij} \xi_i \xi_j \leq \max\{1, p-1\} |\xi|^2 := \Theta |\xi|^2 \quad \forall \xi \in \mathbb{R}^n.$$

Then

$$(10) \quad \mathcal{L}^*(z) := \mathcal{A}(z) + Cz^{q+2-p} - D \frac{|\nabla z|^2}{z} - (n-1)B^2 z \leq 0 \quad \text{in } M_+.$$

The next lemma is a local estimate.

Lemma 1. *Let $B_R(a) \subset M^n(g)$ be the geodesic ball of radius $R > 0$ and center a . Assume that $Ricc_g \geq -(n-1)B^2$ and either $Sec_g \geq -S^2$ for some $S^2 := S_R^2$ in $B_R(a)$ if $p > 2$, or $Sec_g \leq 0$ if $1 < p < 2$. Then there exists $c = c(n, p, q) > 0$ such that the function*

$$(11) \quad w(x) = \lambda (R^2 - r^2(x))^{-\frac{2}{q+1-p}} + \mu \quad \text{with } r = r(x) = d(x, a),$$

satisfies $\mathcal{L}^*(w) \geq 0$ in $B_R(a)$, provided

$$(12) \quad \lambda = c \max \left\{ (R^4 B^2)^{\frac{1}{q+1-p}}, ((1+B+(p-2)_+ S) R^3)^{\frac{1}{q+1-p}} \right\}$$

and

$$(13) \quad \mu = ((n-1)B^2)^{\frac{1}{q+1-p}}.$$

Proof. We recall that $\Delta_2 w = w'' + w' \Delta_2 r$ and by [6, Lemma 1]

$$\Delta_2 r \leq (n-1)B \coth(Br) \leq \frac{n-1}{r} (1+Br).$$

Then

$$(14) \quad \Delta_2 w \leq \frac{4}{q+1-p} (R^2 - r^2)^{-\frac{2(q+2-p)}{q+1-p}} \left(\frac{2r^2(q+3-p)}{q+1-p} + (R^2 - r^2)(1 + (n-1)(1+Br)) \right).$$

Moverover [3, Chap 2, p. 23]

$$(15) \quad D^2 w = w'' dr \otimes dr + w' D^2 r.$$

If $0 \geq \text{Sec}_g(x) \geq -S^2$, there holds

$$(16) \quad 0 \leq D^2 r \leq S \coth(Sr) g \leq \frac{S}{r} (1+Sr) g.$$

Therefore, if $p \geq 2$ and $\text{Sec}_g \geq -S^2$, we get

$$(17) \quad \frac{\langle D^2 w(\nabla u), \nabla u \rangle}{|\nabla u|^2} \leq \frac{4}{q+1-p} (R^2 - r^2)^{-\frac{2(q+2-p)}{q+1-p}} \left(\frac{2r^2(q+3-p)}{q+1-p} + (R^2 - r^2)(2+Sr) \right).$$

while, if $p \leq 2$ and $\text{Sec}_g \leq 0$,

$$(18) \quad \frac{\langle D^2 w(\nabla u), \nabla u \rangle}{|\nabla u|^2} \leq \frac{4}{q+1-p} (R^2 - r^2)^{-\frac{2(q+2-p)}{q+1-p}} \left(\frac{2r^2(q+3-p)}{q+1-p} + 2(R^2 - r^2) \right).$$

As a consequence

$$(19) \quad \begin{aligned} \mathcal{A}(w) &= -\Delta w - (p-2) \frac{\langle D^2 w(\nabla u), \nabla u \rangle}{|\nabla u|^2} \\ &\geq -k \lambda (R^2 - r^2)^{-\frac{2(q+2-p)}{q+1-p}} (R^2 + (R^2 - r^2) B_p r) \end{aligned}$$

for some $k = k(n, p, q)$, where $B_p = B + (p-2)_+ S$. Since

$$w^{q+2-p} \geq \lambda^{q+2-p} (R^2 - r^2)^{-\frac{2(q+1-p)}{q+1-p}} + \mu^{q+2-p}.$$

we have

$$(20) \quad \begin{aligned} \mathcal{L}^*(w) &\geq \lambda (R^2 - r^2)^{-\frac{2(q+2-p)}{q+1-p}} \left(-k (R^2 + (R^2 - r^2) B_p r) - D \frac{16}{(q+1-p)^2} r^2 + C \lambda^{q+1-p} \right) \\ &\quad + \mu^{q+2-p} - (n-1) B^2 \lambda (R^2 - r^2)^{-\frac{2}{q+1-p}} - (n-1) B^2 \mu. \end{aligned}$$

We first take

$$(21) \quad \mu = ((n-1)B^2)^{\frac{1}{q+1-p}}.$$

Next we choose λ in order to have, uniformly for $0 \leq r < R$,

$$2^{-1}C\lambda^{q+1-p} \geq k(R^2 + (R^2 - r^2)B_p r) + \frac{16Dr^2}{(q+1-p)^2}$$

and

$$2^{-1}C\lambda^{q+2-p}(R^2 - r^2)^{-\frac{2(q+2-p)}{q+1-p}} \geq (n-1)B^2\lambda(R^2 - r^2)^{-\frac{2}{q+1-p}}.$$

There exists $c = c(n, p, q)$ such that, if

$$(22) \quad \lambda = c \max \left\{ (R^4 B^2)^{\frac{1}{q+1-p}}, ((1 + B_p)R^3)^{\frac{1}{q+1-p}} \right\},$$

then $\mathcal{L}^*(w) \geq 0$ holds.

Lemma 2. *Under the assumptions of Lemma 1, any C^1 solution of (2) in M satisfies*

$$(23) \quad |\nabla u(x)| \leq c_{n,p,q} \max \left\{ B^{\frac{1}{q+1-p}}, (1 + B_p)^{\frac{1}{2(q+1-p)}} (d(x, \partial\Omega))^{-\frac{1}{2(q+1-p)}} \right\} \quad \forall x \in \Omega,$$

for all open domain $\Omega \subset M$, where $B_p = B + (p-2)_+ S$ and $S = S_{d(x, \partial\Omega)}$.

Proof. Assume $a \in \Omega$, with $R < d(a, \partial\Omega)$. Let w be as in Lemma 1, then in any connected component G of $\{x \in B_R(a) : z(x) - w(x) > 0\}$ we have

$$(24) \quad \mathcal{A}(z-w) + C(z^{q+2-p} - w^{q+2-p}) - (n-1)B^2(z-w) - D \left(\frac{|\nabla z|^2}{z} - \frac{|\nabla w|^2}{w} \right) \leq 0.$$

By the mean value theorem and since $w(a)$ is the minimum of w , there holds

$$(25) \quad C(z^{q+2-p} - w^{q+2-p}) - (n-1)B^2(z-w) > 0,$$

provided $C(q+2-p)(w(a))^{q+1-p} > (n-1)B^2$. Since $w(a) > \mu = ((n-1)B^2)^{\frac{1}{q+1-p}}$ and $q+2-p > 1$, this condition is fulfilled, up to replacing μ by $A\mu$ for some $A = A(p, q) > 1$. If $x_0 \in G$ is such that $z-w$ is maximal at x_0 , we derive that

$$\mathcal{A}(z-w) + C(z^{q+2-p} - w^{q+2-p}) - (N-1)B^2(z-w) - D \left(\frac{|\nabla z|^2}{z} - \frac{|\nabla w|^2}{w} \right) \leq 0$$

if $x = x_0$, which is a contradiction. Thus $G = \emptyset$, $z \leq w$ and (23) follows.

The proof of Theorem 1 and Corollary 2 follows by taking $\Omega = B_R(x)$ and letting $R \rightarrow \infty$.

Proof of Theorem 3. We take $q = p$ and assume that v is p -harmonic and positive. If we write $v = e^{-\frac{u}{p-1}}$, then u satisfies

$$-\Delta_p u + |\nabla u|^p = 0.$$

If $\text{Ric}_g(x) \geq 0$, u is constant by Corollary 2, and so is v . If $\inf\{\text{Ric}_g(x) : x \in M\} = (1-n)B^2 < 0$ we apply (23) to ∇u . If γ is a minimizing geodesic from a to x , then $|\gamma'(t)| = 1$ and

$$u(x) - u(a) = \int_0^{d(x,a)} \frac{d}{dt} u \circ \gamma(t) dt = \int_0^{d(x,a)} \langle \nabla u \circ \gamma(t), \gamma'(t) \rangle dt$$

Since

$$|\langle \nabla u \circ \gamma(t), \gamma'(t) \rangle| \leq |\nabla u \circ \gamma(t)| \leq c_{n,p,p} B$$

we obtain

$$(26) \quad u(a) - c_{n,p,p} B \text{dist}(x, a) \leq u(x) \leq u(a) + c_{n,p,p} B \text{dist}(x, a) \quad \forall x \in M,$$

Then (6) follows since $u = (1-p) \ln v$.

RÉFÉRENCES

- [1] S. Y. Cheng and S.-T. Yau : *Differential equations on Riemannian manifolds and their geometric applications*, **Comm. Pure Appl. Math.** **28**, 333–354 (1975).
- [2] T. H. Colding and W.P. Minicozzi : *An excursion into geometric analysis*, Grigor'yan, Alexander (ed.) et al., *Surveys in differential geometry*. Somerville, MA : International Press, **Surveys in Differential Geometry** **9**, 83-146 (2004).
- [3] R.E. Greene and H. Wu/ *Function Theory on Manifolds Which Possess a Pole*, **Lecture Notes in Mathematics** **699**, Springer-Verlag Berlin Heidelberg (1979).
- [4] B. L. Kotschwar and L. Ni : *Local gradient estimates of p -harmonic functions, $1/H$ -flow, and an entropy formula*, **Ann. Sci. École. Norm. Sup.** **42**, 1–36 (2009).
- [5] A. Porretta and L. Véron : *Separable p -harmonic functions in a cone and related quasilinear equations on manifolds*, **J. Eur. Math. Soc.** **11**, 1285-1305 (2009).
- [6] A. Ratto, M. Rigoli and L. Véron *Conformal immersion of complete Riemannian manifolds and extensions of the Schwarz lemma*, **Duke Math. J.** **74**, 223-236 (1994).