



HAL
open science

Multi-solitons for nonlinear Klein-Gordon equations

Raphaël Côte, Claudio Muñoz

► **To cite this version:**

Raphaël Côte, Claudio Muñoz. Multi-solitons for nonlinear Klein-Gordon equations. Forum of Mathematics, Sigma, 2014, 2, pp.e15. 10.1017/fms.2014.13 . hal-00832756

HAL Id: hal-00832756

<https://hal.science/hal-00832756v1>

Submitted on 14 Feb 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

MULTI-SOLITONS FOR NONLINEAR KLEIN-GORDON EQUATIONS

RAPHAËL CÔTE AND CLAUDIO MUÑOZ

ABSTRACT. In this paper we consider the existence of multi-soliton structures for the nonlinear Klein-Gordon equation (NLKG) in \mathbb{R}^{1+d} . We prove that, independently of the unstable character of (NLKG) solitons, it is possible to construct a N -soliton family of solutions to (NLKG), of dimension $2N$, globally well-defined in the energy space $H^1 \times L^2$ for all large positive times. The method of proof involves the generalization of previous works on supercritical NLS and gKdV equations by Martel, Merle and the first author [3] to the wave case, where we replace the unstable mode associated to the linear NLKG operator by two generalized directions that are controlled without appealing to modulation theory. As a byproduct, we generalize the linear theory described in Grillakis-Shatah-Strauss [10] and Duyckaerts-Merle [7] to the case of boosted solitons, and provide new solutions to be studied using the recent Nakanishi-Schlag [24] theory.

1. INTRODUCTION

In this paper we are interested in the problem of constructing multi-soliton solutions for some well-known scalar field equations. Let $f = f(s)$ be a real-valued \mathcal{C}^1 -function. We consider the nonlinear Klein-Gordon equation (NLKG) in \mathbb{R}^{1+d} , $d \geq 1$,

$$\partial_{tt}u - \Delta u + u - f(u) = 0, \quad u(t, x) \in \mathbb{R}, \quad (t, x) \in \mathbb{R} \times \mathbb{R}^d. \quad (\text{NLKG})$$

This equation arises in Quantum Field Physics as a model for a self-interacting, nonlinear scalar field, invariant under Lorentz transformations (see below).

Let F be the standard integral of f :

$$F(s) := \int_0^s f(\sigma) d\sigma. \quad (1)$$

We will assume that for some fixed constant $C > 0$,

- (A) If $d = 1$,
 - (i) f is odd, and $f(0) = f'(0) = 0$, and
 - (ii) There exists $s_0 > 0$ such that $F(s_0) - \frac{1}{2}s_0^2 > 0$.
- (B) If $d \geq 2$, f is a pure power H^1 -subcritical nonlinearity: $f(u) = \lambda|u|^{p-1}u$, where $\lambda > 0$, $p \in (1, 1 + \frac{4}{d-2})$.

2010 *Mathematics Subject Classification.* 35Q51,35L71,35Q40.

Key words and phrases. Klein-Gordon equation, soliton, construction, instability, multi-soliton.

The first author wishes to thank the University of Chicago for its hospitality during the academic year 2011-12, and acknowledges support from the European Research Council through the project BLOWDISOL.

Prescribing f to the above class of focusing nonlinearities ensures that the corresponding Cauchy problem for (NLKG) is locally well-posed in $H^s(\mathbb{R}^d) \times H^{s-1}(\mathbb{R}^d)$, for any $s \geq 1$: we refer to Ginibre-Velo [12] and Nakamura-Ozawa [23] (when $d = 2$) for more details.

Also under the above conditions, the Energy and Momentum (every integral is taken over \mathbb{R}^d)

$$E[u, u_t](t) = \frac{1}{2} \int [|\partial_t u(t, x)|^2 + |\nabla u(t, x)|^2 + |u(t, x)|^2 - 2F(u(t, x))] dx, \quad (2)$$

$$P[u, u_t](t) = \frac{1}{2} \int \partial_t u(t, x) \nabla u(t, x) dx, \quad (3)$$

are conserved along the flow.

Another important feature of equation (NLKG), still under the previous conditions, is the fact that it admits stationary solutions of the form $u(t, x) = U(x)$ (i.e., with no dependence on t). Among them, we are interested in the ground-state $Q = Q(x)$, where Q is a positive solution of the elliptic PDE

$$\Delta Q - Q + f(Q) = 0, \quad Q > 0, \quad Q \in H^1(\mathbb{R}^d). \quad (4)$$

The existence of this solution goes back to Berestycki-Lions [1], provided the above conditions (in particular (ii)) hold. Additionally, it is well-known that Q is radial and exponentially decreasing, along with its first and second derivatives (Gidas-Nirenberg [9]), and *unique* up to definition of the origin (see Kwong [14], Serrin and Tang [27]).

In fact, our main result written below could be extended to more general nonlinearity under an additional assumption of spectral nature, namely that the linearized operator around Q has a standard simple spectrum. More precisely, Theorem 1 holds, as soon as f satisfies (i), (ii) and:

- (iii) If $d = 2$, $|f'(s)| \leq C|s|^p e^{\kappa s^2}$, for some $p \geq 0$, $\kappa > 0$ and all $s \in \mathbb{R}$.
- (iv) If $d \geq 3$, $|f'(s)| \leq C(1 + |s|^{p-1})$ for some $p < 1 + \frac{4}{d-2}$ and all $s \in \mathbb{R}$.
- (v) $-\Delta z + z - f'(Q)z$ has a unique simple negative eigenvalue, and its kernel is given by $\{x \cdot \nabla Q | x \in \mathbb{R}^d\}$ and it is nondegenerate.

Assumption (v) has been checked in the cases (A) and (B) (using ODE analysis), and is believed to hold for a wide class of functions f . (See Lemma 4.)

Since (NLKG) is invariant under *Lorentz boosts*, we can define a *boosted ground state* (a *soliton* from now on) with relative velocity $\beta \in \mathbb{R}^d$. More precisely, let $\beta = (\beta_1, \dots, \beta_d) \in \mathbb{R}^d$, with $|\beta| < 1$ (we denote $|\cdot|$ the euclidian norm on \mathbb{R}^d), the corresponding Lorentz boost is given by the $(d+1) \times (d+1)$ matrix

$$\Lambda_\beta := \begin{pmatrix} \gamma & -\beta_1 \gamma & \cdots & \beta_d \gamma \\ -\beta_1 \gamma & & & \\ \vdots & & \text{Id}_d + \frac{(\gamma-1)}{|\beta|^2} \beta \beta^T & \\ -\beta_d \gamma & & & \end{pmatrix} \quad \text{where} \quad \gamma := \frac{1}{\sqrt{1-|\beta|^2}}, \quad (5)$$

$(\beta\beta^T$ is the $d \times d$ rank 1 matrix with coefficient of index (i, j) $\beta_i\beta_j$). Then the boosted soliton with velocity β is

$$Q_\beta(t, x) := Q\left(\Lambda_\beta\begin{pmatrix} t \\ x \end{pmatrix}\right), \quad (6)$$

where with a slight abuse of notation we say that $Q(t, x) = Q(x)$ (namely we project on the last d coordinates). Also notice that (NLKG) is invariant by space translation (shifts). Hence the general family of solitons is parametrized by speed $\beta \in \mathbb{R}^d$ and shift (translation) $x_0 \in \mathbb{R}^d$:

$$(t, x) \mapsto Q_\beta(t, x - x_0).$$

This family is the orbit of $\{Q\}$ under all the symmetries of (NLKG) (general Lorentz transformation, time and space shifts), in particular it is invariant under these transformations: see the Appendix A for further details.

In the rest of this work, it will be convenient to work with vector data $(u, \partial_t u)^T$. For notational purposes, we use upper-case letters to denote vector valued functions and lower-case letters for scalar functions (except for the scalar field Q_β).

We will work in the energy space $H^1(\mathbb{R}^d) \times L^2(\mathbb{R}^d)$ endowed with the following scalar product: denote $U = (u_1, u_2)^T$, $V = (v_1, v_2)^T$, we define

$$\begin{aligned} \langle U|V \rangle &= \left\langle \begin{pmatrix} u_1 \\ u_2 \end{pmatrix} \middle| \begin{pmatrix} v_1 \\ v_2 \end{pmatrix} \right\rangle := (u_1|v_1) + (u_2|v_2) = \int (u_1 v_1 + u_2 v_2), \quad (7) \\ &\text{where } (u|v) := \int uv, \end{aligned}$$

and the energy norm

$$\|U\|^2 := \langle U|U \rangle + (\nabla u_1|\nabla u_1) = \|u_1\|_{H^1}^2 + \|u_2\|_{L^2}^2. \quad (8)$$

It is well known (see e.g. Grillakis-Shatah-Strauss [10]) that $(Q, 0)$ is unstable¹ in the energy space. The instability properties of Q and solution with energy slightly above $E[(Q, 0)]$ have recently been further studied by Nakanishi and Schlag, see [24] and subsequent works. Their ideas are further developments of the primary idea introduced in Duyckaerts-Merle [6], in the context of the energy critical nonlinear wave equation (where the relevant nonlinear object is the stationary function W which solves $\Delta W + W^{1+4/(d-2)} = 0$).

In this paper, we want to understand the dynamics of large, quantized energy solutions. More precisely, we address the question whether is it possible to construct a multi-soliton solution for (NLKG), i.e. a solution u to (NLKG) defined on a semi-infinite interval of time, such that

$$u(t, x) \sim \sum_{j=1}^N Q_{\beta_j}(t, x - x_j) \quad \text{as } t \rightarrow +\infty.$$

Such solutions were constructed for the nonlinear Schrödinger equation and the generalized Korteweg-de Vries equation, first in the L^2 -critical and subcritical case by Merle [18], Martel [16] and Martel-Merle [19]. These results followed from the stability and asymptotic stability theory that these authors developed.

¹This result was known in the physics literature as the Derrick's Theorem [5].

The existence of multi-solitons was then extended by Martel-Merle and the first author [3] to the L^2 supercritical case: in this latter case, each single soliton is unstable, hence the multi-soliton is a highly unstable solution. It turns out that this is also the case for scalar field equations as (NLKG). We prove that, regardless of the instability of the soliton, one can construct large mass multi-solitons, on the whole range of parameters $\beta_1, \dots, \beta_N \in \mathbb{R}^d$ distinct, with $|\beta_j| < 1$ and $x_1, \dots, x_N \in \mathbb{R}^d$. More precisely, the main result of this paper is the following.

Theorem 1. *Assume (A) or (B), and let $\beta_1, \beta_2, \dots, \beta_N \in \mathbb{R}^d$ be a set of different velocities*

$$\forall i \neq j, \quad \beta_i \neq \beta_j, \quad \text{and} \quad |\beta_j| < 1,$$

and $x_1, x_2, \dots, x_N \in \mathbb{R}^d$ shift parameters.

Then there exist a time $T_0 \in \mathbb{R}$, constants $C > 0$, and $\gamma_0 > 0$, only depending on the sets $(\beta_j)_j$, $(x_j)_j$, and a solution $(u, \partial_t u) \in \mathcal{C}([T_0, +\infty), H^1(\mathbb{R}^d) \times L^2(\mathbb{R}^d))$ of (NLKG), globally defined for forward times and satisfying

$$\forall t \geq T_0, \quad \left\| (u, \partial_t u)(t, x) - \sum_{j=1}^N (Q_{\beta_j}, \partial_t Q_{\beta_j})(t, x - x_j) \right\| \leq C e^{-\gamma_0 t}.$$

We remark that this is the first multi-soliton result for wave-type equations. Although the nonlinear object under consideration is the same as for (NLS) for example, the structure of the flow is different (recall that all solitons are unstable for (NLKG), irrespective of the nonlinearity). Hence we need to work in a more general framework, the one given by a matrix description of (NLKG).

Let us describe the main steps of the proof. We first revisit the standard spectral theory of linearized operators around the soliton, and the second order derivative of the energy-momentum functional (see H in (14)) [10]. Since solitons are unstable objects, it is clear that such a theory will not be enough to describe the dynamics of several solitons. However, a slight variation of this functional (see \mathcal{H} in (24)) turns out to be the key element to study. We describe its spectrum in great detail, in particular we prove that this operator has three eigenvalues: the kernel zero, and two opposite sign eigenvalues, with associated eigenfunctions Z_{\pm} . After some work we are able to prove a coercivity property for the operator \mathcal{H} modulo the two directions Z_+ and Z_- . This analysis was first conducted by Pego and Weinstein [25] in the context of generalized KdV equations.

The rest of the work is devoted to the study of the dynamics of small perturbations of the sum of N solitons, in particular how the two directions associated to Z_{\pm} evolve. Using a topological argument, we can show the existence of suitable initial data for (NLKG) such that both directions remain controlled for all large positive time, proving the main theorem. We remark that this method is general and does not require the study of the linear evolution at large, but also a deep understanding of suitable alternative directions of the linearized operator. A nice open question should be the extension of this result to the nonlinear wave case, where the soliton decays polynomially.

For the sake of easiness and clarity, we present the detailed computations in the one dimensional case $d = 1$. This case encompass all difficulties, the higher dimension case adding only indices and notational inconvenience: we will briefly describe the corresponding differences at the end of each section.

Organization of this paper. In Section 2, we develop spectral aspects of the linearized flow around Q_β , which are more subtle than in the (NLS) or (gKdV) case. In Section 3, we construct approximate N -soliton solutions in Proposition 3, which we do by estimation backward in time as in [18, 16, 19]. There we present the nonlinear argument, relying *in fine* on a topological argument as in [3]. The Lyapunov functional has to be chosen carefully, as we cannot allow mixed derivatives of the form $\partial_{tx}u$. Finally in Section 4, we prove Theorem 1, relying on the previously proved Proposition 3 and a compactness procedure.

Acknowledgment

We would like to thank Wilhelm Schlag for pointing us this problem and for enlightening discussions. We are deeply indebted to the anonymous referee, who we thank for his thorough reading and comments which improved the manuscript significantly.

2. SPECTRAL THEORY

In this section we describe and solve two spectral problems related to (NLKG). We will work with functions independent of time, unless specified explicitly. The main result of this section is Proposition 2.

2.1. Coercivity of the Hessian. First of all, we recall the structure of the Hessian of the energy around Q . Given $Q = Q(x)$ ground state of (4) and $Q_\beta(x) = Q(\gamma x)$, where $\gamma = (1 - \beta^2)^{-1/2}$, we define the operators

$$L^+ := -\partial_{xx} + \text{Id} - f'(Q), \quad \text{and} \quad L_\beta^+ = -\gamma^{-2}\partial_{xx} + \text{Id} - f'(Q_\beta), \quad (9)$$

so that L_β^+ is a rescaled version of L^+ :

$$L_\beta^+(v(\gamma x)) = (L^+v)(\gamma x).$$

As a consequence of the Sturm-Liouville theory and the previous identity, we have the following spectral properties for L^+ , and therefore for L_β^+ .

Lemma 1. *The unbounded operator L^+ , defined in $L^2(\mathbb{R})$ with domain $H^2(\mathbb{R})$, is self-adjoint, has a unique negative eigenvalue $-\lambda_0 < 0$ (with corresponding L^2 -normalized eigenfunction Q^-) and its kernel is spanned by $\partial_x Q$. Moreover, the continuous spectrum is $[1, +\infty)$, and 0 is an isolated eigenvalue.*

We recall that from standard elliptic theory, Q^- is smooth, even and exponentially decreasing in space: there exists $c_0 > 0$ such that

$$\forall k \in \mathbb{N}, \forall x \in \mathbb{R}, \exists C_k, \quad |\partial_x^k Q^-(x)| \leq C_k e^{-c_0|x|}. \quad (10)$$

It is not difficult to check that one can take any c_0 satisfying $0 < c_0 \leq \sqrt{1 + \lambda_0}$.

Another consequence of Lemma 1 is the following fact: L_β^+ has a unique negative eigenvalue $-\lambda_0$ with (even) eigenfunction $Q_\beta^-(x) := Q^-(\gamma x)$, its kernel is spanned by $\partial_x Q_\beta$ and has continuous spectrum $[1, +\infty)$. Additionally, we have

Corollary 1. *There exists $\nu_0 \in (0, 1)$ such that, if $v \in H^1(\mathbb{R})$ satisfies $(v|Q_\beta^-) = (v|\partial_x Q_\beta) = 0$, then $(L_\beta^+v|v) \geq \nu_0 \|v\|_{H^1}^2$.*

We introduce now suitable matrix operators associated to the dynamics around a soliton. These operators will be dependent on the velocity parameter β , but for simplicity of notation, we will omit the subscript β when there is no ambiguity. Define²

$$T = T_\beta := -\partial_{xx} + \text{Id} - f'(Q_\beta) = L_\beta^+ - \beta^2 \partial_{xx}, \quad (11)$$

$$J := \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \quad (12)$$

$$L := \begin{pmatrix} T & 0 \\ 0 & \text{Id} \end{pmatrix}, \quad (13)$$

and

$$H := L - J \begin{pmatrix} \beta \partial_x & 0 \\ 0 & \beta \partial_x \end{pmatrix} = \begin{pmatrix} T & -\beta \partial_x \\ \beta \partial_x & \text{Id} \end{pmatrix}. \quad (14)$$

The operator H is the standard second order derivative of the functional for which the vector soliton $R = (Q_\beta, \partial_t Q_\beta)^T$ is an associated local minimizer. Later we will discuss in detail this assertion. The following Proposition describes the main spectral properties of H . Recall that $\langle \cdot | \cdot \rangle$ and $(\cdot | \cdot)$ denote the symmetric bilinear forms on $H^1(\mathbb{R}) \times L^2(\mathbb{R})$ and $L^2(\mathbb{R})$ respectively, introduced in (7), and $\| \cdot \|$ is the energy norm defined in (8).

Proposition 1. *Let $\beta \in \mathbb{R}$, $|\beta| < 1$. The matrix operator H , defined in $L^2(\mathbb{R}) \times L^2(\mathbb{R})$ with dense domain $H^2(\mathbb{R}) \times H^1(\mathbb{R})$, is a self-adjoint operator. Furthermore, there exist $\alpha_0 > 0$, $\Phi_0, \Phi_- \in \mathcal{S}(\mathbb{R})^2$ (with exponential decay, along with their derivatives) such that*

$$H\Phi_0 = 0, \quad \langle \Phi_0 | \Phi_- \rangle = 0, \quad (15)$$

$$\langle H\Phi_- | \Phi_- \rangle < 0, \quad (16)$$

and the following coercivity property holds. Let $V = (v_1, v_2)^T \in H^1(\mathbb{R}) \times L^2(\mathbb{R})$. Then,

$$\text{if } \langle V | \Phi_0 \rangle = \langle V | H\Phi_- \rangle = 0 \text{ one has } \langle HV | V \rangle \geq \alpha_0 \|V\|^2. \quad (17)$$

A stronger version of this result was stated by Grillakis, Shatah and Strauss in [10, Lemma 6.2], but the proof given there contained a gap, as noted in the errata at the end of [11, page 347]. As a replacement, the Proposition above (weaker than the original Grillakis-Shatah-Strauss result, but adequate for our purposes) was proposed in the errata [11], without proof. We have not found a clear definition and meaning of the function Φ_- in [11], so therefore, for the convenience of the reader, we write the details of the proof in the following lines.

Proof of Proposition 1. It is easy to check that H is indeed a self-adjoint operator. On the other hand, let $V = (v_1, v_2)^T$. We have from (14),

$$\begin{aligned} \langle HV | V \rangle &= \left\langle \begin{pmatrix} Tv_1 - \beta \partial_x v_2 \\ \beta \partial_x v_1 + v_2 \end{pmatrix} \middle| \begin{pmatrix} v_1 \\ v_2 \end{pmatrix} \right\rangle \\ &= (Tv_1 | v_1) - \beta (\partial_x v_2 | v_1) + \beta (\partial_x v_1 | v_2) + (v_2 | v_2) \\ &= (L_\beta^+ v_1 | v_1) + \beta^2 (\partial_x v_1 | \partial_x v_1) + 2\beta (v_2 | \partial_x v_1) + (v_2 | v_2) \\ &= (L_\beta^+ v_1 | v_1) + (\beta \partial_x v_1 + v_2 | \beta \partial_x v_1 + v_2). \end{aligned} \quad (18)$$

²Do not confuse with the transpose symbol $(\cdot)^T$.

Recalling the notation of Corollary 1, we define

$$\Phi_0 := \begin{pmatrix} \partial_x Q_\beta \\ -\beta \partial_{xx} Q_\beta \end{pmatrix}, \quad \Phi_- := \begin{pmatrix} Q_\beta^- \\ -\beta \partial_x Q_\beta^- \end{pmatrix}, \quad (19)$$

One can check from (18) that $\langle \Phi_0 | \Phi_0 \rangle \neq 0$ and $\langle H\Phi_0 | \Phi_0 \rangle = 0$, since $L_\beta^+ \partial_x Q_\beta = 0$. Note additionally that by parity $\langle \Phi_- | \Phi_0 \rangle = 0$. Therefore, (15) is directly satisfied. Also notice that

$$H\Phi_- = -\lambda_0 \begin{pmatrix} Q_\beta^- \\ 0 \end{pmatrix}. \quad (20)$$

We now prove (17). Let $V = (v_1, v_2)^T \in H^1(\mathbb{R}) \times L^2(\mathbb{R})$ be satisfying the orthogonality properties

$$\langle V | \Phi_0 \rangle = \langle V | H\Phi_- \rangle = 0.$$

Let us decompose v_1 in terms of the nonpositive spectral elements of L_β^+ , and L^2 -orthogonally:

$$v_1 = aQ_\beta^- + b\partial_x Q_\beta + q, \quad (q | Q_\beta^-) = (q | \partial_x Q_\beta) = 0.$$

From the orthogonality conditions in (17), we have

$$\left\langle \begin{pmatrix} v_1 \\ v_2 \end{pmatrix} \middle| \begin{pmatrix} Q_\beta^- \\ 0 \end{pmatrix} \right\rangle = 0,$$

so that $a = 0$, and hence from Corollary 1,

$$\langle HV | V \rangle = (L_\beta^+ q | q) + (\beta \partial_x v_1 + v_2 | \beta \partial_x v_1 + v_2) \geq \nu_0 \|q\|_{H^1}^2 \geq 0. \quad (21)$$

We now argue by contradiction. Assume that there exists a normalized sequence $V^n = (v_1^n, v_2^n)^T \in H^1(\mathbb{R}) \times L^2(\mathbb{R})$ that satisfies the orthogonality properties

$$\langle V^n | \Phi_0 \rangle = \langle V^n | H\Phi_- \rangle = 0, \quad \|V^n\|^2 = 1, \quad \text{and such that } \langle HV^n | V^n \rangle \rightarrow 0. \quad (22)$$

Let us write the L^2 -orthogonal decomposition for each v_1^n :

$$v_1^n = b_n \partial_x Q_\beta + q_n, \quad (q_n | \partial_x Q_\beta) = 0.$$

Then in view of (21) and (22) applied this time to the sequence V^n , $q_n \rightarrow 0$ in H^1 and $\beta \partial_x v_1^n + v_2^n \rightarrow 0$ in L^2 . Now we compute

$$\begin{aligned} 0 &= \langle V^n | \Phi_0 \rangle = \int (v_1^n \partial_x Q_\beta - v_2^n \beta \partial_{xx} Q_\beta) \\ &= \int v_1^n \partial_x Q_\beta + \beta \int (\beta \partial_x v_1^n + o_{L^2}(1)) \partial_{xx} Q_\beta \\ &= b_n \|\partial_x Q_\beta\|_{L^2}^2 + \beta^2 \int (b_n \partial_{xx} Q_\beta + \partial_x q_n) \partial_{xx} Q_\beta + o(1) \\ &= b_n (\|\partial_x Q_\beta\|_{L^2}^2 + \beta^2 \|\partial_{xx} Q_\beta\|_{L^2}^2) - \beta^2 \int q_n \partial_{xxx} Q_\beta + o(1) \end{aligned}$$

Now $q_n \rightarrow 0$ in L^2 , so that $(q_n | \partial_{xxx} Q_\beta) \rightarrow 0$, and hence $b_n \rightarrow 0$ as $n \rightarrow +\infty$. But in this case, $v_1^n = b_n \partial_x Q_\beta + q_n \rightarrow 0$ in H^1 and $v_2^n = \beta \partial_x v_1^n + o_{L^2}(1) \rightarrow 0$ in L^2 . Hence $\|V^n\|^2 = \|v_1^n\|_{H^1}^2 + \|v_2^n\|_{L^2}^2 \rightarrow 0$, a contradiction to (22).

It remains to show that $\langle H\Phi_- | \Phi_- \rangle < 0$, namely (16). Indeed,

$$\langle H\Phi_- | \Phi_- \rangle = -\lambda_0 \left\langle \begin{pmatrix} Q_\beta^- \\ 0 \end{pmatrix} \middle| \begin{pmatrix} Q_\beta^- \\ \partial_x Q_\beta^- \end{pmatrix} \right\rangle = -\lambda_0 \|Q_\beta^-\|_{L^2}^2 < 0.$$

□

2.2. Eigenfunctions of the linearized flow and Hessian. It is still unclear whether or not the coercivity property (17) – a key point in the proof of any stability result – is useful for us, since solitons are actually unstable. It turns out that for our purposes, we need a different version of Proposition 1, for the linearized operator of the **flow around Q** . In order to state such a result, we introduce some additional notation.

Let $\beta \in \mathbb{R}$, $|\beta| < 1$ be a Lorentz parameter, and consider the operators T , J , L and H defined in (11)-(14). Let

$$\mathcal{L} = \mathcal{L}(\beta) = JL = \begin{pmatrix} 0 & \text{Id} \\ -T & 0 \end{pmatrix}, \quad (23)$$

and

$$\mathcal{H} = \begin{pmatrix} -\beta\partial_x & -T \\ \text{Id} & -\beta\partial_x \end{pmatrix} = -HJ. \quad (24)$$

Concerning this last operator, we prove the following result.

Lemma 2. *Let $\beta \in \mathbb{R}$, $|\beta| < 1$, $\gamma = (1 - \beta^2)^{-1/2}$ and λ_0 from Lemma 1. There are functions $Z_0 = Z_{0,\beta}$, and $Z_{\pm} = Z_{\pm,\beta}$, with components exponentially decreasing in space, satisfying the spectral equations*

$$\mathcal{H}Z_0 = 0, \quad \text{and} \quad \mathcal{H}Z_{\pm} = \pm \frac{\sqrt{\lambda_0}}{\gamma} Z_{\pm}. \quad (25)$$

Moreover, by the nondegeneracy of the kernel spanned by Φ_0 , we can assume $\Phi_0 = JZ_0$.

Proof. The proof is similar to that of [10]. In particular, we obtain explicit expressions for Z_0 and Z_{\pm} in the following lines.

The eigenvalue problem $\mathcal{H}Z = \lambda Z$ reads now, with $Z(x) = (Z_1(\gamma x), Z_2(\gamma x))^T$,

$$TZ_2 + \beta(Z_1)_x + \lambda Z_1 = 0, \quad Z_1 - \beta(Z_2)_x - \lambda Z_2 = 0, \quad (26)$$

Replacing Z_1 in the first equation above, we get in the variable $s = \gamma x$ (recall that $Q_{\beta}(x) = Q(\gamma x)$),

$$-\gamma^2 Z_2'' + Z_2 - f'(Q)Z_2 + \beta\gamma(\lambda Z_2' + \beta\gamma Z_2'') + \lambda(\beta\gamma Z_2' + \lambda Z_2) = 0,$$

namely

$$-Z_2'' + Z_2 - f'(Q)Z_2 + 2\beta\gamma\lambda Z_2' = -\lambda^2 Z_2. \quad (27)$$

Performing the transformation $Z_2(s) := \tilde{Z}_2(s)e^{\beta\gamma\lambda s}$, where $s \in \mathbb{R}$, we get

$$-\tilde{Z}_2'' + \tilde{Z}_2 - f'(Q)\tilde{Z}_2 = -(\beta^2\gamma^2 + 1)\lambda^2 \tilde{Z}_2 = -\lambda^2\gamma^2 \tilde{Z}_2.$$

Therefore, by virtue of Lemma 1 we can take $\tilde{Z}_2 = Q^-(s)$ and $\lambda_{\pm}\gamma = \pm\sqrt{\lambda_0}$, where $-\lambda_0 < 0$ is the first eigenvalue of the standard Schrödinger operator L^+ , defined in (9). Thus,

$$Z_{\pm,2}(s) = Q^-(s)e^{\pm\beta\sqrt{\lambda_0}s}.$$

Note that from (10), $Z_{\pm,2}$ decreases exponentially at both sides of the origin, since $|\beta| < 1$ and $\beta\sqrt{\lambda_0} - \sqrt{1 + \lambda_0} < 0$.

From (26), we have

$$\begin{aligned} Z_{\pm,1}(s) &= \beta\gamma Z_{\pm,2}'(s) + \lambda_{\pm} Z_{\pm,2}(s) \\ &= \left[\beta\gamma(Q^-)_s \pm \beta^2\gamma\sqrt{\lambda_0}Q^- \pm \frac{\sqrt{\lambda_0}}{\gamma}Q^- \right] e^{\pm\beta\sqrt{\lambda_0}s} \end{aligned}$$

$$= \gamma[\beta(Q^-)_s \pm \sqrt{\lambda_0}Q^-]e^{\pm\beta\sqrt{\lambda_0}s}.$$

By the same reasons as above, $Z_{\pm,1}$ is an exponentially decreasing function. From these identities, we have

$$\begin{aligned} Z_{\pm}(x) &= \begin{pmatrix} \gamma\beta(Q^-)_s(\gamma x) \pm \gamma\sqrt{\lambda_0}Q^-(\gamma x) \\ Q^-(\gamma x) \end{pmatrix} e^{\pm\beta\sqrt{\lambda_0}\gamma x} \\ &= \begin{pmatrix} \beta(Q^-)_x(x) \pm \gamma\sqrt{\lambda_0}Q^-_x(x) \\ Q^-_x(x) \end{pmatrix} e^{\pm\beta\sqrt{\lambda_0}\gamma x}. \end{aligned} \quad (28)$$

Now, we consider the computation of Z_0 . Replacing $\lambda = 0$ in (27), we can choose

$$Z_{0,2}(s) = Q'(s), \quad \text{and} \quad Z_{0,1} = \beta\gamma Q''(s),$$

from which we get

$$Z_0(x) = \gamma \begin{pmatrix} Z_{0,1}(x) \\ Z_{0,2}(x) \end{pmatrix} = \gamma \begin{pmatrix} \beta\gamma Q''(\gamma x) \\ Q'(\gamma x) \end{pmatrix} = \begin{pmatrix} \beta Q''_{\beta}(x) \\ Q'_{\beta}(x) \end{pmatrix}. \quad (29)$$

It is clear that $\mathcal{H}Z_0 = 0$. Similarly, we have $\mathcal{H}Z_{\pm} = \pm \frac{\sqrt{\lambda_0}}{\gamma}Z_{\pm}$, which proves (25). \square

In order to prove Proposition 2, we need to prove the existence of two additional functions, both associated to Z_{\pm} .

Lemma 3. *There exist unique functions Y_{\pm} , with components exponentially decreasing in space, such that*

$$HY_{\pm} = Z_{\pm}, \quad \langle \Phi_0 | Y_{\pm} \rangle = 0.$$

Moreover, Y_{\pm} satisfy the additional orthogonality conditions $\langle Y_{\pm} | HY_{\pm} \rangle = 0$.

Proof. Let us prove the existence of Y_{\pm} . It is well-known that a necessary and sufficient condition for existence is the following condition: it suffices to check that Z_{\pm} are orthogonal to Φ_0 , the generator of the kernel of H . Indeed, we have from (25), (24), the self-adjointness of H and Proposition 1,

$$\langle \Phi_0 | Z_{\pm} \rangle = \pm \frac{\gamma}{\sqrt{\lambda_0}} \langle \Phi_0 | \mathcal{H}Z_{\pm} \rangle = \mp \frac{\gamma}{\sqrt{\lambda_0}} \langle \Phi_0 | HJZ_{\pm} \rangle = 0.$$

However, we need some additional estimates on Y_{\pm} . In what follows, we write down explicitly the equation $HY_{\pm} = Z_{\pm}$. It is not difficult to check that $Y_{\pm} = (Y_{\pm,1}, Y_{\pm,2})^T$ satisfies the equations

$$TY_{\pm,1} - \beta(Y_{\pm,2})_x = Z_{\pm,1}, \quad \beta(Y_{\pm,1})_x + Y_{\pm,2} = Z_{\pm,2}.$$

Replacing the second equation in the first one, we get (cf. (9))

$$L_{\beta}^+ Y_{\pm,1} = \beta(Z_{\pm,2})_x + Z_{\pm,1}.$$

Note that $(\beta(Z_{\pm,2})_x + Z_{\pm,1} | \partial_x Q_{\beta}) = 0$. Therefore, $Y_{\pm,1}$ exists and it is exponentially decreasing, with the same rate as $Z_{\pm,1}$ and $Z_{\pm,2}$. A similar conclusion follows for $Y_{\pm,2}$.

Since Y_{\pm} is unique modulo the addition of a constant times Φ_0 , it is clear that we can choose Y_{\pm} such that $\langle \Phi_0 | Y_{\pm} \rangle = 0$. On the other hand, from Lemma 2,

$$\langle Y_{\pm} | HY_{\pm} \rangle = \langle Y_{\pm} | Z_{\pm} \rangle = \pm \frac{\gamma}{\sqrt{\lambda_0}} \langle Y_{\pm} | \mathcal{H}Z_{\pm} \rangle = \mp \frac{\gamma}{\sqrt{\lambda_0}} \langle HY_{\pm} | JZ_{\pm} \rangle$$

$$= \mp \frac{\gamma}{\sqrt{\lambda_0}} \langle Z_{\pm} | JZ_{\pm} \rangle = 0.$$

□

The main result of this section is the following alternative to Proposition 1.

Proposition 2. *There exists $\mu_0 > 0$ such that the following holds. Let $V \in H^1 \times L^2$ such that $\langle \Phi_0 | V \rangle = 0$. Then*

$$\langle HV | V \rangle \geq \mu_0 \|V\|^2 - \frac{1}{\mu_0} \left[\langle Z_+ | V \rangle^2 + \langle Z_- | V \rangle^2 \right].$$

Proof. It is enough to prove that $\langle \Phi_0 | V \rangle = \langle Z_+ | V \rangle = \langle Z_- | V \rangle = 0$ imply

$$\langle HV | V \rangle \geq \mu_0 \|V\|^2,$$

for some $\mu_0 > 0$, independently of V . In order to prove this assertion, we first assume $\beta \neq 0$ and decompose orthogonally V and Y_{\pm} (cf. the previous Lemma) as follows

$$V = \tilde{V} + \alpha_- \Phi_- + \alpha_0 \Phi_0, \quad Y_{\pm} = \tilde{Y}_{\pm} + \delta_0 \Phi_0 + \delta_{\pm} \Phi_-, \quad (30)$$

with

$$\langle \tilde{V} | \Phi_0 \rangle = \langle \tilde{Y}_{\pm} | \Phi_0 \rangle = \langle \tilde{V} | H\Phi_- \rangle = \langle \tilde{Y}_{\pm} | H\Phi_- \rangle = 0. \quad (31)$$

Since $\langle \Phi_0 | \Phi_- \rangle = \langle \Phi_0 | V \rangle = \langle \Phi_0 | Y_{\pm} \rangle = 0$ and $\langle \Phi_- | H\Phi_- \rangle < 0$, it is clear that $\alpha_0 = \delta_0 = 0$ and α_-, δ_{\pm} are well-defined. Moreover,

Claim. For all $\beta \in (-1, 1) \setminus \{0\}$, \tilde{Y}_+ and \tilde{Y}_- are linearly independent as $L^2(\mathbb{R})^2$ vector-valued functions with real coefficients.

Indeed, to see this, assume that there is $\tilde{\lambda} \neq 0$ such that $\tilde{Y}_+ = \tilde{\lambda} \tilde{Y}_-$. Then, from the previous decomposition and Lemma 3,

$$Z_+ - \tilde{\lambda} Z_- = H(Y_+ - \tilde{\lambda} Y_-) = (\delta_+ - \tilde{\lambda} \delta_-) H\Phi_-. \quad (32)$$

This identity contradicts (28) and (19), which establish that Z_+ and Z_- have essentially different rates of decay at infinity, different to that of $H\Phi_-$, for all $\beta \neq 0$, which makes (32) impossible.

The analysis is now similar to that in [7, Lemma 5.2]. We have from (30),

$$\langle HV | V \rangle = \langle H\tilde{V} + \alpha_- H\Phi_- | \tilde{V} + \alpha_- \Phi_- \rangle = \langle H\tilde{V} | \tilde{V} \rangle + \alpha_-^2 \langle H\Phi_- | \Phi_- \rangle. \quad (33)$$

On the other hand, since $\langle Z_{\pm} | V \rangle = 0$, we have from Lemma 3,

$$0 = \langle Y_{\pm} | HV \rangle = \langle \tilde{Y}_{\pm} + \delta_{\pm} \Phi_- | H\tilde{V} + \alpha_- H\Phi_- \rangle = \langle \tilde{Y}_{\pm} | H\tilde{V} \rangle + \alpha_- \delta_{\pm} \langle H\Phi_- | \Phi_- \rangle.$$

Similarly,

$$0 = \langle HY_{\pm} | Y_{\pm} \rangle = \langle H\tilde{Y}_{\pm} | \tilde{Y}_{\pm} \rangle + \delta_{\pm}^2 \langle H\Phi_- | \Phi_- \rangle.$$

We get then

$$\langle HV | V \rangle = \langle H\tilde{V} | \tilde{V} \rangle - \frac{\langle \tilde{Y}_- | H\tilde{V} \rangle \langle \tilde{Y}_+ | H\tilde{V} \rangle}{\sqrt{\langle H\tilde{Y}_+ | \tilde{Y}_+ \rangle \langle H\tilde{Y}_- | \tilde{Y}_- \rangle}}. \quad (34)$$

Consider

$$a := \sup_{W \in \text{Span}(\tilde{Y}_+, \tilde{Y}_-) \setminus \{0\}} \left| \frac{\langle \tilde{Y}_+ | H\tilde{W} \rangle}{\sqrt{\langle H\tilde{Y}_+ | \tilde{Y}_+ \rangle \langle HW | W \rangle}} \cdot \frac{\langle \tilde{Y}_- | H\tilde{W} \rangle}{\sqrt{\langle H\tilde{Y}_- | \tilde{Y}_- \rangle \langle HW | W \rangle}} \right|.$$

Recall $\langle H \cdot | \cdot \rangle$ is positive definite on $\text{Span}(\Phi_0, \Phi_-)^\perp$. Hence apply Cauchy Schwarz's inequality to both terms of the product: it transpires that $a \leq 1$. Furthermore, if $a = 1$ (as $\text{Span}(\tilde{Y}_+, \tilde{Y}_-)$ is finite dimensional), there exists W of norm 1 such that both terms are in the equality case in the Cauchy-Schwarz inequality, i.e W and \tilde{Y}_+ are linearly dependent, and W and \tilde{Y}_- are also linearly dependent. But it would then follow that \tilde{Y}_+ and \tilde{Y}_- are linearly dependent, a contradiction the above claim. This proves $a < 1$.

Now using H -orthogonal decomposition on $\text{Span}(\Phi_0, \Phi_-)^\perp$, we deduce that

$$\forall W \in \text{Span}(\Phi_0, \Phi_-)^\perp, \quad \left| \frac{\langle \tilde{Y}_- | HW \rangle \langle \tilde{Y}_+ | HW \rangle}{\sqrt{\langle H\tilde{Y}_+ | \tilde{Y}_+ \rangle \langle H\tilde{Y}_- | \tilde{Y}_- \rangle}} \right| \leq a \langle HW | W \rangle.$$

By (34), (31) and (17), we get

$$\langle HV | V \rangle \geq (1 - a) \langle H\tilde{V} | \tilde{V} \rangle \geq \alpha_0 (1 - a) \|\tilde{V}\|^2 \geq 0.$$

and so (33) implies $\langle H\tilde{V} | \tilde{V} \rangle \geq \alpha_-^2 |\langle H\Phi_- | \Phi_- \rangle|$.

We then conclude that, for $C = \frac{4}{(1-a)} \max\left(\frac{1}{\alpha_0}, \frac{\|\Phi_-\|^2}{|\langle H\Phi_- | \Phi_- \rangle|}\right)$,

$$\begin{aligned} C \langle HV | V \rangle &\geq C(1 - a) \langle H\tilde{V} | \tilde{V} \rangle \\ &\geq \frac{C(1 - a)}{2} (\langle H\tilde{V} | \tilde{V} \rangle + \alpha_-^2 |\langle H\Phi_- | \Phi_- \rangle|) \\ &\geq 2\|\tilde{V}\|^2 + 2\alpha_-^2 \|\Phi_-\|^2 \geq \|\tilde{V} + \alpha_- \Phi_-\|^2 = \|V\|^2. \end{aligned}$$

Finally, if $\beta = 0$, we proceed as follows. First of all, we have from (28) and (19),

$$Z_\pm = Q^- \begin{pmatrix} \pm\sqrt{\lambda_0} \\ 1 \end{pmatrix}, \quad \Phi_0 = Q' \begin{pmatrix} 1 \\ 0 \end{pmatrix},$$

so that $\langle \Phi_0 | V \rangle = \langle Z_\pm | V \rangle = 0$ imply $(v_1 | Q') = (v_1 | Q^-) = (v_2 | Q^-) = 0$, where $V = (v_1, v_2)^T$. Therefore,

$$\langle HV | V \rangle = (L^+ v_1 | v_1) + (v_2 | v_2) \geq \nu_0 \|V\|^2.$$

□

2.3. Extension to higher dimensions. The equivalent of Lemma 1 (and therefore assumption (iv) of the Introduction) in dimension $d \geq 2$ has the form

Lemma 4. *Assume $d \geq 2$ and assumption (B) holds. L^+ has exactly one negative eigenvalue, and its kernel is spanned by $(\partial_{x_i} Q)_{i=1, \dots, d}$. Its continuous spectrum is $[1, +\infty)$.*

Proof. See Maris [15] and McLeod [17]. □

As mentioned in the Introduction, this result is open for general nonlinearity f . In that case, we need to assume that it holds, i.e. assumption (v).

The null directions for H are now the d -dimensional vector space spanned by the functions $\Phi_{0,i} = \begin{pmatrix} \partial_i Q \\ \beta \cdot \nabla \partial_i Q \end{pmatrix}$. In the proof of Lemma 2, one should rather perform the transformation $\tilde{Z}_2 = Z_2 e^{-\gamma \lambda \beta \cdot x}$. The rest of the arguments is dimension insensitive.

3. CONSTRUCTION OF APPROXIMATE N -SOLITONS

In this section we prove Theorem 1. Again, we will give a detailed proof in the one dimensional case $d = 1$, and point out how to extend the proof in higher dimension, which is done in a similar fashion as in [16].

3.1. The topological argument. We continue with the same notation as in the previous section. In particular, we fix $\beta \in (-1, 1)$ and consider now the time-dependent, boosted soliton given by

$$Q_\beta(t, x) = Q(\gamma(x - \beta t)), \quad \gamma = (1 - \beta^2)^{-1/2}.$$

Additionally, we suppose given N different velocities $\beta_1, \dots, \beta_N \in (-1, 1)$, already arranged in such a way that

$$-1 < \beta_1 < \beta_2 < \dots < \beta_N < 1, \tag{35}$$

and N translation parameters $x_1, \dots, x_N \in \mathbb{R}$, such that $Q_{\beta_j}(t, x - x_j)$ is the associated soliton solution of velocity β_j and shift x_j , $j = 1, \dots, N$.

Finally, we introduce some notation. Given \mathcal{B} a real Banach space, $x \in B$ and $r \geq 0$, we denote

$$B_{\mathcal{B}}(x, r) = \{y \in \mathcal{B} \mid \|x - y\|_{\mathcal{B}} \leq r\}$$

the closed ball in \mathcal{B} centered at x of radius r and $\|\cdot\|_{\mathcal{B}}$ is the associated Banach norm on \mathcal{B} .

Lemma 5 (Modulation). *There exist $L_0 > 0$ and $\varepsilon_0 > 0$ such that the following holds for some $C > 0$. For any $L \geq L_0$ and $0 < \varepsilon < \varepsilon_0$, $t_0 \in \mathbb{R}$, if $U \in H^1(\mathbb{R}) \times L^2(\mathbb{R})$ is sufficiently near a sum of solitons whose centers are sufficiently far apart,*

$$\left\| U - \sum_{j=1}^N \begin{pmatrix} Q_{\beta_j} \\ \partial_t Q_{\beta_j} \end{pmatrix} (t_0, \cdot - y_j) \right\| \leq \varepsilon, \quad \min \{|y_j - y_i| \mid i \neq j\} \geq L,$$

then there exist shifts $\tilde{y}_j = \tilde{y}_j(\beta_j, t_0)$ such that if we define

$$\tilde{R}_j(x) := \begin{pmatrix} Q_{\beta_j} \\ \partial_t Q_{\beta_j} \end{pmatrix} (0, x - \tilde{y}_j), \tag{36}$$

$$\tilde{R}(x) := \sum_{j=1}^N \tilde{R}_j(x), \tag{37}$$

$$V(x) := U(x) - \tilde{R}(x), \tag{38}$$

then

$$\|V\| \leq C\varepsilon, \quad \text{and} \quad \langle V | (\tilde{R}_j)_x \rangle = 0. \tag{39}$$

Also, the map $U \mapsto (V, (\tilde{y}_j)_j)$ is a \mathcal{C}^1 -diffeomorphism around $\sum_{j=1}^N \begin{pmatrix} Q_{\beta_j} \\ \partial_t Q_{\beta_j} \end{pmatrix} (t_0, x - y_j)$. In such case, we say that U can be modulated into $(V, (\tilde{y}_j)_j)$.

Proof. This is the classical modulation result, stated as in [3, Lemma 2]. See [30, 31] for more details. \square

In what follows, we introduce additional notation. We assume that U can be modulated into $(V, (\tilde{y}_j)_j)$.

For any $j = 1, \dots, N$ (cf. Proposition 1 and Lemma 2 for the definitions), let

$$\begin{cases} Z_{\pm, j}(s) := Z_{\pm}(\gamma_j(s - \tilde{y}_j)), & Z_{0, \beta_j}(s) := Z_0(\gamma_j(s - \tilde{y}_j)), \\ \Phi_{0, j}(s) := \Phi_0(\gamma_j(s - \tilde{y}_j)), & \Phi_{-, j}(s) := \Phi_{-}(\gamma_j(s - \tilde{y}_j)), \end{cases} \quad (40)$$

where $\gamma_j := (1 - \beta_j^2)^{-1/2}$, and

$$a_{\pm, j} := \langle V | Z_{\mp, j} \rangle, \quad (41)$$

along with the vectors

$$\mathbf{a}_+ = (a_{+, j})_j, \quad \mathbf{a}_- = (a_{-, j})_j, \quad \text{and} \quad \tilde{\mathbf{y}} = (\tilde{y}_j)_j. \quad (42)$$

Finally, we fix a constant γ_0 given by

$$\begin{aligned} \gamma_0 := \min \left\{ \frac{1}{4} \sqrt{\lambda_0} \min \left\{ \frac{1}{\gamma_1}, \frac{1}{\gamma_2}, \dots, \frac{1}{\gamma_N} \right\}, \right. \\ \left. \frac{1}{4} \min \{ \gamma_1, \gamma_2, \dots, \gamma_N \} \min \{ \beta_1, \beta_2 - \beta_1, \dots, \beta_N - \beta_{N-1} \} \right\} > 0. \end{aligned} \quad (43)$$

Assume now $\varepsilon \in (0, \varepsilon_0)$ and $L \geq L_0$, where ε_0 and L_0 are obtained in by Lemma 5. Given $t \in \mathbb{R}$, let us consider the *centers*

$$y_j = y_j(t) := \beta_j t + x_j, \quad j = 1, \dots, N,$$

where the velocities β_j and the shifts x_j are given by (35). It is clear that there exists $T_0 \in \mathbb{R}$ such that, for all $t \geq T_0$, the y_j satisfy

$$\min \{ |y_j - y_i| \mid i \neq j \} \geq L.$$

From now on, we fix $t \geq T_0$. Consider the corresponding sum of solitons $R(t, x)$ associated to these parameters, namely

$$R(t, x) := \sum_{j=1}^N R_j(t, x) = \sum_{j=1}^N \left(\begin{array}{c} Q_{\beta_j} \\ \partial_t Q_{\beta_j} \end{array} \right) (0, x - y_j). \quad (44)$$

Then, according to Lemma 5, if $U \in H^1(\mathbb{R}) \times L^2(\mathbb{R})$ satisfies $\|U - R(t)\| \leq \varepsilon$, then U can be modulated. Moreover, up to increasing T_0 , we can assume that $e^{-\gamma_0 T_0} < \varepsilon_0$. Thus we can define our shrinking set.

Definition 1 (Shrinking set $\mathcal{V}(t)$). For $t \geq T_0$, we define the set

$$\mathcal{V}(t) \subset B_{H^1 \times L^2}(R(t), \varepsilon_0)$$

in the following way: $U \in \mathcal{V}(t)$ if and only if U can be modulated into $(V, \tilde{\mathbf{y}})$ where (cf. (36) and (42))

$$V = U - \sum_{j=1}^N \tilde{R}_j(t),$$

with

$$\|V\| \leq e^{-\gamma_0 t}, \quad |\tilde{y}_j(t) - \beta_j t - x_j| \leq e^{-\gamma_0 t}, \quad (45)$$

$$\|\mathbf{a}_+\|_{\ell^2} \leq e^{-3\gamma_0 t/2}, \quad \|\mathbf{a}_-\|_{\ell^2} \leq e^{-3\gamma_0 t/2}. \quad (46)$$

Definition 2. We denote by $\varphi = (u, \partial_t u)^T$ the flow of the (NLKG) equation, that is, given $S_0 \in \mathbb{R}$ and $U_0 \in H^1(\mathbb{R}) \times L^2(\mathbb{R})$,

$$t \mapsto \varphi(S_0, t, U_0) \quad (47)$$

is the solution to (NLKG) with initial data U_0 at time S_0 (with values in $H^1 \times L^2$).

In most of what we do, we will have $t \leq S_0$ so that U_0 can be thought of as a final data, and we work backwards in time. The key result of this section is the following construction of an approximate N -soliton.

Proposition 3 (Approximate N -soliton). *There exist $T_0 > 0$ such that the following holds. For any $S_0 \geq T_0$, there exist a final data U_0 such that*

$$\forall t \in [T_0, S_0], \quad \varphi(S_0, t, U_0) \in \mathcal{V}(t).$$

At this point, the solution $\phi(S_0, t, U_0)$ depends on S_0 . To prove Theorem 1, we will need to derive such a solution independent of S_0 , which we will do via a compactness argument in the next (and last) section 4.

Our goal is now to prove Proposition 3.

Fix $S_0 \geq T_0$. Consider an initial data U_0 at time S_0 such that $U_0 \in \mathcal{V}(S_0)$. Due to the blow-up criterion for (NLKG), and the fact that $R(t)$ defined in (44) is bounded in $H^1(\mathbb{R}) \times L^2(\mathbb{R})$, we have that $\varphi(S_0, t, U_0)$ is defined at least as long as it belongs to $B_{H^1 \times L^2}(R(t), 1)$. In particular, $\varphi(S_0, t, U_0)$ does not blow-up as long as it belongs to $\mathcal{V}(t)$, and we can define the (backward) exit time

$$T^*(U_0) := \inf \{T \in [T_0, S_0] \mid \forall t \in [T, S_0], \varphi(S_0, t, U_0) \in \mathcal{V}(t)\}.$$

Notice that we could have $T^*(U_0) = S_0$. Our goal is to find $U_0 \in \mathcal{V}(T_0)$ such that $T^*(U_0) = T_0$.

In order to show such an assertion, we will only consider some very specific initial data, namely $U_0 \in \mathcal{V}(S_0)$ such that (see (42))

- $U_0 \in R(S_0) + \text{Span}((Z_{\pm, j})_{j=1, \dots, N})$,
- $\mathbf{a}_-(S_0) = 0$, and
- $\mathbf{a}_+(S_0) \in B_{\mathbb{R}^N}(0, e^{-3\gamma_0 S_0/2})$.

These conditions can be satisfied due to the almost orthogonality of $Z_{\pm, j}$, and this is the content of the following

Lemma 6 (Modulated final data). *Let $S_0 \geq T_0$ be large enough. There exist a \mathcal{C}^1 map $\Theta : B_{\mathbb{R}^N}(0, 1) \rightarrow \mathcal{V}(S_0)$ as follows. Given $\mathbf{a}_+ = (\mathbf{a}_{+, j})_j \in B_{\mathbb{R}^N}(0, 1)$, $U_0 := \Theta(\mathbf{a}_+) \in \mathcal{V}(S_0)$ such that U_0 can be modulated into $(V_0, \tilde{\mathbf{y}})$ and the associated parameters (42) satisfy*

$$\mathbf{a}_+(S_0) = e^{-3\gamma_0 T_0/2} \mathbf{a}_+, \quad \mathbf{a}_-(S_0) = 0. \quad (48)$$

Moreover,

$$\|V_0\| \leq C e^{-3\gamma_0 T_0/2}. \quad (49)$$

Proof. The main idea is to consider the map $B_{\mathbb{R}^{2N}}(0, 1) \rightarrow B_{\mathbb{R}^{2N}}(0, 1)$, $\mathbf{b}_{\pm} \mapsto \mathbf{a}_{\pm}$, where \mathbf{a}_{\pm} corresponds to the data $U_0 = R(S_0) + \sum_{\pm, j} \mathbf{b}_{\pm, j} Z_{\pm, j}$, and to invoke the implicit mapping theorem. We refer to [3, Lemma 3] and its proof in [3, Appendix A] for full details. \square

If $T^* := T^*(U_0) > T_0$, by maximality, we also have that for the function $\varphi(S_0, T^*, U_0)$, at least one of the inequalities in the definition of $\mathcal{V}(T^*)$ is actually an equality. It turns out that the equality is achieved by $\mathbf{a}_+(T_0)$ only, and that the rescaled quantity $e^{3\gamma_0 T^*/2} \mathbf{a}_+(t)$ is *transverse* to the sphere at $t = T^*$. This is at the heart of the proof and is the content of the following

Proposition 4. *Let $\mathbf{a}_+ \in B_{\mathbb{R}^N}(0, 1)$, and assume that its maximal exit time is (strictly) greater than T_0 :*

$$T^* = T^*(\Theta(\mathbf{a}_+)) > T_0.$$

Denote, for all $t \in [T^, S_0]$, the associated modulation $(V(t), \tilde{\mathbf{y}}(t))$ of $\varphi(t, T_0, \Theta(\mathbf{a}_+))$, defined in (47). Then, for all $t \in [T^*, S_0]$,*

$$\|V(t)\| \leq \frac{1}{2} e^{-\gamma_0 t}, \quad |\tilde{y}_j(t) - \beta_j t - x_j| \leq \frac{1}{2} e^{-\gamma_0 t}, \quad (50)$$

$$\|\mathbf{a}_-(t)\|_{\ell^2} \leq \frac{1}{2} e^{-3\gamma_0 t/2}, \quad (51)$$

and

$$\|\mathbf{a}_+(T^*)\|_{\ell^2} = e^{-3\gamma_0 T^*/2}. \quad (52)$$

Furthermore, $\mathbf{a}_+(T^*)$ is transverse to the sphere, i.e.,

$$\left. \frac{d}{dt} (e^{3\gamma_0 t} \|\mathbf{a}_+(t)\|_{\ell^2}^2) \right|_{t=T^*} < 0.$$

For the sake of continuity, we postpone the proof of Proposition 4 until the next paragraph, and conclude the proof of Proposition 3 here, assuming Proposition 4.

Let us state a few direct consequences of Proposition 4, (their proofs will also be done in the next paragraph 3.2).

Corollary 2. *We have the following properties.*

- (1) *The set of final data which give rise to solutions which exit strictly after T_0*

$$\Omega := \{\mathbf{a}_+ \in B_{\mathbb{R}^N}(0, 1) \mid T^*(\Theta(\mathbf{a}_+)) > T_0\}$$

is open (in $B_{\mathbb{R}^N}(0, 1)$).

- (2) *The map $\Omega \rightarrow \mathbb{R}$, $\mathbf{a}_+ \mapsto T^*(\Theta(\mathbf{a}_+)) \in \mathbb{R}$ is continuous (we emphasize that the data belong to Ω).*

- (3) *The exit is instantaneous on the sphere:*

$$\text{if } \|\mathbf{a}_+\|_{\ell^2} = 1, \quad \text{then } T^*(\mathbf{a}_+) = S_0. \quad (53)$$

We are now in a position to complete the proof of Theorem 1.

End of the proof of Proposition 3. We argue by contradiction. Assume that all possible $\mathbf{a}_+ \in B_{\mathbb{R}^N}(0, 1)$ give rise to initial data $U_0 = \Theta(\mathbf{a}_+) \in \mathcal{V}(S_0)$ and corresponding solutions $\varphi(S_0, t, U_0)$ that exit $\mathcal{V}(t)$ strictly after T_0 , i.e.

$$\text{assume that } \Omega = B_{\mathbb{R}^N}(0, 1). \quad (54)$$

Given $U_0 \in \mathcal{V}(T_0)$, we denote $\Phi(U_0)$ the rescaled projection of the *exit spot*

$$\Phi(U_0) = e^{3\gamma_0 T^*(U_0)/2} \mathbf{a}_+(T^*(U_0)),$$

so that $\Phi(U_0) \in B_{\mathbb{R}^N}(0, 1)$. Let us finally consider the rescaled projection of the exit spot Ψ , defined as follows:

$$\Psi : B_{\mathbb{R}^N}(0, 1) \rightarrow \mathbb{B}_{\mathbb{R}^N}(0, 1), \quad \mathbf{a}_+ \mapsto \Psi(\mathbf{a}_+) = \Phi \circ \Theta(\mathbf{a}_+).$$

Corollary 2 then translates into the following properties for Ψ :

- $\Psi : B_{\mathbb{R}^N}(0, 1) \rightarrow \mathbb{S}^{N-1}$ is continuous (like T^* , Φ and Θ);
- If $\|\mathbf{a}_+\|_{\ell^2} = 1$, $\Psi(\mathbf{a}_+) = \mathbf{a}_+$ (cf. (53) and (48)); i.e $\Psi|_{\mathbb{S}^{N-1}} = \text{Id}$.

These two affirmations contradict the Brouwer's Theorem. Hence our assumption (54) is wrong, and there exists \mathbf{a}^+ such that the solution $U(t) = \varphi(S_0, t, \Theta(\mathbf{a}_+))$ satisfies $T^+(\Theta(\mathbf{a}_+)) = T_0$. In particular $U(t) \in \mathcal{V}(t)$ for all $t \in [T_0, S_0]$, and $U_0 := U(S_0) = \Theta(\mathbf{a}_+)$ satisfies the conditions of Proposition 3. \square

3.2. Bootstrap estimates. This paragraph is devoted to the last remaining results needed to complete Proposition 3: Proposition 4 and Corollary 2

Proof of Proposition 4. Step 1. First, we introduce some notation. Consider the flow $\varphi(t) = \varphi(S_0, t, \Theta(\mathbf{a}_+))$ given by Proposition 4, and valid for all $t \in [T^*, S_0]$. From Lemma 5, we have

$$\varphi(t) = \tilde{R}(t) + V(t), \quad (55)$$

where

$$\tilde{R}(t, x) = \sum_{j=1}^N \tilde{R}_j(t, x), \quad \tilde{R}_j(t, x) = (Q_{\beta_j}, \partial_t Q_{\beta_j})^T (x - \tilde{y}_j(t)), \quad (56)$$

$$\tilde{y}_j(t) = \beta_j t + \tilde{x}_j(t), \quad (57)$$

and

$$V(t) = (v_1(t), v_2(t))^T.$$

Additionally, from the equation satisfied by φ , we have

$$\varphi_t = \begin{pmatrix} 0 & \text{Id} \\ \partial_x^2 - \text{Id} & 0 \end{pmatrix} \varphi + \begin{pmatrix} 0 \\ f(u) \end{pmatrix},$$

where $\varphi = (u, u_t)^T$. Replacing the decomposition (55), we have

$$V_t = \begin{pmatrix} 0 & \text{Id} \\ \partial_x^2 - \text{Id} + f'(Q_{\beta_j}) & 0 \end{pmatrix} V + \text{Rem}(t) = \mathcal{L}_j V + \text{Rem}(t), \quad (58)$$

with $\mathcal{L}_j := \mathcal{L}(\beta_j)$ defined in (23),

$$\begin{aligned} \text{Rem}(t) &:= \begin{pmatrix} 0 & \text{Id} \\ \partial_x^2 - \text{Id} & 0 \end{pmatrix} \tilde{R} - \tilde{R}_t + \begin{pmatrix} 0 \\ f(u) - f'(Q_{\beta_j})v_1 \end{pmatrix} \\ &= \sum_{k=1}^N \tilde{x}'_k(t) \partial_x \begin{pmatrix} Q_{\beta_k} \\ -\beta_k \partial_{xx} Q_{\beta_k} \end{pmatrix} \\ &\quad + \begin{pmatrix} 0 \\ f(\sum_{k=1}^N Q_{\beta_k} + v_1) - \sum_{k=1}^N f(Q_{\beta_k}) - f'(Q_{\beta_j})v_1 \end{pmatrix}. \end{aligned}$$

First of all, note that from (19) we have $\begin{pmatrix} \partial_x Q_{\beta_k} \\ -\beta_k \partial_{xx} Q_{\beta_k} \end{pmatrix} = \Phi_{0,k}$. If we take the scalar product of (58) with $(\tilde{R}_j)_x$, then the orthogonality (39) (coming from modulation) leads to the estimate

$$|\tilde{x}'_j(t)| \leq C(\|V(t)\| + e^{-3\gamma_0 t}), \quad (59)$$

valid for all $j = 1, \dots, N$. Indeed, we have

$$\langle (\tilde{R}_j)_x | V_t \rangle = \langle (\tilde{R}_j)_x | \mathcal{L}_j V \rangle + \langle (\tilde{R}_j)_x | \text{Re}(t) \rangle.$$

Note that from (57)

$$\langle (\tilde{R}_j)_x | V_t \rangle = -\langle (\tilde{R}_j)_{xt} | V \rangle = (\beta_j + \tilde{x}'_j(t)) \langle \partial_{xx} \tilde{R}_j | V \rangle.$$

Consequently

$$|\langle (\tilde{R}_j)_x | V_t \rangle| \leq C(1 + |\tilde{x}'_j(t)|) \|V(t)\|.$$

On the other hand,

$$\langle (\tilde{R}_j)_x | \mathcal{L}_j V \rangle = \left\langle \begin{pmatrix} \partial_x Q_{\beta_j} \\ \partial_{xt} Q_{\beta_j} \end{pmatrix} \middle| \begin{pmatrix} v_2 \\ -T_{\beta_j} v_1 \end{pmatrix} \right\rangle = \left\langle \begin{pmatrix} \partial_x Q_{\beta_j} \\ -T_{\beta_j} \partial_{xt} Q_{\beta_j} \end{pmatrix} \middle| \begin{pmatrix} v_2 \\ v_1 \end{pmatrix} \right\rangle,$$

so that

$$|\langle (\tilde{R}_j)_x | \mathcal{L}_j V \rangle| \leq C \|V(t)\|.$$

Finally, we deal with the term $\langle (\tilde{R}_j)_x | Rem(t) \rangle$. From the definition of $Rem(t)$ we have

$$\begin{aligned} \langle (\tilde{R}_j)_x | Rem(t) \rangle &= \sum_{k=1}^N \tilde{x}'_k(t) \langle (\tilde{R}_j)_x | \Phi_{0,k} \rangle \\ &\quad + \left(\partial_{xt} Q_{\beta_j} \middle| f \left(\sum_{k=1}^N Q_{\beta_k} + v_1 \right) - \sum_{k=1}^N f(Q_{\beta_k}) - f'(Q_{\beta_j}) v_1 \right). \end{aligned}$$

Since $(\tilde{R}_j)_x = \Phi_{0,j}$, we get

$$\langle (\tilde{R}_j)_x | \Phi_{0,j} \rangle = \|\Phi_{0,j}\|^2,$$

and if $k \neq j$,

$$|\langle (\tilde{R}_j)_x | \Phi_{0,k} \rangle| = |\langle \Phi_{0,j} | \Phi_{0,k} \rangle| \leq C e^{-3\gamma_0 t}.$$

Now if $x \in [m_j t, m_{j+1} t]$, then for all $p \neq j$ (see (43)),

$$|Q_{\beta_p}(t, x)| \leq C e^{-3\gamma_0 t}.$$

Therefore, inside this region (note that if $d \geq 2$ then f is a pure power nonlinearity)

$$\left\| f \left(\sum_{k=1}^N Q_{\beta_k} + v_1 \right) - \sum_{k=1}^N f(Q_{\beta_k}) - f'(Q_{\beta_j}) v_1 \right\|_{L^2} \leq C e^{-3\gamma_0 t} + C \|V(t)\|^2.$$

On the other hand, if $x \notin [m_j t, m_{j+1} t]$

$$|\partial_{xt} Q_{\beta_j}| \leq C e^{-3\gamma_0 t}.$$

In conclusion, we have

$$\left(\partial_{xt} Q_{\beta_j} \middle| f \left(\sum_{k=1}^N Q_{\beta_k} + v_1 \right) - \sum_{k=1}^N f(Q_{\beta_k}) - f'(Q_{\beta_j}) v_1 \right) \leq C e^{-3\gamma_0 t} + C \|V(t)\|^2. \quad (60)$$

Collecting the preceding estimates we get (59).

Step 2. Control of degenerate directions. The next step of the proof is to consider the dynamics of the associated scalar products $a_{\pm,j}(t)$ and $a_{0,j}(t)$ introduced in (41).

Lemma 7. *Let $a_{\pm,j}(t)$ and $a_{0,j}(t)$ be as defined in (41). There is a constant $C > 0$, independent of S_0 and $T^* \geq T_0$, such that for all $t \in [T^*, S_0]$,*

$$\left| a'_{\pm,j}(t) \pm \frac{\sqrt{\lambda_0}}{\gamma_j} a_{\pm,j}(t) \right| \leq C \|V(t)\|^2 + C e^{-3\gamma_0 t}. \quad (61)$$

Proof. We prove the case of $a_{-,j}(t)$. The other case is similar. We compute the time derivative of $a_{-,j}$ using (56) and (58), and we choose $\gamma_0 > 0$ as small as needed, but fixed.

$$\begin{aligned} a'_{-,j}(t) &= -\tilde{y}'_j(t)\langle (Z_{+,j})_x | V(t) \rangle + \langle Z_{+,j} | V_t(t) \rangle \\ &= -\tilde{x}'_j\langle (Z_{+,j})_x | V(t) \rangle + \langle (\mathcal{L}_j^* - \beta_j \partial_x) Z_{+,j} | V(t) \rangle + \sum_{k=1}^N x'_k \langle \Phi_{0,k} | Z_{+,j} \rangle \\ &\quad + O(\|V(t)\|^2 + e^{-3\gamma_0 t}). \end{aligned}$$

From Lemma 3 we have $\langle \Phi_{0,j} | Z_{+,j} \rangle = 0$. Therefore, since $\mathcal{L}_j^* - \beta_j \partial_x = \mathcal{H}_j$, where $\mathcal{H}_j := \mathcal{H}(Q_{\beta_j})$ (cf. (24)), we have from Lemma 2 and (59),

$$\begin{aligned} a'_{-,j}(t) &= \frac{\sqrt{\lambda_0}}{\gamma_j} a_{-,j}(t) + O(|x'_j| \|V(t)\| + \|V(t)\|^2 + e^{-3\gamma_0 t}) \\ &= \frac{\sqrt{\lambda_0}}{\gamma_j} a_{-,j}(t) + O(\|V(t)\|^2 + e^{-3\gamma_0 t}). \quad \square \end{aligned}$$

Step 3. Lyapunov functional. Let $L_0 > 0$ be a large constant to be chosen later. Let $(\phi_j)_{j=1,\dots,N}$ be a partition of the unity of \mathbb{R} placed at the midpoint between two solitons. More precisely, let

$$\phi \in C^\infty(\mathbb{R}), \quad \phi' > 0, \quad \lim_{-\infty} \phi = 0, \quad \lim_{+\infty} \phi = 1. \quad (62)$$

We have³, for all $L > L_0$,

$$\sum_{j=1}^N \phi_j(t, x) \equiv 1, \quad \phi_j(t, x) = \phi\left(\frac{x - m_j t}{L}\right) - \phi\left(\frac{x - m_{j+1} t}{L}\right), \quad (63)$$

where $m_j := \frac{1}{2}(\beta_j + \beta_{j-1})$, with $j = 2, \dots, N-1$, and $m_1 := -\infty$, $m_N = +\infty$. We introduce the j -th portion of momentum

$$P_j[\varphi](t) := \frac{1}{2} \int \phi_j u_t u_x \, dx, \quad \varphi = (u, u_t)^T, \quad (64)$$

and the modified *Lyapunov functional*

$$\mathcal{F}[\varphi](t) := E[\varphi](t) + 2 \sum_{j=1}^N \beta_j P_j[\varphi](t), \quad (65)$$

with $E[\varphi]$ being the energy defined in (2). Our first result is a suitable decomposition of $\mathcal{F}[u]$ around the multi-soliton solution.

Lemma 8. *Let $V(t) = (v_1(t), v_2(t))^T$ be the error function defined in Proposition 4. There is a positive constant $C > 0$ such that*

$$\left| \mathcal{F}[\varphi](t) - \sum_{j=1}^N \langle H_j V | V \rangle \right| \leq C \|V(t)\|^3 + \frac{C}{L} e^{-2\gamma_0 t}, \quad (66)$$

where

$$\langle H_j V | V \rangle := \int \phi_j (v_2^2 + (v_1)_x^2 + v_1^2 - f'(Q_{\beta_j}) v^2 + 2\beta_j v_2 (v_1)_x). \quad (67)$$

³Do not confuse the constant L in (63) with the operator L in (13).

Proof. From the decomposition

$$\varphi(t) = (u, u_t)(t) = (\tilde{R}_1, \tilde{R}_2)^T + (v_1, v_2)^T(t), \quad (68)$$

we have

$$\begin{aligned} \mathcal{F}[\varphi](t) &= \frac{1}{2} \int (u_t^2 + u_x^2 + u^2 - F(u)) + \sum_{j=1}^N \beta_j \int \phi_j u_t u_x \\ &= \frac{1}{2} \int (\tilde{R}_2^2 + (\tilde{R}_1)_x^2 + \tilde{R}_1^2 - 2F(\tilde{R}_1)) + \sum_{j=1}^N \beta_j \int \tilde{R}_2 (\tilde{R}_1)_x \phi_j \\ &\quad + \int [\tilde{R}_2 v_2 + (\tilde{R}_1)_x (v_1)_x + \tilde{R}_1 v_1 - f(\tilde{R}_1) v_1 + \sum_{j=1}^N \beta_j (\tilde{R}_2 (v_1)_x + v_2 (\tilde{R}_1)_x) \phi_j] \\ &\quad + \frac{1}{2} \int (v_2^2 + (v_1)_x^2 + v_1^2 - f'(\tilde{R}_1) v_1^2) + \sum_{j=1}^N \beta_j \int v_2 (v_1)_x \phi_j \\ &\quad - \int (F(\tilde{R}_1 + v_1) - F(\tilde{R}_1) - f(\tilde{R}_1) v_1 - \frac{1}{2} f'(\tilde{R}_1) v_1^2) \\ &=: I_1 + I_2 + I_3 + I_4. \end{aligned}$$

Let us consider the term I_1 . Since $\tilde{R}_2 = -\sum_{j=1}^N \beta_j (Q_{\beta_j})_x$ and $(\tilde{R}_1)_x = \sum_{j=1}^N (Q_{\beta_j})_x$, one has

$$\begin{aligned} I_1 &= \frac{1}{2} \sum_{j=1}^N \int [\beta_j^2 (Q_{\beta_j})_x^2 + (Q_{\beta_j})_x^2 + Q_{\beta_j}^2 - 2F(Q_{\beta_j}) - 2\beta_j^2 (Q_{\beta_j})_x^2] + O(e^{-3\gamma_0 t}) \\ &= \frac{1}{2} \int [Q_x^2 + Q^2 - 2F(Q)] \sum_{j=1}^N \frac{1}{\gamma_j} + O(e^{-3\gamma_0 t}). \end{aligned}$$

Now we consider I_2 . Integrating by parts, we have

$$\begin{aligned} I_2 &= \int v_2 \left[\tilde{R}_2 + (\tilde{R}_1)_x \sum_{j=1}^N \beta_j \phi_j \right] \\ &\quad - \int v_1 \left[(\tilde{R}_1)_{xx} - \tilde{R}_1 + f(\tilde{R}_1) + (\tilde{R}_2)_x \sum_{j=1}^N \beta_j \phi_j \right] - \sum_{j=1}^N \beta_j \int v_1 \tilde{R}_2 (\phi_j)_x. \end{aligned}$$

Note that

$$\tilde{R}_2 + (\tilde{R}_1)_x \sum_{j=1}^N \beta_j \phi_j = \sum_{k=1}^N [-\beta_k (Q_{\beta_k})_x + (Q_{\beta_k})_x \sum_{j=1}^N \beta_j \phi_j] = \sum_{k=1}^N (Q_{\beta_k})_x \sum_{j \neq k}^N \beta_j \phi_j.$$

Hence

$$\int v_2 \left[\tilde{R}_2 + (\tilde{R}_1)_x \sum_{j=1}^N \beta_j \phi_j \right] = O(e^{-3\gamma_0 t}).$$

On the other hand,

$$\int v_1 \left[(\tilde{R}_1)_{xx} - \tilde{R}_1 + f(\tilde{R}_1) + (\tilde{R}_2)_x \sum_{j=1}^N \beta_j \phi_j \right] = O(e^{-3\gamma_0 t}).$$

Finally,

$$\left| \sum_{j=1}^N \beta_j \int v_1 \tilde{R}_2(\phi_j)_x \right| \leq C \|v_1\|_{L^2(\mathbb{R})} e^{-2\gamma_0 t}.$$

Gathering the above estimates, we get

$$|I_2| \leq C e^{-3\gamma_0 t}.$$

Let us consider the integral I_3 . Since $\sum_j \phi_j = 1$, we have

$$\begin{aligned} I_3 &= \frac{1}{2} \sum_{j=1}^N \int \phi_j (v_2^2 + (v_1)_x^2 + v_1^2 - f'(\tilde{R}_1)v_1^2 + 2\beta_j v_2(v_1)_x) \\ &= \frac{1}{2} \sum_{j=1}^N \int \phi_j (v_2^2 + (v_1)_x^2 + v_1^2 - f'(Q_{\beta_j})v_1^2 + 2\beta_j v_2(v_1)_x) - \frac{1}{2} \sum_{k \neq j} \int \phi_j f'(Q_k)v_1^2 \\ &\quad - \frac{1}{2} \sum_{j=1}^N \int \phi_j \left(f' \left(\sum_{k=1}^N Q_{\beta_k} \right) - \sum_{k=1}^N f'(Q_{\beta_k}) \right) v_1^2. \end{aligned}$$

Fix $\ell \in \{1, \dots, N-1\}$. If $x \in [m_\ell t, m_{\ell+1}t]$, then for all $p \neq \ell$,

$$|Q_{\beta_p}(t, x)| \leq C e^{-2\gamma_0 t}.$$

Therefore, for all $x \in [m_\ell t, m_{\ell+1}t]$,

$$\left| f' \left(\sum_{k=1}^N Q_{\beta_k}(t, x) \right) - \sum_{k=1}^N f'(Q_{\beta_k}(t, x)) \right| \leq C e^{-\gamma_0 t}.$$

Repeating the same argument for each ℓ , and using (50), we get

$$I_3 = \frac{1}{2} \langle H_j V | V \rangle + O(e^{-3\gamma_0 t}).$$

Finally, we consider I_4 . It is not difficult to check that

$$|I_4| \leq C \|V\|^3.$$

Collecting the above results, we get finally (66). \square

Our next result describes the variation of the momentum P_j .

Lemma 9. *There exists $C > 0$ independent of time and L , such that for all $t \in [T^*, S_0]$,*

$$|P_j[\varphi](t) - P_j[\varphi](S_0)| \leq \frac{C}{L} e^{-2\gamma_0 t}. \quad (69)$$

Proof. A simple computation using (NLKG) shows that

$$\begin{aligned} \partial_t P_j[\varphi](t) &= -\frac{1}{4} \int u_t^2(\phi_j)_x - \frac{1}{4} \int u_x^2(\phi_j)_x + \frac{1}{4} \int u^2(\phi_j)_x \\ &\quad - \frac{1}{2} \int F(u)(\phi_j)_x + \frac{1}{2} \int u_t u_x(\phi_j)_t. \end{aligned} \quad (70)$$

Indeed, one has

$$\begin{aligned} \partial_t P_j[\varphi](t) &= \frac{1}{2} \int u_t u_x(\phi_j)_t + \frac{1}{2} \int u_t u_{tx} \phi_j + \frac{1}{2} \int u_{tt} u_x \phi_j \\ &= \frac{1}{2} \int u_t u_x(\phi_j)_t + \frac{1}{4} \int (u_t^2)_x \phi_j + \frac{1}{2} \int (u_{xx} - u + f(u)) u_x \phi_j \end{aligned}$$

$$= \frac{1}{2} \int u_t u_x (\phi_j)_t - \frac{1}{4} \int u_t^2 (\phi_j)_x - \frac{1}{4} \int (u_x^2 - u^2 + 2F(u)) (\phi_j)_x,$$

as desired. Now, from the decomposition (68) we replace above to obtain (compare with (60))

$$\begin{aligned} & |\partial_t P_j[\varphi](t)| \\ & \leq C \left(\frac{e^{-3\gamma_0 t}}{L} + \int v_2^2 (\phi_j)_x + \int (v_1)_x^2 (\phi_j)_x + \int v_1^2 (\phi_j)_x + \int F(v_1) (\phi_j)_x \right). \end{aligned}$$

From the smallness condition of v , we get finally

$$|\partial_t P_j[\varphi](t)| \leq \frac{C}{L} e^{-2\gamma_0 t},$$

as desired. The conclusion follows after integration in time. \square

The previous Lemma and the energy conservation law imply the following

Corollary 3. *There exists $C > 0$ independent of time and $L > 0$ such that, for all $t \in [T^*, S_0]$,*

$$|\mathcal{F}[\varphi](t) - \mathcal{F}[\varphi](S_0)| \leq \frac{C}{L} e^{-2\gamma_0 t}. \quad (71)$$

Now we use the coercivity associated to H_j . A standard localization argument (see e.g. [21]), Proposition 2 and (41) give

$$\sum_{j=1}^N \langle H_j V | V \rangle \geq \nu_0 \|V(t)\|^2 - \frac{1}{\nu_0} (\|\mathbf{a}_+\|_{\ell^2}^2 + \|\mathbf{a}_-\|_{\ell^2}^2),$$

for an independent constant $\nu_0 > 0$. From this coercivity estimate, using (71) and (66), the initial bound (49), and bounding the terms in \mathbf{a}_\pm by (46), we get that for some $C > 0$

$$\forall t \in [T^*, S_0], \quad \|V(t)\| \leq \frac{C}{\sqrt{L}} e^{-\gamma_0 t} + C e^{-3/2\gamma_0 t}.$$

Therefore, for $L \geq 4C^2$, we improve the first condition in (45), to get (50). We can now integrate of the modulation equation (59) for $\tilde{x}'_j(t)$ we get the second estimates in (50) (by increasing L is necessary).

Now, using (61) on $\mathbf{a}'_-(t)$ and integrating in time, we improve in a similar way the conditions in (46), to obtain (51). In conclusion, (52) must be satisfied.

Step 4. Transversality. For notation, let $\mathcal{N}(\mathbf{a}_+, t) := e^{3\gamma_0 t} \|\mathbf{a}_+(t)\|_{\ell^2}^2$. Using the expansion (61), we compute

$$\begin{aligned} \frac{d}{dt} \mathcal{N}(\mathbf{a}_+, t) &= \sum_{j=1}^N e^{3\gamma_0 t} a'_{+,j}(t) a_{+,j}(t) + 3\gamma_0 \mathcal{N}(\mathbf{a}^+, t) \\ &= -e^{3\gamma_0 t} \sqrt{\lambda_0} \sum_{j=1}^N \frac{|a_{+,j}(t)|^2}{\gamma_j} + O(e^{3\gamma_0 t} (\|V(t)\|_{L^2}^2 + e^{-2\gamma_0 t}) \|\mathbf{a}_+(t)\|_{\ell^2}) \\ &\quad + 3\gamma_0 \mathcal{N}(\mathbf{a}^+, t) \\ &\leq -(2c_0 - 3\gamma_0) \mathcal{N}(\mathbf{a}^+, t) - O(e^{3\gamma_0 t} (\|V(t)\|_{L^2}^2 + e^{-2\gamma_0 t}) \|\mathbf{a}_+(t)\|_{\ell^2}), \end{aligned}$$

where $c_0 = \frac{1}{2}\sqrt{\lambda_0} \min_i \{1/\gamma_i\} > 0$. Note that from (43) we have $2c_0 - 3\gamma_0 > \gamma_0 > 0$. Now, at time $T^* = T^*(\mathbf{a}^+)$, $\|V(T^*)\|_{L^2} = O(e^{-\gamma_0 T^*})$, whereas $\|\mathbf{a}_+(T^*)\|_{\ell^2} = e^{-3\gamma_0 T^*/2}$, i.e. $\mathcal{N}(\mathbf{a}^+, T^*) = 1$, hence

$$\left. \frac{d}{dt} \mathcal{N}(\mathbf{a}^+, t) \right|_{t=T^*(\mathbf{a}^+)} \leq -(2c_0 - 3\gamma_0) + O(e^{-\gamma_0 T^*/2}).$$

Choosing T_0 larger if necessary, and as $T^* \geq T_0$ for all \mathbf{a}_+ , we get

$$\left. \frac{d}{dt} \mathcal{N}(\mathbf{a}_+, t) \right|_{t=T^*(\mathbf{a}_+)} \leq -\frac{1}{2}\gamma_0 < 0. \quad (72)$$

This concludes the proof of Proposition 4. \square

We end this paragraph with the proof of Corollary 2.

Proof of Corollary 2. Let us now show that Ω is open and that the mapping $\mathbf{a}_+ \mapsto T^*(\mathbf{a}_+)$ is continuous. Let $\mathbf{a}_+ \in \Omega$. We recall that $\mathcal{N}(\mathbf{a}_+, t) = e^{3\gamma_0 t} \|\mathbf{a}_+(t)\|_{\ell^2}^2$. By (72), for all $\varepsilon > 0$ small, there exists $\delta > 0$ such that

- $\mathcal{N}(\mathbf{a}_+, T^*(\mathbf{a}_+) - \varepsilon) > 1 + \delta$, and
- for all $t \in [T^*(\mathbf{a}_+) + \varepsilon, S_0]$ (possibly empty), $\mathcal{N}(\mathbf{a}_+, t) < 1 - \delta$.

By continuity of the flow of the (NLKG) equation, it follows that there exists $\eta > 0$ such that the following holds. For all $\tilde{\mathbf{a}}_+ \in B_{\mathbb{R}^N}(0, 1)$ such that $\|\tilde{\mathbf{a}}_+ - \mathbf{a}_+\| \leq \eta$, then $|\mathcal{N}(\tilde{\mathbf{a}}_+, t) - \mathcal{N}(\mathbf{a}_+, t)| \leq \delta/2$ for all $t \in [T^*(\mathbf{a}_+) - \varepsilon, S_0]$. In particular, $\tilde{\mathbf{a}}_+ \in \Omega$ and

$$T^*(\mathbf{a}_+) - \varepsilon \leq T^*(\tilde{\mathbf{a}}_+) \leq T^*(\mathbf{a}_+) + \varepsilon.$$

This exactly means that Ω contains a neighbourhood of \mathbf{a}_+ , hence is open, and that $\mathbf{a}_+ \mapsto T^*(\mathbf{a}_+)$ continuous.

Finally, let us show that the exit is instantaneous on the sphere. If $\|\mathbf{a}_+\|_{\ell^2} = 1$, then $\mathcal{N}(\mathbf{a}_+, S_0) = 1$, hence by (72), $\mathcal{N}(\mathbf{a}_+, t) > 1$ for all $t < S_0$ in a neighborhood of S_0 . This means that $T^*(\mathbf{a}_+) = S_0$. \square

3.3. Extension to higher dimension. The main part of the proof remains unchanged. One has to work only for the definition of the Lyapunov functional. The key point is to notice that one can find a suitable direction as in [16]. The set

$$\mathcal{M} = \{\beta \in \mathbb{R}^d \mid \forall j, \beta \cdot \beta_j = 0\},$$

is of zero measure: let $\bar{\beta} \notin \mathcal{M}$; up to rescaling, we can assume $|\bar{\beta}| = 1$. Without loss of generality we can assume that the indexes j satisfy

$$-1 < (\bar{\beta} \cdot \beta_1) < (\bar{\beta} \cdot \beta_2) < \dots < (\bar{\beta} \cdot \beta_N) < 1.$$

We use again the 1d cut-off function ϕ defined at Step 3 of the previous to define the new cut-off functions

$$\psi_j(x) = \phi\left(\frac{\bar{\beta} \cdot x - m_j t}{L}\right) - \phi\left(\frac{\bar{\beta} \cdot x - m_{j+1} t}{L}\right), \quad \text{where } m_j = \frac{1}{2}(\beta_j + \beta_{j-1}) \cdot \bar{\beta}.$$

Then all the computations of Step 3 of Section 3.2 follow unchanged. We refer to [16] (Claim 1 and what follows) for further details.

4. PROOF OF THEOREM 1

The proof of Theorem 1 follows from Proposition 3 in a standard fashion, see e.g. [16]. The main point is continuity of the flow for the weak $H^1 \times L^2$ topology. More precisely:

Lemma 10. *The (NLKG) flow is continuous for the weak $H^1 \times L^2$ topology. More precisely, let $U_n \in \mathcal{C}([0, T], H^1 \times L^2)$ be a sequence of solutions to (NLKG), and assume that for some $M > 0$,*

$$U_n(0) \rightharpoonup U^* \quad \text{in } H^1 \times L^2 \text{ - weak, and } \forall n, \quad \|U_n(t)\|_{\mathcal{C}([0, T], H^1 \times L^2)} \leq M.$$

Define $U \in \mathcal{C}([0, T^+(U)], H^1 \times L^2)$ be the solution to (NLKG) with initial data $U(0) = U^$. Then $T^+(U) > T$ and*

$$\forall t \in [0, T], \quad U_n(t) \rightharpoonup U(t) \quad \text{in } H^1 \times L^2 \text{ - weak.}$$

Proof. This is a simple consequence of the local well posedness of (NLKG) in $H^s \times \dot{H}^{s-1}$ for some $s < 1$. More precisely, we have

Theorem (Local wellposedness). *There exist $0 \leq s_{f,d} < 1$ such that for all $s \geq s_{f,g}$, the following holds. Given any data $U_0 = (u_0, u_1) \in H^s \times \dot{H}^{s-1}$, there exist a unique solution $U \in \mathcal{C}([0, T^+(U)], H^s \times \dot{H}^{s-1})$ to (NLKG) such that $U(0) = U_0$. Furthermore,*

- (1) *The maximal time of existence $T^+(U)$ is the same in all $H^\sigma \times \dot{H}^{\sigma-1}$ for $\sigma \in [s_{f,d}, s]$. If finite, it is characterized by*

$$\lim_{t \rightarrow T^+(U)} \|U(t)\|_{H^s \times \dot{H}^{s-1}} = +\infty.$$

- (2) *The flow is continuous, in the sense that if U_n is a sequence of solution to (NLKG) such that $U_n(0) \rightarrow U(0)$ in $H^s \times \dot{H}^{s-1}$, then $T^+(U) \geq \liminf_n T^+(U_n)$ and*

$$\forall t \in [0, T^+(U)), \quad \|U_n - U\|_{\mathcal{C}([0, t], H^s \times \dot{H}^{s-1})} \rightarrow 0 \quad \text{as } n \rightarrow +\infty.$$

We refer to [29, Theorem 1.2] and the Remark following it for a proof and the precise value of $s_{f,d}$ (which is not important for us).

Fix $0 < s < 1$ be such that the Theorem holds. Let $t \in [0, T]$ such that $t < T^+(U)$. Let $V \in (\mathcal{D}(\mathbb{R}^d))^2$ and $R > 0$ such that $\text{Supp } V \subset B_{\mathbb{R}^d}(0, R)$.

As $U_n(0) \rightharpoonup U(0)$ weakly in $H^1 \times L^2(\mathbb{R}^d)$, there holds by Sobolev compact embedding

$$\|U_n(0) - U(0)\|_{H^s \times \dot{H}^{s-1}(B_{\mathbb{R}^d}(0, R+t))} \rightarrow 0.$$

It follows by finite speed of propagation and the continuity of the flow in the local well posedness Theorem that

$$\|U_n(t) - U(t)\|_{H^s \times \dot{H}^{s-1}(B_{\mathbb{R}^d}(0, R))} \rightarrow 0.$$

Hence denoting $U_n = (u_n, \partial_t u_n)$ and $V = (v_0, v_1)$,

$$\begin{aligned} & |\langle U_n(t) - U(t), V \rangle| \\ &= \left| \int_{|x| \leq R} \left((\partial_t u_n(t, x) - \partial_t u(t, x)) v_1(x) \right. \right. \\ & \quad \left. \left. + \nabla(u_n - u) \cdot \nabla v_0 + (u_n(t, x) - u(t, x)) v_0(x) \right) dx \right| \end{aligned}$$

$$\leq \|U_n(t) - U(t)\|_{H^s \times \dot{H}^{s-1}(B_{\mathbb{R}^d}(0,R))} \|V\|_{H^{2-s} \times \dot{H}^{1-s}} \rightarrow 0.$$

Therefore $U_n(t) \rightharpoonup U(t)$ in \mathcal{D}' , and by the $H^1 \times L^2$ bound, $U_n(t) \rightharpoonup U(t)$ weakly in $H^1 \times L^2$.

In particular $\|U(t)\|_{H^1 \times L^2} \leq \liminf_{n \rightarrow \infty} \|U_n(t)\| \leq M$. From there, a continuity argument shows that $T^+(U) > T$. \square

We can now prove Theorem 1. Let $(S_n)_{n \geq 1} \subset \mathbb{R}$ be a sequence that satisfies $S_n > S_0$, S_n increasing and $S_n \rightarrow +\infty$. From Proposition 3 there exists a sequence of final data functions $U_{0,n} \in H^1 \times L^2$ such that

$$\forall t \in [T_0, S_n], \quad U_n(t) := \varphi(S_n, t, U_{0,n}) \in \mathcal{V}(t). \quad (73)$$

(We recall that φ denotes the flow and is defined in (47)). Note that T_0 does not depend on S_n , and observe that there exists M independent of n such that

$$\forall t \in [T_0, S_n], \quad \|U_n(t) - R(t)\|_{H^1 \times L^2} \leq M e^{-\gamma_0 t}. \quad (74)$$

Let U_0^* be a weak limit in $H^1 \times L^2$ of the bounded sequence $U_n(T_0)$, and define

$$U^*(t) = \varphi(t, T_0, U_0^*).$$

Fix $t \geq T_0$. Then the previous Lemma applies on $[T_0, t]$ and shows that $T^+(U^*) > t$ and $U_n(t) \rightharpoonup U^*(t)$ weakly in $H^1 \times L^2$. Hence (74) yields

$$\|U^*(t) - R(t)\|_{H^1 \times L^2} \leq \liminf_n \|U_n(t) - R(t)\|_{H^1 \times L^2} \leq M e^{-\gamma_0 t}.$$

Therefore, $T^+(U^*) = +\infty$ and U^* is the desired multi-soliton.

APPENDIX A. THE ORBIT OF Q UNDER GENERAL LORENTZ TRANSFORMATIONS

In this appendix we prove that the orbit of Q under the group generated by space and time translations, and general Lorentz transforms is

$$\mathcal{F} := \{(t, x) \mapsto Q_\beta(t, x - x_0) \mid \beta, x_0 \in \mathbb{R}^d, |\beta| < 1\}.$$

We recall that we consider Q as a function of time with the slight abuse of notation $Q(t, x) = Q(x)$.

The map $\beta \mapsto \Lambda_\beta$ (see (5)) is a group homomorphism from $(B_{\mathbb{R}^d}(0, 1), \oplus)$ to $(M_{d+1}(\mathbb{R}), \circ)$, where \oplus denotes Einstein's velocity addition

$$x \oplus y = \frac{1}{1 + x \cdot y} \left(y + \frac{x \cdot y}{|y|^2} y + \sqrt{1 - |y|^2} \left(x - \frac{x \cdot y}{|y|^2} y \right) \right).$$

In particular, $\Lambda_{-\beta} \Lambda_\beta = \text{Id}_{d+1}$.

A general Lorentz transform is an element of $O(1, d) \simeq \mathbb{R}^d \rtimes O(d)$, hence can be written in the form

$$\Lambda_{U, \beta} := \begin{pmatrix} 1 & 0 & \cdots & 0 \\ 0 & & & \\ \vdots & & U & \\ 0 & & & \end{pmatrix} \Lambda_\beta, \quad \text{where } U \in SO(d), \text{ i.e. } UU^T = \text{Id}_d.$$

As $Q(x)$ is radially symmetric, it follows that

$$Q \left(\Lambda_{U, \beta} \begin{pmatrix} t \\ x \end{pmatrix} \right) = Q \left(\Lambda_\beta \begin{pmatrix} t \\ x \end{pmatrix} \right) = Q_\beta(x),$$

hence the orbit of $\{Q\}$ under general Lorentz transform is simply $\{Q_\beta|\beta \in \mathbb{R}^d\}$. We now want to parametrize the other invariances of (NLKG), that is time and space shifts. Fortunately, the former reduce to the latter. Indeed, notice that

$$\beta\beta^T \sim \begin{pmatrix} |\beta|^2 & 0 & \cdots & 0 \\ 0 & & & \\ \vdots & & 0 & \\ 0 & & & \end{pmatrix}, \quad \text{so that} \quad \text{Id}_d + \frac{\gamma-1}{|\beta|^2} \beta\beta^T \sim \begin{pmatrix} \gamma & 0 & \cdots & 0 \\ 0 & & & \\ \vdots & & \text{Id}_{d-1} & \\ 0 & & & \end{pmatrix}.$$

(Here \sim indicates similarity of matrices). In particular, $\text{Id}_d + \frac{\gamma-1}{|\beta|^2} \beta\beta^T$ is invertible. Then time translations for Q_β can be rethought as an adequate space shift:

$$\begin{aligned} Q_\beta(t+t_0, x) &= Q\left(\Lambda_\beta \begin{pmatrix} t+t_0 \\ x \end{pmatrix}\right) = Q\left(\Lambda_\beta \begin{pmatrix} t \\ x \end{pmatrix} + t_0 \begin{pmatrix} \gamma \\ -\beta \end{pmatrix}\right) \\ &= Q\left(\Lambda_\beta \begin{pmatrix} t \\ x \end{pmatrix} - t_0 \begin{pmatrix} 0 \\ \beta \end{pmatrix}\right) = Q_\beta\left(x - t_0(\text{Id}_d + \frac{\gamma-1}{|\beta|^2} \beta\beta^T)^{-1}(\beta)\right). \end{aligned}$$

It follows that \mathcal{F} is stable through all general Lorentz transform, time and space shifts, hence it is the orbit of Q through the group generated by these transformations.

REFERENCES

- [1] H. Berestycki and P.-L. Lions, *Nonlinear scalar field equations. I. Existence of a ground state*, Arch. Rational Mech. Anal. **82**, (1983) 313–345.
- [2] J. Bourgain, *Global solutions of nonlinear Schrödinger equations*, AMS Colloquium publications **46**, AMS, Providence, RI, 1999.
- [3] R. Côte, Y. Martel, and F. Merle, *Construction of multi-soliton solutions for the L^2 -supercritical gKdV and NLS equations*, Rev. Mat. Iberoamericana **27** (2011), no. 1, 273–302.
- [4] R. Côte and H. Zaag, *Construction of a multi-soliton blow-up solution to the semilinear wave equation in one space dimension*, Comm. Pure Appl. Math. **66** (2013), no. 10, 1541–1581.
- [5] G.H. Derrick, *Comments on Nonlinear Wave Equations as Models for Elementary Particles*, J. Math. Phys. **5**, 1252 (1964); doi: 10.1063/1.1704233.
- [6] T. Duyckaerts, and F. Merle, *Dynamic of threshold solutions for energy-critical NLS*, GAFA Vol. **18** (2008), 1787–1840.
- [7] T. Duyckaerts, and F. Merle, *Dynamics of threshold solutions for energy-critical wave equation*, Int. Math. Res. Pap. IMRP **2008**, Art ID rpn002, 67 pp.
- [8] T. Duyckaerts, C. E. Kenig, and F. Merle, *Classification of radial solutions of the focusing, energy-critical wave equation*, preprint, arXiv:1204.0031.
- [9] B. Gidas, W.-M. Ni, and L. Nirenberg, *Symmetry and related properties via the maximum principle*, Comm. Math. Phys. **68** (1979), no. 3, 209–243.
- [10] M. Grillakis, J. Shatah, and W. Strauss, *Stability theory of solitary waves in the presence of symmetry. I*. J. Funct. Anal. **74** (1987), no. 1, 160–197.
- [11] M. Grillakis, J. Shatah, and W. Strauss, *Stability theory of solitary waves in the presence of symmetry. II*. J. Funct. Anal. **94** (1990), no. 2, 308–348.

- [12] J. Ginibre, and G. Velo, *The global Cauchy problem for the nonlinear Klein-Gordon equation*. Math. Z. **189** (1985), no. 4, 487–505.
- [13] J. Krieger, K. Nakanishi, and W. Schlag, *Global dynamics above the ground state energy for the one-dimensional NLKG equation*, preprint.
- [14] Kwong, M. K., *Uniqueness of positive solutions of $\Delta u - u + u^p = 0$ in \mathbb{R}^n* , Arch. Rational Mech. Anal. **105** (1989), no. 3, 243–266.
- [15] M. Maris, *Existence of nonstationary bubbles in higher dimension*, J. Math. Pures Appl. **81** (2002) 1207–1239.
- [16] Y. Martel, *Asymptotic N -soliton-like solutions of the subcritical and critical generalized Korteweg-de Vries equations*, Amer. J. Math. **127** (2005), no. 5, 1103–1140.
- [17] K. McLeod, *Uniqueness of positive radial solutions of $\Delta u + f(u) = 0$ in \mathbb{R}^n . II*, Trans. Amer. Math. Soc. **339** (1993) 495–505.
- [18] F. Merle, *Construction of solutions with exactly k blow-up points for the Schrödinger equation with critical nonlinearity*, Comm. Math. Phys., **129** (1990), no. 2, 223–240.
- [19] Y. Martel, and F. Merle, *Multi solitary waves for nonlinear Schrödinger equations*, Ann. IHP Nonlinear Anal. **23** (2006), 849–864.
- [20] Y. Martel and F. Merle, *Stability of two soliton collision for nonintegrable $gKdV$ equations*, Comm. Math. Phys. **286** (2009), 39–79.
- [21] Y. Martel, F. Merle, and T. P. Tsai, *Stability in H^1 of the sum of K solitary waves for some nonlinear Schrödinger equations*, Duke Math. Journal, **133** (2006), no. 3, 405–466.
- [22] F. Merle, and H. Zaag, *Existence and classification of characteristic points at blow-up for a semilinear wave equation in one space dimension*. Amer. J. Math. **134** (2012), no. 3, 581–648.
- [23] M. Nakamura, and T. Ozawa, *The Cauchy problem for nonlinear Klein-Gordon equations in the Sobolev spaces*. Publ. Res. Inst. Math. Sci. **37** (2001), no. 3, 255–293.
- [24] K. Nakanishi, and W. Schlag, *Invariant manifolds and dispersive Hamiltonian evolution equations*, Zurich Lectures in Advanced Mathematics. European Mathematical Society (EMS), Zürich, 2011, vi+253 pp.
- [25] R. L. Pego, and M. I. Weinstein, *Eigenvalues, and instabilities of solitary waves*. Philos. Trans. Roy. Soc. London Ser. A **340** (1992), no. 1656, 47–94.
- [26] K. Nakanishi, and W. Schlag, *Global dynamics above the ground state energy for the focusing nonlinear Klein-Gordon equation*, J. Differential Equations **250** (2011), no. 5, 2299–2333.
- [27] J. Serrin, M. Tang, *Uniqueness of ground states for quasilinear elliptic equations*, Indiana Univ. Math. J. **49** (2000), no. 3, 897–923.
- [28] J. Shatah, and W. Strauss, *Walter, Instability of nonlinear bound states*, Comm. Math. Phys. **100** (1985), no. 2, 173–190.
- [29] T. Tao *Low regularity semi-linear wave equations*. Comm. PDE, **24** (1999), 599–630.
- [30] M.I. Weinstein, *Modulational stability of ground states of nonlinear Schrödinger equations*, SIAM J. Math. Anal. **16** (1985), no. 3, 472–491.

- [31] M.I. Weinstein, *Lyapunov stability of ground states of nonlinear dispersive evolution equations*, Comm. Pure Appl. Math. **39**, (1986) 51–68.

RAPHAËL CÔTE

CNRS and École polytechnique
Centre de Mathématiques Laurent Schwartz UMR 7640
Route de Palaiseau, 91128 Palaiseau cedex, France
cote@math.polytechnique.fr

CLAUDIO MUÑOZ

Department of Mathematics, University of Chicago
5734 South University Avenue, Chicago, IL 60615, U.S.A.
cmunoz@math.uchicago.edu