



ASYMPTOTIC PROPERTIES OF FIBONACCI CUBES AND LUCAS CUBE

Sandi Klavzar, Michel Mollard

► To cite this version:

Sandi Klavzar, Michel Mollard. ASYMPTOTIC PROPERTIES OF FIBONACCI CUBES AND LUCAS CUBE. 2013. hal-00836788v1

HAL Id: hal-00836788

<https://hal.science/hal-00836788v1>

Preprint submitted on 21 Jun 2013 (v1), last revised 5 Sep 2013 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Asymptotic properties of Fibonacci cubes and Lucas cubes

Sandi Klavžar

Faculty of Mathematics and Physics

University of Ljubljana, Slovenia

and

Faculty of Natural Sciences and Mathematics

University of Maribor, Slovenia

and

Institute of Mathematics, Physics and Mechanics, Ljubljana

`sandi.klavzar@fmf.uni-lj.si`

Michel Mollard

CNRS Université Joseph Fourier

Institut Fourier, BP 74

100 rue des Maths, 38402 St Martin d'Hères Cedex, France

`michel.mollard@ujf-grenoble.fr`

Abstract

It is proved that the asymptotic average eccentricity and the asymptotic average degree of Fibonacci cubes and Lucas cubes are $(5 + \sqrt{5})/10$ and $(5 - \sqrt{5})/5$, respectively. A new labeling of the leaves of Fibonacci trees is introduced and proved that the eccentricity of a vertex of a given Fibonacci cube is equal to the depth of the associated leaf in the corresponding Fibonacci tree. Hypercube density is also introduced and studied. The hypercube density of both Fibonacci cubes and Lucas cubes is shown to be $(1 - 1/\sqrt{5})/\log_2 \varphi$, where φ is the golden ratio, and the Cartesian product of graphs is used to construct families of graphs with a fixed, non-zero hypercube density. It is also proved that the limit normed sum of ratios of Fibonacci words and Lucas words with fixed coordinate 0 and 1, respectively, is φ^2 .

Key words: Fibonacci cube; Lucas cube; convergence of sequences; average eccentricity; Fibonacci tree; average distance; hypercube density

AMS Subj. Class.: 05C12, 40A05, 05A15

1 Introduction

Fibonacci cubes [8] and Lucas cubes [15] form appealing infinite families of graphs extensively investigated by now; see the recent survey [10]. These cubes are subgraphs

of hypercubes and on one hand inherit many of the fine properties of hypercubes, while on the other hand their size grows significantly slower than that of hypercubes. Moreover, Fibonacci cubes and Lucas cubes found several applications, for instance in theoretical chemistry; see [21] for a use of Fibonacci cubes and [20] for a use of Lucas cubes. We also mention that in [16] an investigation of the on-line routing of linear permutations on these cubes is performed.

In the last years large graphs (and/or complex networks) became a topic of great interest—not only in mathematics but elsewhere—hence it seems justified to consider the asymptotic behaviour of applicable families of graphs, such as Fibonacci and Lucas cubes. The average normed distance of these graphs was proved in [12] to be $2/5$. In this paper we study the limit behaviour of the average eccentricity of these cubes, the limit behaviour of their average degree, and some related topics. For some general properties of the average eccentricity see [3, 9], while [6] gives the the average eccentricity of Sierpiński graphs.

We proceed as follows. In the rest of this section concepts needed in this paper are formally introduced. In Section 2 we determine the limit average eccentricity of Fibonacci and Lucas cubes using related generating functions. In the subsequent section we then connect Fibonacci cubes with Fibonacci trees in a rather surprising way. Using a new labeling of the leaves of Fibonacci trees we prove that the eccentricity of a vertex of a given Fibonacci cube is equal to the depth of the associated leaf in the corresponding Fibonacci tree. Then, in Section 4, we obtain the limit average fraction between the Fibonacci and Lucas words with coordinates fixed to 0 and 1, respectively; see Theorem 4.1 for the precise statement of the result. In the final section we first compute the limit average degree of these cubes. Then we introduce the hypercube density of a family of subgraphs of hypercubes and prove that it is equal to $(1 - 1/\sqrt{5})/\log_2 \varphi$ for both Fibonacci cubes and Lucas cubes. We conclude the paper by demonstrating that the Cartesian product of graphs can be applied to construct families of graphs with a fixed, non-zero hypercube density.

The distance $d_G(u, v)$ between vertices u and v of a graph G is the number of edges on a shortest u, v -path. The *eccentricity* $\text{ecc}_G(u)$ of $u \in V(G)$ is the maximum distance between u and any other vertex of G . We will shortly write $d(u, v)$ and $\text{ecc}(u)$ when G will be clear from the context. The *average eccentricity* and the *average degree* of a graph G are respectively defined as:

$$\begin{aligned}\overline{\text{ecc}}(G) &= \frac{1}{|V(G)|} \sum_{u \in V(G)} \text{ecc}(u), \\ \overline{\text{deg}}(G) &= \frac{1}{|V(G)|} \sum_{u \in V(G)} \text{deg}(u).\end{aligned}$$

The vertex set of the n -cube Q_n is the set of all binary strings of length n , two vertices being adjacent if they differ in precisely one position. A *Fibonacci string* of length n is a binary string $b_1 \dots b_n$ with $b_i \cdot b_{i+1} = 0$ for $1 \leq i < n$. The *Fibonacci cube* Γ_n ($n \geq 1$) is the subgraph of Q_n induced by the Fibonacci strings of length n . A Fibonacci string $b_1 \dots b_n$ is a *Lucas string* if in addition $b_1 \cdot b_n = 0$ holds. The

Lucas cube Λ_n ($n \geq 1$) is the subgraph of Q_n induced by the Lucas strings of length n . For convenience we also consider the empty string and set $\Gamma_0 = K_1 = \Lambda_0$.

Let $\{F_n\}$ be the *Fibonacci numbers*: $F_0 = 0$, $F_1 = 1$, $F_n = F_{n-1} + F_{n-2}$ for $n \geq 2$. Recall that $\lim_{n \rightarrow \infty} F_{n+1}/F_n = \varphi$, where $\varphi = (1 + \sqrt{5})/2$ is the *golden ratio*. More generally, if k is a given integer, then $\lim_{n \rightarrow \infty} F_{n+k}/F_n = \varphi^k$. Let $\{L_n\}$ be the *Lucas numbers*: $L_0 = 2$, $L_1 = 1$, $L_n = L_{n-1} + L_{n-2}$ for $n \geq 2$. Recall finally that $|V(\Gamma_n)| = F_{n+2}$ and $|V(\Lambda_n)| = L_n = F_{n-1} + F_{n+1}$.

2 Average eccentricity

In this section we determine the limit average eccentricity of Fibonacci and Lucas cubes. It is intuitively rather “obvious” that in both cases the result should be the same, however, the proofs are somehow different. We begin with:

Theorem 2.1

$$\lim_{n \rightarrow \infty} \frac{\overline{\text{ecc}}(\Gamma_n)}{n} = \frac{5 + \sqrt{5}}{10}.$$

Proof. Let $f_{n,k}$ be the number of vertices of Γ_n with eccentricity k . It is proved in [2, Theorem 4.3] that the corresponding generating function is

$$F(x, y) = \sum_{n, k \geq 0} f_{n,k} x^n y^k = \frac{1 + xy}{1 - x(x+1)y}. \quad (1)$$

If e_n is the sum of the eccentricities of all vertices of Γ_n ,

$$e_n = \sum_{x \in V(\Gamma_n)} \text{ecc}(x),$$

then

$$\left. \frac{\partial F(x, y)}{\partial y} \right|_{y=1} = \sum_{n, k \geq 0} k f_{n,k} x^n = \sum_{n \geq 0} e_n x^n.$$

On the other hand,

$$\left. \frac{\partial F(x, y)}{\partial y} \right|_{y=1} = \frac{2x + x^2}{(1 - x - x^2)^2}.$$

Since $\sum_{n \geq 0} F_n x^n = \frac{x}{1 - x - x^2}$, we have

$$\begin{aligned} \sum_{n \geq 0} F_{n+1} x^n &= \frac{1}{1 - x - x^2}, \\ \sum_{n \geq 0} n F_{n+1} x^n &= \frac{x + 2x^2}{(1 - x - x^2)^2}, \text{ and} \\ \sum_{n \geq 0} n F_n x^n &= \frac{x + x^3}{(1 - x - x^2)^2}. \end{aligned}$$

Notice that

$$\frac{2x + x^2}{(1 - x - x^2)^2} = \frac{1}{5} \left(3 \frac{x}{1 - x - x^2} + 4 \frac{x + 2x^2}{(1 - x - x^2)^2} + 3 \frac{x + x^3}{(1 - x - x^2)^2} \right).$$

Therefore,

$$\sum_{n \geq 0} e_n x^n = \frac{1}{5} \left(3 \sum_{n \geq 0} F_n x^n + 4 \sum_{n \geq 0} n F_{n+1} x^n + 3 \sum_{n \geq 0} n F_n x^n \right)$$

and thus

$$e_n = \frac{3F_n + 4nF_{n+1} + 3nF_n}{5} = \frac{3F_n + nF_{n+1} + 3nF_{n+2}}{5}.$$

Therefore,

$$\overline{\text{ecc}}(\Gamma_n) = \frac{3F_n + nF_{n+1} + 3nF_{n+2}}{5F_{n+2}}.$$

We conclude that

$$\lim_{n \rightarrow \infty} \frac{\overline{\text{ecc}}(\Gamma_n)}{n} = \frac{3}{5} + \lim_{n \rightarrow \infty} \frac{1}{5} \frac{F_{n+1}}{F_{n+2}} = \frac{3}{5} + \frac{1}{5} \varphi^{-1} = \frac{5 + \sqrt{5}}{10}.$$

□

Note that $(5 + \sqrt{5})/10 \approx 0.7236$ which should be compared with the (trivial) fact that

$$\lim_{n \rightarrow \infty} \frac{\overline{\text{ecc}}(Q_n)}{n} = 1.$$

We next give the parallel result for Lucas cubes:

Theorem 2.2

$$\lim_{n \rightarrow \infty} \frac{\overline{\text{ecc}}(\Lambda_n)}{n} = \frac{5 + \sqrt{5}}{10}.$$

Proof. The proof proceeds along the same lines as the proof of Theorem 2.1, but the computations are much different, hence we give a sketch of the proof. Let $f'_{n,k}$ be the number of vertices of Λ_n with eccentricity k . We start from the generating function of this sequence, obtained in [2, Theorem 5.16]:

$$G(x, y) = \sum_{n, k \geq 0} f'_{n,k} x^n y^k = \frac{1 + x^2 y}{1 - xy - x^2 y} + \frac{1}{1 + xy} - \frac{1 - x}{1 - x^2 y}. \quad (2)$$

Let $e'_n = \sum_{x \in V(\Lambda_n)} \text{ecc}(x)$ be the sum of the eccentricities of all vertices of Λ_n . We deduce from (2) that the generating function of the sequence $\{e'_n\}$ is

$$\sum_{n \geq 0} e'_n x^n = \frac{\partial G(x, y)}{\partial y} \Big|_{y=1} = \frac{x + 2x^2}{(1 - x - x^2)^2} - \frac{x}{(1 + x)^2} - \frac{x^2}{(1 + x)(1 - x^2)}.$$

The first term is the generating function of $n F_{n+1}$, and developing the other terms we obtain

$$e'_n = n F_{n+1} + (-1)^n n + (-1)^{n+1} \left\lfloor \frac{n}{2} \right\rfloor.$$

Since $|V(\Lambda_n)| = F_{n-1} + F_{n+1}$, we conclude that

$$\lim_{n \rightarrow \infty} \frac{\overline{\text{ecc}}(\Lambda_n)}{n} = \lim_{n \rightarrow \infty} \frac{F_{n+1}}{F_{n-1} + F_{n+1}} = \frac{\varphi^2}{1 + \varphi^2} = \frac{5 + \sqrt{5}}{10}.$$

□

3 Fibonacci trees and eccentricity

In the previous section we considered the sequence e_n , where $e_n = \sum_{x \in V(\Gamma_n)} \text{ecc}(x)$. The sequence e_{n+1} starts with 2, 5, 12, 25, 50, 96, ... This is also the start of the sequence [17, Sequence A067331] described as the sum of the depth of leaves in Fibonacci trees. The two sequences indeed coincide since they have the same generating function but the connection seems mysterious. In this section we give a bijective proof that the two sequences coincide and along the way propose a new labeling of the Fibonacci trees.

Fibonacci trees have been introduced in computer science for efficient search algorithm [7, 13, 19]. They are binary trees defined recursively as follow:

- T_0 and T_1 are trees with only one vertex—the root.
- T_n , $n \geq 2$, is the rooted whose left subtree is T_{n-1} and whose right subtree is T_{n-2} .

Clearly, for any n the number of leaves of T_n is F_{n+1} .

We can recursively construct a labeling of the leaves of T_n with Fibonacci strings of length $n - 1$ as follows. Let \mathcal{F}_n be the set of Fibonacci strings of length n . Let \mathcal{F}_n^0 and \mathcal{F}_n^1 be the sets of Fibonacci strings ending with 0 and 1, respectively. We then have, for $n \geq 2$, $\mathcal{F}_n = \{s0; s \in \mathcal{F}_{n-1}\} \uplus \{s01; s \in \mathcal{F}_{n-2}\}$, where \uplus is the disjoint union of sets. First label T_1 and T_2 and assume $n \geq 3$. We add 01 to the labels of the right leaves already labeled as leaves of T_{n-2} , and 0 to the labels of the left leaves already labeled as leaves of T_{n-1} . In Fig. 1 the construction is presented for the first three non-trivial Fibonacci trees, where the currently attached strings are underlined.

The described standard labeling of Fibonacci trees does not respect the equality between depth and eccentricity. For example, the depth of the leaf of T_3 labeled 01 is 1, but $\text{ecc}_{\Gamma_2}(01) = 2$. Nevertheless, the sum of the depths of leaves of T_3 is 5 like the sum of the eccentricities of vertices of Γ_2 . We next construct a new labeling that respects the equality vertex by vertex.

Notice first that, for $n \geq 2$, $\mathcal{F}_n = \{s00; s \in \mathcal{F}_{n-2}\} \uplus \{s0; s \in \mathcal{F}_{n-1}^1\} \uplus \{s1; s \in \mathcal{F}_{n-1}^0\}$. Label T_1 with the empty string and T_2 according to Fig. 2. Assume that




Figure 1: Fibonacci trees equipped with the standard labeling

$n \geq 3$ and that the leaves of T_{n-1} and T_{n-2} were already labeled. We add 00 to the label of the right leaves. For the left leaves add 0 to the label of a leaf with a label ending with 1; otherwise append label 1. Let θ denote this labeling of the leaves of T_n by vertices of Γ_{n-1} . It is shown in Fig. 2, again for the first three non-trivial Fibonacci trees.




Figure 2: Fibonacci trees equipped with the labeling θ

The main result of this section now reads as follows:

Theorem 3.1 *Let $n \geq 1$ and let $u \in V(\Gamma_n)$. Then*

$$\text{ecc}_{\Gamma_n}(u) = \text{depth}_{T_{n+1}}(\theta^{-1}(u)).$$

Proof. We proceed by induction on n , the cases $n = 1, 2$ being trivial. Let $n \geq 3$ and let $u \in V(\Gamma_n)$. Consider the following three cases.

Suppose first that $u = v00$, where $v \in \mathcal{F}_{n-2}$. Then we claim that $\text{ecc}_{\Gamma_n}(u) = \text{ecc}_{\Gamma_{n-2}}(v) + 1$. Let u' be a vertex of Γ_n with $d_{\Gamma_n}(u, u') = \text{ecc}(u)$. Let $u' = wab$, where $w \in \mathcal{F}_{n-2}$. Since $ab \neq 11$, $d_{\Gamma_n}(u, u') \neq d_{\Gamma_{n-2}}(v, w) + 2$. Then

$$\text{ecc}_{\Gamma_n}(u) = d_{\Gamma_n}(u, u') \leq d_{\Gamma_{n-2}}(v, w) + 1 \leq \text{ecc}(v)_{\Gamma_{n-2}} + 1.$$

Conversely, let $w \in \mathcal{F}_{n-2}$, such that $d_{\Gamma_{n-2}}(v, w) = \text{ecc}(v)$. Then $w01 \in \mathcal{F}_n$ and $d_{\Gamma_n}(u, w01) = \text{ecc}_{\Gamma_{n-2}}(v) + 1$. It follows that $\text{ecc}_{\Gamma_n}(u) \geq \text{ecc}(v)_{\Gamma_{n-2}} + 1$. This proves the claim.

Suppose next that $u = v1$, where $v \in \mathcal{F}_{n-1}^0$. Now we claim that $\text{ecc}_{\Gamma_n}(u) = \text{ecc}_{\Gamma_{n-1}}(v) + 1$. The inequality $\text{ecc}_{\Gamma_n}(u) \leq \text{ecc}_{\Gamma_{n-1}}(v) + 1$ follows by an argument similar as in the first case. Conversely, let $w \in \mathcal{F}_{n-1}$, such that $d_{\Gamma_{n-1}}(v, w) = \text{ecc}(v)$. Then $w0 \in \mathcal{F}_n$ and $d_{\Gamma_n}(u, w0) = \text{ecc}_{\Gamma_{n-1}}(v) + 1$. It follows that $\text{ecc}_{\Gamma_n}(u) \geq \text{ecc}_{\Gamma_{n-1}}(v) + 1$.

Suppose finally that $u = v0$, where $v \in \mathcal{F}_{n-1}^1$. We claim again that $\text{ecc}_{\Gamma_n}(u) = \text{ecc}_{\Gamma_{n-1}}(v) + 1$. Again, the inequality $\text{ecc}_{\Gamma_n}(u) \leq \text{ecc}_{\Gamma_{n-1}}(v) + 1$ follows as above. Conversely, let $w \in \mathcal{F}_{n-1}$, such that $d_{\Gamma_{n-1}}(v, w) = \text{ecc}_{\Gamma_{n-1}}(v)$. Then w ends with 0, because otherwise w would not be an eccentric vertex of v . Indeed, if w would end with 1, then the word w' obtained from w by changing its last bit to 0 would lie in \mathcal{F}_{n-1} and hence $d_{\Gamma_{n-1}}(w', v) > d_{\Gamma_{n-1}}(w, v) = \text{ecc}_{\Gamma_{n-1}}(v)$, a contradiction. It follows that $w1 \in \mathcal{F}_n$ and $d_{\Gamma_n}(u, w1) = d_{\Gamma_{n-1}}(w, v) + 1 = \text{ecc}_{\Gamma_{n-1}}(v) + 1$.

We have thus proved that for any $u \in \mathcal{F}_n$, $\text{ecc}_{\Gamma_n}(u)$ increases by 1 with respect to the word v to which a suffix has been added to obtain u . By the construction of T_n and by the induction hypothesis,

$$\text{depth}_{T_{n+1}}(\theta^{-1}(u)) = \text{depth}_{T_{n-1}}(\theta^{-1}(v)) + 1 = \text{ecc}_{\Gamma_{n-2}}(v) + 1 = \text{ecc}_{\Gamma_n}(u)$$

in the first case, and

$$\text{depth}_{T_{n+1}}(\theta^{-1}(u)) = \text{depth}_{T_n}(\theta^{-1}(v)) + 1 = \text{ecc}_{\Gamma_{n-1}}(v) + 1 = \text{ecc}_{\Gamma_n}(u)$$

in the last two cases. \square

4 Average fractional weights

Let G be a subgraph of Q_n , so that the vertices of G are binary strings of length n . Then for $i = 1, \dots, n$ and $\chi = 0, 1$, let

$$W_{(i, \chi)}(G) = \{u = u_1 \dots u_n \in V(G) \mid u_i = \chi\}.$$

In this section we prove the following result which might be of independent interest:

Theorem 4.1

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \frac{|W_{(i,0)}(\Gamma_n)|}{|W_{(i,1)}(\Gamma_n)|} = \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \frac{|W_{(i,0)}(\Lambda_n)|}{|W_{(i,1)}(\Lambda_n)|} = \lim_{n \rightarrow \infty} \frac{|W_{(i,0)}(\Lambda_n)|}{|W_{(i,1)}(\Lambda_n)|} = \varphi^2.$$

To prove it, we will make use of the following result, see [14, Exercise 3.9.13]:

Lemma 4.2 *If $\{a_n\}$ is a convergent complex sequence with limit ℓ , then*

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n a_i a_{n+1-i} = \ell^2.$$

Proof. (of Theorem 4.1) Set $w_{i,0}^{(n)} = |W_{(i,0)}(\Gamma_n)|$, $w_{i,1}^{(n)} = |W_{(i,1)}(\Gamma_n)|$, and $x_i^{(n)} = w_{i,0}^{(n)} / w_{i,1}^{(n)}$. Notice first that the vertices of $W_{(i,0)}(\Gamma_n)$ are the string $u0v$ where u and v are arbitrary vertices of $V(\Gamma_{i-1})$ and $V(\Gamma_{n-i})$, respectively. Similarly the vertices of $W_{(i,1)}(\Gamma_n)$ are the string $u010v$, where u and v are arbitrary vertices of $V(\Gamma_{i-2})$ and $V(\Gamma_{n-i-1})$, respectively. Then, having in mind that $|V(\Gamma_n)| = F_{n+2}$, we have

$$x_i^{(n)} = \frac{F_{i+1} \cdot F_{n-i+2}}{F_i \cdot F_{n-i+1}}.$$

Setting $a_n = F_{n+1}/F_n$, recalling that $\{a_n\} \rightarrow \varphi$, and using Lemma 4.2, we get

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n x_i^{(n)} = \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n a_i a_{n-i+1} = \varphi^2,$$

which proves the result for Fibonacci cubes.

The proof for Lucas cubes proceeds along the same lines as the above proof for Fibonacci cubes. The only difference is that now $x_i^{(n)} = F_{n+1}/F_{n-1}$, and hence $\lim_{n \rightarrow \infty} x_i^{(n)} = \varphi^2$. Then the classical Cesàro Means Theorem (it asserts that if $\lim_{n \rightarrow \infty} a_n = \ell$, then $\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n a_i = \ell$ as well) can be applied instead of Lemma 4.2. \square

5 Average degree and density

In this section we first compute the limit average degree of the considered graphs. The result then motivates us to introduce the hypercube density, to determine it for the cubes, and to show that the Cartesian product of graphs is useful in the context.

Theorem 5.1

$$\lim_{n \rightarrow \infty} \frac{\overline{\deg}(\Gamma_n)}{n} = \lim_{n \rightarrow \infty} \frac{\overline{\deg}(\Lambda_n)}{n} = \frac{5 - \sqrt{5}}{5}.$$

Proof. It was proved in [15] that $|E(\Gamma_n)| = (nF_{n+1} + 2(n+1)F_n)/5$, hence

$$\overline{\deg}(\Gamma_n) = \frac{1}{|V(\Gamma_n)|} 2|E(\Gamma_n)| = \frac{2}{5F_{n+2}} (nF_{n+1} + 2(n+1)F_n).$$

Therefore,

$$\lim_{n \rightarrow \infty} \frac{\overline{\deg}(\Gamma_n)}{n} = \lim_{n \rightarrow \infty} \frac{2}{5} \left(\frac{F_{n+1}}{F_{n+2}} + \frac{2F_n}{F_{n+2}} + \frac{2}{n} \frac{F_n}{F_{n+2}} \right) = \frac{2}{5} (\varphi^{-1} + 2\varphi^{-2}) = \frac{5 - \sqrt{5}}{5}.$$

For the Lucas cubes there is no such nice expression for the number of edges as the above for Fibonacci cubes. Hence we use the “weighted approach” (as in the proof of Theorem 4.1) to prove the result for Lucas cubes.

Let $w_{i,1}^{(n)} = |W_{(i,1)}(\Lambda_n)|$. If $uv \in E(\Lambda_n)$, then u and v differ in exactly one coordinate, say in the i -th, and assume without loss of generality that $u_i = 1$ (and so $v_i = 0$). Then assign the bit $u_i = 1$ to the edge uv and note that this assignment is bijective. It follows that $|E(\Lambda_n)| = \sum_{i=1}^n w_{i,1}^{(n)}$ and therefore

$$\overline{\deg}(\Lambda_n) = \frac{1}{|V(\Lambda_n)|} 2|E(\Lambda_n)| = \frac{2}{L_n} \sum_{i=1}^n w_{i,1}^{(n)}.$$

Observe that $w_{i,1}^{(n)} = |V(\Gamma_{n-3})| = F_{n-1}$ holds for any i . Hence, using the fact that $L_n = F_{n-1} + F_{n+1}$, we get:

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{\overline{\deg}(\Lambda_n)}{n} &= \lim_{n \rightarrow \infty} \frac{2}{nL_n} \sum_{i=1}^n w_{i,1}^{(n)} \\ &= \lim_{n \rightarrow \infty} \frac{2F_{n-1}}{F_{n-1} + F_{n+1}} \\ &= 2(1 + \varphi^2)^{-1} = \frac{5 - \sqrt{5}}{5}. \end{aligned}$$

□

Graham [4] proved the following fundamental property of subgraphs of hypercubes (see [5, Lemma 3.2] for an alternative proof of it):

Lemma 5.2 (Density Lemma) *Let G be a subgraph of a hypercube. Then*

$$|E(G)| \leq \frac{1}{2} |V(G)| \cdot \log_2 |V(G)|.$$

Moreover, equality holds if and only if G is a hypercube.

The lemma has important consequences, in particular for fast recognition algorithms for classes of subgraphs of hypercubes; see [18] for the case of Fibonacci cubes. The Density Lemma also asserts that hypercubes have the largest density among all subgraphs of hypercubes. We therefore introduce the following concept.

If G is a subgraph of a hypercube, then let

$$\rho(G) = \frac{\overline{\deg}(G)}{\log_2 |V(G)|}.$$

Let $\mathcal{G} = \{G_k\}_{k \geq 1}$ be an *increasing family* of subgraphs of hypercubes, that is, a family with $|V(G_{n+1})| > |V(G_n)|$ for $n \geq 1$. Then the *hypercube density* of \mathcal{G} is

$$\rho(\mathcal{G}) = \limsup_{k \rightarrow \infty} \rho(G_k).$$

By the Density Lemma, $0 \leq \rho(\{G_k\}) \leq 1$ holds for any family $\{G_k\}$ and

$$\rho(\{Q_k\}) = 1.$$

For Fibonacci cubes and Lucas cubes we have:

Corollary 5.3

$$\rho(\{\Gamma_n\}) = \rho(\{\Lambda_n\}) = \frac{5 - \sqrt{5}}{5 \log_2 \varphi}.$$

Proof. The result easily follows from Theorem 5.1 together with the facts that $F_n \rightarrow \varphi^n / \sqrt{5}$ and that $\Lambda_n \rightarrow \varphi^n$. \square

Hence $\rho(\{\Gamma_n\}) \approx 0.7962$ which is in particular interesting because the hypercube density of many other important families of hypercube subgraphs is 0. For instance, if WB_k denotes the bipartite wheel with k spokes, then an easy calculation shows that $\rho(\{WB_k\}) = 0$. (Cf. [1] for the role of bipartite wheels among subgraphs of hypercubes.) For another example consider the subdivision $S(K_k)$ of the complete graph K_k , that is, the graph obtained from K_k by subdividing each of its edges precisely once. These graphs embed isometrically into hypercubes (cf. [11]) and

$$\rho(\{S(K_k)\}) = \lim_{k \rightarrow \infty} \rho(S(K_k)) = \lim_{k \rightarrow \infty} \frac{2 \left(2^{\binom{k}{2}} \right)}{\left(k + \binom{k}{2} \right) \log_2 \left(k + \binom{k}{2} \right)} = 0.$$

Families of graphs with bounded degree also have density equal to 0. More precisely:

Proposition 5.4 *Let $\{G_k\}$ be an increasing family of hypercube subgraphs. If there exists a constant M such that $\Delta(G_k) \leq M$ for any $k \geq 1$, then $\rho(\{G_k\}) = 0$.*

Proof. For $k \geq 1$ set $n_k = |V(G_k)|$ and $m_k = |E(G_k)|$. Then from the Handshaking Lemma it follows that $2m_k = \sum_{u \in V(G_k)} \deg(u) \leq n_k M$ and hence $2m_k/n_k \leq M$. The assertion then follows because $\log_2(n_k) \rightarrow \infty$. \square

On the other hand, the Cartesian product of graphs can be used to obtain families with positive hypercube density. If G^k denotes the k -tuple Cartesian product of G , then we have:

Proposition 5.5 *Let G be a hypercube subgraph with $\rho(G) = c$. Then $\rho(\{G^k\}) = c$.*

Proof. Let $n = |V(G)|$ and $m = |E(G)|$. Then it is easily shown by induction that $|V(G^k)| = n^k$ and $|E(G^k)| = kn^{k-1}m$. Consequently,

$$\rho(\{G^k\}) = \lim_{k \rightarrow \infty} \rho(G^k) = \lim_{k \rightarrow \infty} \frac{2kn^{k-1}m}{n^k \log_2(n^k)} = \frac{2m}{n \log_2(n)} = \rho(G).$$

\square

Suppose that a family $\{G_k\}$ is given with $\rho(\{G_k\}) = c$. Then we can again use the Cartesian product of graphs to obtain an infinite number of families with the same density.

Proposition 5.6 *Let $\rho(\{G_k\}) = c$ and let G be a fixed subgraph of a hypercube. Then $\rho(\{G_k \square G\}) = c$.*

Proof. Let $n = |V(G)|$, $m = |E(G)|$, and for $k \geq 1$, let $n_k = |V(G_k)|$ and $m_k = |E(G_k)|$. Then we have

$$\begin{aligned}
\rho(\{G_k \square G\}) &= \lim_{k \rightarrow \infty} \rho(G_k \square G) \\
&= \lim_{k \rightarrow \infty} \frac{2(n m_k + n_k m)}{n_k n \log_2(n_k n)} \\
&= \lim_{k \rightarrow \infty} \frac{2m_k}{n_k \log_2(n_k n)} + \lim_{k \rightarrow \infty} \frac{2m}{n \log_2(n_k n)} \\
&= \lim_{k \rightarrow \infty} \left(\frac{2m_k}{n_k \log_2(n_k)} \cdot \frac{\log_2(n_k)}{\log_2(n_k) + \log_2(n)} \right) \\
&= \lim_{k \rightarrow \infty} \frac{2m_k}{n_k \log_2(n_k)} = \rho(\{G_k\}).
\end{aligned}$$

□

Acknowledgments

This work was supported in part by the Proteus project BI-FR/12-139-PROTEUS-008, by ARRS Slovenia under the grant P1-0297, and within the EUROCORES Programme EUROGIGA/GReGAS of the European Science Foundation.

References

- [1] H.-J. Bandelt, V. Chepoi, Graphs of acyclic cubical complexes, *European J. Combin.* 17 (1996) 113–120.
- [2] A. Castro, M. Mollard, The eccentricity sequences of Fibonacci and Lucas cubes, *Discrete Math.* 312 (2012) 1025–1037.
- [3] P. Dankelmann, W. Goddard, C.S. Swart, The average eccentricity of a graph and its subgraphs, *Util. Math.* 65 (2004) 41–51.
- [4] R. L. Graham, On primitive graphs and optimal vertex assignments, *Ann. New York Acad. Sci.* 175 (1970) 170–186.
- [5] R. Hammack, W. Imrich, S. Klavžar, *Handbook of Product Graphs*, Second Edition, CRC Press, Boca Raton, FL, 2011.
- [6] A.M. Hinz, D. Parisse, The average eccentricity of Sierpiński graphs, *Graphs Combin.* 28 (2012) 671–686.

- [7] Y. Horibe, An entropy view of Fibonacci trees, *Fibonacci Quart.* 20 (1982) 168–178.
- [8] W.-J. Hsu, Fibonacci cubes—a new interconnection technology, *IEEE Trans. Parallel Distrib. Syst.* 4 (1993) 3–12.
- [9] A. Ilić, On the extremal properties of the average eccentricity, *Comp. Math. Appl.* 64 (2012) 2877–2885.
- [10] S. Klavžar, Structure of Fibonacci cubes: a survey, *J. Comb. Optim.* 25 (2013) 505–522.
- [11] S. Klavžar, A. Lipovec, Partial cubes as subdivision graphs and as generalized Petersen graphs, *Discrete Math.* 263 (2003) 157–165.
- [12] S. Klavžar, M. Mollard, Wiener index and Hosoya polynomial of Fibonacci and Lucas cubes, *MATCH Commun. Math. Comput. Chem.* 68 (2012) 311–324.
- [13] D.E. Knuth, *The Art of Computer Programming*, Vol. 3, 2nd edition, Addison-Wesley, Reading, MA, 1998, p. 417.
- [14] J.-M. Monier, *Analyse*, Tome 1, Dunod, 1990.
- [15] E. Munarini, C. Perelli Cippo, N. Zagaglia Salvi, On the Lucas cubes, *Fibonacci Quart.* 39 (2001) 12–21.
- [16] M. Ramras, Routing linear permutations on Fibonacci and Lucas cubes, manuscript, arXiv:1207.1518v1 [math.CO].
- [17] N.J.A. Sloane, The On-Line Encyclopedia of Integer Sequences, Published electronically at <http://oeis.org>, 2013.
- [18] A. Taranenko, A. Vesel, Fast recognition of Fibonacci cubes, *Algorithmica* 49 (2007) 81–93.
- [19] S.G. Wagner, The Fibonacci number of Fibonacci trees and a related family of polynomial recurrence systems, *Fibonacci Quart.* 45 (2007) 247–253.
- [20] P. Žigert Pleteršek, M. Berlič, Resonance graphs of armchair nanotubes cyclic polypyrenes and amalgams of Lucas cubes, *MATCH Commun. Math. Comput. Chem.* 70 (2013) 533–543.
- [21] H. Zhang, L. Ou, H. Yao, Fibonacci-like cubes as Z -transformation graphs, *Discrete Math.* 309 (2009) 1284–1293.