



HAL
open science

Cremona groups of real surfaces

Jérémy Blanc, Frédéric Mangolte

► **To cite this version:**

Jérémy Blanc, Frédéric Mangolte. Cremona groups of real surfaces. Automorphisms in Birational and Affine Geometry - Levico Terme, Italy, October 2012, 79, Springer, pp.35–58, 2014, Springer Proceedings in Mathematics & Statistics. hal-00838522

HAL Id: hal-00838522

<https://hal.science/hal-00838522>

Submitted on 25 Jun 2013

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

CREMONA GROUPS OF REAL SURFACES

JÉRÉMY BLANC AND FRÉDÉRIC MANGOLTE

ABSTRACT. We give an explicit set of generators for various natural subgroups of the real Cremona group $\text{Bir}_{\mathbb{R}}(\mathbb{P}^2)$. This completes and unifies former results by several authors.

MSC 2000: 14E07, 14P25, 14J26

Keywords: real algebraic surface, rational surface, birational geometry, algebraic automorphism, Cremona transformation

1. INTRODUCTION

1.1. **On the real Cremona group** $\text{Bir}_{\mathbb{R}}(\mathbb{P}^2)$. The classical Noether-Castelnuovo Theorem (1917) gives generators of the group $\text{Bir}_{\mathbb{C}}(\mathbb{P}^2)$ of birational transformations of the complex projective plane. The group is generated by the biregular automorphisms, which form the group $\text{Aut}_{\mathbb{C}}(\mathbb{P}^2) \cong \text{PGL}(3, \mathbb{C})$ of projectivities, and by the standard quadratic transformation

$$\sigma_0: (x : y : z) \dashrightarrow (yz : xz : xy).$$

This result does not work over the real numbers. Indeed, recall that a *base point* of a birational transformation is a (possibly infinitely near) point of indeterminacy; and note that the base points of the quadratic involution

$$\sigma_1: (x : y : z) \dashrightarrow (y^2 + z^2 : xy : xz)$$

are not real. Thus σ_1 cannot be generated by projectivities and σ_0 . More generally, we cannot generate this way maps having non real base-points. Hence the group $\text{Bir}_{\mathbb{R}}(\mathbb{P}^2)$ of birational transformations of the real projective plane is not generated by $\text{Aut}_{\mathbb{R}}(\mathbb{P}^2) \cong \text{PGL}(3, \mathbb{R})$ and σ_0 .

The first result of this note is that $\text{Bir}_{\mathbb{R}}(\mathbb{P}^2)$ is generated by $\text{Aut}_{\mathbb{R}}(\mathbb{P}^2)$, σ_0 , σ_1 , and a family of birational maps of degree 5 having only non real base-points.

Theorem 1.1. *The group $\text{Bir}_{\mathbb{R}}(\mathbb{P}^2)$ is generated by $\text{Aut}_{\mathbb{R}}(\mathbb{P}^2)$, σ_0 , σ_1 , and the standard quintic transformations of \mathbb{P}^2 (defined in Example 3.1).*

The proof of this result follows the so-called Sarkisov program, which amounts to decompose a birational map between Mori fiber spaces as a sequence of simple maps, called *Sarkisov links*. The description of all possible links has been done in [Isko96] for perfect fields, and in [Poly97] for real surfaces. We recall it in Section 2 and show how to deduce Theorem 1.1 from the list of Sarkisov links.

Let X be an algebraic variety defined over \mathbb{R} , we denote as usual by $X(\mathbb{R})$ the set of real points endowed with the induced algebraic structure. The topological space $\mathbb{P}^2(\mathbb{R})$ is then the real projective plane, letting $\mathbb{F}_0 := \mathbb{P}^1 \times \mathbb{P}^1$, the space $\mathbb{F}_0(\mathbb{R})$

First author supported by the SNSF grant no PP00P2_128422 /1.

This research was partially supported by ANR Grant "BirPol" ANR-11-JS01-004-01.

is the torus $\mathbb{S}^1 \times \mathbb{S}^1$ and letting $Q_{3,1} = \{(w : x : y : z) \in \mathbb{P}^3 \mid w^2 = x^2 + y^2 + z^2\}$, the real locus $Q_{3,1}(\mathbb{R})$ is the sphere \mathbb{S}^2 .

Recall that an *automorphism* of $X(\mathbb{R})$ is a birational transformation $\varphi \in \text{Bir}_{\mathbb{R}}(X)$ such that φ and φ^{-1} are defined at all real points of X . The set of such maps form a group $\text{Aut}(X(\mathbb{R}))$, and we have natural inclusions

$$\text{Aut}_{\mathbb{R}}(X) \subset \text{Aut}(X(\mathbb{R})) \subset \text{Bir}_{\mathbb{R}}(X).$$

The strategy used to prove Theorem 1.1 allows us to treat similarly the case of natural subgroups of $\text{Bir}_{\mathbb{R}}(\mathbb{P}^2)$, namely the groups $\text{Aut}(\mathbb{P}^2(\mathbb{R}))$, $\text{Aut}(Q_{3,1}(\mathbb{R}))$ and $\text{Aut}(\mathbb{F}_0(\mathbb{R}))$ of the three *minimal* real rational surfaces (see 2.8). This way, we give a unified treatment to prove three theorems on generators, the first two of them already proved in a different way in [RV05] and [KM09].

Observe that $\text{Aut}(Q_{3,1}(\mathbb{R}))$ and $\text{Aut}(\mathbb{F}_0(\mathbb{R}))$ are not really subgroups of $\text{Bir}_{\mathbb{R}}(\mathbb{P}^2)$, but each of them is isomorphic to a subgroup which is determined up to conjugation. Indeed, for any choice of a birational map $\psi: \mathbb{P}^2 \dashrightarrow X$ ($X = Q_{3,1}$ or \mathbb{F}_0), $\psi^{-1} \text{Aut}(X(\mathbb{R}))\psi \subset \text{Bir}_{\mathbb{R}}(\mathbb{P}^2)$.

Theorem 1.2 ([RV05]). *The group $\text{Aut}(\mathbb{P}^2(\mathbb{R}))$ is generated by $\text{Aut}_{\mathbb{R}}(\mathbb{P}^2) = \text{PGL}(3, \mathbb{R})$ and by standard quintic transformations.*

Theorem 1.3 ([KM09]). *The group $\text{Aut}(Q_{3,1}(\mathbb{R}))$ is generated by $\text{Aut}_{\mathbb{R}}(Q_{3,1}) = \text{PO}(3, 1)$ and by standard cubic transformations.*

Theorem 1.4. *The group $\text{Aut}(\mathbb{F}_0(\mathbb{R}))$ is generated by $\text{Aut}_{\mathbb{R}}(\mathbb{F}_0) = \text{PGL}(2, \mathbb{R})^2 \rtimes \mathbb{Z}/2\mathbb{Z}$ and by the involution*

$$\tau_0 : ((x_0 : x_1), (y_0 : y_1)) \dashrightarrow ((x_0 : x_1), (x_0 y_0 + x_1 y_1 : x_1 y_0 - x_0 y_1)).$$

The proof of theorems 1.1, 1.2, 1.3, 1.4 is given in Sections 4, 3, 5, 6, respectively. Section 7 is devoted to present some related recent results on birational geometry of real projective surfaces.

In the sequel, surfaces and maps are assumed to be real. In particular if we consider that a real surface is a complex surface endowed with a Galois-action of $G := \text{Gal}(\mathbb{C}|\mathbb{R})$, a map is G -equivariant. On the contrary, points and curves are not assumed to be real a priori.

2. MORI THEORY FOR REAL RATIONAL SURFACES AND SARKISOV PROGRAM

We work with the tools of Mori theory. A good reference in dimension 2, over any perfect field, is [Isko96]. The theory, applied to smooth projective real rational surfaces, becomes really simple. The description of Sarkisov links between real rational surfaces has been done in [Poly97], together with a study of relations between these links. In order to state this classification, we first recall the following classical definitions (which can be found in [Isko96]).

Definition 2.1. A smooth projective real rational surface X is said to be *minimal* if any birational morphism $X \rightarrow Y$, where Y is another smooth projective real surface, is an isomorphism.

Definition 2.2. A *Mori fibration* is a morphism $\pi: X \rightarrow W$ where X is a smooth projective real rational surface and one of the following occurs

- (1) $\rho(X) = 1$, W is a point (usually denoted $\{*\}$), and X is a del Pezzo surface;
- (2) $\rho(X) = 2$, $W = \mathbb{P}^1$ and the map π is a conic bundle.

Note that for an arbitrary surface, the curve W in the second case should be any smooth curve, but we restrict ourselves to rational surfaces which implies that W is isomorphic to \mathbb{P}^1 .

Proposition 2.3. *Let X be a smooth projective real rational surface. If X is minimal, then it admits a morphism $\pi: X \rightarrow W$ which is a Mori fibration.*

Proof. Follows from [Isko79]. See also [Mori82]. \square

Definition 2.4. A *Sarkisov link* between two Mori fibrations $\pi_1: X_1 \rightarrow W_1$ and $\pi_2: X_2 \rightarrow W_2$ is a birational map $\varphi: X_1 \dashrightarrow X_2$ of one of the following four types, where each of the diagrams is commutative:

(1) LINK OF TYPE I

$$\begin{array}{ccc} X_1 & \xrightarrow{\varphi} & X_2 \\ \pi_1 \downarrow & & \downarrow \pi_2 \\ \{*\} = W_1 & \xleftarrow{\tau} & W_2 = \mathbb{P}^1 \end{array}$$

where $\varphi^{-1}: X_2 \rightarrow X_1$ is a birational morphism, which is the blow-up of either a real point or two imaginary conjugate points of X_1 , and where τ is the contraction of $W_2 = \mathbb{P}^1$ to the point W_1 .

(2) LINK OF TYPE II

$$\begin{array}{ccccc} X_1 & \xleftarrow{\sigma_1} & Z & \xrightarrow{\sigma_2} & X_2 \\ \pi_1 \downarrow & & \downarrow \varphi & & \downarrow \pi_2 \\ W_1 & \xrightarrow{\tau} & & & W_2 \end{array}$$

where $\sigma_i: Z \rightarrow X_i$ is a birational morphism, which is the blow-up of either a real point or two imaginary conjugate points of X_i , and where τ is an isomorphism between W_1 and W_2 .

(3) LINK OF TYPE III

$$\begin{array}{ccc} X_1 & \xrightarrow{\varphi} & X_2 \\ \pi_1 \downarrow & & \downarrow \pi_2 \\ \mathbb{P}^1 = W_1 & \xrightarrow{\tau} & W_2 = \{*\} \end{array}$$

where $\varphi: X_1 \rightarrow X_2$ is a birational morphism, which is the blow-up of either a real point or two imaginary conjugate points of X_2 , and where τ is the contraction of $W_1 = \mathbb{P}^1$ to the point W_2 . (*It is the inverse of a link of type I.*)

(4) LINK OF TYPE IV

$$\begin{array}{ccc} X_1 & \xrightarrow{\varphi} & X_2 \\ \pi_1 \downarrow & & \downarrow \pi_2 \\ \mathbb{P}^1 = W_1 & & W_2 = \mathbb{P}^1 \end{array}$$

where $\varphi: X_1 \rightarrow X_2$ is an isomorphism and $\pi_1, \pi_2 \circ \varphi$ are conic bundles on X_1 with distinct fibres.

Note that the morphism τ is important only for links of type II, between two surfaces with a Picard group of rank 2 (in higher dimension τ is important also for other links).

Definition 2.5. If $\pi: X \rightarrow W$ and $\pi': X' \rightarrow W'$ are two (Mori) fibrations, an isomorphism $\psi: X \rightarrow X'$ is called an *isomorphism of fibrations* if there exists an isomorphism $\tau: W \rightarrow W'$ such that $\pi'\psi = \tau\pi$.

Note that the composition $\alpha\varphi\beta$ of a Sarkisov link φ with some automorphisms of fibrations α and β is again a Sarkisov link. We have the following fundamental result:

Proposition 2.6. *If $\pi: X \rightarrow W$ and $\pi': X' \rightarrow W'$ are two Mori fibrations, then any birational map $\psi: X \dashrightarrow X'$ is either an isomorphism of fibrations or decomposes into Sarkisov links.*

Proof. Follows from [Isko96]. See also [Cort95]. \square

Theorem 2.7 ([Com12] (see also [Isko79])). *Let X be a real rational surface, if X is minimal, then it is isomorphic to one of the following:*

- (1) \mathbb{P}^2 ,
- (2) the quadric $Q_{3,1} = \{(w : x : y : z) \in \mathbb{P}^3 \mid w^2 = x^2 + y^2 + z^2\}$,
- (3) a Hirzebruch surface $\mathbb{F}_n = \{((x : y : z), (u : v)) \in \mathbb{P}^2 \times \mathbb{P}^1 \mid yv^n = zu^n\}$ with $n \neq 1$.

By [Mang06], if $n - n' \equiv 0 \pmod{2}$, $\mathbb{F}_n(\mathbb{R})$ is isomorphic to $\mathbb{F}_{n'}(\mathbb{R})$, we get:

Corollary 2.8. *Let $X(\mathbb{R})$ be the real locus of a real rational surface. If X is minimal, then $X(\mathbb{R})$ is isomorphic to one of the following:*

- (1) $\mathbb{P}^2(\mathbb{R})$,
- (2) $Q_{3,1}(\mathbb{R}) \sim \mathbb{S}^2$,
- (3) $\mathbb{F}_0(\mathbb{R}) \sim \mathbb{S}^1 \times \mathbb{S}^1$.

We give a list of Mori fibrations on real rational surfaces, and will show that, up to isomorphisms of fibrations, this list is exhaustive.

Example 2.9. The following morphisms $\pi: X \rightarrow W$ are Mori fibrations on the plane, the sphere, the Hirzebruch surfaces, and a particular Del Pezzo surface of degree 6.

- (1) $\mathbb{P}^2 \rightarrow \{*\}$;
- (2) $Q_{3,1} = \{(w : x : y : z) \in \mathbb{P}_{\mathbb{R}}^3 \mid w^2 = x^2 + y^2 + z^2\} \rightarrow \{*\}$;
- (3) $\mathbb{F}_n = \{((x : y : z), (u : v)) \in \mathbb{P}^2 \times \mathbb{P}^1 \mid yv^n = zu^n\} \rightarrow \mathbb{P}^1$ for $n \geq 0$ (the map is the projection on the second factor and \mathbb{F}_n is the n -th Hirzebruch surface);
- (4) $\mathcal{D}_6 = \{(w : x : y : z), (u : v) \in Q_{3,1} \times \mathbb{P}^1 \mid wv = xu\} \rightarrow \mathbb{P}^1$ (the map is the projection on the second factor).

Example 2.10. The following maps between the surfaces of Example 2.9 are Sarkisov links:

- (1) The contraction of the exceptional curve of \mathbb{F}_1 (or equivalently the blow-up of a real point of \mathbb{P}^2), is a link $\mathbb{F}_1 \rightarrow \mathbb{P}^2$ of type III. Note that the converse of this link is of type I.

- (2) The stereographic projection from the North pole $p_N = (1 : 0 : 0 : 1)$, $\varphi: Q_{3,1} \dashrightarrow \mathbb{P}^2$ given by $\varphi: (w : x : y : z) \dashrightarrow (x : y : w - z)$ and its inverse $\varphi^{-1}: \mathbb{P}^2 \dashrightarrow Q_{3,1}$ given by $\varphi^{-1}: (x : y : z) \dashrightarrow (x^2 + y^2 + z^2 : 2xz : 2yz : x^2 + y^2 - z^2)$ are both Sarkisov links of type II.
- The map φ decomposes into the blow-up of p_N , followed by the contraction of the strict transform of the curve $z = w$ (intersection of $Q_{3,1}$ with the tangent plane at p_N), which is the union of two imaginary conjugate lines. The map φ^{-1} decomposes into the blow-up of the two imaginary points $(1 : \pm i : 0)$, followed by the contraction of the strict transform of the line $z = 0$.
- (3) The projection on the first factor $\mathcal{D}_6 \rightarrow Q_{3,1}$ which contracts the two disjoint conjugate imaginary (-1) -curves $(0 : 0 : 1 : \pm i) \times \mathbb{P}^1 \subset \mathcal{D}_6$ onto the two conjugate imaginary points $(0 : 0 : 1 : \pm i) \in Q_{3,1}$ is a link of type III.
- (4) The blow-up of a real point $q \in \mathbb{F}_n$, lying on the exceptional section if $n > 0$ (or any point if $n = 0$), followed by the contraction of the strict transform of the fibre passing through q onto a real point of \mathbb{F}_{n+1} not lying on the exceptional section is a link $\mathbb{F}_n \dashrightarrow \mathbb{F}_{n+1}$ of type II.
- (5) The blow-up of two conjugate imaginary points $p, \bar{p} \in \mathbb{F}_n$ lying on the exceptional section if $n > 0$, or on the same section of self-intersection 0 if $n = 0$, followed by the contraction of the strict transform of the fibres passing through p, \bar{p} onto two imaginary conjugate points of \mathbb{F}_{n+2} not lying on the exceptional section is a link $\mathbb{F}_n \dashrightarrow \mathbb{F}_{n+2}$ of type II.
- (6) The blow-up of two conjugate imaginary points $p, \bar{p} \in \mathbb{F}_n$, $n \in \{0, 1\}$ not lying on the same fibre (or equivalently not lying on a real fibre) and not on the same section of self-intersection $-n$ (or equivalently not lying on a real section of self-intersection $-n$), followed by the contraction of the fibres passing through p, \bar{p} onto two imaginary conjugate points of \mathbb{F}_n having the same properties is a link $\mathbb{F}_n \dashrightarrow \mathbb{F}_n$ of type II.
- (7) The exchange of the two components $\mathbb{P}^1 \times \mathbb{P}^1 \rightarrow \mathbb{P}^1 \times \mathbb{P}^1$ is a link $\mathbb{F}_0 \rightarrow \mathbb{F}_0$ of type IV.
- (8) The blow-up of a real point $p \in \mathcal{D}_6$, not lying on a singular fibre (or equivalently $p \neq ((1 : 1 : 0 : 0), (1 : 1))$, $p \neq ((1 : -1 : 0 : 0), (1 : -1))$), followed by the contraction of the strict transform of the fibre passing through p onto a real point of \mathcal{D}_6 , is a link $\mathcal{D}_6 \dashrightarrow \mathcal{D}_6$ of type II.
- (9) The blow-up of two imaginary conjugate points $p, \bar{p} \in \mathcal{D}_6$, not lying on the same fibre (or equivalently not lying on a real fibre), followed by the contraction of the strict transform of the fibres passing through p, \bar{p} onto two imaginary points of \mathcal{D}_6 is a link $\mathcal{D}_6 \dashrightarrow \mathcal{D}_6$ of type II.

Remark 2.11. Note that in the above list, the three links $\mathbb{F}_n \dashrightarrow \mathbb{F}_m$ of type II can be put in one family, and the same is true for the two links $\mathcal{D}_6 \dashrightarrow \mathcal{D}_6$. We distinguished here the possibilities for the base points to describe more precisely the geometry of each link. The two links $\mathcal{D}_6 \dashrightarrow \mathcal{D}_6$ could also be arranged into extra families, by looking if the base points belong to the two exceptional sections of self-intersection -1 , but go in any case from \mathcal{D}_6 to \mathcal{D}_6 .

Proposition 2.12. *Any Mori fibration $\pi: X \rightarrow W$, where X is a smooth projective real rational surface, belongs to the list of Example 2.9.*

Any Sarkisov link between two such Mori fibrations is equal to $\alpha\varphi\beta$, where φ or φ^{-1} belongs to the list described in Example 2.10 and where α and β are isomorphisms of fibrations.

Proof. Since any birational map between two surfaces with Mori fibrations decomposes into Sarkisov links and that all links of Example 2.10 involve only the Mori fibrations of Example 2.9, it suffices to check that any link starting from one of the Mori fibrations of 2.9 belongs to the list 2.10. This is an easy case-by-case study; here are the steps.

Starting from a Mori fibration $\pi: X \rightarrow W$ where W is a point, the only links we can perform are links of type I or II centered at a real point or two conjugate imaginary points. From 2.7, the surface X is either $Q_{3,1}$ or \mathbb{P}^2 , and both are homogeneous under the action of $\text{Aut}(X)$, so the choice of the point is not relevant. Blowing-up a real point in \mathbb{P}^2 or two imaginary points in $Q_{3,1}$ gives rise to a link of type I to \mathbb{F}_1 or \mathcal{D}_6 . The remaining cases correspond to the stereographic projection $Q_{3,1} \dashrightarrow \mathbb{P}^2$ and its converse.

Starting from a Mori fibration $\pi: X \rightarrow W$ where $W = \mathbb{P}^1$, we have additional possibilities. If the link is of type IV, then X admits two conic bundle structures and by 2.7, the only possibility is $\mathbb{F}_0 = \mathbb{P}^1 \times \mathbb{P}^1$. If the link is of type III, then we contract a real (-1) -curve of X or two disjoint conjugate imaginary (-1) -curves. The only possibilities for X are respectively \mathbb{F}_1 and \mathcal{D}_6 , and the image is respectively \mathbb{P}^2 and $Q_{3,1}$ (these are the inverses of the links described before). The last possibility is to perform a link a type II, by blowing up a real point or two conjugate imaginary points, on respectively one or two smooth fibres, and to contract the strict transform. We go from \mathcal{D}_6 to \mathcal{D}_6 or from \mathbb{F}_m to $\mathbb{F}_{m'}$ where $m' - m \in \{-2, -1, -0, 1, 1\}$. All possibilities are described in Example 2.10. \square

We end this section by reducing the number of links of type II needed for the classification. For this, we introduce the notion of standard links.

Definition 2.13. The following links of type II are called *standard*:

- (1) links $\mathbb{F}_m \dashrightarrow \mathbb{F}_n$, with $m, n \in \{0, 1\}$;
- (2) links $\mathcal{D}_6 \dashrightarrow \mathcal{D}_6$ which do not blow-up any point on the two exceptional section of self-intersection -1 .

The other links of type II will be called *special*.

The following result allows us to simplify the set of generators of our groups.

Lemma 2.14. Any Sarkisov link of type IV decomposes into links of type I, III, and standard links of type II.

Proof. Note that a link of type IV is, up to automorphisms preserving the fibrations, equal to the following automorphism of $\mathbb{P}^1 \times \mathbb{P}^1$

$$\tau: ((x_1 : x_2), (y_1 : y_2)) \mapsto ((y_1 : y_2), (x_1 : x_2)).$$

We denote by $\psi: \mathbb{P}^2 \dashrightarrow \mathbb{P}^1 \times \mathbb{P}^1$ the birational map $(x : y : z) \dashrightarrow ((x : y), (x : z))$ and observe that $\tau\psi = \psi\sigma$, where $\sigma \in \text{Aut}_{\mathbb{R}}(\mathbb{P}^2)$. Hence, $\tau = \psi\tau\psi^{-1}$. Observing that ψ decomposes into the blow-up of the point $(0 : 0 : 1)$, which is a link of type III, followed by a standard link of type II, we get the result. \square

Lemma 2.15. Let $\pi: X \rightarrow \mathbb{P}^1$ and $\pi': X' \rightarrow \mathbb{P}^1$ be two Mori fibrations, where X, X' belong to the list $\mathbb{F}_0, \mathbb{F}_1, \mathcal{D}_6$. Let $\psi: X \dashrightarrow X'$ be a birational map, such

that $\pi'\psi = \alpha\pi$ for some $\alpha \in \text{Aut}_{\mathbb{R}}(\mathbb{P}^1)$. Then, ψ is either an automorphism or $\psi = \varphi_n \cdots \varphi_1$, where each φ_i is a sequence of standard links of type II. Moreover, if ψ is an isomorphism on the real points (i.e. is an isomorphism $X(\mathbb{R}) \rightarrow X'(\mathbb{R})$), the standard links φ_i can also be chosen to be isomorphisms on the real points.

Proof. We first show that $\psi = \varphi_n \cdots \varphi_1$, where each φ_i is a sequence of links of type II, not necessarily standard. This is done by induction on the number of base-points of ψ . If ψ has no base-point, it is an isomorphism. If q is a real proper base-point, or q, \bar{q} are two proper imaginary base-points (here proper means not infinitely near), we denote by φ_1 a Sarkisov link of type II centered at q (or q, \bar{q}). Then, $(\varphi_1)^{-1}\psi$ has less base-points than ψ . The result follows then by induction. Moreover, if ψ is an isomorphism on the real points, i.e. if ψ and ψ^{-1} have no real base-point, then so are all φ_i .

Let $\varphi: \mathcal{D}_6 \dashrightarrow \mathcal{D}_6$ be a special link of type II. Then, it is centered at two points p_1, \bar{p}_1 lying on the (-1) -curves E_1, \bar{E}_1 . We choose then two general imaginary conjugate points q_1, \bar{q}_1 , and let $q_2 := \varphi(q_1)$ and $\bar{q}_2 := \varphi(\bar{q}_1)$. For $i = 1, 2$, we denote by $\varphi_i: \mathcal{D}_6 \dashrightarrow \mathcal{D}_6$ a standard link centered at q_i, \bar{q}_i . The image by φ_2 of E_1 is a curve of self-intersection 1. Hence, $\varphi_2\varphi(\varphi_1)^{-1}$ is a standard link of type II.

It remains to consider the case where each φ_i is a link $\mathbb{F}_{n_i} \dashrightarrow \mathbb{F}_{n_{i+1}}$. We denote by N the maximum of the integers n_i . If $N \leq 1$, we are done because all links of type II between \mathbb{F}_i and $\mathbb{F}_{i'}$ with $i, i' \leq 1$ are standard. We can thus assume $N \geq 2$, which implies that there exists i such that $n_i = N, n_{i-1} < N, n_{i+1} < N$. We choose two general imaginary points $q_1, \bar{q}_1 \in \mathbb{F}_{n_{i-1}}$, and write $q_2 = \varphi_{i-1}(q_1), q_3 = \varphi_i(q_2)$. For $i = 1, 2, 3$, we denote by $\tau_i: \mathbb{F}_{n_i} \dashrightarrow \mathbb{F}_{n'_i}$ a Sarkisov link centered at q_i, \bar{q}_i . We obtain then the following commutative diagram

$$\begin{array}{ccccc} \mathbb{F}_{n_{i-1}} & \xrightarrow{\varphi_{i-1}} & \mathbb{F}_{n_i} & \xrightarrow{\varphi_i} & \mathbb{F}_{n_{i+1}} \\ | \tau_1 & & | \tau_2 & & | \tau_3 \\ \mathbb{F}_{n'_{i-1}} & \xrightarrow{\varphi'_{i-1}} & \mathbb{F}_{n'_i} & \xrightarrow{\varphi'_i} & \mathbb{F}_{n'_{i+1}}, \end{array}$$

where $\varphi'_{i-1}, \varphi'_i$ are Sarkisov links. By construction, $n'_{i-1}, n'_i, n'_{i+1} < N$, we can then replace $\varphi_i\varphi_{i-1}$ with $(\tau_3)^{-1}\varphi'_i\varphi'_{i-1}\tau_1$ and "avoid" \mathbb{F}_N . Repeating this process if needed, we end up with a sequence of Sarkisov links passing only through \mathbb{F}_1 and \mathbb{F}_0 . Moreover, since this process does not add any real base-point, it preserves the regularity at real points. \square

Corollary 2.16. *Let $\pi: X \rightarrow W$ and $\pi': X' \rightarrow W'$ be two Mori fibrations, where X, X' are either $\mathbb{F}_0, \mathbb{F}_1, \mathcal{D}_6$ or \mathbb{P}^2 . Any birational map $\psi: X \dashrightarrow X'$ is either an isomorphism preserving the fibrations or decomposes into links of type I, III, and standard links of type II.*

Proof. Follows from Proposition 2.6, Lemmas 2.14 and 2.15, and the description of Example 2.10. \square

3. GENERATORS OF THE GROUP $\text{Aut}(\mathbb{P}^2(\mathbb{R}))$

We start this section by describing three kinds of elements of $\text{Aut}(\mathbb{P}^2(\mathbb{R}))$, which are birational maps of \mathbb{P}^2 of degree 5. These maps are associated to three pairs of conjugate imaginary points; the description is then analogue to the description of quadratic maps, which are associated to three points.

Example 3.1. Let $p_1, \bar{p}_1, p_2, \bar{p}_2, p_3, \bar{p}_3 \in \mathbb{P}^2$ be three pairs of imaginary points of \mathbb{P}^2 , not lying on the same conic. Denote by $\pi: X \rightarrow \mathbb{P}^2$ the blow-up of the six points, which is an isomorphism $X(\mathbb{R}) \rightarrow \mathbb{P}^2(\mathbb{R})$. Note that X is isomorphic to a smooth cubic of \mathbb{P}^3 . The set of strict transforms of the conics passing through five of the six points corresponds to three pairs of imaginary (-1) -curves (or lines on the cubic), and the six curves are disjoint. The contraction of the six curves gives a birational morphism $\eta: X \rightarrow \mathbb{P}^2$, inducing an isomorphism $X(\mathbb{R}) \rightarrow \mathbb{P}^2(\mathbb{R})$, which contracts the curves onto three pairs of imaginary points $q_1, \bar{q}_1, q_2, \bar{q}_2, q_3, \bar{q}_3 \in \mathbb{P}^2$; we choose the order so that q_i is the image of the conic not passing through p_i . The map $\psi = \eta\pi^{-1}$ is a birational map $\mathbb{P}^2 \dashrightarrow \mathbb{P}^2$ inducing an isomorphism $\mathbb{P}^2(\mathbb{R}) \rightarrow \mathbb{P}^2(\mathbb{R})$.

Let $L \subset \mathbb{P}^2$ be a general line of \mathbb{P}^2 . The strict transform of L on X by π^{-1} has self-intersection 1 and intersects the six curves contracted by η into 2 points (because these are conics). The image $\psi(L)$ has then six singular points of multiplicity 2 and self-intersection 25; it is thus a quintic passing through the q_i with multiplicity 2. The construction of ψ^{-1} being symmetric as the one of ψ , the linear system of ψ consists of quintics of \mathbb{P}^2 having multiplicity 2 at $p_1, \bar{p}_1, p_2, \bar{p}_2, p_3, \bar{p}_3$.

One can moreover check that ψ sends the pencil of conics through $p_1, \bar{p}_1, p_2, \bar{p}_2$ onto the pencil of conics through $q_1, \bar{q}_1, q_2, \bar{q}_2$ (and the same holds for the two other real pencil of conics, through $p_1, \bar{p}_1, p_3, \bar{p}_3$ and through $p_2, \bar{p}_2, p_3, \bar{p}_3$).

Example 3.2. Let $p_1, \bar{p}_1, p_2, \bar{p}_2 \in \mathbb{P}^2$ be two pairs of imaginary points of \mathbb{P}^2 , not on the same line. Denote by $\pi_1: X_1 \rightarrow \mathbb{P}^2$ the blow-up of the four points, and by $E_2, \bar{E}_2 \subset X_1$ the curves contracted onto p_2, \bar{p}_2 respectively. Let $p_3 \in E_2$ be a point, and $\bar{p}_3 \in \bar{E}_2$ its conjugate. We assume that there is no conic of \mathbb{P}^2 passing through $p_1, \bar{p}_1, p_2, \bar{p}_2, p_3, \bar{p}_3$ and let $\pi_2: X_2 \rightarrow X_1$ be the blow-up of p_3, \bar{p}_3 .

On X , the strict transforms of the two conics C, \bar{C} of \mathbb{P}^2 , passing through $p_1, \bar{p}_1, p_2, \bar{p}_2, p_3$ and $p_1, \bar{p}_1, p_2, \bar{p}_2, \bar{p}_3$ respectively, are imaginary conjugate disjoint (-1) curves. The contraction of these two curves gives a birational morphism $\eta_2: X_2 \rightarrow Y_1$, contracting C, \bar{C} onto two points q_3, \bar{q}_3 . On Y_1 , we find two pairs of imaginary (-1) -curves, all four curves being disjoint. These are the strict transforms of the exceptional curves associated to p_2, \bar{p}_2 , and of the conics passing through $p_1, p_2, \bar{p}_2, p_3, \bar{p}_3$ and $\bar{p}_1, p_2, \bar{p}_2, p_3, \bar{p}_3$ respectively. The contraction of these curves gives a birational morphism $\eta_1: Y_1 \rightarrow \mathbb{P}^2$, and the images of the four curves are points $q_2, \bar{q}_2, q_1, \bar{q}_1$ respectively. Note that the four maps $\pi_1, \pi_2, \eta_1, \eta_2$ are blow-ups of imaginary points, so the birational map $\psi = \eta_1\eta_2(\pi_1\pi_2)^{-1}: \mathbb{P}^2 \dashrightarrow \mathbb{P}^2$ induces an isomorphism $\mathbb{P}^2(\mathbb{R}) \rightarrow \mathbb{P}^2(\mathbb{R})$.

In the same way as in Example 3.1, we find that the linear system of ψ is of degree 5, with multiplicity 2 at the points p_i, \bar{p}_i . The situation is similar for ψ^{-1} , with the six points q_i, \bar{q}_i in the same configuration: $q_1, \bar{q}_1, q_2, \bar{q}_2$ lie on the plane and q_3, \bar{q}_3 are infinitely near to q_2, \bar{q}_2 respectively.

One can moreover check that ψ sends the pencil of conics through $p_1, \bar{p}_1, p_2, \bar{p}_2$ onto the pencil of conics through $q_1, \bar{q}_1, q_2, \bar{q}_2$ and the pencil of conics through $p_2, \bar{p}_2, p_3, \bar{p}_3$ onto the pencil of conics through $q_2, \bar{q}_2, q_3, \bar{q}_3$. But, contrary to Example 3.1, there is no pencil of conics through $q_1, \bar{q}_1, q_3, \bar{q}_3$.

Example 3.3. Let p_1, \bar{p}_1 be a pair of two conjugate imaginary points of \mathbb{P}^2 . We choose a point p_2 in the first neighbourhood of p_1 , and a point p_3 in the first neighbourhood of p_2 , not lying on the exceptional divisor of p_1 . We denote by $\pi: X \rightarrow \mathbb{P}^2$ the blow-up of $p_1, \bar{p}_1, p_2, \bar{p}_2, p_3, \bar{p}_3$. We denote by $E_i, \bar{E}_i \subset X$ the irreducible exceptional curves corresponding to the points p_i, \bar{p}_i , for $i = 1, 2, 3$. The strict transforms

of the two conics through $p_1, \bar{p}_1, p_2, \bar{p}_2, p_3$ and $p_1, \bar{p}_1, p_2, \bar{p}_2, \bar{p}_3$ respectively are disjoint (-1) -curves on X , intersecting the exceptional curves $E_1, \bar{E}_1, E_2, \bar{E}_2$ similarly as E_3, \bar{E}_3 . Hence, there exists a birational morphism $\eta: X \rightarrow \mathbb{P}^2$ contracting the strict transforms of the two conics and the curves $E_1, \bar{E}_1, E_2, \bar{E}_2$.

As in Examples 3.1 and 3.2, the linear system of $\psi = \eta\pi^{-1}$ consists of quintics with multiplicity two at the six points $p_1, \bar{p}_1, p_2, \bar{p}_2, p_3, \bar{p}_3$.

Definition 3.4. The birational maps of \mathbb{P}^2 of degree 5 obtained in Example 3.1 will be called *standard quintic transformations* and those of Example 3.2 and Example 3.3 will be called *special quintic transformations* respectively.

Lemma 3.5. *Let $\psi: \mathbb{P}^2 \dashrightarrow \mathbb{P}^2$ be a birational map inducing an isomorphism $\mathbb{P}^2(\mathbb{R}) \rightarrow \mathbb{P}^2(\mathbb{R})$. The following hold:*

- (1) *The degree of ψ is $4k + 1$ for some integer $k \geq 0$.*
- (2) *Every multiplicity of the linear system of ψ is even.*
- (3) *Every curve contracted by ψ is of even degree.*
- (4) *If ψ has degree 1, it belongs to $\text{Aut}_{\mathbb{R}}(\mathbb{P}^2) = \text{PGL}(3, \mathbb{R})$.*
- (5) *If ψ has degree 5, then it is a standard or special quintic transformation, described in Examples 3.1, 3.2 or 3.3, and has thus exactly 6 base-points.*
- (6) *If ψ has at most 6 base-points, then ψ has degree 1 or 5.*

Remark 3.6. Part (1) is [RV05, Teorema 1].

Proof. Denote by d the degree of ψ and by m_1, \dots, m_k the multiplicities of the base-points of ψ . The Noether equalities yield $\sum_{i=1}^k m_i = 3(d-1)$ and $\sum_{i=1}^k (m_i)^2 = d^2 - 1$.

Let C, \bar{C} be a pair of two curves contracted by ψ . Since $C \cap \bar{C}$ does not contain any real point, the degree of C and \bar{C} is even. This yields (2), and implies that all multiplicities of the linear system of ψ^{-1} are even, giving (2).

In particular, $3(d-1)$ is a multiple of 4 (all multiplicities come by pairs of even integers), which implies that $d = 4k + 1$ for some integer k . Hence (1) is proved.

If the number of base-points is at most $k = 6$, then by Cauchy-Schwartz we get

$$9(d-1)^2 = \left(\sum_{i=1}^k m_i \right)^2 \leq k \sum_{i=1}^k (m_i)^2 = k(d^2 - 1) = 6(d^2 - 1)$$

This yields $9(d-1) \leq 6(d+1)$, hence $d \leq 5$.

If $d = 1$, all m_i are zero, and $\psi \in \text{Aut}_{\mathbb{R}}(\mathbb{P}^2)$, so we get (4).

If $d = 5$, the Noether equalities yield $k = 6$ and $m_1 = m_2 = \dots = m_6 = 2$. Hence, the base-points of ψ consist of three pairs of conjugate imaginary points $p_1, \bar{p}_1, p_2, \bar{p}_2, p_3, \bar{p}_3$. Moreover, if a conic passes through 5 of the six points, its free intersection with the linear system is zero, so it is contracted by ψ , and there is no conic through the six points.

(a) If the six points belong to \mathbb{P}^2 , the map is a standard quintic transformation, described in Example 3.1.

(b) If two points are infinitely near, the map is a special quintic transformation, described in Example 3.2.

(c) If four points are infinitely near, the map is a special quintic transformation, described in Example 3.3. \square

Before proving Theorem 1.2, we will show that all quintic transformations are generated by linear automorphisms and standard quintic transformations:

Lemma 3.7. *Every quintic transformation $\psi \in \text{Aut}(\mathbb{P}^2(\mathbb{R}))$ belongs to the group generated by $\text{Aut}_{\mathbb{R}}(\mathbb{P}^2)$ and standard quintic transformations.*

Proof. By Lemma 3.5, we only need to show the result when ψ is a special quintic transformation as in Example 3.2 or Example 3.3.

We first assume that ψ is a special quintic transformation as in Example 3.2, with base-points $p_1, \bar{p}_1, p_2, \bar{p}_2, p_3, \bar{p}_3$, where p_3, \bar{p}_3 are infinitely near to p_2, \bar{p}_2 . For $i = 1, 2$, we denote by $q_i \in \mathbb{P}^2$ the point which is the image by ψ of the conic passing the five points of $\{p_1, \bar{p}_1, p_2, \bar{p}_2, p_3, \bar{p}_3\} \setminus \{p_i\}$. Then, the base-points of ψ^{-1} are $q_1, \bar{q}_1, q_2, \bar{q}_2, q_3, \bar{q}_3$, where q_3, \bar{q}_3 are points infinitely near to q_2, \bar{q}_2 respectively (see Example 3.2). We choose a general pair of conjugate imaginary points $p_4, \bar{p}_4 \in \mathbb{P}^2$, and write $q_4 = \psi(p_4)$, $\bar{q}_4 = \psi(\bar{p}_4)$. We denote by φ_1 a standard quintic transformation having base-points at $p_1, \bar{p}_1, p_2, \bar{p}_2, p_4, \bar{p}_4$, and by φ_2 a standard quintic transformation having base-points at $q_1, \bar{q}_1, q_2, \bar{q}_2, q_4, \bar{q}_4$. We now prove that $\varphi_2\psi(\varphi_1)^{-1}$ is a standard quintic transformation; this will yield the result. Denote by p'_i, \bar{p}'_i the base-points of $(\varphi_1)^{-1}$, with the order associated to the p_i , which means that p'_i is the image by φ_i of a conic not passing through p_i (see Example 3.1). Similarly, we denote by q'_i, \bar{q}'_i the base-points of $(\varphi_2)^{-1}$. We obtain the following commutative of birational maps, where the arrows indexed by points are blow-ups of these points:

$$\begin{array}{ccccccc}
 & & Y_1 & & Y_2 & & Y_3 & & \\
 & \swarrow & & \searrow & \swarrow & \searrow & \swarrow & \searrow & \\
 & p'_4, \bar{p}'_4 & & p_4, \bar{p}_4 & p_3, \bar{p}_3 & & q_3, \bar{q}_3 & q_4, \bar{q}_4 & & q'_4, \bar{q}'_4 \\
 & & X_1 & \xleftarrow{\hat{\varphi}_1} & X_2 & \xrightarrow{\hat{\psi}} & X_3 & \xrightarrow{\hat{\varphi}_2} & X_4 & \\
 & \downarrow & & & \downarrow & & \downarrow & & \downarrow & \\
 & p'_1, \bar{p}'_1 & & & p_1, \bar{p}_1 & & q_1, \bar{q}_1 & & q'_1, \bar{q}'_1 & \\
 & p'_2, \bar{p}'_2 & & & p_2, \bar{p}_2 & & q_2, \bar{q}_2 & & q_2, \bar{q}_2 & \\
 & & \mathbb{P}^2 & \xleftarrow{\varphi_1} & \mathbb{P}^2 & \xrightarrow{\psi} & \mathbb{P}^2 & \xrightarrow{\varphi_2} & \mathbb{P}^2 &
 \end{array}$$

Each of the surfaces X_1, X_2, X_3, X_4 admits a conic bundle structure $\pi_i : X_i \rightarrow \mathbb{P}^1$, which fibres corresponds to the conics passing through the four points blown-up on \mathbb{P}^2 to obtain X_i . Moreover, $\hat{\varphi}_1, \hat{\psi}, \hat{\varphi}_2$ preserve these conic bundle structures. The map $(\hat{\varphi}_1)^{-1}$ blows-up p_4, \bar{p}_4 and contract the fibres associated to them, then $\hat{\psi}$ blows-up p_3, \bar{p}_3 and contract the fibres associated to them. The map $\hat{\varphi}_2$ blow-ups the points q_4, \bar{q}_4 , which correspond to the image of the curves contracted by $(\hat{\varphi}_1)^{-1}$, and contracts their fibres, corresponding to the exceptional divisors of the points p_4, \bar{p}_4 . Hence, $\hat{\varphi}_2\hat{\psi}\hat{\varphi}_1$ is the blow-up of two imaginary points $p'_3, \bar{p}'_3 \in X_1$, followed by the contraction of their fibres. We obtain the following commutative diagram


$$\begin{array}{ccc}
 & Z & \\
 & \swarrow & \searrow \\
 & p'_3, \bar{p}'_3 & q'_3, \bar{q}'_3 \\
 & & \\
 & X_1 & \xrightarrow{\hat{\varphi}_2\hat{\psi}(\hat{\varphi}_1)^{-1}} & X_4 \\
 & \downarrow & & \downarrow \\
 & p'_1, \bar{p}'_1 & & q'_1, \bar{q}'_1 \\
 & p'_2, \bar{p}'_2 & & q_2, \bar{q}_2 \\
 & & \mathbb{P}^2 & \xrightarrow{\varphi_2\psi(\varphi_1)^{-1}} & \mathbb{P}^2,
 \end{array}$$

and the points p'_3, \bar{p}'_3 correspond to point of \mathbb{P}^2 , hence $\varphi_2\psi(\varphi_1)^{-1}$ is a standard quintic transformation.

The remaining case is when ψ is a special quintic transformation as in Example 3.3, with base-points with base-points $p_1, \bar{p}_1, p_2, \bar{p}_2, p_3, \bar{p}_3$, where p_3, \bar{p}_3 are infinitely near to p_2, \bar{p}_2 and these latter are infinitely near to p_1, \bar{p}_1 . The map ψ^{-1} has base-points $q_1, \bar{q}_1, q_2, \bar{q}_2, q_3, \bar{q}_3$, having the same configuration (see Example 3.3). We choose a general pair of conjugate imaginary points $p_4, \bar{p}_4 \in \mathbb{P}^2$, and write $q_4 = \psi(p_4)$, $\bar{q}_4 = \psi(\bar{p}_4)$. We denote by φ_1 a special quintic transformation having base-points at $p_1, \bar{p}_1, p_2, \bar{p}_2, p_4, \bar{p}_4$, and by φ_2 a special quintic transformation having base-points at $q_1, \bar{q}_1, q_2, \bar{q}_2, q_4, \bar{q}_4$. The maps φ_1, φ_2 have four proper base-points, and are thus given in Example 3.2. The same proof as before implies that $\varphi_2\psi(\varphi_1)^{-1}$ is a special quintic transformation with four base-points. This gives the result. \square

Lemma 3.8. *Let $\varphi: \mathbb{P}^2 \dashrightarrow \mathbb{P}^2$ be a birational map, that decomposes as $\varphi = \varphi_5 \cdots \varphi_1$, where $\varphi_i: X_{i-1} \dashrightarrow X_i$ is a Sarkisov link for each i , where $X_0 = \mathbb{P}^2$, $X_1 = Q_{3,1}$, $X_2 = X_3 = \mathcal{D}_6$, $X_4 = \mathcal{S}_2$, $X_5 = \mathbb{P}^2$. If φ_2 is an automorphism of $\mathcal{D}_6(\mathbb{R})$ and $\varphi_4\varphi_3\varphi_2$ sends the base-point of $(\varphi_1)^{-1}$ onto the base-point of φ_5 , then φ is an automorphism of $\mathbb{P}^2(\mathbb{R})$ of degree 5.*

Proof. We have the following commutative diagram, where each π_i is the blow-up of two conjugate imaginary points and each η_i is the blow-up of one real point. The two maps $(\varphi_2)^{-1}$ and φ_4 are also blow-ups of imaginary points.



The only real base-points are those blown-up by η_1 and η_2 . Since η_2 blows-up the image by $\varphi_4\varphi_3\varphi_2$ of the real point blown-up by η_1 , the map φ has at most 6 base-points, all being imaginary, and the same holds for φ^{-1} . Hence, φ is an automorphism of $\mathbb{P}^2(\mathbb{R})$ with at most 6 base-points. We can moreover see that $\varphi \notin \text{Aut}_{\mathbb{R}}(\mathbb{P}^2)$, since the two curves of Y_2 contracted by π_2 are sent by $\varphi_4\pi_3$ onto conics of $Q_{3,1}$, which are therefore not contracted by φ_5 .

Lemma 3.5 implies that ψ has degree 5. \square

Proposition 3.9. *The group $\text{Aut}(\mathbb{P}^2(\mathbb{R}))$ is generated by $\text{Aut}_{\mathbb{R}}(\mathbb{P}^2)$ and by elements of $\text{Aut}(\mathbb{P}^2(\mathbb{R}))$ of degree 5.*

Proof. Let us prove that any $\varphi \in \text{Aut}(\mathbb{P}^2(\mathbb{R}))$ is generated by $\text{Aut}_{\mathbb{R}}(\mathbb{P}^2)$ and elements of $\text{Aut}(\mathbb{P}^2(\mathbb{R}))$ of degree 5. We can assume that $\varphi \notin \text{Aut}_{\mathbb{R}}(\mathbb{P}^2)$, decompose it into Sarkisov links: $\varphi = \varphi_r \circ \cdots \circ \varphi_1$.

It follows from the construction of the links (see [Cort95]) that if φ_i is of type I or II, each base-point of φ_i is a base-point of $\varphi_r \cdots \varphi_i$ (and in fact of maximal multiplicity).

We proceed by induction on r .

Since φ has no real base-point, the first link φ_1 is then of type II from \mathbb{P}^2 to $Q_{3,1}$, and $\varphi_r \cdots \varphi_2$ has a unique real base-point $r \in Q_{3,1}$, which is the base-point of $(\varphi_1)^{-1}$. If φ_2 blows-up this point, then $\varphi_2\varphi_1 \in \text{Aut}_{\mathbb{R}}(\mathbb{P}^2)$. We can then assume that φ_2 is a link of type I from $Q_{3,1}$ to \mathcal{D}_6 . The map $\varphi_2\varphi_1$ can be written as $\eta\pi^{-1}$, where $\pi: X \rightarrow \mathbb{P}^2$ is the blow-up of two pairs of imaginary points, say $p_1, \bar{p}_1, p_2, \bar{p}_2$ and $\eta: X \rightarrow \mathcal{D}_6$ is the contraction of the strict transform of the real line passing through p_1, \bar{p}_1 , onto a real point $q \in \mathcal{D}_6$. Note that p_1, \bar{p}_1 are proper points of \mathbb{P}^2 , blown-up by φ_1 and p_2, \bar{p}_2 either are proper base-points or are infinitely near to p_1, \bar{p}_1 .

The fibration $\mathcal{D}_6 \rightarrow \mathbb{P}^1$ corresponds to conics through $p_1, \bar{p}_1, p_2, \bar{p}_2$. If φ_3 is a link of type III, then $\varphi_3\varphi_2$ is an automorphism of $Q_{3,1}$ and we decrease r . Then, φ_3 is of type II. If q is a base-point of φ_3 , then $\varphi_3 = \eta'\eta^{-1}$, where $\eta': X \rightarrow \mathcal{D}_6$ is the contraction of the strict transform of the line through p_2, \bar{p}_2 . We can then write $\varphi_3\varphi_2\varphi_1$ into only two links, exchanging p_1 with p_2 and \bar{p}_1 with \bar{p}_2 . The remaining case is when φ_3 is the blow-up of two imaginary points p_3, \bar{p}_3 of \mathcal{D}_6 , followed by the contraction of the strict transforms of their fibres.

We denote by $q' \in \mathcal{D}_6(\mathbb{R})$ the image of q by φ_3 , consider $\psi_4 = (\varphi_2)^{-1}: \mathcal{D}_6 \rightarrow Q_{3,1}$, which is a link of type III, and write $\psi_5: Q_{3,1} \dashrightarrow \mathbb{P}^2$ the stereographic projection by $\psi_4(q')$, which is a link of type II centered at $\psi_4(q')$. By Lemma 3.8, the map $\chi = \psi_5\psi_4\varphi_3\varphi_2\varphi_1$ is an element of $\text{Aut}(\mathbb{P}^2(\mathbb{R}))$ of degree 5. Since $\varphi\chi^{-1}$ decomposes into one link less than φ , this concludes the proof by induction. \square

Proof of Theorem 1.2. By Proposition 3.9, $\text{Aut}(\mathbb{P}^2(\mathbb{R}))$ is generated by $\text{Aut}_{\mathbb{R}}(\mathbb{P}^2)$ and by elements of $\text{Aut}(\mathbb{P}^2(\mathbb{R}))$ of degree 5. Thanks to Lemma 3.7, $\text{Aut}(\mathbb{P}^2(\mathbb{R}))$ is indeed generated by projectivities and standard quintic transformations. \square

4. GENERATORS OF THE GROUP $\text{Bir}_{\mathbb{R}}(\mathbb{P}^2)$

Lemma 4.1. *Let $\varphi: Q_{3,1} \dashrightarrow Q_{3,1}$ be a birational map, that decomposes as $\varphi = \varphi_3\varphi_2\varphi_1$, where $\varphi_i: X_{i-1} \dashrightarrow X_i$ is a Sarkisov link for each i , where $X_0 = Q_{3,1} = X_2$, $X_1 = \mathcal{D}_6$. If φ_2 has a real base-point, then, φ can be written as $\varphi = \psi_2\psi_1$, where $\psi_1, (\psi_2)^{-1}$ are links of type II from $Q_{3,1}$ to \mathbb{P}^2 .*

Proof. We have the following commutative diagram, where each of the maps η_1, η_2 blow-ups a real point, and each of the maps $(\varphi_1)^{-1}, \varphi_3$ is the blow-up of two conjugate imaginary points.

$$\begin{array}{ccc}
 & Y & \\
 \eta_1 \swarrow & & \searrow \eta_2 \\
 \mathcal{D}_6 & \overset{\varphi_2}{\dashrightarrow} & \mathcal{D}_6 \\
 \downarrow (\varphi_1)^{-1} & & \downarrow \varphi_3 \\
 Q_{3,1} & \overset{\varphi}{\dashrightarrow} & Q_{3,1}
 \end{array}$$

The map φ has thus exactly three base-points, two of them being imaginary and one being real; we denote them by p_1, \bar{p}_1, q . The fibres of the Mori fibration $\mathcal{D}_6 \rightarrow \mathbb{P}^1$ correspond to conics of $Q_{3,1}$ passing through the points p_1, \bar{p}_1 . The real curve contracted by η_2 is thus the strict transform of the conic C of $Q_{3,1}$ passing through

p_1, \bar{p}_1 and q . The two curves contracted by φ_3 are the two imaginary sections of self-intersection -1 , which corresponds to the strict transforms of the two imaginary lines L_1, L_2 of $Q_{3,1}$ passing through q .

We can then decompose φ as the blow-up of p_1, p_2, q , followed by the contraction of the strict transforms of C, L_1, L_2 . Denote by $\psi_1: Q_{3,1} \dashrightarrow \mathbb{P}^2$ the link of type II centered at q , which is the blow-up of q followed by the contraction of the strict transform of L_1, L_2 , or equivalently the stereographic projection centered at q . The curve $\psi_1(C)$ is a real line of \mathbb{P}^2 , which contains the two points $\psi_1(p_1), \psi_1(\bar{p}_1)$. The map $\psi_2 = \varphi(\psi_1)^{-1}: \mathbb{P}^2 \dashrightarrow Q_{3,1}$ is then the blow-up of these two points, followed by the contraction of the line passing through both of them. It is then a link of type II. \square

Proof of Theorem 1.1. Let us prove that any $\varphi \in \text{Bir}_{\mathbb{R}}(\mathbb{P}^2)$ is in the group generated by $\text{Aut}_{\mathbb{R}}(\mathbb{P}^2)$, σ_0, σ_1 , and standard quintic transformations of \mathbb{P}^2 . We can assume that $\varphi \notin \text{Aut}_{\mathbb{R}}(\mathbb{P}^2)$, decompose it into Sarkisov links: $\varphi = \varphi_r \circ \dots \circ \varphi_1$. By Corollary 2.16, we can assume that all the φ_i are links of type I, III, or standard links of type II.

We proceed by induction on r , the case $r = 0$ being obvious.

Note that φ_1 is either a link of type I from \mathbb{P}^2 to \mathbb{F}_1 , or a link of type II from \mathbb{P}^2 to $Q_{3,1}$. We now study the possibilities for the base-points of φ_1 and the next links:

(1) Suppose that $\varphi_1: \mathbb{P}^2 \dashrightarrow \mathbb{F}_1$ is a link of type I, and that φ_2 is a link $\mathbb{F}_1 \dashrightarrow \mathbb{F}_1$. Then, φ_2 blows-up two imaginary base-points of \mathbb{F}_1 , not lying on the exceptional curve. Hence, $\psi = (\varphi_1)^{-1}\varphi_2\varphi_1$ is a quadratic transformation of \mathbb{P}^2 with three proper base-points, one real and two imaginary. It is thus equal to $\alpha\sigma_1\beta$ for some $\alpha, \beta \in \text{Aut}_{\mathbb{R}}(\mathbb{P}^2)$. Replacing φ with $\varphi\psi^{-1}$, we obtain a decomposition with less Sarkisov links, and conclude by induction.

(2) Suppose that $\varphi_1: \mathbb{P}^2 \dashrightarrow \mathbb{F}_1$ is a link of type I, and that φ_2 is a link $\mathbb{F}_1 \dashrightarrow \mathbb{F}_0$. Then, $\varphi_2\varphi_1$ is the blow-up of two real points p_1, p_2 of \mathbb{P}^2 followed by the contraction of the line through p_1, p_2 . The exceptional divisors of p_1, p_2 are two (0)-curves of $\mathbb{F}_0 = \mathbb{P}^1 \times \mathbb{P}^1$, intersecting at one real point.

(2a) Suppose first that φ_3 has a base-point which is real, and not lying on E_1, E_2 . Then, $\psi = (\varphi_1)^{-1}\varphi_3\varphi_2\varphi_1$ is a quadratic transformation of \mathbb{P}^2 with three proper base-points, all real. It is thus equal to $\alpha\sigma_0\beta$ for some $\alpha, \beta \in \text{Aut}_{\mathbb{R}}(\mathbb{P}^2)$. Replacing φ with $\varphi\psi^{-1}$, we obtain a decomposition with less Sarkisov links, and conclude by induction.

(2b) Suppose now that φ_3 has imaginary base-points, which are q, \bar{q} . Since φ_3 is a standard link of type II, it goes from \mathbb{F}_0 to \mathbb{F}_0 , so q and \bar{q} do not lie on a (0)-curve, and then do not belong to the curves E_1, E_2 . We can then decompose $\varphi_2\varphi_3: \mathbb{F}_1 \dashrightarrow \mathbb{F}_2$ into a Sarkisov link centered at two imaginary points, followed by a Sarkisov link centered at a real point. This reduces to case (1), already treated before.

(2c) The remaining case (for (2)) is when φ_3 has a base-point p_3 which is real, but lying on E_1 or E_2 . We choose a general real point $p_4 \in \mathbb{F}_0$, and denote by $\theta: \mathbb{F}_0 \dashrightarrow \mathbb{F}_1$ a Sarkisov link centered at p_4 . We observe that $\psi = (\varphi_1)^{-1}\theta\varphi_2\varphi_1$ is a quadratic map as in case (2a), and that $\varphi\psi^{-1} = \varphi_n \dots \varphi_3\theta^{-1}\varphi_1$ admits now a decomposition of the same length, but which is in case (2a).

(3) Suppose now that $\varphi_1: \mathbb{P}^2 \dashrightarrow Q_{3,1}$ is a link of type II and that φ_2 is a link of type II from S to \mathbb{P}^2 . If φ_2 and $(\varphi_1)^{-1}$ have the same real base-point, the map

$\varphi_2\varphi_1$ belongs to $\text{Aut}_{\mathbb{R}}(\mathbb{P}^2)$. Otherwise, $\varphi_2\varphi_1$ is a quadratic map with one unique real base-point q and two imaginary base-points. It is then equal to $\alpha\sigma_0\beta$ for some $\alpha, \beta \in \text{Aut}_{\mathbb{R}}(\mathbb{P}^2)$. We conclude as before by induction hypothesis.

(4) Suppose that $\varphi_1: \mathbb{P}^2 \dashrightarrow Q_{3,1}$ is a link of type II and φ_2 is a link of type I from S to \mathcal{D}_6 . If φ_3 is a Sarkisov link of type III, then $\varphi_3\varphi_2$ is an automorphism of $Q_{3,1}$, so we can decrease the length. We only need to consider the case where φ_3 is a link of type II from \mathcal{D}_6 to \mathcal{D}_6 . If φ_3 has a real base-point, we apply Lemma 4.1 to write $(\varphi_2)^{-1}\varphi_3\varphi_2 = \psi_2\psi_1$ where $\psi_1, (\psi_2)^{-1}$ are links $Q_{3,1} \dashrightarrow \mathbb{P}^2$. By (3), the map $\chi = \psi_1\varphi_1$ is generated by $\text{Aut}_{\mathbb{R}}(\mathbb{P}^2)$ and σ_0 . We can then replace φ with $\varphi\chi^{-1} = \varphi_r \cdots \varphi_3\varphi_2(\psi_1)^{-1} = \varphi_r \cdots \varphi_4\varphi_2\psi_2$, which has a shorter decomposition. The last case is when φ_3 has two imaginary base-points. We denote by $q \in Q_{3,1}$ the real base-point of $(\varphi_1)^{-1}$, write $q' = (\varphi_2)^{-1}\varphi_3\varphi_2(q) \in Q_{3,1}$ and denote by $\psi: Q_{3,1} \dashrightarrow \mathbb{P}^2$ the stereographic projection centered at q' . By Lemma 3.8, the map $\chi = \psi(\varphi_2)^{-1}\varphi_3\varphi_2\varphi_1$ is an automorphism of $\mathbb{P}^2(\mathbb{R})$ of degree 5, which is generated by $\text{Aut}_{\mathbb{R}}(\mathbb{P}^2)$ and standard automorphisms of $\mathbb{P}^2(\mathbb{R})$ of degree 5 (Lemma 3.7). We can thus replace φ with $\varphi\chi^{-1}$, which has a decomposition of shorter length. \square

5. GENERATORS OF THE GROUP $\text{Aut}(Q_{3,1}(\mathbb{R}))$

Example 5.1. Let $p_1, \bar{p}_1, p_2, \bar{p}_2 \in Q_{3,1} \subset \mathbb{P}^3$ be two pairs of conjugate imaginary points, not on the same plane of \mathbb{P}^3 . Let $\pi: X \rightarrow Q_{3,1}$ be the blow-up of these points. The imaginary plane of \mathbb{P}^3 passing through $p_1, \bar{p}_2, \bar{p}_2$ intersects $Q_{3,1}$ onto a conic, having self-intersection 2: two general different conics on $Q_{3,1}$ are the trace of hyperplanes, and intersect then into two points, being on the line of intersection of the two planes. The strict transform of this conic on S is thus a (-1) -curve on S . Doing the same for the other conics passing through 3 of the points $p_1, \bar{p}_1, p_2, \bar{p}_2$, we obtain four disjoint (-1) -curves on X , that we can contract in order to obtain a birational morphism $\eta: X \rightarrow Q_{3,1}$; note that the target is $Q_{3,1}$ because it is a smooth projective rational surface of Picard rank 1. We obtain then a birational map $\psi = \eta\pi^{-1}: Q_{3,1} \dashrightarrow Q_{3,1}$ inducing an isomorphism $Q_{3,1}(\mathbb{R}) \rightarrow Q_{3,1}(\mathbb{R})$.

Denote by $H \subset Q_{3,1}$ a general hyperplane section. The strict transform of H on X by π^{-1} has self-intersection 2 and has intersection 2 with the 4 curves contracted. The image $\psi(H)$ has thus multiplicity 2 and self-intersection 18; it is then the trace of a cubic section. The construction of ψ and ψ^{-1} being similar, the linear system of ψ consists of cubic sections with multiplicity 2 at $p_1, \bar{p}_1, p_2, \bar{p}_2$.

Example 5.2. Let $p_1, \bar{p}_1 \in Q_{3,1} \subset \mathbb{P}^3$ be two conjugate imaginary points and let $\pi_1: X_1 \rightarrow Q_{3,1}$ be the blow-up of the two points. Denote by $E_1, \bar{E}_1 \subset X_1$ the curves contracted onto p_1, \bar{p}_1 respectively. Let $p_2 \in E_1$ be a point, and $\bar{p}_2 \in \bar{E}_1$ its conjugate. We assume that there is no conic of $Q_{3,1} \subset \mathbb{P}^3$ passing through $p_1, \bar{p}_1, p_2, \bar{p}_2$ and let $\pi_2: X_2 \rightarrow X_1$ be the blow-up of p_2, \bar{p}_2 .

On X , the strict transforms of the two conics C, \bar{C} of \mathbb{P}^2 , passing through p_1, \bar{p}_1, p_2 and $p_1, \bar{p}_1, \bar{p}_2$ respectively, are imaginary conjugate disjoint (-1) curves. The contraction of these two curves gives a birational morphism $\eta_2: X_2 \rightarrow Y_1$. On this latter surface, we find two disjoint conjugate imaginary (-1) -curves. These are the strict transforms of the exceptional curves associated to p_1, \bar{p}_1 . The contraction of these curves gives a birational morphism $\eta_1: Y_1 \rightarrow Q_{3,1}$. The birational map $\psi = \eta_1\eta_2(\pi_1\pi_2)^{-1}: Q_{3,1} \dashrightarrow Q_{3,1}$ induces an isomorphism $Q_{3,1}(\mathbb{R}) \rightarrow Q_{3,1}(\mathbb{R})$.

Definition 5.3. The birational maps of $Q_{3,1}$ of degree 3 obtained in Example 5.1 will be called *standard cubic transformations* and those of Example 5.2 will be called *special cubic transformations*.

Note that since $\text{Pic}(Q_{3,1}) = \mathbb{Z}H$, where H is an hyperplane section, we can associate to any birational map $Q_{3,1} \dashrightarrow Q_{3,1}$, an integer d , which is the *degree of the map*, such that $\psi^{-1}(H) = dH$.

Lemma 5.4. *Let $\psi: Q_{3,1} \dashrightarrow Q_{3,1}$ be a birational map inducing an isomorphism $Q_{3,1}(\mathbb{R}) \rightarrow Q_{3,1}(\mathbb{R})$. The following hold:*

- (1) *The degree of ψ is $2k + 1$ for some integer $k \geq 0$.*
- (2) *If ψ has degree 1, it belongs to $\text{Aut}_{\mathbb{R}}(Q_{3,1}) = \text{PO}(3,1)$.*
- (3) *If ψ has degree 3, then it is a standard or special cubic transformation, described in Examples 5.1 and 5.2, and has thus exactly 4 base-points.*
- (4) *If ψ has at most 4 base-points, then ψ has degree 1 or 3.*

Proof. Denote by d the degree of ψ and by a_1, \dots, a_n the multiplicities of the base-points of ψ . Denote by $\pi: X \rightarrow Q_{3,1}$ the blow-up of the base-points, and by $E_1, \dots, E_n \in \text{Pic}(X)$ the divisors being the total pull-back of the exceptional (-1) -curves obtained after blowing-up the points. Writing $\eta: X \rightarrow Q_{3,1}$ the birational morphism $\psi\pi$, we obtain

$$\begin{aligned} \eta^*(H) &= d\pi^*(H) - \sum_{i=1}^n a_i E_i \\ K_X &= \pi^*(-2H) + \sum_{i=1}^n E_i. \end{aligned}$$

Since H corresponds to a smooth rational curve of self-intersection 2, we have $(\eta^*(H))^2$ and $\eta^*(H) \cdot K_X = -4$. We find then

$$\begin{aligned} 2 &= (\eta^*(H))^2 = 2d^2 - \sum_{i=1}^n (a_i)^2 \\ 4 &= -K_X \cdot \eta^*(H) = 4d - \sum_{i=1}^n a_i. \end{aligned}$$

Since multiplicities come by pairs, $n = 2m$ for some integer m and we can order the a_i so that $a_i = a_{n+1-i}$ for $i = 1, \dots, m$. This yields

$$\begin{aligned} d^2 - 1 &= \sum_{i=1}^m (a_i)^2 \\ 2(d-1) &= \sum_{i=1}^m a_i \end{aligned}$$

Since $(a_i)^2 \equiv a_i \pmod{2}$, we find $d^2 - 1 \equiv 2(d-1) \equiv 0 \pmod{2}$, hence d is odd. This gives (1).

If the number of base-points is at most 4, we can choose $m = 2$, and obtain by Cauchy-Schwartz

$$4(d-1)^2 = \left(\sum_{i=1}^m a_i \right)^2 \leq m \sum_{i=1}^m (a_i)^2 = m(d^2 - 1) = 2(d^2 - 1).$$

This yields $2(d-1) \leq d+1$, hence $d \leq 3$.

If $d = 1$, all a_i are zero, and $\psi \in \text{Aut}_{\mathbb{R}}(Q_{3,1})$.

If $d = 3$, we get $\sum_{i=1}^m (a_i)^2 = 8$, $\sum_{i=1}^m a_i = 4$, so $m = 2$ and $a_1 = a_2 = 2$. Hence, the base-points of ψ consist of two pairs of conjugate imaginary points $p_1, \bar{p}_1, p_2, \bar{p}_2$. Moreover, if a conic passes through 3 of the points, its free intersection with the linear system is zero, so it is contracted by ψ , and there is no conic through the four points.

(a) If the four points belong to $Q_{3,1}$, the map is a standard cubic transformation, described in Example 5.1.

(b) If two points are infinitely near, the map is a special cubic transformation, described in Example 5.2. \square

Lemma 5.5. *Let $\varphi: Q_{3,1} \dashrightarrow Q_{3,1}$ be a birational map, that decomposes as $\varphi = \varphi_3\varphi_2\varphi_1$, where $\varphi_i: X_{i-1} \dashrightarrow X_i$ is a Sarkisov link for each i , where $X_0 = Q_{3,1} = X_2$, $X_1 = \mathcal{D}_6$. If φ_2 is an automorphism of $\mathcal{D}_6(\mathbb{R})$ then φ is a cubic automorphism of $Q_{3,1}(\mathbb{R})$ of degree 3 described in Examples 5.1 and 5.2. Moreover, φ is a standard cubic transformation if and only if the link φ_2 of type II is a standard link of type II.*

Proof. We have the following commutative diagram, where each of the maps $\pi_1, \pi_2, (\varphi_1)^{-1}, \varphi_3$ is the blow-up of two conjugate imaginary points.

$$\begin{array}{ccc}
 & Y & \\
 \pi_1 \swarrow & & \searrow \pi_2 \\
 \mathcal{D}_6 & \overset{\varphi_2}{\dashrightarrow} & \mathcal{D}_6 \\
 \downarrow (\varphi_1)^{-1} & & \downarrow \varphi_3 \\
 Q_{3,1} & \overset{\varphi}{\dashrightarrow} & Q_{3,1}.
 \end{array}$$

Hence, φ is an automorphism of $\mathbb{P}^2(\mathbb{R})$ with at most 4 base-points. We can moreover see that $\varphi \notin \text{Aut}_{\mathbb{R}}(Q_{3,1})$, since the two curves of Y_2 contracted blown-up by π_2 are sent by $\varphi_3\pi_3$ onto conics of $Q_{3,1}$, contracted by φ^{-1} .

Lemma 3.5 implies that φ is cubic automorphism of $Q_{3,1}(\mathbb{R})$ of degree 3 described in Examples 5.1 and 5.2. In particular, φ has exactly four base-points, blown-up by $(\varphi_1)^{-1}\pi_1$. Moreover, φ is a standard cubic transformation if and only if these four points are proper base-points of $Q_{3,1}$. This corresponds to saying that the two base-points of φ_2 do not belong to the exceptional curves contracted by $(\varphi_1)^{-1}$, and is thus the case exactly when φ_2 is a standard link of type II. \square

Proof of Theorem 1.3. Let us prove that any $\varphi \in \text{Aut}(Q_{3,1}(\mathbb{R}))$ is generated by $\text{Aut}_{\mathbb{R}}(Q_{3,1})$ and standard cubic transformations of $\text{Aut}(Q_{3,1}(\mathbb{R}))$ of degree 3. We can assume that $\varphi \notin \text{Aut}_{\mathbb{R}}(\mathbb{P}^2)$, decompose it into Sarkisov links: $\varphi = \varphi_r \circ \dots \circ \varphi_1$.

As already explained in the proof of Proposition 3.9, if φ_i is of type I or II, each base-point of φ_i is a base-point of $\varphi_r \dots \varphi_i$. This implies that all links are either of type I, from $Q_{3,1}$ to \mathcal{D}_6 , of type II from \mathcal{D}_6 to \mathcal{D}_6 with imaginary base-points, or of type III from \mathcal{D}_6 to $Q_{3,1}$. Moreover, by Lemma 2.15, we can assume that all links of type II are standard.

We proceed by induction on r .

Since φ has no real base-point, the first link φ_1 is then of type I from $Q_{3,1}$ to \mathcal{D}_6 . If φ_2 is of type III, then $\varphi_2\varphi_1 \in \text{Aut}_{\mathbb{R}}(Q_{3,1})$. We replace these two links and conclude by induction. If φ_2 is a standard link of type II, then $\psi = (\varphi_1)^{-1}\varphi_2\varphi_1$ is a standard cubic transformation. Replacing φ with $\varphi\psi^{-1}$ decreases the number of links, so we conclude by induction. \square

Twisting maps and factorisation. Choose a real line $L \subset \mathbb{P}^3$, which does not meet $Q_{3,1}(\mathbb{R})$. The projection from L gives a morphism $\pi_L: Q_{3,1}(\mathbb{R}) \rightarrow \mathbb{P}^1(\mathbb{R})$, which induces a conic bundle structure on the blow-up $\tau_L: \mathcal{D}_6 \rightarrow Q_{3,1}$ of the two imaginary points of $L \cap Q_{3,1}$.

We denote by $T(Q_{3,1}, \pi_L) \subset \text{Aut}(Q_{3,1}(\mathbb{R}))$ the group of elements $\varphi \in \text{Aut}(Q_{3,1}(\mathbb{R}))$ such that $\pi_L \varphi = \pi_L$ and such that the lift $(\tau_L)^{-1} \varphi \tau_L \in \text{Aut}(\mathcal{D}_6(\mathbb{R}))$ preserves the set of two imaginary (-1) -curves which are sections of the conic bundle $\pi_L \tau_L$.

Any element $\varphi \in T(Q_{3,1}, \pi_L)$ is called a *twisting map of $Q_{3,1}$ with axis L* .

Choosing the line $w = x = 0$ for L , we can get the more precise description given in [HM09a, KM09]: the twisting maps corresponds in local coordinates $(x, y, z) \mapsto (1 : x : y : z)$ to

$$\varphi_M : (x, y, z) \mapsto (x, (y, z) \cdot M(x))$$

where $M : [-1, 1] \rightarrow O(2) \subset \text{PGL}(2, \mathbb{R}) = \text{Aut}(\mathbb{P}^1)$ is a real algebraic map.

Proposition 5.6. *Any twisting map with axis L is a composition of twisting maps with axis L , of degree 1 and 3.*

Proof. We can assume that L is the line $y = z = 0$.

The projection $\tau : \mathcal{D}_6 \rightarrow Q_{3,1}$ is a link of type III, described in Example 2.10(3), which blows-up two imaginary points of $Q_{3,1}$. The fibres of the Mori Fibration $\pi : \mathcal{D}_6 \rightarrow \mathbb{P}^1$ correspond then, via τ , to the fibres of $\pi_L : Q_{3,1}(\mathbb{R}) \rightarrow \mathbb{P}^1(\mathbb{R})$. Hence, a twisting map of $Q_{3,1}$ corresponds to a map of the form $\tau \varphi \tau^{-1}$, where $\varphi : \mathcal{D}_6 \dashrightarrow \mathcal{D}_6$ is a birational map such that $\pi \varphi = \pi$, and which preserves the set of two (-1) -curves. This implies that φ has all its base-points on the two (-1) -curves. It remains to argue as in Lemma 2.15, and decompose φ into links that have only base-points on the set of two (-1) -curves. \square

6. GENERATORS OF THE GROUP $\text{Aut}(\mathbb{F}_0(\mathbb{R}))$

Proof of Theorem 1.4. Let us prove that any $\varphi \in \text{Aut}(\mathbb{F}_0(\mathbb{R}))$ is generated by $\text{Aut}_{\mathbb{R}}(\mathbb{F}_0)$ and by the the involution

$$\tau_0 : ((x_0 : x_1), (y_0 : y_1)) \dashrightarrow ((x_0 : x_1), (x_0 y_0 + x_1 y_1 : x_1 y_0 - x_0 y_1)).$$

Observe that τ_0 is a Sarkisov link $\mathbb{F}_0 \dashrightarrow \mathbb{F}_0$ that is the blow-up of the two imaginary points $p = ((\mathbf{i} : 1), (\mathbf{i} : 1))$, $\bar{p} = ((-\mathbf{i} : 1), (-\mathbf{i} : 1))$, followed by the contraction of the two fibres of the first projection $\mathbb{F}_0 \rightarrow \mathbb{P}^1$ passing through p, \bar{p} .

We can assume that $\varphi \notin \text{Aut}_{\mathbb{R}}(\mathbb{P}^2)$, decompose it into Sarkisov links: $\varphi = \varphi_r \circ \dots \circ \varphi_1$. As already explained, if φ_i is of type I or II, each base-point of φ_i is a base-point of $\varphi_r \dots \varphi_i$. This implies that all links are either of type IV, or of II, from \mathbb{F}_{2d} to $\mathbb{F}_{2d'}$, with exactly two imaginary base-points. Moreover, by Lemma 2.15, we can assume that all links of type II are standard, so all go from \mathbb{F}_0 to \mathbb{F}_0 .

Each link of type IV is an element of $\text{Aut}_{\mathbb{R}}(\mathbb{F}_0)$.

Each link φ_i of type II consists of the blow-up of two imaginary points q, \bar{q} , followed by the contraction of the fibres of the first projection $\mathbb{F}_0 \rightarrow \mathbb{P}^1$ passing through q, \bar{q} . Since the two points do not belong to the same fibre by any projection, we have $q = ((a + \mathbf{i}b : 1), (c + \mathbf{i}d : 1))$, for some $a, b, c, d \in \mathbb{R}$, $bd \neq 0$. There exists thus an element $\alpha \in \text{Aut}_{\mathbb{R}}(\mathbb{F}_0)$ that sends q onto p and then \bar{q} onto \bar{p} . In consequence, $\tau_0 \alpha (\varphi_i)^{-1} \in \text{Aut}_{\mathbb{R}}(\mathbb{P}^2)$. This yields the result. \square

7. OTHER RESULTS

Infinite transitivity on surfaces. The group of automorphisms of a complex algebraic variety is small: indeed, it is finite in general. Moreover, the group of automorphisms is 3-transitive only if the variety is \mathbb{P}^1 . On the other hand, it was proved in [HM09a] that for a real rational surface X , the group of automorphisms

$\text{Aut}(X(\mathbb{R}))$ acts n -transitively on $X(\mathbb{R})$ for any n . The next theorem determines all real algebraic surfaces X having a group of automorphisms which acts infinitely transitively on $X(\mathbb{R})$.

Definition 7.1. Let G be a topological group acting continuously on a topological space M . We say that two n -tuples of distinct points (p_1, \dots, p_n) and (q_1, \dots, q_n) are *compatible* if there exists a homeomorphism $\psi: M \rightarrow M$ such that $\psi(p_i) = q_i$ for each i . The action of G on M is then said to be *infinitely transitive* if for any pair of compatible n -tuples of points (p_1, \dots, p_n) and (q_1, \dots, q_n) of M , there exists an element $g \in G$ such that $g(p_i) = q_i$ for each i . More generally, the action of G is said to be *infinitely transitive on each connected component* if we require the above condition only in case, for each i , p_i and q_i belong to the same connected component of M .

Theorem 7.2. [BM10] *Let X be a nonsingular real projective surface. The group $\text{Aut}(X(\mathbb{R}))$ is then infinitely transitive on each connected component if and only if X is geometrically rational and $\#X(\mathbb{R}) \leq 3$.*

Density of automorphisms in diffeomorphisms. In [KM09], it is proved that $\text{Aut}(X(\mathbb{R}))$ is dense in $\text{Diff}(X(\mathbb{R}))$ for the C^∞ -topology when X is a geometrically rational surface with $\#X(\mathbb{R}) = 1$ (or equivalently when X is rational). In the cited paper, it is said that $\#X(\mathbb{R}) = 2$ is probably the only other case where the density holds. The following collect the known results in this direction.

Theorem 7.3. [KM09, BM10] *Let X be a smooth real projective surface.*

- *If X is not a geometrically rational surface, then $\overline{\text{Aut}(X(\mathbb{R}))} \neq \text{Diff}(X(\mathbb{R}))$;*
- *If X is a geometrically rational surface, then*
 - *If $\#X(\mathbb{R}) \geq 5$, then $\overline{\text{Aut}(X(\mathbb{R}))} \neq \text{Diff}(X(\mathbb{R}))$;*
 - *if $\#X(\mathbb{R}) = 3, 4$, then for most X , $\overline{\text{Aut}(X(\mathbb{R}))} \neq \text{Diff}(X(\mathbb{R}))$;*
 - *if $\#X(\mathbb{R}) = 1$, then $\overline{\text{Aut}(X(\mathbb{R}))} = \text{Diff}(X(\mathbb{R}))$.*

Here the closure is taken in the C^∞ -topology.

REFERENCES

- [BM10] J. Blanc, F. Mangolte, *Geometrically rational real conic bundles and very transitive actions*, Compos. Math. **147** (2011), no. 1, 161–187.
- [Com12] A. Comessatti, *Fondamenti per la geometria sopra superficie razionali dal punto di vista reale*, Math. Ann. **73** (1912) 1–72.
- [Cort95] A. Corti, *Factoring birational maps of threefolds after Sarkisov*, J. Algebraic Geom. **4** (1995), no. 2, 223–254.
- [HM09a] J. Huisman, F. Mangolte, *The group of automorphisms of a real rational surface is n -transitive*, Bull. London Math. Soc. **41**, 563–568 (2009).
- [Isko79] V.A. Iskovskikh, *Minimal models of rational surfaces over arbitrary fields*, Izv. Akad. Nauk SSSR Ser. Mat. **43** (1979), no 1, 19–43, 237.
- [Isko96] V.A. Iskovskikh, *Factorization of birational mappings of rational surfaces from the point of view of Mori theory*, Uspekhi Mat. Nauk **51** (1996) no 4 (310), 3–72.
- [KM09] J. Kollár, F. Mangolte, *Cremona transformations and diffeomorphisms of surfaces*, Adv. in Math. **222**, 44–61 (2009).
- [Mang06] F. Mangolte, *Real algebraic morphisms on 2-dimensional conic bundles*, Adv. Geom. **6** (2006), 199–213.
- [Mori82] S. Mori, *Threefolds whose canonical bundles are not numerically effective*, Ann. of Math. (2) **116** (1982), 133–176.
- [Poly97] Yu. Polyakova, *Factorization of birational mappings of rational surfaces over the field of real numbers*, Fundam. Prikl. Mat. **3** (1997), no. 2, 519–547.

- [RV05] F. Ronga, T. Vust, *Diffeomorfismi birazionali del piano proiettivo reale*, Comm. Math. Helv. **80** (2005), 517–540.

JÉRÉMY BLANC, MATHEMATISCHES INSTITUT, UNIVERSITÄT BASEL, RHEINSPRUNG 21, CH-4051 BASEL, SCHWEIZ

E-mail address: `Jeremy.Blanc@unibas.ch`

FRÉDÉRIC MANGOLTE, LUNAM UNIVERSITÉ, LAREMA, UNIVERSITÉ D'ANGERS, BD. LAVOISIER, 49045 ANGERS CEDEX 01, FRANCE

E-mail address: `frederic.mangolte@univ-angers.fr`