



HAL
open science

Symmetric powers of $\text{Nat } \mathfrak{sl}(2, \mathbb{K})$

Adrien Deloro

► **To cite this version:**

Adrien Deloro. Symmetric powers of $\text{Nat } \mathfrak{sl}(2, \mathbb{K})$. Communications in Algebra, 2016, 44 (5), pp.1981-2008. 10.1080/00927872.2014.900690 . hal-00840201v2

HAL Id: hal-00840201

<https://hal.science/hal-00840201v2>

Submitted on 22 Jan 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Symmetric powers of $\text{Nat } \mathfrak{sl}_2(\mathbb{K})$

Adrien Deloro

18th February 2015

C'est une série logarithmique.

Abstract

We identify the spaces of homogeneous polynomials in two variables $\mathbb{K}[Y^k, XY^{k-1}, \dots, X^k]$ among representations of the Lie ring $\mathfrak{sl}_2(\mathbb{K})$. This amounts to constructing a compatible \mathbb{K} -linear structure on some abstract $\mathfrak{sl}_2(\mathbb{K})$ -modules, where $\mathfrak{sl}_2(\mathbb{K})$ is viewed as a Lie ring.

The present article comes immediately next to [1] but may be read independently. For the sake of self-containedness we shall briefly recast the main motivation; should the reader desire a more complete and speculative introduction we would direct him or her to [1].

The problem addressed in the present ongoing series of articles is to understand to which extent the group-theoretic constraints of the abstract group $G = \mathbb{G}_{\mathbb{K}}$ of \mathbb{K} -points of some algebraic group \mathbb{G} actually determine the representation theory of G , that is to which extent representations of G over \mathbb{K} are characterizable among abstract G -modules. It is the author's belief that something can in the long run be achieved or at least formulated.

At this point it seems natural to ask about the Lie \mathbb{K} -algebra $(\text{Lie } \mathbb{G})_{\mathbb{K}}$ as well. Parallel to the difference between an algebraic group and the abstract group of \mathbb{K} -points runs the difference between the Lie algebra and the abstract Lie *ring* \mathfrak{g} of \mathbb{K} -points, that is the underlying $(+, [\cdot, \cdot])$ -structure with no reference to a vector space structure. As a matter of fact the relationships between \mathfrak{g} -modules and $\mathbb{K}\mathfrak{g}$ -modules are far from clear. The question may deserve interest although it appears not to have been asked before.

In the present article we shall deal with a fundamental case and characterize the symmetric powers $\text{Sym}^k \text{Nat } \mathfrak{sl}_2(\mathbb{K})$ among representations of the Lie ring $\mathfrak{sl}_2(\mathbb{K})$.

Remember that the modules $\text{Sym}^k \text{Nat } \mathfrak{sl}_2(\mathbb{K})$ ($k \geq 1$ an integer) are the irreducible, finite-dimensional representations of the Lie \mathbb{K} -algebra $\mathfrak{sl}_2(\mathbb{K})$. It is convenient to realize them as the spaces of homogeneous polynomials of degree k in two variables $\mathbb{K}[Y^k, XY^{k-1}, \dots, X^k]$ equipped with the standard action.

Such are the irreducible finite-dimensional representations of the Lie \mathbb{K} -algebra. But when one sees $\mathfrak{sl}_2(\mathbb{K})$ as a mere Lie *ring*, the question is more general. For one simply lets $\mathfrak{sl}_2(\mathbb{K})$ act on an arbitrary abelian group V in such a way that $\mathfrak{sl}_2(\mathbb{K}) \rightarrow \text{End}(V)$ be a morphism of Lie rings. In this abstract setting V need not be a \mathbb{K} -vector space, which leaves us quite far from weight theory. The present article thus deals like [1] with linearizing abstract modules, that is constructing a \mathbb{K} -vector space structure compatible with the given action of an algebraic structure, here the Lie ring $\mathfrak{sl}_2(\mathbb{K})$.

It is not surprising to turn the assumption that V is finite-dimensional (which is a priori meaningless since there is no \mathbb{K} -linear structure to start with) into an assumption on the length of the action: $x^n \cdot V = 0$, where h, x, y form the standard \mathbb{K} -basis of $\mathfrak{sl}_2(\mathbb{K})$. We thus extend the results of [1] which considered the simpler, quadratic case $x^2 \cdot V = 0$. Whenever we write $A \simeq \oplus_I B$ we merely mean that A is isomorphic to a direct sum of copies of B indexed by some (possibly finite) set I . Our main result is the following scalar extension principle.

Keywords: Lie ring $\mathfrak{sl}(2, \mathbb{K})$, Lie ring representation
MSC 2010: 17B10, 16G99, 16S30

Adrien Deloro, Institut de Mathématiques de Jussieu, Université Paris 6
Case 247, 4 place Jussieu, 75252 Paris Cedex – FRANCE
adrien.deloro@imj-prg.fr

Variation n°19. Let $n \geq 2$ be an integer and \mathbb{K} be a field of characteristic 0 or $\geq n$. Let $\mathfrak{g} = \mathfrak{sl}_2(\mathbb{K})$ viewed as a Lie ring and V be a \mathfrak{g} -module. Let \mathbb{K}_1 be the prime subfield of \mathbb{K} and $\mathfrak{g}_1 = \mathfrak{sl}_2(\mathbb{K}_1)$. Suppose that V is a \mathbb{K}_1 -vector space such that $V \simeq \bigoplus_I \text{Sym}^{n-1} \text{Nat } \mathfrak{g}_1$ as $\mathbb{K}_1 \mathfrak{g}_1$ -modules.

Then V bears a compatible \mathbb{K} -vector space structure for which $V \simeq \bigoplus_J \text{Sym}^{n-1} \text{Nat } \mathfrak{g}$ as $\mathbb{K} \mathfrak{g}$ -modules.

One may actually say a little more under an extra hypothesis which we shall later call of *coherence* of the action, in the sense that the kernels and/or images of the nilpotent operators must obey some global behaviour.

Variations n°20, n°21 and n°22. Let $n \geq 2$ be an integer and \mathbb{K} be a field. Let $\mathfrak{g} = \mathfrak{sl}_2(\mathbb{K})$ viewed as a Lie ring and V be a \mathfrak{g} -module. If the characteristic of \mathbb{K} is 0 one requires V to be torsion-free. Suppose either that $x^n = 0$ in $\text{End}(V)$ and the characteristic of \mathbb{K} is 0 or $\geq 2n + 1$, or that $x^n = y^n = 0$ in $\text{End}(V)$ and the characteristic of \mathbb{K} is $\geq n + 1$. Then the following hold.

- If for all λ , one has $\ker x \leq \ker x_\lambda$, then there is a series $\text{Ann}_V(\mathfrak{g}) = V_0 \leq V_1 \leq \dots \leq V_{n-1} = V$ of \mathfrak{g} -submodules such that for all $k = 1, \dots, n-1$, V_k/V_{k-1} bears a compatible \mathbb{K} -vector space structure for which $V_k/V_{k-1} \simeq \bigoplus_{I_k} \text{Sym}^k \text{Nat } \mathfrak{g}$ as $\mathbb{K} \mathfrak{g}$ -modules.
- If for all $\lambda \in \mathbb{K}$, one has $\text{im } x_\lambda \leq \text{im } x$, then there is a series $0 = V_0 \leq V_1 \leq \dots \leq V_{n-1} = \mathfrak{g} \cdot V$ of \mathfrak{g} -submodules such that for all $k = 1, \dots, n-1$, V_k/V_{k-1} bears a compatible \mathbb{K} -vector space structure for which $V_k/V_{k-1} \simeq \bigoplus_{I_{n-k}} \text{Sym}^{n-k} \text{Nat } \mathfrak{g}$ as $\mathbb{K} \mathfrak{g}$ -modules.
- If for all λ , one has $\ker x \leq \ker x_\lambda$ and $\text{im } x_\lambda \leq \text{im } x$, then our series split: $V = \text{Ann}_V(\mathfrak{g}) \oplus \mathfrak{g} \cdot V$, and $\mathfrak{g} \cdot V$ bears a compatible \mathbb{K} -vector space structure for which $\mathfrak{g} \cdot V \simeq \bigoplus_{k=1}^{n-1} \bigoplus_{I_k} \text{Sym}^k \text{Nat } \mathfrak{g}$ as $\mathbb{K} \mathfrak{g}$ -modules.

One should in particular note the immediate consequence:

if V is a simple \mathfrak{g} -module with n minimal such that $x^n \cdot V = 0$, and – either for all $\lambda \in \mathbb{K}$, $\ker x \leq \ker x_\lambda$ – or for all $\lambda \in \mathbb{K}$, $\text{im } x_\lambda \leq \text{im } x$, then $V \simeq \text{Sym}^{n-1} \text{Nat } \mathfrak{g}$.

Without simplicity the statement we gave is quite clumsier due to the trouble one has controlling the cohomology of representations of a Lie ring. The reader will observe that we cannot prove in general that $\mathfrak{g} \cdot V / \text{Ann}_{\mathfrak{g} \cdot V}(\mathfrak{g})$ bears a compatible \mathbb{K} -vector space structure under either assumption $\ker x \leq \ker x_\lambda$ or $\text{im } x_\lambda \leq \text{im } x$: we seem to need both.

The present article starts with a few notations (§1) and basic remarks on length (§2). Section 3 then studies the actions of $\mathfrak{sl}_2(\mathbb{K}_1)$ for a *prime* field \mathbb{K}_1 . Everything is as expected in large enough characteristic (§3.1). In §3.2 we somehow digress by lowering a little the characteristic which results in creating pathologies. These are removed in §3.3 under the assumption that the action is decently “two-sided”. At this point we leave prime fields for the general case and move to Section 4. Our main result Variation n°19 describes the extension of the linear structure from the combinatorial skeleton (i.e., the action at the level of the prime subfield \mathbb{K}_1) to the scalar flesh (i.e., the action at the level of the field \mathbb{K}); it is proved in §4.1. In §4.2 the inclusions $\ker x \leq \ker x_\lambda$ and $\text{im } x_\lambda \leq \text{im } x$ finally appear. A few ideas on their possible meaning are put forth in Section 5.

Technically speaking the only tool is patience; the reader should expect tedious computations. Some results on prime fields may be known or accessible by less naive methods and the article might seem longer than necessary. But it is the philosophy of the present series to do things as explicitly (not to say clumsily) as possible. Finally, studying $\mathfrak{sl}_2(\mathbb{K})$ as a Lie ring may sound somehow arbitrary and the author has no illusions on his results. The real purpose of the study was to prepare him for the future case of the group $\text{SL}_2(\mathbb{K})$.

This work was finished during a visit to the French-Russian “Poncelet” Mathematics Laboratory in Moscow. The author warmly thanks everyone involved, with a special thought for the young lady who likes Belgian chocolates.

1 The setting

This section contains notations and very basic facts which will be used with no reference.

1.1 The Lie ring

\mathbb{K}, \mathfrak{g}

Notation. Let \mathbb{K} be a field and \mathfrak{g} be the Lie ring $\mathfrak{sl}_2(\mathbb{K})$.

Literature on Lie rings looks scarce when compared to other topics. Fortunately we deal with a concrete and familiar Lie ring, so any reference on Lie algebras such as [3] will do. We simply use the group law $+$, the bracket $[\cdot, \cdot]$, and forget about the \mathbb{K} -linear structure on \mathfrak{g} .

$\mathbb{K}_1, \mathfrak{g}_1$

Notation. Let \mathbb{K}_1 be the prime subfield of \mathbb{K} and \mathfrak{g}_1 be the Lie ring $\mathfrak{sl}_2(\mathbb{K}_1)$; one has $\mathfrak{g}_1 \leq \mathfrak{g}$.

$h_\lambda, x_\lambda, y_\lambda$

Notation. For $\lambda \in \mathbb{K}$ let

$$h_\lambda = \begin{pmatrix} \lambda & 0 \\ 0 & -\lambda \end{pmatrix}, \quad x_\lambda = \begin{pmatrix} 0 & \lambda \\ 0 & 0 \end{pmatrix}, \quad y_\lambda = \begin{pmatrix} 0 & 0 \\ \lambda & 0 \end{pmatrix}$$

h, x, y

One simply writes $h = h_1, x = x_1, y = y_1$.

$\mathfrak{b}, \mathfrak{u}, \mathfrak{t}$

Notation. Let \mathfrak{b} be the Borel subring generated by the h_λ 's and the x_μ 's, and $\mathfrak{u} = \{x_\mu : \mu \in \mathbb{K}\}$ be its unipotent radical. Let \mathfrak{t} be the Cartan subring $\{h_\lambda : \lambda \in \mathbb{K}\}$.

Relations.

- $[h_\lambda, x_\mu] = 2x_{\lambda\mu}$;
- $[h_\lambda, y_\nu] = -2y_{\lambda\nu}$;
- $[x_\mu, y_\nu] = h_{\mu\nu}$.

\mathbb{K} will never have characteristic 2; as a consequence \mathfrak{g} will always be perfect. One should be careful that $[\mathfrak{g}, \mathfrak{g}]$ will merely denote the *additive subgroup* of \mathfrak{g} generated by all brackets since we forget about the \mathbb{K} -linear structure on \mathfrak{g} . It is however the case that $\mathfrak{g} = [\mathfrak{g}, \mathfrak{g}]$ which is the definition of perfectness.

We shall sometimes go to the enveloping (associative) ring which is defined among rings just like the enveloping (associative) \mathbb{K} -algebra is defined among \mathbb{K} -algebras. It enjoys a similar universal property in the broader category of \mathfrak{g} -modules. This simply amounts to viewing $\mathfrak{sl}_2(\mathbb{K})$ as a Lie algebra over the prime ring of \mathbb{K} and taking its enveloping algebra as such, but to prevent confusion we shall always refer to this object as the enveloping ring. The usual enveloping \mathbb{K} -algebra can be retrieved as a quotient of the enveloping Lie ring by relations expressing \mathbb{K} -linearity. It has no reason to play a role since we are a priori not dealing with \mathbb{K} -linear representations.

Relations. One has in the enveloping ring the following equalities which the reader may check by induction:

$$x^i y = y x^i + i(h+1-i)x^{i-1}; \quad (1)$$

$$y^j x = x y^j - j(h+j-1)y^{j-1}; \quad (2)$$

$$y^j x^i = \sum_{k=0}^{\min(i,j)} (-1)^k k! \binom{i}{k} \binom{j}{k} \left(\prod_{\ell=0}^{k-1} (h-i+j+\ell) \right) x^{i-k} y^{j-k}; \quad (3)$$

$$x^i h_\mu = h_\mu x^i - 2i x_\mu x^{i-1}; \quad (4)$$

$$\begin{aligned} x_{\lambda_1} \dots x_{\lambda_i} y_\mu &= y_\mu x_{\lambda_1} \dots x_{\lambda_i} + \sum_k h_{\mu \cdot \lambda_k} x_{\lambda_1} \dots \widehat{x_{\lambda_k}} \dots x_{\lambda_i} \\ &\quad - \sum_{k \neq \ell} x_{\mu \cdot \lambda_k \cdot \lambda_\ell} x_{\lambda_1} \dots \widehat{x_{\lambda_k}} \dots \widehat{x_{\lambda_\ell}} \dots x_{\lambda_i}; \end{aligned} \quad (5)$$

$$\begin{aligned} y_\mu x_{\lambda_1} \dots x_{\lambda_i} &= x_{\lambda_1} \dots x_{\lambda_i} y_\mu - \sum_k x_{\lambda_1} \dots \widehat{x_{\lambda_k}} \dots x_{\lambda_i} h_{\mu \cdot \lambda_k} \\ &\quad - \sum_{k \neq \ell} x_{\lambda_1} \dots \widehat{x_{\lambda_k}} \dots \widehat{x_{\lambda_\ell}} \dots x_{\lambda_i} x_{\mu \cdot \lambda_k \cdot \lambda_\ell} \end{aligned} \quad (6)$$

(The terms in the hats do not appear.)

Be however careful that in the enveloping ring $x_\lambda y \neq y x_\lambda$. So checking the formulas in the enveloping algebra does not suffice in order to establish them in the enveloping ring.

Notation. Let $c_1 = 2xy + 2yx + h^2$ be the Casimir operator.

The Casimir operator is central in the enveloping algebra but not in the enveloping ring; for instance a quick computation yields $[c_1, h_\lambda] = 8(xy_\lambda - x_\lambda y)$ which is non-zero.

However when $\mathbb{K}_1 = \mathbb{F}_p$ and $\mathfrak{g}_1 = \mathfrak{sl}_2(\mathbb{K}_1)$, c_1 is central indeed in the enveloping ring of \mathfrak{g}_1 . This is not quite true over \mathbb{Q} , but it is readily checked that for all z in the enveloping ring there is an integer k with $k[c_1, z] = 0$. It follows that provided $\mathbb{K}_1 = \mathbb{Q}$ and V is a *torsion-free* $\mathfrak{g}_1 = \mathfrak{sl}_2(\mathbb{K}_1)$ -module, the action of c_1 commutes with the action of \mathfrak{g}_1 . This will always be the case when we use c_1 .

1.2 The module

V **Notation.** Let V be a \mathfrak{g} -module.

We shall keep writing $x_\lambda, y_\lambda, h_\mu$ for the images in $\text{End}(V)$ of the corresponding elements of \mathfrak{g} .

$\lambda_u(V)$ **Notation.** The length $\lambda_u(V)$ of the u -module V is the least integer n , if there is one such, with $u^n \cdot V = 0$.

E_i **Notation.** For $i \in \mathbb{Z}$, let $E_i = E_i(V) = \{a \in V : h \cdot a = iv\}$.

Using the familiar relations one sees that h_λ (resp. x_μ , resp. y_ν) maps E_i to E_i (resp. E_{i+2} , resp. E_{i-2}).

We shall in a minute deal with constructing vector space structures on modules. If \mathbb{K}_1 is a prime field then an abelian group V bears at most one linear structure over \mathbb{K}_1 . If it is the case and V is a $\mathfrak{g}_1 = \mathfrak{sl}_2(\mathbb{K}_1)$ -module as well then V is a $\mathbb{K}_1\mathfrak{g}_1$ -module.

Let us also remind the reader why Lie rings do not admit cross-characteristic representations.

Observation. Let \mathbb{K} be a field, \mathbb{K}_1 its prime subfield, $\mathfrak{g} = \mathfrak{sl}_2(\mathbb{K})$ and V be a \mathfrak{g} -module. Then $\mathfrak{g} \cdot V / \text{Ann}_{\mathfrak{g} \cdot V}(\mathfrak{g})$ is a \mathbb{K}_1 -vector space.

Proof of Claim. If $\mathbb{K}_1 = \mathbb{F}_p$ then \mathfrak{g} annihilates pV so $V / \text{Ann}_V(\mathfrak{g})$ has exponent p . Also note that p annihilates $\mathfrak{g} \cdot V$, so $\mathfrak{g} \cdot V$ has exponent p as well. Hence in prime characteristic both $V / \text{Ann}_V(\mathfrak{g})$ and $\mathfrak{g} \cdot V$ are actually \mathbb{K}_1 -vector spaces.

If $\mathbb{K}_1 = \mathbb{Q}$ then $\mathfrak{g} \cdot V$ is divisible and \mathfrak{g} annihilates the torsion submodule of V , so $\mathfrak{g} \cdot V / \text{Ann}_{\mathfrak{g} \cdot V}(\mathfrak{g})$ is torsion-free and divisible: a \mathbb{Q} -vector space. \diamond

This certainly does not prove that V need be a \mathbb{K} -vector space (which is not true in general) but already removes the outmost pathologies.

1.3 Symmetric powers

Let us finally state a few facts about the very modules we try to characterize.

$\text{Nat } \mathfrak{g}$ **Notation.** Let $\text{Nat } \mathfrak{g}$ denote the natural representation of \mathfrak{g} , that is \mathbb{K}^2 equipped with the left action of $\mathfrak{g} = \mathfrak{sl}_2(\mathbb{K})$.

$\text{Sym}^k \text{Nat } \mathfrak{g}$ **Notation.** For $k \geq 1$ an integer, let $\text{Sym}^k \text{Nat } \mathfrak{g}$ denote the k^{th} symmetric power of $\text{Nat } \mathfrak{g}$.

We do not wish to go into tensor algebra, and will more conveniently handle $\text{Sym}^k \text{Nat } \mathfrak{g}$ as follows.

Fact ([3, §II.7]). Let \mathbb{K} be a field of characteristic 0 or $\geq k + 1$.

- $S_k = \text{Sym}^k \text{Nat } \mathfrak{g}$ is isomorphic to $\mathbb{K}[Y^k, XY^{k-1}, \dots, X^k]$ as a $\mathbb{K}\mathfrak{g}$ -module, where x acts as $X \frac{\partial}{\partial Y}$ and y as $Y \frac{\partial}{\partial X}$.
- S_k is an irreducible $\mathbb{K}\mathfrak{g}$ -module; it remains irreducible as a \mathfrak{g} -module.
- $h = [x, y]$ acts on S_k as $X \frac{\partial}{\partial X} - Y \frac{\partial}{\partial Y}$.
- $\mathbb{K} \cdot X^{k-i} Y^i = E_{k-2i}(S_k)$.
- The length of S_k is $k + 1$, meaning that $u^{k+1} \cdot S_k = 0$ and $u^k \cdot S_k \neq 0$.
- The Casimir operator c_1 acts on S_k as multiplication by $k(k + 2)$. In particular in characteristic 0 or $\geq k + 3$, c_1 induces a bijection of S_k .

2 Length

This section contains two minor results on the notion of length as defined in §1. They are fairly straightforward and so are their proofs.

Variation n°15. *Let $n \geq 2$ be an integer and \mathbb{K} be a field of characteristic 0 or $\geq n+1$. Let $\mathfrak{g} = \mathfrak{sl}_2(\mathbb{K})$, $\mathfrak{b} \leq \mathfrak{g}$ be a Borel subring and $\mathfrak{u} = \mathfrak{b}'$ be its radical. Let V be a \mathfrak{u} -module. Suppose that for all $\lambda \in \mathbb{K}$, $x_\lambda^n = 0$ in $\text{End}(V)$. Then V has \mathfrak{u} -length at most n , meaning that $\mathfrak{u}^n \cdot V = 0$.*

Proof. This is a simpler analog of Variation n°6 [1]: only the end of the argument need be reproduced as the induction on the “weights” of monomials is not necessary. Indeed $x_{\lambda+\mu} = x_\lambda + x_\mu$ whence immediately:

$$0 = \sum_{j=1}^{n-1} \binom{n}{j} x_\lambda^j x_\mu^{n-j}$$

One then replaces μ by $i\mu$ for $i = 1 \dots n-1$; this yields the same linear $(n-1)$ by $(n-1)$ system as in Variation n°6 [1]. Hence $dx_\lambda^{n-1} x_\mu = 0$ where d is the determinant of the system; all prime divisors of d divide $n!$. In particular replacing μ by $\frac{\mu}{d}$ in \mathbb{K} , one finds $x_\lambda^{n-1} x_\mu = 0$ in $\text{End}(V)$. Since \mathfrak{u} acts on $\text{im } x_\mu$ one may use induction on n . \square

Next comes an easy generalization of Variation n°9 [1]. (There seems to be no parallel argument in the case of $\text{SL}_2(\mathbb{K})$; the quadratic setting painfully dealt with in Variation n°7 [1] actually required a full $\text{SL}_2(\mathbb{K})$ -module.)

Variation n°16. *Let $n \geq 2$ be an integer and \mathbb{K} be a field of characteristic 0 or $\geq n+1$. Let $\mathfrak{g} = \mathfrak{sl}_2(\mathbb{K})$ and $\mathfrak{b} \leq \mathfrak{g}$ be a Borel subring. Let V be a \mathfrak{b} -module. Suppose that $x^n \cdot V = 0$. Then V has \mathfrak{u} -length at most n , meaning that $\mathfrak{u}^n \cdot V = 0$.*

Proof. We go to $\text{End}(V)$. Let us prove by induction on $i = 0 \dots n$:

$$\forall (\lambda_1, \dots, \lambda_i) \in \mathbb{K}^i, x^{n-i} x_{\lambda_i} \dots x_{\lambda_1} = 0$$

This holds of $i = 0$. Suppose that the result holds of fixed $i < n$; let $(\lambda_1, \dots, \lambda_i, \lambda_{i+1}) \in \mathbb{K}^{i+1}$.

We show by induction on $j = 0 \dots i$:

$$x^{n-i} x_{\lambda_i} \dots x_{\lambda_{j+1}} h_{\lambda_{i+1}} x_{\lambda_j} \dots x_{\lambda_1} = 0$$

- This holds of $j = 0$ by assumption on i .
- Suppose that the result holds of fixed j . Then:

$$\begin{aligned} & x^{n-i} x_{\lambda_i} \dots x_{\lambda_{j+2}} h_{\lambda_{i+1}} x_{\lambda_{j+1}} \dots x_{\lambda_1} \\ &= x^{n-i} x_{\lambda_i} \dots x_{\lambda_{j+2}} ([h_{\lambda_{i+1}}, x_{\lambda_{j+1}}] + x_{\lambda_{j+1}} h_{\lambda_{i+1}}) x_{\lambda_j} \dots x_{\lambda_1} \\ &= 2x^{n-i} x_{\lambda_i} \dots x_{\lambda_{j+2}} x_{\lambda_{i+1} \cdot \lambda_{j+1}} x_{\lambda_j} \dots x_{\lambda_1} \\ &\quad + x^{n-i} x_{\lambda_i} \dots x_{\lambda_{j+1}} h_{\lambda_{i+1}} x_{\lambda_j} \dots x_{\lambda_1} \\ &= 0 \end{aligned}$$

by assumption on j and i (the latter applied with $\lambda'_{j+1} = \lambda_{i+1} \cdot \lambda_{j+1}$). This concludes the induction on j .

With $j = i$, one gets:

$$x^{n-i} h_{\lambda_{i+1}} x_{\lambda_i} \dots x_{\lambda_1} = 0$$

Let us now prove by induction on $k = 0 \dots n-i$:

$$x^{n-(i+k)} h_{\lambda_{i+1}} x^k x_{\lambda_i} \dots x_{\lambda_1} = 2k x^{n-(i+1)} x_{\lambda_{i+1}} \dots x_{\lambda_1}$$

- This holds of $k = 0$ by what we have just shown.
- Suppose that the result holds of fixed k . Then:

$$\begin{aligned} & x^{n-(i+k+1)} h_{\lambda_{i+1}} x^{k+1} x_{\lambda_i} \dots x_{\lambda_1} \\ &= x^{n-(i+k+1)} ([h_{\lambda_{i+1}}, x] + x h_{\lambda_{i+1}}) x^k x_{\lambda_i} \dots x_{\lambda_1} \\ &= 2x^{n-(i+k+1)} x_{\lambda_{i+1}} x^k x_{\lambda_i} \dots x_{\lambda_1} + x^{n-(i+k)} h_{\lambda_{i+1}} x^k x_{\lambda_i} \dots x_{\lambda_1} \\ &= 2x^{n-(i+1)} x_{\lambda_{i+1}} \dots x_{\lambda_1} + 2k x^{n-(i+1)} x_{\lambda_{i+1}} \dots x_{\lambda_1} \end{aligned}$$

This concludes the induction on k .

With $k = n - i$ one gets:

$$h_{\lambda_{i+1}} x^{n-i} x_{\lambda_i} \dots x_{\lambda_1} = 2(n-i) x^{n-(i+1)} x_{\lambda_{i+1}} \dots x_{\lambda_i}$$

but the left-hand side is zero by assumption. If instead of λ_{i+1} we had started with $\frac{\lambda_{i+1}}{2(n-i)}$, which is legitimate by assumption on the characteristic of \mathbb{K} , we would have obtained:

$$x^{n-(i+1)} x_{\lambda_{i+1}} \dots x_{\lambda_1} = 0$$

This concludes the induction on i .

With $i = n$, one has the desired statement. \square

Remark. One cannot use induction on n via $\text{im } x$ since $\text{im } x$ may fail to be $\mathfrak{t} = \{h_\lambda : \lambda \in \mathbb{K}\}$ -invariant; such a configuration will be met in the example illustrating the following.

Remark. The mere existence of a product $x_{\lambda_1} \dots x_{\lambda_n}$ which is zero in $\text{End}(V)$ does not suffice to force the length to be at most n . Take indeed $\mathbb{K} = \mathbb{C}$, $\mathfrak{g} = \mathfrak{sl}_2(\mathbb{C})$, and let φ stand for complex conjugation. Also let $V = \text{Nat } \mathfrak{g} \simeq \mathbb{C}^2$, $V' = {}^\varphi V$ (a copy “twisted” by the field automorphism), and $W = V \otimes V'$. One sees that W has no $\mathbb{C}\mathfrak{g}$ -submodule other than $\{0\}$ and W .

Let (e_1, e_2) be the standard basis of \mathbb{C}^2 ; one has $x \cdot e_1 = 0$ and $x \cdot e_2 = e_1$. Write for simplicity $e_{i,j} = e_i \otimes e_j$. One finds:

$$\begin{cases} x_\lambda \cdot e_{1,1} &= 0 \\ x_\lambda \cdot e_{2,1} &= \lambda e_{1,1} \\ x_\lambda \cdot e_{1,2} &= \varphi(\lambda) e_{1,1} \\ x_\lambda \cdot e_{2,2} &= \lambda e_{1,2} + \varphi(\lambda) e_{2,1} \end{cases}$$

so that $x_\lambda x_\mu \cdot e_{2,2} = (\lambda\varphi(\mu) + \mu\varphi(\lambda))e_{1,1}$. Clearly $x_1^2 \neq 0$ and yet $x_1 x_i = 0$ (where i stands for a root of -1).

One may object that W though simple as a $\mathbb{C}\mathfrak{g}$ -module, is not as a \mathfrak{g} -module; we then go down to $W_0 = \mathbb{R}e_{1,1} \oplus \{\lambda e_{1,2} + \varphi(\lambda) e_{2,1} : \lambda \in \mathbb{C}\} \oplus \mathbb{R}e_{2,2}$, which as a \mathfrak{g} -module is simple; one has $x_1^2 \neq 0$ in $\text{End}(W_0)$.

Remark. By Variation n°16 the u -length is therefore the nilpotence order of x in $\text{End}(V)$; one may wonder whether it is the nilpotence order of y as well. (One should not expect this in full generality: after Variation n°12 [1] we saw that it can be achieved in characteristic 3 that $x^2 = 0 \neq y^2$.)

Here is an unsatisfactory argument in characteristic zero.

We go to the enveloping algebra \mathfrak{A} . Then $\text{Ann}_{\mathfrak{A}}(V)$ is a two-sided ideal containing x^n . But $\text{SL}_2(\mathbb{K})$ acts on \mathfrak{A} and normalizes every two-sided ideal by [2, Proposition 2.4.17]; since the Weyl group exchanges x and y one has $y^n \in \text{Ann}_{\mathfrak{A}}(V)$ as well, whence $y^n = 0$ in $\text{End}(V)$.

The argument is not quite satisfactory: we have been using the \mathbb{K} -algebra \mathfrak{A} . It is a fact that every $\mathbb{K}\mathfrak{g}$ -module is an \mathfrak{A} -module but all we had was a mere \mathfrak{g} -module; turning it into a $\mathbb{K}\mathfrak{g}$ -module is precisely the core of the matter.

More prosaically, a crude computation will show that in characteristic $\geq 2n + 1$ one does have $x^n = 0 \Rightarrow y^n = 0$: we shall see this while proving Variation n°17.

3 Combinatorial skeleton

In this section we focus on $\mathfrak{sl}_2(\mathbb{K}_1)$ -modules of finite length, with \mathbb{K}_1 a prime field. If the characteristic is 0 or large enough, Variation n°17 of §3.1 gives a complete description. But some other objects appear if one tries to lower the characteristic too much (§3.2). Provided one assumes that y has the same order as x , the monsters vanish (Variation n°18, §3.3).

The author cannot believe that the results of this section are new, but found no evidence. We shall give purely computational arguments without going to the algebraic closure. Again, this reflects a methodological line more than pure foolishness.

3.1 Large Enough Characteristic

Variation n°17. Let $n \geq 2$ be an integer and \mathbb{K}_1 be a prime field of characteristic 0 or $\geq 2n + 1$. Let $\mathfrak{g}_1 = \mathfrak{sl}_2(\mathbb{K}_1)$ and V be a \mathfrak{g}_1 -module. If the characteristic of \mathbb{K} is 0 one requires V to be torsion-free. Suppose that $x^n = 0$ in $\text{End}(V)$.

Then $V = \text{Ann}_V(\mathfrak{g}_1) \oplus \mathfrak{g}_1 \cdot V$, and $\mathfrak{g}_1 \cdot V$ is a \mathbb{K}_1 -vector space with $\mathfrak{g}_1 \cdot V \simeq \bigoplus_{k=1}^{n-1} \oplus_{I_k} \text{Sym}^k \text{Nat } \mathfrak{g}_1$ as $\mathbb{K}_1 \mathfrak{g}_1$ -modules.

Proof. Induction on n . When $n = 2$ this is Variation n°12 [1].

All along $c_1 = 2xy + 2yx + h^2$ will be the Casimir operator; the action of c_1 commutes with the action of \mathfrak{g}_1 on V . In characteristic 0 this holds only since we assumed V to be torsion-free.

Step 1 (see Variation n°3 [1]). We may assume $V = \mathfrak{g}_1 \cdot V$ and $\text{Ann}_V(\mathfrak{g}_1) = 0$.

Proof of Claim. Let $W = \mathfrak{g}_1 \cdot V$ and $\overline{W} = W / \text{Ann}_W(\mathfrak{g}_1)$. Let $\bar{\cdot}$ stand for projection modulo $\text{Ann}_W(\mathfrak{g}_1)$. By perfectness of \mathfrak{g}_1 one has $\overline{W} = \mathfrak{g}_1 \cdot \overline{W}$ and $\text{Ann}_{\overline{W}}(\mathfrak{g}_1) = 0$. In particular if $\mathbb{K}_1 = \mathbb{Q}$ then \overline{W} is torsion-free. Suppose that the result holds of \overline{W} ; let us prove it for V : suppose that \overline{W} is a \mathbb{K}_1 -vector space satisfying $\overline{W} \simeq \bigoplus_{k=1}^{n-1} \oplus_{I_k} \text{Sym}^{n-1} \text{Nat } \mathfrak{g}_1$ as $\mathbb{K}_1 \mathfrak{g}_1$ -modules.

One then sees that c_1 is a bijection of \overline{W} . We claim the following:

- $W = c_1 \cdot W + \text{Ann}_W(\mathfrak{g}_1)$. For take $w \in W$. Since c_1 is surjective onto \overline{W} there exists $w' \in W$ with $\bar{w} = c_1 \cdot \bar{w}'$.
- $c_1 \cdot W = W$. Let us apply \mathfrak{g}_1 to the previous equality, bearing in mind perfectness of \mathfrak{g}_1 and centralness of c_1 . One finds $W = \mathfrak{g}_1 \cdot W = \mathfrak{g}_1 c_1 \cdot W + \mathfrak{g}_1 \cdot \text{Ann}_W(\mathfrak{g}_1) = c_1 \mathfrak{g}_1 \cdot W = c_1 \cdot W$.
- $W \cap \ker c_1 = 0$. For take $w \in W \cap \ker c_1$. Then by the previous claim there exists $w' \in W$ with $w = c_1 \cdot w'$. Modulo $\text{Ann}_W(\mathfrak{g}_1)$ one has $0 = c_1 \cdot \bar{w} = c_1^2 \cdot \bar{w}'$. By injectivity of the Casimir operator on \overline{W} it follows $\bar{w}' = 0$, whence $w' \in \text{Ann}_V(\mathfrak{g}_1) \leq \ker c_1$ and $w = c_1 \cdot w' = 0$.
- $\text{Ann}_V(\mathfrak{g}_1) = \ker c_1$. One inclusion is obvious and was just used; if conversely $k \in \ker c_1$ then $\mathfrak{g}_1 \cdot \langle c_1 \cdot k \rangle = 0 = c_1 \cdot \langle \mathfrak{g}_1 \cdot k \rangle$ so $\langle \mathfrak{g}_1 \cdot k \rangle \leq W \cap \ker c_1 = 0$.
- $V = \text{Ann}_V(\mathfrak{g}_1) \oplus W$. The sum is direct indeed as we just saw. Moreover if $v \in V$ then there exists $w \in W$ with $c_1 \cdot v = c_1 \cdot w$; in particular $V \leq W + \ker c_1 = W \oplus \ker c_1 = W \oplus \text{Ann}_V(\mathfrak{g}_1)$.

V therefore has the desired structure. \diamond

We now suppose $V = \mathfrak{g}_1 \cdot V$ and $\text{Ann}_V(\mathfrak{g}_1) = 0$. It follows that $V \simeq (\mathfrak{g}_1 \cdot V) / (\text{Ann}_{\mathfrak{g}_1 \cdot V}(\mathfrak{g}_1))$ is a \mathbb{K}_1 -vector space.

Step 2. In $\text{End}(V)$, one has $(h - n + 1)(h - n + 2) \dots (h + n - 1) = 0$.

Proof of Claim. Remember that in the enveloping ring, for $i, j \geq 1$:

$$y^j x^i = \sum_{k=0}^{\min(i,j)} \left[(-1)^k k! \binom{i}{k} \binom{j}{k} \left(\prod_{\ell=0}^{k-1} (h - i + j + \ell) \right) x^{i-k} y^{j-k} \right]$$

In the subring of $\text{End}(V)$ generated by the image of \mathfrak{g}_1 one has $x^n = 0$; the formula becomes with $i = n$ and $j \leq n$:

$$\sum_{k=1}^j (-1)^k k! \binom{n}{k} \binom{j}{k} \left(\prod_{\ell=0}^{k-1} (h - n + j + \ell) \right) x^{n-k} y^{j-k} = 0 \quad (F_j)$$

Let us prove by induction on $j = 0 \dots n$:

$$(h - n + 1)(h - n + 2) \dots (h - n + 2j - 1) x^{n-j} = 0$$

When $j = 0$ the (ascending) product is empty: our claim holds by assumption on x . Suppose that the result holds of fixed j and let us prove it for $j + 1 \leq n$. Consider formula (F_{j+1}) multiplied on the left by $(h - n + 1) \dots (h - n + j)$. One gets:

$$\sum_{k=1}^{j+1} (-1)^k k! \binom{n}{k} \binom{j+1}{k} \pi_k x^{n-k} y^{j+1-k} = 0$$

where:

$$\begin{aligned}\pi_k &= (h-n+1)\dots(h-n+j)\cdot(h-n+j+1)\dots(h-n+j+k) \\ &= (h-n+1)\dots(h-n+j+k)\end{aligned}$$

Since $j+k \geq 2k-1$ the term with index k contains $(h-n+1)\dots(h-n+2k-1)x^{n-k}$, which by induction is zero while $k \leq j$. So only remains the term with index $k = j+1$ namely:

$$(-1)^{j+1}(j+1)! \binom{n}{j+1} (h-n+1)\dots(h-n+2j+1)x^{n-(j+1)} = 0$$

By $n!$ -torsion-freeness of V we may remove the coefficients and complete the induction. When $j = n$ one finds $(h-n+1)(h-n+2)\dots(h+n-1) = 0$. \diamond

Step 3. $V = \oplus_{j=1-n}^{n-1} E_j$.

Proof of Claim. Let us first observe that the sum $\oplus_{j=1-n}^{n-1} E_j$ is direct indeed by $(2n-2)!$ -torsion-freeness of V . For the same reason the monomials $X-j$ are pairwise coprime in $\mathbb{K}_1[X]$ for $j = 1-n, \dots, n-1$. Since their product annihilates h in $\text{End}(V)$ one has $V = \oplus_{j=1-n}^{n-1} \ker(h-j) = \oplus_{j=1-n}^{n-1} E_j$. \diamond

Since $n-1+2 = n+1$ and $n-2+2 = n$ are not congruent to any $j \in \{1-n, \dots, n-1\}$ the operator x annihilates E_{n-1} and E_{n-2} . Similarly y annihilates E_{1-n} and E_{2-n} .

Remark. It is now clear that $y^n \cdot V = 0$.

Notation 4. Let $V_\perp = \text{im}(c_1 - n^2 + 1)$ and $V_\top = \ker(c_1 - n^2 + 1)$.

Step 5. V_\perp is a \mathfrak{g}_1 -submodule isomorphic to $\oplus_{k=1}^{n-2} \oplus_{I_k} \text{Sym}^k \text{Nat } \mathfrak{g}_1$.

Proof of Claim. V_\perp is clearly \mathfrak{g}_1 -invariant. But by Step 3 or the proof of Step 2 one has in $\text{End}(V)$ the identity $hx^{n-1} = (n-1)x^{n-1}$. Hence always in $\text{End}(V)$:

$$\begin{aligned}x^{n-1}(c_1 - n^2 + 1) &= x^{n-1}(2xy + 2yx + h^2 - n^2 + 1) \\ &= 2(x^{n-1}y)x + (h+2-2n)^2 x^{n-1} - (n^2-1)x^{n-1} \\ &= 2(yx^{n-1} + (n-1)(h+2-n)x^{n-2})x \\ &\quad + (1-n)^2 x^{n-1} + (1-n^2)x^{n-1} \\ &= 2(n-1)(h+2-n)x^{n-1} + 2(1-n)x^{n-1} \\ &= 0\end{aligned}$$

It follows that x^{n-1} annihilates $\text{im}(c_1 - n^2 + 1) = V_\perp$ and one may apply induction. Since $\text{Ann}_{V_\perp}(\mathfrak{g}_1) \leq \text{Ann}_V(\mathfrak{g}_1) = 0$ there remains only $V_\perp = \mathfrak{g}_1 \cdot V_\perp \simeq \oplus_{k=1}^{n-2} \oplus_{I_k} \text{Sym}^k \text{Nat } \mathfrak{g}_1$. \diamond

Step 6. We may assume $V = V_\top$.

Proof of Claim. We claim that $V = V_\perp \oplus V_\top$. The way the Casimir operator acts on each $\text{Sym}^k \text{Nat } \mathfrak{g}_1$ is known: like multiplication by $k(k+2)$. But for $k = 1, \dots, n-2$, one has $k(k+2) \neq n^2 - 1$ in \mathbb{K}_1 by assumption on the characteristic. It follows that $(c_1 - n^2 + 1)$ induces a bijection of V_\perp . As a consequence $V_\perp \cap V_\top = V_\perp \cap \ker(c_1 - n^2 + 1) = 0$. Moreover for all $v \in V$ there exists $v' \in V_\perp$ with $(c_1 - n^2 + 1) \cdot v = (c_1 - n^2 + 1) \cdot v'$, whence $V = V_\perp + \ker(c_1 - n^2 + 1) = V_\perp + V_\top = V_\perp \oplus V_\top$. If the result were proved for V_\top it would therefore follow for V . \diamond

From now on we suppose $V = V_\top$; in particular $c_1 - n^2 + 1$ annihilates V .

Step 7. $\ker x = E_{n-1}$.

Proof of Claim. We claim that x is injective on $\oplus_{j=1-n}^{n-2} E_j$. For if $a \in E_j$ with $j \in \{1-n, \dots, n-2\}$ satisfies $x \cdot a = 0$, then

$$(n^2-1)a = c_1 \cdot a = (2xy + 2yx + h^2) \cdot a = (2h + h^2) \cdot a = j(j+2)a$$

so either $a = 0$ or $n^2 = j(j+2) + 1 = (j+1)^2$. But the latter equation solves into $j = \pm n - 1$ which is not the case (even in characteristic $p \geq 2n + 1$). \diamond

Step 8. V is isomorphic to $\oplus_{I_{n-1}} \text{Sym}^{n-1} \text{Nat } \mathfrak{g}_1$.

Proof of Claim. We claim that for all $i = 1 \dots n$, one has $E_{n-2i} = 0$. At $i = 1$ this is because $E_{n-2} \leq \ker x = E_{n-1}$ by Step 7. If this is known at i , then $x \cdot E_{n-2(i+1)} \leq E_{n-2i}$ whence $E_{n-2(i+1)} \leq \ker x = E_{n-1}$.

On the other hand observe that for all $i = 1 \dots n$: $yx|_{E_{n+1-2i}} = (i-1)(n+1-i)$ and $xy|_{E_{n+1-2i}} = i(n-i)$. This is actually obvious since $c_1 = 4yx + h^2 + 2h = 4xy + h^2 - 2h$ is constant and equals multiplication by $n^2 - 1$.

It is therefore now clear that for all $a_{n-1} \in E_{n-1} \setminus \{0\}$, the span $\langle \mathfrak{g}_1 \cdot a_{n-1} \rangle$ is a \mathbb{K}_1 -vector space isomorphic to $\text{Sym}^{n-1} \text{Nat } \mathfrak{g}_1$ as a $\mathbb{K}_1 \mathfrak{g}_1$ -module; in particular $b \in \langle \mathfrak{g}_1 \cdot a_{n-1} \rangle \setminus \{0\}$ then $\langle \mathfrak{g}_1 \cdot a_{n-1} \rangle = \langle \mathfrak{g}_1 \cdot b \rangle$. Let $M \leq V$ be a maximal direct sum of such spaces. Then M has the desired structure, and our computations show $V = \bigoplus_{j=1-n}^{n-1} E_j \leq \mathfrak{g}_1 \cdot E_{n-1} \leq M$. \diamond

This finishes the proof. \square

3.2 A Digression: Pathologies in Low Characteristic

As in Variation n°12 [1] the characteristic must be 0 or $\geq 2n + 1$ in order to prove Variation n°17. In this section we suppose the characteristic to lie between n and $2n$. We shall construct counterexamples to Variation n°17 and remove them later in §3.3 under the extra assumption that y has the same order as x in $\text{End}(V)$.

The construction generalizes the one given in characteristic 3 at the end of [1]. Let $n \geq 2$ be an integer and p be a prime number with $n < p < 2n$; let m be such that $n + m = p$. Observe that if $i \in \{1, \dots, n\}$ and $j \in \{1, \dots, m\}$, then $n + 1 - 2i$ and $m + 1 - 2j$ are never congruent modulo p . Hence modulo p , the $n + 1 - 2i$'s and $m + 1 - 2j$'s are all distinct, and their global number is p ; there are n of the former kind and m of the latter.

Construction. Let V_1 and V_2 be two vector spaces over \mathbb{F}_p . Let $\alpha : V_1 \rightarrow V_2$ and $\beta : V_2 \rightarrow V_1$ be two linear maps. Define a \mathfrak{g}_1 -module $\mathbb{S}_{\alpha, \beta}$ as follows.

For each $j \in \{1, \dots, m\}$ let E_{m+1-2j} be a copy of V_1 whose elements we shall denote e_{m+1-2j, v_1} for $v_1 \in V_1$. For each $i \in \{1, \dots, n\}$ let $E_{n+1-2i} = \{e_{n+1-2i, v_2} : v_2 \in V_2\}$ be a copy of V_2 .

The underlying vector space of $\mathbb{S}_{\alpha, \beta}$ is

$$\left(\bigoplus_{j=1}^m E_{m+1-2j} \right) \bigoplus \left(\bigoplus_{i=1}^n E_{n+1-2i} \right)$$

Now define an action of $\mathfrak{g}_1 = \mathfrak{sl}_2(\mathbb{F}_p)$ on $\mathbb{S}_{\alpha, \beta}$ by:

$$\begin{aligned} h \cdot e_{m+1-2j, v_1} &= (m+1-2j) \cdot e_{m+1-2j, v_1} && \text{if } 1 \leq j \leq m \\ x \cdot e_{m+1-2j, v_1} &= (j-1)e_{m+1-2(j-1), v_1} && \text{if } 1 \leq j \leq m \\ y \cdot e_{m+1-2j, v_1} &= (m-j)e_{m+1-2(j+1), v_1} && \text{if } 1 \leq j < m \\ y \cdot e_{1-m, v_1} &= e_{n-1, \alpha(v_1)} \\ h \cdot e_{n+1-2i, v_2} &= (n+1-2i) \cdot e_{n+1-2i, v_2} && \text{if } 1 \leq i \leq n \\ x \cdot e_{n+1-2i, v_2} &= (i-1)e_{n+1-2(i-1), v_2} && \text{if } 1 \leq i \leq n \\ y \cdot e_{n+1-2i, v_2} &= (n-i)e_{n+1-2(i+1), v_2} && \text{if } 1 \leq i < n \\ y \cdot e_{1-n, v_2} &= e_{m-1, \beta(v_2)} \end{aligned}$$

Note that by construction, x annihilates E_{n-1} and E_{m-1} . The following diagram explains the construction:

$$\begin{array}{c} \left. \begin{array}{c} \bigoplus \text{Sym}^{m-1} \\ \text{Nat } \mathfrak{g}_1 \end{array} \right\} \left\{ \begin{array}{c} E_{m-1} \xrightleftharpoons[x]{y} \dots \xrightleftharpoons[x]{y} E_{1-m} \\ \beta \uparrow \qquad \qquad \qquad \downarrow \alpha \\ E_{1-n} \xrightleftharpoons[x]{y} \dots \xrightleftharpoons[x]{y} E_{n-1} \end{array} \right\} \bigoplus \text{Sym}^{n-1} \\ \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \text{Nat } \mathfrak{g}_1 \end{array}$$

Observation. $\mathbb{S}_{\alpha, \beta}$ is a \mathfrak{g}_1 -module annihilated by x^n .

Proof of Claim. It suffices to prove that the defining relations of \mathfrak{g}_1 are satisfied at every vector e_{m+1-2j, v_1} ; the e_{n+1-2i, v_2} 's are treated similarly.

At e_{m+1-2j, v_1} with $1 < j < m$ there is nothing to prove since everything is locally as in $\text{Sym}^{n-1} \text{Nat } \mathfrak{g}_1$; since x annihilates e_{m-1, v_1} , this also holds at e_{m-1, v_1} . Let us now consider a vector e_{1-m, v_1} . One checks:

$$\begin{aligned} x \cdot (y \cdot e_{1-m, v_1}) - y \cdot (x \cdot e_{1-m, v_1}) &= x \cdot e_{n-1, \alpha(v_1)} - y \cdot (m-1)e_{3-m, v_1} \\ &= (1-m)e_{1-m, v_1} \\ &= h \cdot e_{1-m, v_1} \end{aligned}$$

then $h \cdot (x \cdot e_{1-m, v_1}) - x \cdot (h \cdot e_{1-m, v_1}) = 0 = 2x \cdot e_{1-m, v_1}$, and finally:

$$\begin{aligned} h \cdot (y \cdot e_{1-m, v_1}) - y \cdot (h \cdot e_{1-m, v_1}) &= h \cdot e_{n-1, \alpha(v_1)} + (m-1)y \cdot e_{1-m, v_1} \\ &= (n-1)e_{n-1, \alpha(v_1)} + (m-1)e_{n-1, \alpha(v_1)} \\ &= (p-2)e_{n-1, \alpha(v_1)} \\ &= -2y \cdot e_{1-m, v_1} \end{aligned}$$

By construction, $x^n \cdot \mathbb{S}_{\alpha, \beta} = 0$. ◇

On the face of it our construction of $\mathbb{S}_{\alpha, \beta}$ may seem to depend on the bases we chose for V_1 and V_2 . The following shows that up to isomorphism it is not the case.

Observation. $\mathbb{S}_{\alpha, \beta}$ and $\mathbb{S}_{\alpha', \beta'}$ are isomorphic iff the pairs (α, β) and (α', β') are equivalent, that is iff there exist linear isomorphisms $u_1 : V_1 \simeq V_1'$ and $u_2 : V_2 \simeq V_2'$ with $\alpha' = u_2 \alpha u_1^{-1}$ and $\beta' = u_1 \beta u_2^{-1}$.

Proof of Claim. If the pairs (α, β) and (α', β') are equivalent, an isomorphism of \mathfrak{g}_1 -modules $\mathbb{S}_{\alpha, \beta} \simeq \mathbb{S}_{\alpha', \beta'}$ is easily constructed by setting $f(e_{m+1-2j, v_1}) = e'_{m+1-2j, u_1(v_1)}$ with obvious notations, and similarly on the other row.

For the converse suppose that there is such an isomorphism $f : \mathbb{S}_{\alpha, \beta} \simeq \mathbb{S}_{\alpha', \beta'}$. Let $V_1 = E_{1-m}(\mathbb{S}_{\alpha, \beta})$ and $V_2 = E_{1-n}(\mathbb{S}_{\alpha, \beta})$; these are \mathbb{F}_p -vector spaces. One then retrieves $\alpha(v_1) = \frac{1}{(n-1)!} y^n \cdot v_1$ and $\beta(v_2) = \frac{1}{(m-1)!} y^m \cdot v_2$ (which do induce $\mathbb{S}_{\alpha, \beta}$). Proceed similarly on $\mathbb{S}_{\alpha', \beta'}$.

Let $u_1(v_1) = f(v_1)$ and $u_2(v_2) = f(v_2)$. Since f is an isomorphism, these are linear isomorphisms between V_1 and V_1' , resp. V_2 and V_2' . Now for all $v_1 \in V_1$, since f is an isomorphism of \mathfrak{g}_1 -modules,

$$\begin{aligned} u_2 \circ \alpha(v_1) &= u_2 \left(\frac{1}{(n-1)!} y^n \cdot v_1 \right) \\ &= \frac{1}{(n-1)!} f(y^n \cdot v_1) \\ &= \frac{1}{(n-1)!} y^n \cdot f(v_1) \\ &= \alpha'(f(v_1)) \\ &= \alpha' \circ u_1(v_1) \end{aligned}$$

A similar verification can be carried on V_2 , proving that u_1 and u_2 define an equivalence of (α, β) and (α', β') . ◇

Observation. $\mathbb{S}_{\alpha, \beta}$ is non-simple iff there are subspaces $W_1 \leq V_1$ and $W_2 \leq V_2$ not both zero nor both full such that α maps W_1 to W_2 and β maps W_2 to W_1 .

Proof of Claim. We give a correspondence between submodules of $\mathbb{S}_{\alpha, \beta}$ and pairs (W_1, W_2) as in the statement. One direction is clear: if such a pair (W_1, W_2) is given, a \mathfrak{g}_1 -submodule is readily defined.

So let $W \leq V$ be a \mathfrak{g}_1 -submodule. Set $W_1 = \{v_1 \in V_1 : e_{1-m, v_1} \in W\}$ and $W_2 = \{v_2 \in V_2 : e_{1-n, v_2} \in W\}$. We claim that α maps W_1 to W_2 , and that β maps W_2 to W_1 . It suffices to prove it for α . Take indeed $w_1 \in W_1$. Then by construction $e_{1-m, w_1} \in W$ whence $y \cdot e_{1-m, w_1} = e_{n-1, \alpha(w_1)} \in W$. Applying y^{n-1} , one finds up to multiplication by $(n-1)!$ which is coprime to p : $e_{1-n, \alpha(w_1)} \in W$, so by definition, $\alpha(w_1) \in W_2$. ◇

Observation. $\mathbb{S}_{\alpha, \beta}$ is simple iff α and β are isomorphisms and $\beta \circ \alpha$ is irreducible (as an automorphism of V_1).

Proof of Claim. Suppose that $\mathbb{S}_{\alpha,\beta}$ is simple. Take $f \in V_1 \setminus \{0\}$. Consider the sequences $f_r = (\beta\alpha)^r(f)$ and $g_r = \alpha(f_r)$, for $r \geq 0$. These sequences span subspaces $W_1 \leq V_1$ and $W_2 \leq V_2$ mapped one to another by α and β . By simplicity $W_1 = V_1$ and $W_2 = V_2$. Hence α is a bijection. A similar argument holds for β . Now let W_1 be invariant under $\beta \circ \alpha$ and set $W_2 = \alpha(W_1)$. Then α maps W_1 to W_2 and β maps W_2 to W_1 so by simplicity $W_1 = V_1$ or 0 .

Suppose that α and β are isomorphisms such that $\beta \circ \alpha$ is irreducible. If $W_1 \leq V_1$ and $W_2 \leq V_2$ are mapped one to another by α and β then $\beta \circ \alpha(W_1) \leq W_1$ so $W_1 = 0$ or V_1 . In the former case $W_2 = 0$ by injectivity of β , in the latter $W_2 = V_2$ by surjectivity of α . \diamond

Suppose in particular that $\mathbb{S}_{\alpha,\beta}$ is simple. Then for any $f \in V_1 \setminus \{0\}$, the sequence $(f_r)_{r \geq 1}$ as above spans V_1 whence a linear relation $f_0 = \sum_{r=1}^d k_r f_r$. It follows that f_d lies in the span W_1 of $(f_r)_{0 \leq r \leq d-1}$, so $V = W_1$ is finite-dimensional. Moreover the characteristic polynomial of $\beta \circ \alpha$ is irreducible over $\mathbb{F}_p[X]$.

Observation. Let $n \geq 2$ be an integer and \mathbb{K}_1 be the field \mathbb{F}_p with $n < p < 2n$. Let $\mathfrak{g}_1 = \mathfrak{sl}_2(\mathbb{K}_1)$ and V be a simple \mathfrak{g}_1 -module. Suppose that $x^n = 0$ in $\text{End}(V)$.

Then V is some $\mathbb{S}_{\alpha,\beta}$.

Proof of Claim. We may suppose n minimal such that $x^n = 0$. (The reader will observe that had we wished to be fully rigorous we should have written $\mathbb{S}_{\alpha,\beta}^n$ throughout.)

By simplicity $\text{Ann}_V(\mathfrak{g}_1) = 0$ and $\mathfrak{g}_1 \cdot V = V$, so V is a vector space over \mathbb{F}_p ; in particular it is $n!$ -torsion-free and $n!$ -divisible. Now Step 2 of Variation n°17 requires only $n!$ -torsion-freeness, so we get $hx^{n-1} = (n-1)x^{n-1}$ in $\text{End}(V)$ (this is only the first step of the induction fully carried in Step 2 of Variation n°17). As $x^{n-1} \neq 0$, we deduce $E_{n-1} \neq 0$. Since $\bigoplus_{\ell \in \mathbb{Z}/p\mathbb{Z}} E_\ell$ is clearly \mathfrak{g}_1 -invariant, one finds by simplicity $V = \bigoplus_{\ell \in \mathbb{Z}/p\mathbb{Z}} E_\ell$.

We now make the following observation: if for some $\ell \in \mathbb{Z}/p\mathbb{Z}$, one has $E_\ell \cap \ker x \neq 0$, then $E_\ell \leq \ker x$, and likewise with $\ker y$ instead of $\ker x$. We prove it only for x as length plays no role here. Consider $W = \bigoplus_{i=0}^{p-1} y^i \cdot (E_\ell \cap \ker x)$. We claim that W is x -invariant. This is because for $a \in E_\ell \cap \ker x$ and $i \in \{0, \dots, p-1\}$ one has $x \cdot (y^0 \cdot a) = x \cdot a = 0 \in W$ when $i = 0$ and otherwise

$$xy^i \cdot a = y^i x \cdot a + i(h+i-1)y^{i-1} \cdot a \in W$$

We claim that W is y -invariant as well. This is because for $a \in E_\ell \cap \ker x$,

$$xy \cdot (y^{p-1} \cdot a) = xy^p \cdot a = y^p x \cdot a = 0$$

whence $y \cdot (y^{p-1} \cdot a) \in E_\ell \cap \ker x \leq W$. By assumption W is non-trivial, by simplicity of V one has $W = V$ and therefore $E_\ell \leq \ker x$.

Since $0 \neq \text{im } x^{n-1} \leq \ker x \cap E_{n-1}$, one has $E_{n-1} \leq \ker x$. Now the Casimir operator $c_1 = 4yx + h^2 + 2h$ equals $n^2 - 1$ on E_{n-1} ; by simplicity of V , one has $c_1 = n^2 - 1$ everywhere. In particular on E_{m-1} one finds $4yx + m^2 - 1 = n^2 - 1 = m^2 - 1$ so yx annihilates E_{m-1} . If $x \cdot E_{m-1} \neq 0$ then $0 \neq x \cdot E_{m-1} \leq \ker y \cap E_{m+1} = \ker y \cap E_{1-n}$, so by the above observation y annihilates E_{1-n} and one readily sees $V \simeq \text{Sym}^{n-1} \text{Nat } \mathfrak{g}_1$ (a very special case of our construction). If $x \cdot E_{m-1} = 0$ then one retrieves an $\mathbb{S}_{\alpha,\beta}$. \diamond

We thus have described all simple \mathfrak{g}_1 -modules of length n in characteristic $\geq n+1$: they correspond to irreducible polynomials in $\mathbb{F}_p[X]$. There remains one pending question: can one analyze *all* \mathfrak{g}_1 -modules of length n in characteristic $\geq n+1$, in terms of $\mathbb{S}_{\alpha,\beta}$'s? It could be conjectured so but the author wishes to dwell no longer on a subject of disputable interest.

3.3 The Symmetric Case

There is something odd in assuming the characteristic of \mathbb{K}_1 to be $\geq 2n+1$ in length n ; we bring no evidence to support the feeling that a better lower bound should be $n+1$ as it was in Variation n°12 [1].

We know from the previous subsection that lowering the characteristic can result in creating pathologies. Observe how in $\mathbb{S}_{\alpha,\beta}$ the actions of x and y are dissymmetrical as soon as α or β is non-zero. In particular $\mathbb{S}_{\alpha,\beta}$ cannot be made into an $\text{SL}_2(\mathbb{K})$ -module in a ‘‘consistent’’ way since x and y should then have the same order in $\text{End}(V)$ being conjugate under the adjoint action of the Weyl group of $\text{SL}_2(\mathbb{K})$. In short all our previous counterexamples shared the feature that the action of y was quite different from that of x , which is ill-behaved. The minimal decency requirement on an $\mathfrak{sl}_2(\mathbb{K})$ -module V in order to stem from an associated $\text{SL}_2(\mathbb{K})$ -module is that x and y should have the same order in $\text{End}(V)$.

Under this extra symmetry assumption it is possible to classify $\mathfrak{sl}_2(\mathbb{F}_p)$ -modules of two-sided finite length even in low characteristic, as it was in Variation n°14 [1] for a two-sided quadratic module in characteristic 3.

Variation n°18. Let $n \geq 2$ be an integer and \mathbb{K}_1 be the field \mathbb{F}_p with $n < p < 2n$. Let $\mathfrak{g}_1 = \mathfrak{sl}_2(\mathbb{K}_1)$ and V be a \mathfrak{g}_1 -module. Suppose that $x^n = y^n = 0$ in $\text{End}(V)$.

Then $V = \text{Ann}_V(\mathfrak{g}_1) \oplus \mathfrak{g}_1 \cdot V$, and $\mathfrak{g}_1 \cdot V$ is a \mathbb{K}_1 -vector space with $\mathfrak{g}_1 \cdot V \simeq \bigoplus_{k=1}^{n-1} \oplus_{I_k} \text{Sym}^k \text{Nat } \mathfrak{g}_1$ as $\mathbb{K}_1 \mathfrak{g}_1$ -modules.

Proof. Induction on n . When $n = 2$ this is a special case of Variation n°14 [1] with $\mathbb{K} = \mathbb{F}_3$. We shall adapt the proof of Variation n°17. Write $p = n + m$ with $0 < m < n$.

One might desire to assume $\text{Ann}_V(\mathfrak{g}_1) = 0$ and $\mathfrak{g}_1 \cdot V = V$. Actually if $p > n + 1$ the proof given in Variation n°17 remains correct as $k(k+2) \neq 0$ for $k \in \{1, \dots, n-1\}$. But when $p = n + 1$ the Casimir operator c_1 now annihilates $\text{Sym}^{p-2} \text{Nat } \mathfrak{g}_1$ and there may be some subtleties.

Step 1. We may assume that V is a \mathbb{K}_1 -vector space.

Proof of Claim. Suppose the result is known for \mathbb{K}_1 -vector spaces and bear in mind that assumptions on the length of x and y go down to subquotients.

As \mathfrak{g}_1 annihilates pV the factor $\bar{V} = V / \text{Ann}_V(\mathfrak{g}_1)$ is a vector space. By assumption $\bar{V} = \mathfrak{g}_1 \cdot \bar{V} \oplus \text{Ann}_{\bar{V}}(\mathfrak{g}_1)$. Then using perfectness one has $\text{Ann}_{\bar{V}}(\mathfrak{g}_1) = 0$ so $\bar{V} = \mathfrak{g}_1 \cdot \bar{V}$.

As a consequence $V = \mathfrak{g}_1 \cdot V + \text{Ann}_V(\mathfrak{g}_1)$. But p annihilates the submodule $W = \mathfrak{g}_1 \cdot V$ which is therefore another vector space. Still by assumption $W = \mathfrak{g}_1 \cdot W \oplus \text{Ann}_W(\mathfrak{g}_1)$. Then perfectness again yields $\mathfrak{g}_1 \cdot W = W$ so $\text{Ann}_W(\mathfrak{g}_1) = 0$.

In particular $W \cap \text{Ann}_V(\mathfrak{g}_1) = 0$ and $V = \mathfrak{g}_1 \cdot V \oplus \text{Ann}_V(\mathfrak{g}_1)$ has the desired structure since $W = \mathfrak{g}_1 \cdot V$ does. \diamond

Step 2. In $\text{End}(V)$, one has $(h - n + 1)(h - n + 2) \dots (h + n - 1) = 0$.

Proof of Claim. Since $p > n$ the proof given in Variation n°17 remains correct. \diamond

We move to the weight space decomposition. Unfortunately the various E_j 's with $j \in \{1 - n, \dots, n - 1\}$ are no longer pairwise distinct so special attention must be paid. Observe how since V is a $\mathbb{K}_1 = \mathbb{F}_p$ -vector space one should actually talk about the $E_{[j]}$'s (where $[j]$ is the congruence class of j modulo p) in order to prevent confusion. This is what we do from now on.

Step 3. $V = \bigoplus_{j \in \{0, \dots, p-1\}} E_{[j]}$.

Proof of Claim. Bear in mind that $(h - n + 1)(h - n + 2) \dots (h + n - 1) = 0$. If $p = 2n - 1$ the argument of Variation n°17 remains correct since the polynomials $X - j$ with $j \in \{1 - n, \dots, n - 1\}$ are still pairwise coprime and coincide with the polynomials $X - j$ with $j \in \{0, \dots, p - 1\}$. But for $p \leq 2n - 3$ which we now assume it is no longer the case as some appear twice. Let us determine which with care.

As $p \leq 2n - 3$ we have $n - 1 \geq m + 1$. We lift every congruence class modulo p to its canonical representative in $\{0, \dots, p - 1\}$.

$$\begin{array}{cccccccc} \text{class} & [1 - n] & [2 - n] & \dots & [-1] & [0] & [1] & \dots & [n - 1] \\ \text{repr.} & m + 1 & m + 2 & \dots & p - 1 & 0 & 1 & \dots & n - 1 \end{array}$$

Let us partition $I = \{0, \dots, p - 1\}$ into the set $I_1 = \{0, \dots, m\} \cup \{n, \dots, p - 1\}$ of the $2m + 1$ elements occurring once and the set $I_2 = \{m + 1, \dots, n - 1\}$ of the $n - 1 - m$ elements occurring twice:

$$0, 1, \dots, m, \underbrace{m + 1, \dots, n - 1}_{I_2}, n, \dots, p - 1$$

Therefore the polynomial

$$P(X) = \prod_{\ell \in I_1} (X - [\ell]) \cdot \prod_{\ell \in I_2} (X - [\ell])^2$$

annihilates h in $\text{End}(V)$. For $\ell \in I$ let $F_{[\ell]} = \ker(h - [\ell])^2 \geq E_{[\ell]} = \ker(h - [\ell])$. It is readily observed that x , resp. y , maps $F_{[\ell]}$ to $F_{[\ell+2]}$, resp. $F_{[\ell-2]}$. Since all monomials powers in $P(X)$ are pairwise coprime in $\mathbb{F}_p[X]$ one has:

$$V = \bigoplus_{\ell \in I_1} E_{[\ell]} \bigoplus \bigoplus_{\ell \in I_2} F_{[\ell]}$$

Observe that for all $\ell \in I_1$, one has $F_{[\ell]} = E_{[\ell]}$. Our task is to prove it for $\ell \in I_2$ as well. So let $k \in I$ be minimal with $F_{[k]} > E_{[k]}$; $k \in I_2$ so $k \geq m + 1$. We wish to take the least i with $k + 2i \in I_1$. Unfortunately this may fail to exist, for instance when $n = p - 1$ and $k = p - 2$. But there certainly is i minimal with $[k + 2i] \in [I_1]$. Then $i \leq \frac{n-k}{2} + 1$.

Let $W = \bigoplus_{\ell \in I} E_{[\ell]}$ which is clearly \mathfrak{g}_1 -invariant. We shall compute modulo W which we denote by \equiv . Let $v \in F_{[k]}$. Recall that $y \cdot v \in y \cdot F_{[k]} \leq F_{[k-2]} = E_{[k-2]} \leq W$, so $y \cdot v \equiv 0$. Moreover by construction $x^i \cdot v \in F_{[k+2i]} = E_{[k+2i]} \leq W$. Finally by definition $(h - [k]) \cdot v \in \ker(h - [k]) = E_{[k]} \leq W$ so $h \cdot v \equiv kv$. Hence

$$\begin{aligned} 0 &\equiv y^i x^i \cdot v \\ &\equiv \sum_{k=0}^i (-1)^k k! \binom{i}{k} \binom{i}{k}^{k-1} \prod_{\ell=0}^{k-1} (h + [\ell]) x^{i-k} y^{i-k} \cdot v \\ &\equiv (-1)^i i! \prod_{\ell=0}^{i-1} (h + [\ell]) \cdot v \\ &\equiv (-1)^i i! \prod_{\ell=0}^{i-1} [k + \ell] v \end{aligned}$$

Now $k \neq 0$ since $0 \in I_1$ and $k + i - 1 \leq k + \frac{n-k}{2} \leq n < p$. Thus remains $v \equiv 0$ meaning $F_{[k]} \leq W$ and $F_{[k]} = E_{[k]}$. Therefore $V = W$. \diamond

Notation 4. Let $V_{\perp} = \text{im}(c_1 - n^2 + 1)$ and $V_{\top} = \ker(c_1 - n^2 + 1)$. Let also $V_{\perp\perp} = \text{im}(c_1 - n^2 + 1)^2$ and $V_{\top\top} = \ker(c_1 - n^2 + 1)^2$.

Step 5. $V_{\perp\perp}$ is a \mathfrak{g}_1 -submodule isomorphic to $\bigoplus_{k=1}^{n-2} \bigoplus_{I_k} \text{Sym}^k \text{Nat } \mathfrak{g}_1$ if $p = n + 1$ and to $\text{Ann}_{V_{\perp}}(\mathfrak{g}_1) \oplus \bigoplus_{\substack{k=1 \\ k \neq m-1}}^{n-2} \bigoplus_{I_k} \text{Sym}^k \text{Nat } \mathfrak{g}_1$ otherwise.

Proof of Claim. As in Variation n°17, V_{\perp} is a \mathfrak{g}_1 -submodule annihilated by x^{n-1} , and by y^{n-1} similarly. One certainly has $n - 1 < p$. If $p \geq 2(n - 1) + 1$ then we apply Variation n°17. Otherwise $p < 2(n - 1)$ and we apply induction. In any case $V_{\perp} = \text{Ann}_{V_{\perp}}(\mathfrak{g}_1) \oplus \mathfrak{g}_1 \cdot V_{\perp}$ and $\mathfrak{g}_1 \cdot V_{\perp}$ is isomorphic to $\bigoplus_{k=1}^{n-2} \bigoplus_{I_k} \text{Sym}^k \text{Nat } \mathfrak{g}_1$.

The operator $c_1 - n^2 + 1$ is no longer a bijection of V_{\perp} as $k(k + 2) = n^2 - 1$ solves into $k = n - 1$ or $m - 1$ in \mathbb{F}_p . Hence $c_1 - n^2 + 1$ annihilates the component isomorphic to $\bigoplus_{I_{m-1}} \text{Sym}^{m-1} \text{Nat } \mathfrak{g}_1$ but acts bijectively on the other $\bigoplus_{I_k} \text{Sym}^k \text{Nat } \mathfrak{g}_1$'s.

As for the $\text{Ann}_{V_{\perp}}(\mathfrak{g}_1)$ term, there are two possibilities. Either $p = n + 1$ in which case $n^2 - 1 = 0$ and $(c_1 - n^2 + 1)$ annihilates $\text{Ann}_{V_{\perp}}(\mathfrak{g}_1)$, or $p > n + 1$ in which case $n^2 - 1 \neq 0$ and $(c_1 - n^2 + 1)$ is a bijection of $\text{Ann}_{V_{\perp}}(\mathfrak{g}_1)$.

Hence if $p = n + 1$ one has $V_{\perp\perp} = \bigoplus_{k=1}^{n-2} \bigoplus_{I_k} \text{Sym}^k \text{Nat } \mathfrak{g}_1$ whereas if $p > n + 1$ one has $V_{\perp\perp} = \text{Ann}_{V_{\perp}}(\mathfrak{g}_1) \oplus \bigoplus_{\substack{k=1 \\ k \neq m-1}}^{n-2} \bigoplus_{I_k} \text{Sym}^k \text{Nat } \mathfrak{g}_1$. \diamond

We shall simplify notations letting $\text{Sym}^0 \text{Nat } \mathfrak{g}_1$ denote the trivial \mathbb{F}_p -line so that $\text{Ann}_{V_{\perp}}(\mathfrak{g}_1)$ handily rewrites into $\bigoplus_{I_0} \text{Sym}^0 \text{Nat } \mathfrak{g}_1$. We preferred to avoid such notation in general due to possible confusions: for instance when $V = \mathbb{Z}/2\mathbb{Z}$ as a trivial $\mathfrak{g}_1 = \mathfrak{sl}_2(\mathbb{F}_3)$ -module one has $V = \text{Ann}_V(\mathfrak{g}_1)$ but V certainly is no sum of copies of $\text{Sym}^0 \text{Nat } \mathfrak{g}_1 = \mathbb{F}_3$. Here we know from Step 1 that V is a $\mathbb{K}_1 = \mathbb{F}_p$ -vector space and confusion is no longer possible.

As a consequence $V_{\perp\perp}$ is isomorphic to $\bigoplus_{\substack{k=0 \\ k \neq m-1}}^{n-2} \bigoplus_{I_k} \text{Sym}^k \text{Nat } \mathfrak{g}_1$ in either case.

Step 6. We may assume $V = V_{\top\top}$.

Proof of Claim. We claim that $V = V_{\perp\perp} \oplus V_{\top\top}$. Here again $c_1 - n^2 + 1$ is a bijection of $V_{\perp\perp}$ and its square as well whence $V_{\perp\perp} \cap V_{\top\top} = 0$ and $V = V_{\perp\perp} \oplus V_{\top\top}$. \diamond

From now on we suppose $V = V_{\top\top}$; in particular $(c_1 - n^2 + 1)^2$ annihilates V . The assumption that $y^n = 0$ in $\text{End}(V)$ had played no real role up to this point.

Step 7. $\ker x = E_{[n-1]} \oplus E_{[m-1]}$ and $\ker y = E_{[1-n]} \oplus E_{[1-m]}$.

Proof of Claim. We claim that x is injective on $\bigoplus_{\substack{j \in \{0, \dots, p-1\} \\ j \neq m-1, n-1}} E_{[j]}$. Let $j \neq m-1, n-1$ and $a_j \in E_{[j]} \cap \ker x$. Then

$$0 = (c_1 - n^2 + 1)^2 \cdot a_j = (j(j+2) - n^2 + 1)^2 a_j$$

implies $((j+1)^2 - n^2)a_j = 0$ so by assumption on j one has $a_j = 0$.

It remains to prove that x does annihilate all of $E_{[n-1]} \oplus E_{[m-1]}$. First let $a_{m-1} \in E_{[m-1]}$. Do not forget that $hx^{n-1} = (n-1)x^{n-1}$ in $\text{End}(V)$. So $x^{n-1} \cdot a_{m-1} \in E_{[n-3]} \cap E_{[m-1]} = 0$. But x is injective on each of $E_{[1-n]}, \dots, E_{[n-5]}$ which implies $x \cdot a_{m-1} = 0$. Hence x annihilates $E_{[m-1]}$ and by symmetry $y \cdot E_{[1-m]} = 0$ as well.

Now let $a_{1-n} \in E_{[1-n]}$. Then $y^{n-1}x^{n-2} \cdot a_{1-n} \in E_{[1-n]} \cap E_{[-n-1]} = 0$. But also bearing in mind that x annihilates $E_{[m-1]} = E_{[-n-1]}$:

$$\begin{aligned} 0 &= y^{n-1}x^{n-2} \cdot a_{1-n} \\ &= y \cdot \left(\sum_{k=0}^{n-2} (-1)^k k! \binom{n-2}{k} \binom{n-2}{k} \left(\prod_{\ell=0}^{k-1} (h+\ell) \right) x^{n-2-k} y^{n-2-k} \cdot a_{1-n} \right) \\ &= (-1)^{n-2} (n-2)! y \cdot \left(\prod_{\ell=0}^{n-3} (h+\ell) \cdot a_{1-n} \right) \\ &= (-1)^{n-2} (n-2)! \prod_{\ell=0}^{n-3} (1-n+\ell) y \cdot a_{1-n} \\ &= ky \cdot a_{n-1} \end{aligned}$$

where k is non-zero modulo p . Hence $y \cdot a_{1-n} = 0$. By symmetry the analogue holds of x . \diamond

We may conclude as in Variation n°17:

$$V_{\top\top} \simeq \bigoplus_{I_{m-1}} \text{Sym}^{m-1} \text{Nat } \mathfrak{g}_1 \bigoplus \bigoplus_{I_{n-1}} \text{Sym}^{n-1} \text{Nat } \mathfrak{g}_1$$

This finishes the proof. \square

4 Scalar Flesh

When the irreducible $\mathfrak{sl}_2(\mathbb{K}_1)$ -submodules of an $\mathfrak{sl}_2(\mathbb{K})$ -module V are all isomorphic, V bears a compatible \mathbb{K} -vector space structure: §4.1 contains Variation n°19 which is our main result. Otherwise, and always in order to retrieve a linear geometry, one has to make some assumptions on the behaviour of $\ker x_\lambda$ and of $\text{im } x_\lambda$. Under either assumption things work more or less in quotients of a certain composition series (§4.2); should one wish to have a direct sum, one needs *both* assumptions (Variation n°22, §4.3).

4.1 The Separated Case

Variation n°19. *Let $n \geq 2$ be an integer and \mathbb{K} be a field of characteristic 0 or $\geq n$. Let $\mathfrak{g} = \mathfrak{sl}_2(\mathbb{K})$ and V be a \mathfrak{g} -module. Let \mathbb{K}_1 be the prime subfield of \mathbb{K} and $\mathfrak{g}_1 = \mathfrak{sl}_2(\mathbb{K}_1)$. Suppose that V is a \mathbb{K}_1 -vector space such that $V \simeq \bigoplus_I \text{Sym}^{n-1} \text{Nat } \mathfrak{g}_1$ as $\mathbb{K}_1 \mathfrak{g}_1$ -modules.*

Then V bears a compatible \mathbb{K} -vector space structure for which $V \simeq \bigoplus_J \text{Sym}^{n-1} \text{Nat } \mathfrak{g}$ as $\mathbb{K} \mathfrak{g}$ -modules.

Proof.

Notation 1. For $i = 1 \dots n$ let:

$$d_i = \frac{(i-1)! (n-1)!}{(n-i)!} = ((i-1)!)^2 \binom{n-1}{i-1}$$

This is an integer with prime factors $< n$. Moreover $d_{i+1} = i(n-i)d_i$.

Step 2. $V = \bigoplus_{i=1}^n E_{n+1-2i}$. For all $i = 1 \dots n$, one has $(yx)|_{E_{n+1-2i}} = (i-1)(n+1-i)$, also $(xy)|_{E_{n+1-2i}} = i(n-i)$, and $(x^{i-1}y^{i-1})|_{E_{n-1}} = d_i$.

Proof of Claim. All by assumption on V as a \mathfrak{g}_1 -module. \diamond

Notation 3. [see the linear structure in the Theme [1]] Let $1 \leq i \leq n$. Set for $\lambda \in \mathbb{K}$ and $a_{n+1-2i} \in E_{n+1-2i}$:

$$\lambda \cdot a_{n+1-2i} = \frac{1}{n-1} \frac{1}{d_i} y^{i-1} h_\lambda x^{i-1} \cdot a_{n+1-2i}$$

Observe that multiplication by λ normalizes each E_{n+1-2i} . Extend the definition to $V = \bigoplus_{i=1}^n E_{n+1-2i}$.

Remark. One has for all $a \in V$:

$$\lambda \cdot a = \frac{1}{n-1} \frac{1}{((n-1)!)^2} \sum_{i=1}^n \frac{1}{d_i} y^{i-1} h_\lambda x^{n-1} y^{n-1} x^{i-1} \cdot a$$

We shall not use this.

Step 4. V is a \mathbb{K} -vector space.

Proof of Claim. Let us prove that we have defined an action of \mathbb{K} . The construction is well-defined. Additivity in λ and a is obvious. So it suffices to prove multiplicativity. Let $(\lambda, \mu) \in \mathbb{K}^2$ and $a \in E_{n-1}$. By definition $\lambda \cdot a_{n-1} = \frac{1}{n-1} h_\lambda \cdot a_{n-1}$. So by Step 2 applied to $y_\mu \cdot a$ with $i = 2$ one has $xy_\mu \cdot a = (n-1)y_\mu \cdot a$, whence:

$$\begin{aligned} (n-1)^2 \lambda \cdot (\mu \cdot a) &= h_\lambda h_\mu \cdot a \\ &= (x_\lambda y - y x_\lambda)(x y_\mu - y_\mu x) \cdot a \\ &= (x_\lambda y - y x_\lambda) x y_\mu \cdot a \\ &= x_\lambda y x y_\mu \cdot a \\ &= (n-1) x_\lambda y_\mu \cdot a \\ &= (n-1) h_{\lambda\mu} \cdot a \\ &= (n-1)^2 (\lambda\mu) \cdot a \end{aligned}$$

and we obtain multiplicativity on E_{n-1} .

Let now i be any integer in $\{1, \dots, n\}$ and $a \in E_{n+1-2i}$. Let $b = x^{i-1} \cdot a \in E_{n-1}$. Then by definition for any $\lambda \in \mathbb{K}$:

$$\lambda \cdot a = \frac{1}{(n-1)} \frac{1}{d_i} y^{i-1} h_\lambda \cdot b$$

so with Step 2 applied to $h_\mu \cdot b = h_\mu x^{i-1} \cdot a$:

$$\begin{aligned} (n-1)^2 d_i^2 \lambda \cdot (\mu \cdot a) &= y^{i-1} h_\lambda x^{i-1} y^{i-1} h_\mu x^{i-1} \cdot a \\ &= d_i y^{i-1} h_\lambda h_\mu x^{i-1} \cdot a \\ &= (n-1) d_i y^{i-1} h_{\lambda\mu} x^{i-1} \cdot a \\ &= (n-1)^2 d_i^2 (\lambda\mu) \cdot a \end{aligned}$$

and we obtain multiplicativity on E_{n+1-2i} . ◇

Step 5. \mathfrak{g} is linear on V .

Proof of Claim. Let $\lambda \in \mathbb{K}$. Let us first prove linearity of x . It is obvious on E_{n-1} . So let $i \geq 2$ and $a \in E_{n+1-2i}$; one has thanks to Step 2 applied to $y^{i-2} h_\lambda x^{i-1} \cdot a \in E_{n+1-2(i-1)}$ and $i-1$:

$$\begin{aligned} (n-1) d_i x \cdot (\lambda \cdot a) &= x \cdot (y^{i-1} h_\lambda x^{i-1} \cdot a) \\ &= xy \cdot (y^{i-2} h_\lambda x^{i-2} \cdot (x \cdot a)) \\ &= (i-1)(n+1-i)(n-1) d_{i-1} \lambda \cdot (x \cdot a) \\ &= (n-1) d_i \lambda \cdot (x \cdot a) \end{aligned}$$

and we obtain linearity of x .

Linearity of y is very similar. It is obvious on E_{n-1} . Now for $a \in E_{n+1-2i}$ with $1 \leq i < n$ one has by Step 2 $xy \cdot a = i(n-i)a$, whence:

$$\begin{aligned} (n-1) d_{i+1} y \cdot (\lambda \cdot a) &= (n-1) i (n-i) d_i y \cdot (\lambda \cdot y) \\ &= y (y^{i-1} h_\lambda x^{i-1}) \cdot (xy \cdot a) \\ &= y^i h_\lambda x^i \cdot (y \cdot a) \\ &= (n-1) d_{i+1} \lambda \cdot (y \cdot a) \end{aligned}$$

which proves linearity of y .

It remains to prove linearity of h_μ . For $a \in E_{n-1}$ one has:

$$(n-1)h_\mu \cdot (\lambda \cdot a) = h_\mu h_\lambda \cdot a = h_\lambda \cdot (h_\mu \cdot a) = (n-1)\lambda \cdot (h_\mu \cdot a)$$

which proves linearity of h_μ on E_{n-1} . Now let $i \geq 2$, and take $a \in E_{n+1-2i}$, and $b = x^{i-1} \cdot a \in E_{n-1}$. With Step 2 applied to b one finds $y \cdot b = (n-1)x^{i-2} \cdot a$. Now remember that $x^{i-1}h_\mu = h_\mu x^{i-1} - 2(i-1)x_\mu x^{i-2}$. Then using linearity of x :

$$\begin{aligned} (n-1)x^{i-1}h_\mu \cdot a &= (n-1)(h_\mu x^{i-1} - 2(i-1)x_\mu x^{i-2}) \cdot a \\ &= (n-1)h_\mu \cdot b - 2(i-1)x_\mu y \cdot b \\ &= (n-1 - 2(i-1))h_\mu \cdot b \\ &= (n-1)(n+1-2i)\mu \cdot b \\ &= (n-1)x^{i-1} \cdot ((n+1-2i)\mu \cdot a) \end{aligned}$$

Since x^{i-1} is injective on E_{n+1-2i} by Step 2 one derives $h_\mu \cdot a = (n+1-2i)\mu \cdot a$, and this holds of any $a \in E_{n+1-2i}$. In particular by multiplicativity:

$$\lambda \cdot (h_\mu \cdot a) = \lambda \cdot ((n+1-2i)\mu \cdot a) = (n+1-2i)\mu \cdot (\lambda \cdot a) = h_\mu \cdot (\lambda \cdot a)$$

so h_μ is linear. \diamond

V is therefore a $\mathbb{K}\mathfrak{g}$ -module and its structure as such is clear. This finishes the proof. \square

Remark (see Variation n°10 [1]). It is now obvious that for any $\lambda \neq 0$: $\ker x = \ker x_\lambda$ and $\operatorname{im} x = \operatorname{im} x_\lambda$; also $\ker y = \ker y_\lambda$ and $\operatorname{im} y = \operatorname{im} y_\lambda$.

Remark. Although our proof only requires the characteristic to be $\geq n$ it is not possible to apply the method to the modules $\mathbb{S}_{\alpha,\beta}$ obtained in §3.2. All one can get is the following which generalizes Variation n°13 [1].

Let $n \geq 2$ be an integer and \mathbb{K} be a field of characteristic 0 or $\geq n$. Let $\mathfrak{g} = \mathfrak{sl}_2(\mathbb{K})$ and V be a \mathfrak{g} -module. Let \mathbb{K}_1 be the prime subfield of \mathbb{K} and $\mathfrak{g}_1 = \mathfrak{sl}_2(\mathbb{K}_1)$. Suppose that V is a \mathbb{K}_1 -vector space such that $V \simeq \mathbb{S}_{\alpha,\beta}$ as $\mathbb{K}_1\mathfrak{g}_1$ -modules.

Then V bears a \mathbb{K} -vector space structure such that the maps h_λ and x_λ are everywhere linear, but y_λ only on E_ℓ for $\ell \notin \{1-n, 1-m\}$.

Preservation of the linear structure under α and β depends on properties which cannot be prescribed over \mathbb{K}_1 .

4.2 Composition series

We now prove two dual partial results.

Variation n°20. *Let $n \geq 2$ be an integer and \mathbb{K} be a field. Let $\mathfrak{g} = \mathfrak{sl}_2(\mathbb{K})$ and V be a \mathfrak{g} -module. If the characteristic of \mathbb{K} is 0 one requires V to be torsion-free. Suppose either that $x^n = 0$ in $\operatorname{End}(V)$ and the characteristic of \mathbb{K} is 0 or $\geq 2n+1$, or that $x^n = y^n = 0$ in $\operatorname{End}(V)$ and the characteristic of \mathbb{K} is $\geq n+1$.*

Suppose in addition that for all $\lambda \in \mathbb{K}$, one has $\ker x \leq \ker x_\lambda$.

Then there exists a series $\operatorname{Ann}_V(\mathfrak{g}) = V_0 \leq V_1 \leq \dots \leq V_{n-1} = V$ of \mathfrak{g} -submodules such that for all $k = 1, \dots, n-1$, the quotient V_k/V_{k-1} bears a compatible \mathbb{K} -vector space structure for which $V_k/V_{k-1} \simeq \oplus_{I_k} \operatorname{Sym}^k \operatorname{Nat} \mathfrak{g}$ as $\mathbb{K}\mathfrak{g}$ -modules.

Proof. Induction on n . When $n = 2$ this is Variation n°12 [1] and one even has $V = \operatorname{Ann}_V(\mathfrak{g}) \oplus \oplus_{I_1} \operatorname{Nat} \mathfrak{g}$. Let \mathbb{K}_1 denote the prime subfield and $\mathfrak{g}_1 = \mathfrak{sl}_2(\mathbb{K}_1)$. By Variation n°17 or n°18 depending on the assumptions, $V = \operatorname{Ann}_V(\mathfrak{g}_1) \oplus \mathfrak{g}_1 \cdot V$ where $\mathfrak{g}_1 \cdot V \simeq \oplus_{k=1}^{n-1} \oplus_{I_k} \operatorname{Sym}^k \operatorname{Nat} \mathfrak{g}_1$ as $\mathbb{K}_1\mathfrak{g}_1$ -modules.

Let $V_\perp = \operatorname{Ann}_V(\mathfrak{g}_1) \oplus \oplus_{k=1}^{n-2} \oplus_{I_k} \operatorname{Sym}^k \operatorname{Nat} \mathfrak{g}_1$ and $V_\top = \oplus_{I_{n-1}} \operatorname{Sym}^{n-1} \operatorname{Nat} \mathfrak{g}_1$. These are \mathfrak{g}_1 -submodules satisfying $V = V_\perp \oplus V_\top$. One should be careful with the Casimir operator c_1 . Since this operator does not commute with \mathfrak{g} in $\operatorname{End}(V)$, V_\perp and V_\top as defined in Variation n°17 have no reason a priori to be \mathfrak{g} -invariant. Moreover the definition of V_\perp in terms of c_1 fails in characteristic $\leq 2n$ as seen in Variation n°18.

Yet in the present case one sees by inspection in the \mathfrak{g}_1 -module V :

$$V_\perp = (\oplus_{i=1}^n E_{n-2i}) \bigoplus (\oplus_{i=1}^n (E_{n+1-2i} \cap \ker x^{i-1}))$$

Let us now prove that V_\perp is a \mathfrak{g} -submodule. It suffices to show that it is $\mathfrak{t} = \{h_\lambda : \lambda \in \mathbb{K}\}$ -invariant. All E_j 's are h_λ -invariant. But by assumption on the kernels in V , so is $\ker x$: for if $a \in \ker x$ then $x_\lambda \cdot a = 0$ and $xh_\lambda \cdot a = (h_\lambda x - 2x_\lambda) \cdot a = 0$. So the subgroup V_\perp is \mathfrak{g} -invariant; it is a \mathfrak{g} -submodule.

Remark. There is no reason why V_\top should be \mathfrak{g} -invariant as well.

One sees that x^{n-1} acts trivially on $V_\perp = 0$. Moreover V_\perp still enjoys the property $\ker x \leq \ker x_\lambda$; induction provides the desired structure on V_\perp . But $V/V_\perp \simeq V_\top$ as \mathfrak{g}_1 -modules so in the quotient V/V_\perp , one has $\ker x \cap \ker y^{n-1} = 0$. One then applies Variation n°19 to the \mathfrak{g} -module V/V_\perp in order to conclude. \square

Variation n°21. *Let $n \geq 2$ be an integer and \mathbb{K} be a field. Let $\mathfrak{g} = \mathfrak{sl}_2(\mathbb{K})$ and V be a \mathfrak{g} -module. If the characteristic of \mathbb{K} is 0 one requires V to be torsion-free. Suppose either that $x^n = 0$ in $\text{End}(V)$ and the characteristic of \mathbb{K} is 0 or $\geq 2n+1$, or that $x^n = y^n = 0$ in $\text{End}(V)$ and the characteristic of \mathbb{K} is $\geq n+1$.*

Suppose in addition that for all $\lambda \in \mathbb{K}$, one has $\text{im } x_\lambda \leq \text{im } x$.

Then there exists a series $0 = V_0 \leq V_1 \leq \dots \leq V_{n-1} = \mathfrak{g} \cdot V$ of \mathfrak{g} -submodules such that for all $k = 1, \dots, n-1$, the quotient V_k/V_{k-1} bears a compatible \mathbb{K} -vector space structure for which $V_k/V_{k-1} \simeq \bigoplus_{I_{n-k}} \text{Sym}^{n-k} \text{Nat } \mathfrak{g}$ as $\mathbb{K}\mathfrak{g}$ -modules.

Proof. Induction on n . When $n = 2$ this is Variation n°12 [1] and one even has $V = \text{Ann}_V(\mathfrak{g}) \oplus \bigoplus_{I_1} \text{Nat } \mathfrak{g}$. Let \mathbb{K}_1 denote the prime subfield and $\mathfrak{g}_1 = \mathfrak{sl}_2(\mathbb{K}_1)$. By Variation n°17 or n°18 depending on the assumptions, $V = \text{Ann}_V(\mathfrak{g}_1) \oplus \mathfrak{g}_1 \cdot V$ where $\mathfrak{g}_1 \cdot V \simeq \bigoplus_{k=1}^{n-1} \bigoplus_{I_k} \text{Sym}^k \text{Nat } \mathfrak{g}_1$ as $\mathbb{K}_1\mathfrak{g}_1$ -modules.

Let $V_\perp = \text{Ann}_V(\mathfrak{g}_1) \oplus \bigoplus_{k=1}^{n-2} \bigoplus_{I_k} \text{Sym}^k \text{Nat } \mathfrak{g}_1$ and $V_\top = \bigoplus_{I_{n-1}} \text{Sym}^{n-1} \text{Nat } \mathfrak{g}_1$. One sees by inspection in the \mathfrak{g}_1 -module V that:

$$V_\top = \bigoplus_{i=1}^n E_{n+1-2i} \cap \text{im } x^{n-i}$$

Let us then prove that V_\top is a \mathfrak{g} -submodule. It suffices to show that it is $\mathfrak{t} = \{h_\lambda : \lambda \in \mathbb{K}\}$ -invariant. All E_j 's are h_λ -invariant. But by assumption on the images in V , so is $\text{im } x$: for if $a \in \text{im } x$ then writing $a = x \cdot b$ one finds $h_\lambda \cdot a = xh_\lambda \cdot b + 2x_\lambda \cdot b \in \text{im } x$ by assumption. The subgroup V_\top is therefore \mathfrak{g} -invariant: it is a \mathfrak{g} -submodule.

One sees that in the submodule V_\top , $\ker x \cap \ker y^{n-1} = 0$; Variation n°19 provides the desired structure on V_\top . But $V/V_\top \simeq V_\perp$ as \mathfrak{g}_1 -modules so in the quotient V/V_\top , x^{n-1} acts trivially. Moreover V/V_\top still enjoys the property $\text{im } x_\lambda \leq \text{im } x$. One then applies induction to the \mathfrak{g} -module V/V_\top in order to conclude. \square

4.3 Separation

Variation n°22. *Let $n \geq 2$ be an integer and \mathbb{K} be a field. Let $\mathfrak{g} = \mathfrak{sl}_2(\mathbb{K})$ and V be a \mathfrak{g} -module. If the characteristic of \mathbb{K} is 0 one requires V to be torsion-free. Suppose either that $x^n = 0$ in $\text{End}(V)$ and the characteristic of \mathbb{K} is 0 or $\geq 2n+1$, or that $x^n = y^n = 0$ in $\text{End}(V)$ and the characteristic of \mathbb{K} is $\geq n+1$.*

Suppose in addition that for all $\lambda \in \mathbb{K}$, one has $\ker x \leq \ker x_\lambda$ and $\text{im } x_\lambda \leq \text{im } x$.

Then $V = \text{Ann}_V(\mathfrak{g}) \oplus \mathfrak{g} \cdot V$, and $\mathfrak{g} \cdot V$ bears a compatible \mathbb{K} -vector space structure for which $\mathfrak{g} \cdot V \simeq \bigoplus_{k=1}^{n-1} \bigoplus_{I_k} \text{Sym}^{n-1} \text{Nat } \mathfrak{g}$ as $\mathbb{K}\mathfrak{g}$ -modules.

Proof. Induction on n . When $n = 2$ this is Variation n°12 [1]. As in Variations n°20 and n°21, V_\perp and V_\top are \mathfrak{g} -invariant. But the property $\ker x \leq \ker x_\lambda$ clearly goes to submodules, and the property $\text{im } x_\lambda \leq \text{im } x$ clearly goes to quotients. Hence $V_\perp \simeq V/V_\top$ (here as \mathfrak{g} -modules) allows to use induction. \square

5 Lesson: coherence degrees

Notation. *Let V be a \mathfrak{g} -module.*

- $\kappa(V)$ • Let $\kappa(V)$ be the least integer n , if there is one, such that for all $(\lambda_1, \dots, \lambda_n) \in \mathbb{K}^n$, one has $\ker x^n \leq \ker x_{\lambda_1} \dots x_{\lambda_n}$;
- $\iota(V)$ • Let $\iota(V)$ be the least integer n , if there is one, such that for all $(\lambda_1, \dots, \lambda_n) \in \mathbb{K}^n$, one has $\text{im } x_{\lambda_1} \dots x_{\lambda_n} \leq \text{im } x^n$.

The parameters $\kappa(V)$ and $\iota(V)$ may play some role in the rest of the present series of articles. A convenient name would be the *ascending* (resp., *descending*) *coherence degrees* of the action. Be careful that they are *not* the least n such that the kernels (resp. images) of $x_{\lambda_1} \dots x_{\lambda_n}$ do not depend on $(\lambda_1, \dots, \lambda_n)$. They are the least n such that one always has an inclusion.

Observation. *Let V be a \mathfrak{b} -module. Then $\kappa(V) = \min\{n \in \mathbb{N} \cup \{\infty\} : \ker x^n \text{ is } \mathfrak{t}\text{-invariant}\}$, and $\iota(V) = \min\{n \in \mathbb{N} \cup \{\infty\} : \text{im } x^n \text{ is } \mathfrak{t}\text{-invariant}\}$.*

Proof of Claim. We claim that $\ker x^n$ is \mathfrak{t} -invariant iff $\forall (\lambda_1, \dots, \lambda_n) \in \mathbb{K}^n$, one has $\ker x^n \leq \ker x_{\lambda_1} \dots x_{\lambda_n}$:

- if $\ker x^n$ is \mathfrak{t} -invariant then $\ker x^n$ is a \mathfrak{b} -submodule with x -length $\leq n$, so by Variation n°15 it has u -length $\leq n$, meaning $x_{\lambda_1} \dots x_{\lambda_n} = 0$ in $\text{End}(V)$;
- the converse is obvious since if $a \in \ker x^n$ and $\lambda \in \mathbb{K}$ then $x^n h_\lambda \cdot a = h_\lambda x^n \cdot a - 2n x_\lambda x^{n-1} a = 0$.

Similarly $\text{im } x^n$ is \mathfrak{t} -invariant iff $\forall (\lambda_1, \dots, \lambda_n) \in \mathbb{K}^n$, one has $\text{im } x_{\lambda_1} \dots x_{\lambda_n} \leq \text{im } x^n$ (apply Variation n°15 to the quotient module $V/\text{im } x^n$), which proves the second claim. \diamond

We have in Variation n°22 been using an obvious fact.

Observation. *Let V be a \mathfrak{g} -module and $W \leq V$ be a \mathfrak{g} -submodule. Then $\kappa(W) \leq \kappa(V)$ and $\iota(V/W) \leq \iota(V)$.*

Remember that $\lambda(V)$ stands for the length of V as a u -module. One knows from Variation n°16 that $\lambda(V)$ equals the length of V as an x -module, at least provided the characteristic is not too low.

Variation n°23. *Let $n \geq 2$ be an integer and \mathbb{K} be a field of characteristic 0 or $\geq n+1$. Let V be a $\mathfrak{sl}_2(\mathbb{K})$ -module of u -length at most n . Then for all $\lambda_1, \dots, \lambda_{n-1} \in \mathbb{K}$ one has $\ker x^{n-1} \leq \ker(x_{\lambda_1} \dots x_{\lambda_{n-1}})$ and $\text{im}(x_{\lambda_1} \dots x_{\lambda_{n-1}}) \leq \text{im } x^{n-1}$.*

In our notations this writes $\kappa(V) \leq \lambda(V) - 1$ and $\iota(V) \leq \lambda(V) - 1$.

Proof. Let us first deal with the kernels. We shall need the following identity of the enveloping ring (remember that the terms in the hats do not appear):

$$\begin{aligned} x_{\lambda_1} \dots x_{\lambda_i} y_\mu &= y_\mu x_{\lambda_1} \dots x_{\lambda_i} + \sum_j h_{\mu \cdot \lambda_j} x_{\lambda_1} \dots \widehat{x_{\lambda_j}} \dots x_{\lambda_i} \\ &\quad - \sum_{j \neq k} x_{\mu \cdot \lambda_j \cdot \lambda_k} x_{\lambda_1} \dots \widehat{x_{\lambda_j}} \dots \widehat{x_{\lambda_k}} \dots x_{\lambda_i} \end{aligned} \quad (7)$$

We prove by induction on $i = 0 \dots n-1$ that for all $(\lambda_1, \dots, \lambda_i) \in \mathbb{K}^i$, $x^{n-1-i} x_{\lambda_1} \dots x_{\lambda_i}$ annihilates $\ker x^{n-1}$. When $i = 0$ this is obvious. Let us suppose that the property holds of i and prove it of $i+1 \leq n-1$. Let $(\lambda_1, \dots, \lambda_i, \mu)$ be an $(i+1)$ -tuple of \mathbb{K} and set $\pi = x^{n-1-(i+1)} x_{\lambda_1} \dots x_{\lambda_i} x_\mu$. Recall that $2x_\mu = 2xy_\mu x - y_\mu x^2 - x^2 y_\mu$. By assumption on the length all products of the form $x^{n-i} x_{\lambda_1} \dots x_{\lambda_i}$ are zero, whence in $\text{End}(V)$:

$$\begin{aligned} 2\pi &= 2x^{n-1-(i+1)} x_{\lambda_1} \dots x_{\lambda_i} x_\mu \\ &= x^{n-2-i} x_{\lambda_1} \dots x_{\lambda_i} (2xy_\mu x - y_\mu x^2 - x^2 y_\mu) \\ &= 2x^{n-1-i} x_{\lambda_1} \dots x_{\lambda_i} y_\mu x - x^{n-2-i} x_{\lambda_1} \dots x_{\lambda_i} y_\mu x^2 \end{aligned}$$

It remains to move the y_μ 's to the left using equation (7) applied to the various tuples $(x, \dots, x, x_{\lambda_1}, \dots, x_{\lambda_i})$. Let us do it mentally. Terms with a y_μ on the left will end in $x^{n-i} x_{\lambda_1} \dots x_{\lambda_i}$: by assumption they are zero in $\text{End}(V)$. Terms with a h_ν on the left end either in $x^{n-1-i} x_{\lambda_1} \dots x_{\lambda_i}$ or in $x^{n-i} x_{\lambda_1} \dots \widehat{x_{\lambda_j}} \dots x_{\lambda_i}$ for some j : by induction they annihilate $\ker x^{n-1}$. It thus only remains to consider the pure products of x and the various x_ν 's. There are three cases:

- the j^{th} and k^{th} (omitted) terms were among the x_ν 's: the product is of the form $x_{\mu \cdot \nu_1 \cdot \nu_2} x^{n-i} x_{\lambda_1} \dots \widehat{x_{\nu_1}} \dots \widehat{x_{\nu_2}} \dots x_{\lambda_i}$; by induction it annihilates $\ker x^{n-1}$.
- the j^{th} (omitted) term was among the x 's and the k^{th} among the x_ν 's (or vice-versa): the product is of the form $x_{\mu \cdot \nu} x^{n-1-i} x_{\lambda_1} \dots \widehat{x_\nu} \dots x_{\lambda_i}$; it annihilates $\ker x^{n-1}$.

- the j^{th} and k^{th} (omitted) terms were among the x 's: the product is then of the form $x_\mu x^{n-i-2} x_{\lambda_1} \dots x_{\lambda_i} = \pi$.

The latter case is of interest. Paying attention to the signs and coefficients it appears exactly $-4\binom{n-1-i}{2} + 2\binom{n-2-i}{2} = -(n-2-i)(n+1-i)$ times, whence:

$$2\pi = -(n-2-i)(n+1-i)\pi + z$$

where z annihilates $\ker x^{n-1}$, that is $(n-i-1)(n-i)\pi$ annihilates $\ker x^{n-1}$. Now $i \leq n-2$ so if we had started with $\frac{\mu}{(n-i-1)(n-i)}$ we would have found that π annihilates $\ker x^{n-1}$. This completes the induction; with $i = n-1$ one obtains the desired conclusion.

As far as the images are concerned we proceed similarly using the dual formula:

$$\begin{aligned} y_\mu x_{\lambda_1} \dots x_{\lambda_i} &= x_{\lambda_1} \dots x_{\lambda_i} y_\mu - \sum_j x_{\lambda_1} \dots \widehat{x_{\lambda_j}} \dots x_{\lambda_i} h_{\mu \cdot \lambda_j} \\ &\quad - \sum_{j \neq k} x_{\lambda_1} \dots \widehat{x_{\lambda_j}} \dots \widehat{x_{\lambda_k}} \dots x_{\lambda_i} x_{\mu \cdot \lambda_j \cdot \lambda_k} \end{aligned} \quad (8)$$

proving by induction on $i = 0 \dots n-1$ that for all $(\lambda_1, \dots, \lambda_i) \in \mathbb{K}^i$ one has the inclusion $\text{im}(x^{n-1-i} x_{\lambda_1} \dots x_{\lambda_i}) \leq \text{im } x^{n-1}$. When rewriting π use instead:

$$2\pi = (2x y_\mu x - y_\mu x^2 - x^2 y_\mu) x^{n-2-i} x_{\lambda_1} \dots x_{\lambda_i}$$

and move the y_μ 's to the right using formula (8). \square

Let us briefly comment on duality. First notice that if V is a $\mathbb{K}_1[\mathfrak{u}]$ -module, then V has length at most n iff the dual module V^* has, meaning $\lambda(V) = \lambda(V^*)$.

Observation. *Let n be an integer and \mathbb{K} be a field of characteristic 0 or $\geq n$, with prime subfield \mathbb{K}_1 . Let V be a $\mathbb{K}_1[\mathfrak{u}]$ -module of length at most n . Then $\iota(V) = \kappa(V^*)$ and $\kappa(V) = \iota(V^*)$.*

Proof of Claim. This is routine. For a tuple $\mu = (\lambda_1, \dots, \lambda_d) \in \mathbb{K}^d$, let χ_μ (resp., χ_μ^*) stand for the operator $x_{\mu_1} \dots x_{\mu_d}$ in $\text{End}(V)$ (resp., $\text{End}(V^*)$). Also let $\eta = (\mu_d, \dots, \mu_1)$ be the tuple μ in reverse order; in $\text{End}(V)$ one has $\chi_\eta = \chi_\mu$ but for the sake of clarity we shall not use this. For $(v, \delta) \in V \times V^*$, one has $(\chi_\mu^* \cdot \delta)(v) = (-1)^d \delta(\chi_\eta \cdot v)$.

Now for $A \subseteq V$ let $A^\perp = \{\delta \in V^* : \forall a \in A, \delta(a) = 0\}$ and for $\Delta \subseteq V^*$ let $\Delta^\perp = \{v \in V : \forall \delta \in \Delta, \delta(a) = 0\}$. We then observe that $(\ker \chi_\mu)^\perp = \text{im } \chi_\eta^*$ and $(\ker \chi_\mu^*)^\perp = \text{im } \chi_\eta$ as immediate verifications show.

Finally let $\mathbb{1}$ be the tuple $(1, \dots, 1) \in \mathbb{K}^d$. Then $\iota(V) \leq d$ iff $\forall \mu \in \mathbb{K}^d$, $\text{im } \chi_\mu \leq \text{im } \chi_{\mathbb{1}}$ iff $\forall \mu \in \mathbb{K}^d$, $(\text{im } \chi_{\mathbb{1}})^\perp \leq (\text{im } \chi_\mu)^\perp$ iff $\forall \mu \in \mathbb{K}^d$, $\ker \chi_{\mathbb{1}}^* \leq \ker \chi_\mu^*$ iff $\kappa(V^*) \leq d$; the other equality is proved similarly. \diamond

Remark. We can now explain the redundancy in the proof of Variation $n^\circ 23$, in the case of $\mathbb{K}_1[\mathfrak{g}]$ -modules. Check by crude computation, as we did, only one of the two inequalities, say $\kappa(V) \leq \lambda(V)$. Then $\iota(V) = \kappa(V^*) \leq \lambda(V^*) = \lambda(V)$.

The argument requires a \mathbb{K}_1 -vector space structure, since the author does not care for duality arguments over more general rings; it however requires only an action of \mathfrak{u} . But in order to get the first inequality $\kappa(V) \leq \lambda(V)$, one does need an action of \mathfrak{g} .

Remark.

- Equalities may not hold in Variation $n^\circ 23$: remember that in $\text{Nat } \mathfrak{sl}_2(\mathbb{C}) \otimes^\varphi \text{Nat } \mathfrak{sl}_2(\mathbb{C})$ (φ stands for complex conjugation) one has $x^3 = 0$ and $x_i x_1 = 0$ but $x^2 \neq 0$.
- The value $n-1$ is optimal. Take distinct field automorphisms $\varphi_1, \dots, \varphi_n$ and set $V = (\varphi_1 \text{Nat } \mathfrak{sl}_2) \otimes \dots \otimes (\varphi_n \text{Nat } \mathfrak{sl}_2)$. This is an irreducible representation. Its length is $n+1$; in particular $\ker x^n \leq \ker x_{\lambda_1} \dots x_{\lambda_n}$ for all $(\lambda, \dots, \lambda_n) \in \mathbb{K}^n$, but this fails at stage $n-1$. Let indeed $\lambda \in \mathbb{K}$ be such that $\varphi_1(\lambda) \neq \varphi_n(\lambda)$. The standard basis (e_1, e_2) of $\text{Nat } \mathfrak{sl}_2$ being fixed, e_{i_1, \dots, i_n} will denote the pure tensor $e_{i_1} \otimes \dots \otimes e_{i_n}$. Consider $a = e_{2, \dots, 2, 1} - e_{1, 2, \dots, 2}$; one sees that $x^{n-1} \cdot a = 0$ but $x_\lambda x^{n-2} \cdot a = (n-2)! (\varphi_1(\lambda) - \varphi_n(\lambda)) e_{1, \dots, 1} \neq 0$.

One might expect $\kappa(V)$ and $\iota(V)$ to provide an indication of the number of tensor factors; but one would first need to conjecture that every simple \mathfrak{g} -module of finite length is a tensor product of copies, twisted by field automorphisms, of a same representation of \mathfrak{g} as a Lie algebra. The author does not wish to do so even under model-theoretic assumptions. Anyway we have until now been dealing mostly with actions of coherence degree 1, in a sense or the other.

It is not a priori clear that $\kappa(V)$ and $\iota(V)$ need in general be equal and the question deserves to be asked, at least for an action of finite length. Note that one could define the same numbers for the action of y ; perhaps one should not expect a relation with the coherence degrees for x even in the finite length case.

Finally, an alternative indicator could be the nilpotence height of the Casimir operator, that is the least n such that $[\mathfrak{g}, \dots, [\mathfrak{g}, c_1]]$ acts trivially on V . Our results would have been more naive under the assumption that c_1 commutes with the action of \mathfrak{g} since instead of Variations n°20, n°21, and n°22 it would have sufficed to adapt the rather standard techniques of Variation n°17. Besides we found no relation between the nilpotence height of the Casimir operator and the coherence degrees.

One easily imagines how to define λ, κ, ι for an action of $\mathrm{SL}_2(\mathbb{K})$.

Future variations will explore the symmetric powers of $\mathrm{Nat} \mathrm{SL}_2(\mathbb{K})$.

References

- [1] Adrien Deloro. Quadratic actions. *Confluentes Math.*, 5(2):23–41, 2013.
- [2] Jacques Dixmier. *Enveloping algebras*, volume 11 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 1996. Revised reprint of the 1977 translation.
- [3] James E. Humphreys. *Introduction to Lie algebras and representation theory*, volume 9 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1978. Second printing, revised.