



HAL
open science

New steps in walks with small steps in the quarter plane

Irina Kourkova, Kilian Raschel

► **To cite this version:**

Irina Kourkova, Kilian Raschel. New steps in walks with small steps in the quarter plane. 2013.
hal-00840239v1

HAL Id: hal-00840239

<https://hal.science/hal-00840239v1>

Preprint submitted on 2 Jul 2013 (v1), last revised 20 Sep 2014 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

NEW STEPS IN WALKS WITH SMALL STEPS IN THE QUARTER PLANE

I. KURKOVA AND K. RASCHEL

ABSTRACT. In this article we present a new approach for finding the generating function counting (non-singular) walks with small steps in the quarter plane. We obtain its explicit expression in terms of infinite series, while in the literature, its standard expression is given in terms of solutions to boundary value problems. As an illustration of the computational interest of our approach, we work out these new formulæ for Kreweras' model: this leads to an expression for the generating function with Weierstrass Zeta functions, and eventually (using transformation theory of elliptic functions) to the well-known algebraic expression.

1. INTRODUCTION

1.1. **Context.** In the field of enumerative combinatorics, much progress has been recently made in the study of walks in the quarter plane $\mathbf{Z}_+^2 = \{0, 1, \dots\}^2$. The general aim is the following: given a set \mathcal{S} of allowed steps (or jumps), it is a matter of counting the number of walks constructed from these steps, which start from a given point and end at a given point or subdomain of the quarter plane. Without hypotheses on \mathcal{S} , this problem is, still today, out of reach. In this article, following the seminal work of Bousquet-Mélou and Mishna [6], we shall assume that the steps are small: in other words, $\mathcal{S} \subset \{-1, 0, 1\}^2 \setminus \{(0, 0)\}$. See Figures 1, 2, 3 and 6 for examples. There are obviously $2^8 = 256$ models. But one is easily convinced that some models are trivial; some models are equivalent (by diagonal symmetry) to other ones; and finally, some models are equivalent to models of walks confined in a half-plane, for which the general theory already exists [7]. It happens that out of the 256 models, only 79 inherently different ones remain to be studied [6]. Let $q(i, j; n)$ denote the number of paths in \mathbf{Z}_+^2 having length n , starting from $(0, 0)$ and ending at (i, j) . Define their generating function (GF) as

$$(1) \quad Q(x, y; z) = \sum_{i, j, n \geq 0} q(i, j; n) x^i y^j z^n.$$

There are then two usual key challenges:

- (i) Finding an expression for the numbers $q(i, j; n)$, or equivalently for $Q(x, y; z)$;

Date: July 2, 2013.

1991 *Mathematics Subject Classification.* Primary 05A15; Secondary 30F10, 30D05.

Key words and phrases. Walks in the quarter plane; Counting generating function; Holonomy; Group of the walk; Riemann surface; Elliptic functions; Uniformization; Universal covering.




FIGURE 1. Example of model (with an infinite group) considered in this article (note that on the boundary, the jumps are the natural ones: those that would take the walk out \mathbf{Z}_+^2 are discarded)

- (ii) Determining the nature of $Q(x, y; z)$: is it holonomic?¹ And in that event, is it algebraic, or even rational?

The functional equation (2) below served as the basis for all further analyses [2, 6, 9, 10, 18, 19, 23, 24, 25]. It determines $Q(x, y; z)$ through the boundary functions $Q(x, 0; z)$, $Q(0, y; z)$ and $Q(0, 0; z)$, as follows:

$$(2) \quad K(x, y; z)Q(x, y; z) = K(x, 0; z)Q(x, 0; z) + K(0, y; z)Q(0, y; z) - K(0, 0; z)Q(0, 0; z) - xy,$$

where

$$(3) \quad K(x, y; z) = xyz[\sum_{(i,j) \in \mathcal{S}} x^i y^j - 1/z]$$

is called the kernel of the walk. We refer to [6] for the proof of Equation (2). It has been anticipated in [6] and confirmed in the articles [2, 10, 18, 19, 23, 24, 25] that the analysis of both problems (i) and (ii) highlights the notion of the group of the walk, introduced by Malyshev [20, 21, 22]. This is the group

$$(4) \quad \langle \xi, \eta \rangle$$

of birational transformations of $(\mathbf{C} \cup \{\infty\})^2$, which is generated by

$$(5) \quad \xi(x, y) = \left(x, \frac{1}{y} \frac{\sum_{(i,-1) \in \mathcal{S}} x^i}{\sum_{(i,+1) \in \mathcal{S}} x^i} \right), \quad \eta(x, y) = \left(\frac{1}{x} \frac{\sum_{(-1,j) \in \mathcal{S}} y^j}{\sum_{(+1,j) \in \mathcal{S}} y^j}, y \right).$$

Each element of $\langle \xi, \eta \rangle$ leaves invariant $\sum_{(i,j) \in \mathcal{S}} x^i y^j$, the GF of the step set \mathcal{S} . Further, $\xi^2 = \eta^2 = \text{id}$, and $\langle \xi, \eta \rangle$ is a dihedral group of order even and larger than or equal to four. It has been proved in [6] that 23 of the 79 walks have a finite group, while the 56 others admit an infinite group.²

¹In other words, see [12, Appendix B.4], is the vector space over $\mathbf{C}(x, y, z)$ —the field of rational functions in the three variables x, y, z —spanned by the set of all derivatives of $Q(x, y; z)$ finite dimensional?

²Proving that a given model has a finite group is easy: it suffices to compute the group defined in (4). As examples, the models of Figure 2 all have finite groups, of order 4, 6, 8 and 8, respectively. On the other hand, it is much more complicated to prove that a model has an infinite group. To that purpose, methods are presented in [6]; see also [9, 11].




FIGURE 2. Four famous examples, known as simple, Kreweras', Gessel's and Gouyou-Beauchamps' walks, respectively




FIGURE 3. The 5 singular walks studied in [23, 24]

1.2. Existing results in the literature. For 22 of the 23 models with finite group, GFs $Q(x, 0; z)$, $Q(0, y; z)$ and $Q(0, 0; z)$ —and hence $Q(x, y; z)$ by (2)—have been computed in [6] by means of certain (half-)orbit sums of the functional equation (2). For the 23rd model with finite group, known as Gessel's walks (see Figure 2), the GFs have been expressed by radicals in [2] thanks to a guessing-proving method using computer calculations; they were also found in [18] by solving some boundary value problems. All 23 finite group models admit a holonomic GF: 19 walks turn out to have a holonomic but non-algebraic GF, while for 4 walks (among which Kreweras' and Gessel's models), $Q(x, y; z)$ is algebraic. This was first proved in [2, 6]; see also [10, 19] for alternative proofs.

The 5 models with infinite group on Figure 3 are commonly known as singular walks. They are notably distinct from the others, since for them, the genus of the elliptic curve

$$(6) \quad \mathbf{T}_z = \{(x, y) \in (\mathbf{C} \cup \{\infty\})^2 : K(x, y; z) = 0\}$$

is 0, while for all other 74 models the genus of \mathbf{T}_z is 1, see [9, 25]. These 5 models are studied in detail in [23, 24]; they all have non-holonomic GFs.

At this step, there remain $51 = 79 - 23 - 5$ models. In [25] the problem (i) was resolved for all these 51 models—and in fact for all the 74 non-singular walks. This was done via a unified approach: integral representations were obtained for GFs $Q(x, 0; z)$, $Q(0, y; z)$ and $Q(0, 0; z)$ in certain domains, by solving boundary value problems (of Riemann-Carleman type). However, these complicated explicit expressions have not been helpful for solving problem (ii), that is for determining the nature of the GFs for the 51 non-singular walks with infinite group.

This problem has been finally solved in [19], as follows. Since the transformations ξ and η of $(\mathbf{C} \cup \{\infty\})^2$ leave $\sum_{(i,j) \in \mathcal{S}} x^i y^j$ invariant, one can consider a group $\langle \xi, \eta \rangle_{\mathbf{T}_z}$ of automorphisms of the elliptic curve (6) generated by its automorphisms ξ and η . Due to the obvious inclusion $\mathbf{T}_z \subset (\mathbf{C} \cup \{\infty\})^2$, it may happen that the group $\langle \xi, \eta \rangle_{\mathbf{T}_z}$ is finite

for some z , while the group $\langle \xi, \eta \rangle$ on $(\mathbf{C} \cup \{\infty\})^2$ is infinite. Let

$$(7) \quad \mathcal{H} = \{z \in]0, 1/|\mathcal{S}[[: |\langle \xi, \eta \rangle|_{\mathbf{T}_z}| < \infty\}.$$

Clearly $\mathcal{H} =]0, 1/|\mathcal{S}[[$ for any of the 23 models with finite group $\langle \xi, \eta \rangle$. The following result is proved in [19].³

Theorem 1 ([19]). *For all 51 non-singular models with infinite group:*

- (i) *The subsets \mathcal{H} and $]0, 1/|\mathcal{S}[[\setminus \mathcal{H}$ are both dense in $]0, 1/|\mathcal{S}[[$;*
- (ii) *For all $z \in \mathcal{H}$, the GFs $x \mapsto Q(x, 0; z)$ and $y \mapsto Q(0, y; z)$ are holonomic;*
- (iii) *For all $z \in]0, 1/|\mathcal{S}[[\setminus \mathcal{H}$, $x \mapsto Q(x, 0; z)$ and $y \mapsto Q(0, y; z)$ are non-holonomic.*

Assertion (iii) of Theorem 1 combined with usual properties of holonomic functions (see, e.g., [12, Appendix B.4]) implies the non-holonomy of the trivariate GF $Q(x, y; z)$ for all 51 models and therefore proves Bousquet-Mélou's and Mishna's conjecture stated in [6]. It has been remarked in [19] that assertion (ii) of the same theorem suggests a promising start in order to achieve better understanding and easier representations of the GFs, than those in [19, 25] mentioned above. This is the subject of the present paper.

1.3. Aim, methods and structure of this article. Here we analyze the GFs $Q(x, 0; z)$ and $Q(0, y; z)$ for all 74 non-singular models and all z such that $z \in \mathcal{H}$. More precisely, we compute the GFs (together with all their infinitely many branches) as sums of absolutely convergent series, where all terms are made explicit via an analysis of points of \mathbf{T}_z where $x = \infty$ or $y = \infty$. This result provides an alternative and completely different representation of these functions than the one given in [25] in terms of solutions to boundary value problems. It seems us more explicit and in this sense more satisfactory. Further, as an illustration of our approach, we work out this representation for Kreweras' model (see Figure 2 for its step set). In this way we obtain new expressions for the GFs of this model in terms of Weierstrass Zeta functions with known periods.

Our approach is deeply different that the ones used in the recent papers [10, 11, 18, 19, 25]. It has been first suggested by Malyshev in [20, § 6 in Chapter 5], at that time for studying the stationary probability generating functions of ergodic random walks in \mathbf{Z}_+^2 . The main idea consists in using, in a constructive way, Mittag-Leffler's theorem.⁴ Specifically, we first find the poles (and the principal parts at these poles) of the lifted GFs on the universal covering \mathbf{C} of \mathbf{T}_z ; then we show how to express the GFs as an infinite series of principal parts at its poles.

Another possible method for obtaining the GFs for $z \in \mathcal{H}$ is based on [9, Chapter 4]. It is commented in Appendix A.

³The key tool for the proof in [19] is the following: for any $z \in]0, 1/|\mathcal{S}[[$, GFs $Q(x, 0; z)$ and $Q(0, y; z)$ can be continued on the whole of \mathbf{C} as multi-valued functions with infinitely many (and explicit) meromorphic branches. Then the set of poles of all these branches is proved to be infinite for any $z \in]0, 1/|\mathcal{S}[[\setminus \mathcal{H}$, which leads to the non-holonomy of the GFs.

⁴Roughly speaking, Mittag-Leffler's theorem states that it is possible to construct a meromorphic function with prescribed (discrete) set of poles and prescribed principal parts at these poles, see [26, Section 4.13].

Our paper has the following organization: Sections 2–6 are devoted to the general theory for the 74 models of non-singular walks. In Section 2, functions $Q(x, 0; z)$ and $Q(0, y; z)$ are lifted on the elliptic curve \mathbf{T}_z , and then on its universal covering \mathbf{C} . They are then continued as meromorphic functions on the whole of \mathbf{C} . This continuation procedure, which is described in detail in [19], is based on ideas of [9]. We only sketch it here, restricting ourselves with notations and details necessary for the next sections. In Sections 3–5, for all 74 models and any $z \in \mathcal{H}$, all branches of $Q(x, 0; z)$ and $Q(0, y; z)$ are computed as sums of absolutely convergent series, the main result of the paper is stated in Theorem 7. Section 6 proposes some refinements in several directions. Sections 7–9 propose an illustration of our results on the example of Kreweras' model. In Section 7 we obtain an expression for the generating function in terms of Weierstrass ζ -functions. In Sections 8–9 we show how to derive the well-known expressions for $Q(0, 0; z)$ and $Q(x, 0; z) = Q(0, x; z)$ by the theory of transformation of elliptic functions. Finally, in Appendix A we present another possible approach.

In what follows, most of the time, we shall drop the dependence of all quantities with respect to the variable z , writing, for example, $Q(0, 0)$ instead of $Q(0, 0; z)$.

Part 1. GFs for all non singular models of walks

2. MEROMORPHIC CONTINUATION OF THE GFs ON THE UNIVERSAL COVERING

Branch points. Let us fix $z \in]0, 1[\setminus \mathcal{S}[$. The kernel $K(x, y)$ defined in (3) is a polynomial of the second degree in both x and y . The algebraic function $X(y)$ defined by $K(X(y), y) = 0$ has thus two branches and four branch points, that we call y_i , $i \in \{1, \dots, 4\}$.⁵ The latter are the roots of the discriminant of the second degree equation $K(x, y) = 0$ in the variable x :

$$(8) \quad \tilde{d}(y) = (zy^2\delta_{0,1} - y + z\delta_{0,-1})^2 - 4z^2(y^2\delta_{1,1} + y\delta_{1,0} + \delta_{1,-1})(y^2\delta_{-1,1} + y\delta_{-1,0} + \delta_{-1,-1}).$$

Above and throughout, we note

$$\delta_{i,j} = \begin{cases} 1 & \text{if } (i, j) \in \mathcal{S}, \\ 0 & \text{if } (i, j) \notin \mathcal{S}. \end{cases}$$

We refer to [19, Section 2.1] for the numbering and for further properties of the y_i 's. For any non-singular model, the Riemann surface of $X(y)$ is a torus (i.e., a Riemann surface of genus 1) \mathbf{T}_y composed of two complex spheres $\mathbf{C} \cup \{\infty\}$ which are properly glued together along the cuts $[y_1, y_2]$ and $[y_3, y_4]$. The analogous statement is true for the algebraic function $Y(x)$ defined by $K(x, Y(x)) = 0$, and for its four branch points that are roots of

$$(9) \quad d(x) = (zx^2\delta_{1,0} - x + z\delta_{-1,0})^2 - 4z^2(x^2\delta_{1,1} + x\delta_{0,1} + \delta_{-1,1})(x^2\delta_{1,-1} + x\delta_{0,-1} + \delta_{-1,-1}),$$

and its Riemann surface which is a torus \mathbf{T}_x . Since \mathbf{T}_x and \mathbf{T}_y are equivalent, in the sequel we consider a single Riemann surface \mathbf{T} with two different coverings $x, y : \mathbf{T} \rightarrow \mathbf{S}$.

⁵That is, four points such that the two branches of $X(y)$ are equal.




FIGURE 4. The universal covering and the location of $\Pi_{0,0} = \mathcal{M}_{0,0} \cup \mathcal{M}_{1,0}$ on it

Galois automorphisms. Any point $s \in \mathbf{T}$ admits the two coordinates $(x(s), y(s))$, which by construction satisfy $K(x(s), y(s)) = 0$. For any $s \in \mathbf{T}$, there exists a unique s' (resp. s'') such that $x(s') = x(s)$ (resp. $y(s'') = y(s)$). The values $x(s), x(s')$ (resp. $y(s), y(s'')$) are the two roots of the equation of the second degree $K(x, y(s)) = 0$ (resp. $K(x(s), y) = 0$) in x (resp. y). We define the automorphisms $\xi : \mathbf{T} \rightarrow \mathbf{T}$ (resp. $\eta : \mathbf{T} \rightarrow \mathbf{T}$) such that $\xi s = s'$ (resp. $\eta s = s''$) and call them Galois automorphisms, following the terminology of [9]. Clearly $\xi s = s$ (resp. $\eta s = s$) if and only if $x(s) = x_i$, $i \in \{1, \dots, 4\}$ (resp. $y(s) = y_i$, for some $i \in \{1, \dots, 4\}$).

Universal covering and uniformization. The torus \mathbf{T} is isomorphic to a certain quotient space $\mathbf{C}/(\omega_1\mathbf{Z} + \omega_2\mathbf{Z})$, where ω_1, ω_2 are complex numbers linearly independent on \mathbf{R} . This set can obviously be thought as the (fundamental) parallelogram $[0, \omega_1] + [0, \omega_2]$, whose opposed edges are identified. Up to a unimodular transform, ω_1, ω_2 are unique and found in [9, Lemma 3.3.2]:

$$(10) \quad \omega_1 = i \int_{x_1}^{x_2} \frac{dx}{\sqrt{-d(x)}}, \quad \omega_2 = \int_{x_2}^{x_3} \frac{dx}{\sqrt{d(x)}}.$$

For an upcoming use, we define

$$g_x(t) = \begin{cases} d''(x_4)/6 + d'(x_4)/[t - x_4] & \text{if } x_4 \neq \infty, \\ d''(0)/6 + d'''(0)t/6 & \text{if } x_4 = \infty, \end{cases}$$

as well as

$$g_y(t) = \begin{cases} \tilde{d}''(y_4)/6 + \tilde{d}'(y_4)/[t - y_4] & \text{if } y_4 \neq \infty, \\ \tilde{d}''(0)/6 + \tilde{d}'''(0)t/6 & \text{if } y_4 = \infty, \end{cases}$$

and finally we introduce $\wp(\omega; \omega_1, \omega_2)$, the Weierstrass elliptic function with periods ω_1, ω_2 . Throughout, we shall write $\wp(\omega)$ for $\wp(\omega; \omega_1, \omega_2)$. By definition (see, e.g., [15, 28]), we have

$$(11) \quad \wp(\omega) = \frac{1}{\omega^2} + \sum_{(\ell_1, \ell_2) \in \mathbf{Z}^2 \setminus \{(0,0)\}} \left[\frac{1}{(\omega - \ell_1\omega_1 - \ell_2\omega_2)^2} - \frac{1}{(\ell_1\omega_1 + \ell_2\omega_2)^2} \right].$$




FIGURE 5. The fundamental parallelogram $\Pi_{0,0} = \omega_1[0, 1[+ \omega_2[0, 1[$, and important points and domains on it

The universal covering of \mathbf{T} has the form (\mathbf{C}, λ) , where \mathbf{C} is the usual complex plane (that can be viewed as the union of infinitely many parallelograms $\Pi_{m,n} = \omega_1[m, m + 1[+ \omega_2[n, n + 1[$ glued together for $m, n \in \mathbf{Z}$) and $\lambda : \mathbf{C} \rightarrow \mathbf{T}$ is a non-branching covering map; see Figure 4. For any $\omega \in \mathbf{C}$ such that $\lambda\omega = s \in \mathbf{T}$, we have $x(\lambda\omega) = x(s)$ and $y(\lambda\omega) = y(s)$. The uniformization formulæ [9, Lemma 3.3.1] are

$$(12) \quad \begin{cases} x(\lambda\omega) = g_x^{-1}(\wp(\omega)), \\ y(\lambda\omega) = g_y^{-1}(\wp(\omega - \omega_3/2)), \end{cases}$$

where

$$(13) \quad \omega_3 = \int_{X(y_1)}^{x_1} \frac{dx}{\sqrt{d(x)}}.$$

Throughout, we shall write $x(\lambda\omega) = x(\omega)$ and $y(\lambda\omega) = y(\omega)$. These are elliptic functions on \mathbf{C} with periods ω_1, ω_2 . Clearly

$$(14) \quad K(x(\omega), y(\omega)) = 0, \quad \forall \omega \in \mathbf{C}.$$

Furthermore, since each parallelogram $\Pi_{m,n}$ represents a torus \mathbf{T} composed of two complex spheres, the function $x(\omega)$ (resp. $y(\omega)$) takes each value of $\mathbf{C} \cup \infty$ twice within this parallelogram, except for the branch points $x_i, i \in \{1, \dots, 4\}$ (resp. $y_i, i \in \{1, \dots, 4\}$). Points $\omega_{x_i} \in \Pi_{0,0}$ such that $x(\omega_{x_i}) = x_i, i \in \{1, \dots, 4\}$ are pictured on Figure 5. They are equal to $\omega_{x_4} = 0, \omega_{x_1} = \omega_2/2, \omega_{x_3} = \omega_1/2$ and $\omega_{x_2} = (\omega_1 + \omega_2)/2$. Points ω_{y_i} such that $y(\omega_{y_i}) = y_i$ are just the shifts of ω_{x_i} by the real vector $\omega_3/2$: $\omega_{y_i} = \omega_{x_i} + \omega_3/2$ for $i \in \{1, \dots, 4\}$, see also Figure 5.

The automorphisms ξ and η on \mathbf{T} are lifted on the universal covering \mathbf{C} in such a way that ω_{x_2} and ω_{y_2} stay their fixed points, respectively:

$$(15) \quad \widehat{\xi}\omega = -\omega + 2\omega_{x_2}, \quad \widehat{\eta}\omega = -\omega + 2\omega_{y_2}, \quad \forall \omega \in \mathbf{C}.$$

Clearly, one has

$$(16) \quad x(\widehat{\xi}\omega) = x(\omega), \quad y(\widehat{\eta}\omega) = y(\omega), \quad \forall \omega \in \mathbf{C}.$$

Meromorphic continuation of the GFs. The domains

$$\{\omega \in \mathbf{C} : |x(\omega)| < 1\}, \quad \{\omega \in \mathbf{C} : |y(\omega)| < 1\}$$

consist of infinitely many curvilinear strips, which differ from translations by a multiple of ω_2 . We denote by Δ_x (resp. Δ_y) the strip that is within $\cup_{n \in \mathbf{Z}} \Pi_{0,n}$ (resp. $\cup_{n \in \mathbf{Z}} \Pi_{0,n} + \omega_3/2$). The domain Δ_x (resp. Δ_y) is centered around the straight line $L_{x_1}^{x_2} = \{\omega \in \mathbf{C} : \Re \omega = \Re \omega_{x_2}\}$ (resp. $L_{y_1}^{y_2} = \{\omega \in \mathbf{C} : \Re \omega = \Re \omega_{y_2}\}$), see Figure 5. Note that the function $Q(x(\omega), 0)$ (resp. $Q(0, y(\omega))$) is well defined in Δ_x (resp. Δ_y), by (1). Let us put

$$(17) \quad r_x(\omega) = K(x(\omega), 0)Q(x(\omega), 0), \quad \forall \omega \in \Delta_x, \quad r_y(\omega) = K(0, y(\omega))Q(0, y(\omega)), \quad \forall \omega \in \Delta_y.$$

The domain $\Delta_x \cap \Delta_y$ is a non-empty open strip. It follows from (2), (14) and (17) that

$$(18) \quad r_x(\omega) + r_y(\omega) - Q(0, 0)K(0, 0) - x(\omega)y(\omega) = 0, \quad \forall \omega \in \Delta_x \cap \Delta_y.$$

Let

$$(19) \quad \Delta = \Delta_x \cup \Delta_y.$$

Then the functions $r_x(\omega)$ and $r_y(\omega)$ can be continued as meromorphic functions on the whole of Δ , setting

$$(20) \quad r_x(\omega) = -r_y(\omega) + Q(0, 0)K(0, 0) + x(\omega)y(\omega), \quad \forall \omega \in \Delta_y,$$

$$(21) \quad r_y(\omega) = -r_x(\omega) + Q(0, 0)K(0, 0) + x(\omega)y(\omega), \quad \forall \omega \in \Delta_x.$$

The following theorem holds true, see [19] or [9].

Theorem 2 ([19],[9]). *We have*

$$(22) \quad \bigcup_{n \in \mathbf{Z}} (\Delta + n\omega_3) = \mathbf{C}.$$

The functions $r_x(\omega)$ and $r_y(\omega)$ can be continued meromorphically from Δ to the whole of \mathbf{C} via the following formulæ:

$$(23) \quad r_x(\omega - \omega_3) = r_x(\omega) + y(\omega)[x(-\omega + 2\omega_{y_2}) - x(\omega)], \quad \forall \omega \in \mathbf{C},$$

$$(24) \quad r_y(\omega + \omega_3) = r_y(\omega) + x(\omega)[y(-\omega + 2\omega_{x_2}) - y(\omega)], \quad \forall \omega \in \mathbf{C}.$$

Further, we have, for any $\omega \in \mathbf{C}$,

$$(25) \quad r_x(\omega) + r_y(\omega) - K(0,0)Q(0,0) - x(\omega)y(\omega) = 0,$$

$$(26) \quad r_x(\widehat{\xi}\omega) = r_x(\omega),$$

$$(27) \quad r_y(\widehat{\eta}\omega) = r_y(\omega),$$

$$(28) \quad r_x(\omega + \omega_1) = r_x(\omega),$$

$$(29) \quad r_y(\omega + \omega_1) = r_y(\omega).$$

The restrictions of $r_x(\omega)/K(x(\omega), 0)$ on

$$(30) \quad \mathcal{M}_{k,0} = \omega_1[0, 1[+ \omega_2[k/2, (k+1)/2[$$

for $k \in \{1, 2, \dots\}$ provide all branches on $\mathbf{C} \setminus ([x_1, x_2] \cup [x_3, x_4])$ of $Q(x, 0)$ as follows:

$$(31)$$

$$Q(x, 0) = \{r_x(\omega)/K(x(\omega), 0) : \omega \text{ is the (unique) element of } \mathcal{M}_{k,0} \text{ such that } x(\omega) = x\}.$$

Indeed, due to the ω_1 -periodicity of $r_x(\omega)$ and $x(\omega)$, the restrictions of these functions on $\mathcal{M}_{k,\ell} = \omega_1[\ell, \ell+1[+ \omega_2[k/2, (k+1)/2[$ do not depend on $\ell \in \mathbf{Z}$, and therefore determine the same branch as on $\mathcal{M}_{k,0}$ for any ℓ . Further, by (26) the restrictions of $r_x(\omega)/K(x(\omega), 0)$ on $\mathcal{M}_{-k+1,0}$ and on $\mathcal{M}_{k,0}$ lead to the same branches for any $k \in \mathbf{Z}$. See Figure 4. The analogous statement holds for the restrictions of $r_y(\omega)/K(0, y(\omega))$ on

$$(32) \quad \mathcal{N}_{k,0} = \omega_3/2 + \omega_1[0, 1[+ \omega_2[k/2, (k+1)/2[$$

for $k \in \{1, 2, \dots\}$.

3. EXPRESSION OF THE GFs IN TERMS OF THE PRINCIPAL PARTS AT THEIR POLES

Although the dynamic of the meromorphic continuation procedure (see (20), (21), (23) and (24)) is fairly easy, the initial values (17) of $r_x(\omega)$ in Δ_x and of $r_y(\omega)$ in Δ_y can be obtained only via complicated integral representations of $Q(x, 0)$ and $Q(0, y)$ given in [25], by solving certain boundary value problems. In this section we give another representation of $r_x(\omega)$ and $r_y(\omega)$ for any $\omega \in \mathbf{C}$. It is valid for any $z \in \mathcal{H}$ (see (7) for its definition). We remind that for all 23 models with finite group, $\mathcal{H} =]0, 1/\mathcal{S}[$, while for all 51 non-singular models with infinite group, \mathcal{H} is dense in $]0, 1/\mathcal{S}[$, but not equal to the latter, see Theorem 1. In [19], the set \mathcal{H} has been also characterized as

$$\mathcal{H} = \{z \in]0, 1/\mathcal{S}[: \omega_3/\omega_2 \text{ is rational}\}.$$

Therefore, for any fixed $z \in \mathcal{H}$, one has

$$(33) \quad \frac{\omega_3}{\omega_2} = \frac{k}{\ell}$$

for some integers $\ell > k > 0$ without common divisors. In the sequel we work under this assumption.

The other representation of the GFs $r_x(\omega)$ and $r_y(\omega)$ that we shall obtain here is based on a constructive application of Mittag-Leffler's theorem. Specifically, the latter theorem states that it is possible to construct a meromorphic function with prescribed (discrete) set of poles and prescribed principal parts at these poles (see Definition 3 below), see [26,

Section 4.13]. Our main result in this section (Theorem 4) states that the GFs are the limits of the sums of the principal parts at their poles along certain sequences of parallelograms.

Definition 3. *Let $g(\omega)$ be a function meromorphic in the neighborhood of $\omega_0 \in \mathbf{C}$, where it has a pole of order p , such that $g(\omega) = \sum_{k=-p}^{\infty} g_k(\omega - \omega_0)^k$. Its principal part at ω_0 is given by $\sum_{k=-p}^{-1} g_k(\omega - \omega_0)^k$.*

Therefore, we first need to construct a sequence of parallelograms $(P_m)_{m \geq 1}$ on the universal covering \mathbf{C} with the following properties:

- (i) $P_1 \subset P_2 \subset \dots$;
- (ii) $\cup_{m \geq 1} P_m = \mathbf{C}$;
- (iii) All the P_m 's are centered around ω_{y_2} ;
- (iv) There exist constants $C_1, C_2 > 0$ such that for all $m \geq 1$,

$$(34) \quad \sup_{\omega \in \partial P_m} |r_y(\omega)| \leq C_1 m,$$

$$(35) \quad \inf_{\omega \in \partial P_m} |\omega - \omega_{y_2}| \geq C_2 m,$$

where ∂P_m denotes the boundary of P_m .

The angle stone of this construction is the equality (33), under which there are many possibilities to perform it; below we just give one.

Construction of a family $(P_m)_{m \geq 1}$ satisfying (i), (ii), (iii) and (iv). Let

$$(36) \quad f_y(\omega) = x(\omega)[y(-\omega + 2\omega_{x_2}) - y(\omega)]$$

be the function used in the meromorphic continuation procedure (24). It has at most six distinct poles in the main parallelogram $\Pi_{0,0} + \omega_3/2$ for r_y , we denote them by $p_1^{(0)}, \dots, p_6^{(0)}$. (It is worth noting that for some of the 51 non-singular models with infinite group, some of these poles may coincide, so that their number may be smaller—see [19] and also Remark 13 in this article.) Let $n \in \{1, \dots, \ell - 1\}$, $i \in \{1, \dots, 6\}$, and consider $p_i^{(0)} - n\omega_3$. There exists a unique $m_{i,n} \in \mathbf{Z}_+$ such that $p_i^{(n)} = p_i^{(0)} - n\omega_3 + m_{i,n}\omega_2 \in \Pi_{0,0} + \omega_3/2$. Let

$$\mathcal{R} = \{\omega \in \Pi_{0,0} + \omega_3/2 : \Re \omega \neq p_i^{(n)}, \forall i \in \{1, \dots, 6\}, \forall n \in \{1, \dots, \ell - 1\}\}.$$

In other words, \mathcal{R} is the domain $\Pi_{0,0} + \omega_3/2$ without the set of at most $6(\ell - 1)$ vertical segments. With (24) combined with (33), one deduces that for any $\omega \in \mathcal{R}$ and any $m \in \mathbf{Z}$, $\omega + m\omega_3$ is not a pole of $r_y(\omega)$.

Let us now fix $\omega^+, \omega^- \in \mathcal{R}$ symmetric with respect to ω_{y_2} and such that $\Re \omega^+ > \Re \omega^-$, and define the vertical straight lines

$$V_m^+ = \{\omega \in \mathbf{C} : \Re \omega = \Re \omega^+ + m\omega_3\}, \quad V_m^- = \{\omega \in \mathbf{C} : \Re \omega = \Re \omega^- - m\omega_3\}.$$

Then by Equations (24), (29) and by the assumption (33) the function $|r_y(\omega)|$ on them can be estimated as follows:

$$(37) \quad \sup_{\omega \in V_m^+ \cup V_m^-} |r_y(\omega)| \leq C_3 + mC_4,$$

where

$$(38) \quad C_3 = \sup_{\substack{\omega \in \Pi_{0,0} + \omega_3/2 \\ \Re\omega = \omega^+ \text{ or } \Re\omega = \omega^-}} |r_y(\omega)| < \infty$$

and

$$C_4 = \max_{n \in \{0, \dots, \ell-1\}} \left\{ \sup_{\substack{\omega \in \Pi_{0,0} + \omega_3/2, \\ \Re\omega = \omega^+}} |f_y(\omega + n\omega_3)|, \sup_{\substack{\omega \in \Pi_{0,0} + \omega_3/2, \\ \Re\omega = \omega^-}} |f_y(\omega - n\omega_3)| \right\} < \infty.$$

Let us fix any $\omega^*, \omega^{**} \in \Pi_{0,0} + \omega_3/2$ symmetric with respect to ω_{y_2} , $\Im\omega^* > \Im\omega^{**}$, and such that for all $i \in \{1, \dots, 6\}$, $\Im\omega^* \neq \Im p_i^{(0)}$ and $\Im\omega^{**} \neq \Im p_i^{(0)}$. Then due to ω_2 -periodicity of f_y ,

$$(39) \quad C_5 = \sup_{\Im\omega = \Im\omega^* \text{ or } \Im\omega = \Im\omega^{**}} |f_y(\omega)| < \infty.$$

It follows from (24) that $r_y(\omega)$ does not have poles at points $\omega \in \mathbf{C}$ such that $\Im\omega \in \{\Im\omega^*, \Im\omega^{**}\}$, and in particular

$$(40) \quad C_6 = \sup_{\substack{\Im\omega = \Im\omega^* \text{ or } \Im\omega = \Im\omega^{**}, \\ \Re\omega^- \leq \Re\omega \leq \Re\omega^+}} |r_y(\omega)| < \infty.$$

Define now the horizontal straight lines

$$H_p^* = \{\omega \in \mathbf{C} : \Im\omega = \Im\omega^* + p|\omega_1|\}, \quad H_p^{**} = \{\omega \in \mathbf{C} : \Im\omega = \Im\omega^{**} - p|\omega_1|\}$$

for any $p \in \mathbf{Z}_+$. Let us estimate $|r_y(\omega)|$ on their segments bounded by the intersection points with V_m^+ and V_m^- . By (24), (29), (39) and (40) the following bound holds true for any $p \in \mathbf{Z}_+$:

$$(41) \quad \sup_{\substack{\omega \in H_p^* \cup H_p^{**}, \\ \Re\omega^- - m\omega_3 \leq \Re\omega \leq \Re\omega^+ + m\omega_3}} |r_y(\omega)| = \sup_{\substack{\Im\omega = \Im\omega^* \text{ or } \Im\omega = \Im\omega^{**}, \\ \Re\omega^- - m\omega_3 \leq \Re\omega \leq \Re\omega^+ + m\omega_3}} |r_y(\omega)| \leq C_6 + mC_5 < \infty.$$

Let $(p_m)_{m \geq 1}$ be any sequence of integers strictly increasing to infinity as $m \rightarrow \infty$ and such that $p_m \geq m$. Let us construct the parallelograms P_m 's bounded by V_m^+ , V_m^- , $H_{p_m}^*$ and $H_{p_m}^{**}$, which are defined above. Properties (i), (ii) and (iii) are trivially satisfied for these P_m 's. The estimate (34) is ensured by (37) and (41). Note that (33) was crucial to make (37) valid. Furthermore,

$$\inf_{\omega \in \partial P_m} |\omega - \omega_{y_2}| \geq m \min\{\omega_3, |\omega_1|\},$$

which shows the estimate (35). This concludes the construction of the P_m 's with required properties (i), (ii), (iii) and (iv). \square

We are now ready to state and prove the main result of this section.

Theorem 4. *Let d_i be the poles of the function $r_y(\omega)$ on \mathbf{C} and let $R_{d_i,y}(\omega)$ be the principal parts of $r_y(\omega)$ at these poles. Then for any family of parallelograms $(P_m)_{m \geq 1}$ satisfying (i), (ii), (iii) and (iv), one has*

$$(42) \quad r_y(\omega) - r_y(\omega_{y_2}) = \lim_{m \rightarrow \infty} \sum_{d_i \in P_m} R_{d_i,y}(\omega) - R_{d_i,y}(\omega_{y_2}), \quad \forall \omega \in \mathbf{C}.$$

Proof. We follow [8, Chapter V, proof of Theorem 2], as suggested in [20, page 163]. We have

$$(43) \quad \frac{1}{2\pi i} \int_{\partial P_m} \frac{r_y(\zeta)}{\zeta - \omega} d\zeta = \operatorname{res}_\omega \frac{r_y(\zeta)}{\zeta - \omega} + \sum_{d_i \in P_m} \operatorname{res}_{d_i} \frac{r_y(\zeta)}{\zeta - \omega} = r_y(\omega) + \sum_{d_i \in P_m} \operatorname{res}_{d_i} \frac{\sum_{d_i \in P_m} R_{d_i, y}(\zeta)}{\zeta - \omega}.$$

The rational function

$$G(\zeta) = \frac{\sum_{d_i \in P_m} R_{d_i, y}(\zeta)}{\zeta - \omega}$$

has a residue at infinity equal to zero (if there is at least one pole of r_y in P_m , that holds for all m large enough). Then $\sum_{d_i \in P_m} \operatorname{res}_{d_i} G(\zeta) + \operatorname{res}_\omega G(\zeta) = 0$. Hence

$$\sum_{d_i \in P_m} \operatorname{res}_{d_i} G(\zeta) = -\operatorname{res}_\omega G(\zeta) = -\sum_{d_i \in P_m} R_{d_i, y}(\omega),$$

so that

$$(44) \quad \frac{1}{2\pi i} \int_{\partial P_m} \frac{r_y(\zeta)}{\zeta - \omega} d\zeta = r_y(\omega) - \sum_{d_i \in P_m} R_{d_i, y}(\omega).$$

In particular, for $\omega = \omega_{y_2}$ we have

$$(45) \quad \frac{1}{2\pi i} \int_{\partial P_m} \frac{r_y(\zeta)}{\zeta - \omega_{y_2}} d\zeta = r_y(\omega_{y_2}) - \sum_{d_i \in P_m} R_{d_i, y}(\omega_{y_2}).$$

Moreover, taking the derivative in (44) at $\omega = \omega_{y_2}$ leads to

$$(46) \quad \frac{1}{2\pi i} \int_{\partial P_m} \frac{r_y(\zeta)}{(\zeta - \omega_{y_2})^2} d\zeta = r'_y(\omega_{y_2}) - \sum_{d_i \in P_m} R'_{d_i, y}(\omega_{y_2}).$$

Note that since $r_y(\omega) = r_y(-\omega + 2\omega_{y_2})$ by (26), then for each d_i pole of r_y , $-d_i + 2\omega_{y_2}$ is also a pole of r_y , and further $R'_{d_i, y}(\omega_{y_2}) = -R'_{-d_i + 2\omega_{y_2}, y}(\omega_{y_2})$. Since P_m is centered in ω_{y_2} , then either both $d_i, -d_i + 2\omega_{y_2}$ belong to P_m , or both do not. Hence $\sum_{d_i \in P_m} R'_{d_i, y}(\omega_{y_2}) = 0$.

Subtracting from (44) the identity (45) and also (46) multiplied by the quantity $(\omega - \omega_{y_2})$, we find

$$(47) \quad \begin{aligned} & \frac{1}{2\pi i} \int_{P_m} r_y(\zeta) \frac{(\omega - \omega_{y_2})^2}{(\zeta - \omega_{y_2})^2 (\zeta - \omega)} d\zeta \\ &= r_y(\omega) - \sum_{d_i \in P_m} R_{d_i, y}(\omega) - \left(r_y(\omega_{y_2}) - \sum_{d_i \in P_m} R_{d_i, y}(\omega_{y_2}) \right) - r'_y(\omega_{y_2})(\omega - \omega_{y_2}). \end{aligned}$$

By the assumption (34), one has

$$(48) \quad \left| \frac{1}{2\pi i} \int_{P_m} r_y(\zeta) \frac{(\omega - \omega_{y_2})^2}{(\zeta - \omega_{y_2})^2 (\zeta - \omega)} d\zeta \right| \leq \frac{|\omega - \omega_{y_2}|^2}{2\pi} \frac{C_1 m}{\inf_{\zeta \in \partial P_m} |\zeta - \omega|} \int_{\partial P_m} \frac{1}{|\zeta - \omega_{y_2}|^2} d\zeta.$$

By the assumption (35), one has

$$\inf_{\zeta \in \partial P_m} |\zeta - \omega| \geq \inf_{\zeta \in \partial P_m} |\zeta - \omega_{y_2}| - |\omega_{y_2} - \omega| \geq C_2 m - |\omega_{y_2} - \omega|.$$

We can also estimate

$$\int_{\partial P_m} \frac{1}{|\zeta - \omega_{y_2}|^2} d\zeta \leq \frac{4\pi}{\inf_{\zeta \in \partial P_m} |\zeta - \omega_{y_2}|} \leq \frac{4\pi}{C_2 m}.$$

It follows that (48) equals $O(1/m)$ as $m \rightarrow \infty$. Thus

$$r_y(\omega) - r_y(\omega_{y_2}) = \lim_{m \rightarrow \infty} \left(\sum_{d_i \in P_m} R_{d_i, y}(\omega) - \sum_{d_i \in P_m} R_{d_i, y}(\omega_{y_2}) \right) + r'_y(\omega_{y_2})(\omega - \omega_{y_2}).$$

The identity $r_y(\omega) = r_y(-\omega + 2\omega_{y_2})$ due to (26) implies $R_{d_i, y}(\omega) = R_{-d_i + 2\omega_{y_2}, y}(-\omega + 2\omega_{y_2})$. Hence for any $m \geq 1$,

$$\sum_{d_i \in P_m} R_{d_i, y}(\omega) = \sum_{d_i \in P_m} R_{d_i, y}(-\omega + 2\omega_{y_2}).$$

Then for any $\omega \in \mathbf{C}$, $r'_y(\omega_{y_2})(\omega - \omega_{y_2}) = r'_y(\omega_{y_2})(-\omega + \omega_{y_2})$, thus $r'_y(\omega_{y_2}) = 0$. This finishes the proof of the theorem. \square

The result of Theorem 4 has to be improved for several reasons: first, it would be important to identify the poles of r_y and the principal parts at them; second, it would be more convenient to represent r_y as an absolutely convergent series, independent of the parallelograms P_m 's; third, it would be useful to find the unknown constant $r_y(\omega_{y_2})$. These remarks give the structure of the following sections. In Section 4 we compute the principal parts at the poles of r_y in terms of the principal parts at the poles of f_y . In Section 5 we represent r_y as the sum of an absolutely convergent series, where all terms are expressed via the principal parts at the (at most) six poles of f_y in the parallelogram $\Pi_{0,0} - \omega_{x_1}$. In Section 5 we also compute $r_y(\omega_{y_2})$.

4. COMPUTATION OF THE POLES OF THE GFs AND OF THE PRINCIPAL PARTS AT THEM

In this section we study the poles of r_y and the principal parts at them. The first lemma deals with the poles of r_y in the domain Δ , which is defined in (19). Let f_y be the function defined in (36). We denote by $F_{d,y}(\omega)$ the principal part of $f_y(\omega)$ at the pole d .

Lemma 5. *A point d is a pole of r_y in Δ if and only if it is a pole of the function f_y in Δ with $\Re d < \Re \omega_{x_1}$. Furthermore, at such a point d ,*

$$(49) \quad R_{d,y}(\omega) = -F_{d,y}(\omega), \quad \forall \omega \in \mathbf{C}.$$

A point d' is a pole of $r_y + f_y$ in Δ if and only if it is a pole of f_y in Δ with $\Re d' > \Re \omega_{x_1}$.

Proof. We first prove the following facts:

- (i) $\Delta \cap \{\omega \in \mathbf{C} : y(\omega) = \infty\} \subset \{\omega \in \mathbf{C} : \Re \omega < \Re \omega_{x_1}\} \cup (\omega_{x_1} + \omega_1 \mathbf{Z})$;
- (ii) $\Delta \cap \{\omega \in \mathbf{C} : y(\widehat{\xi}\omega) = \infty\} \subset \{\omega \in \mathbf{C} : \Re \omega > \Re \omega_{x_1}\} \cup (\omega_{x_1} + \omega_1 \mathbf{Z})$;
- (iii) $\Delta \cap \{\omega \in \mathbf{C} : x(\omega) = \infty\} \subset \{\omega \in \mathbf{C} : \Re \omega > \Re \omega_{x_1}\}$.

Consider first the models with $y_4 < 0$. For them, points $\omega \in \mathbf{C}$ such that $y(\omega) = \infty$ are located on the vertical lines $\{\omega \in \mathbf{C} : \Re \omega = \Re \omega_{y_4} + m\omega_2\}$, $m \in \mathbf{Z}$. Only those on the line $\{\omega \in \mathbf{C} : \Re \omega = \Re \omega_{y_4}\}$ can be in Δ , and on this line we have $\Re \omega_{y_4} < \Re \omega_{x_1}$. This proves (i) for these models. Points ω with $y(\widehat{\xi}\omega) = \infty$ are on the lines $\widehat{\xi}\{\omega \in \mathbf{C} : \Re \omega = \Re \omega_{y_4} + m\omega_2\}$,

$m \in \mathbf{Z}$. None of them can be in Δ , except for those on $\widehat{\xi}\{\omega \in \mathbf{C} : \Re\omega = \Re\omega_{y_4}\}$. But we have $\Re\widehat{\xi}\omega_{y_4} > \Re\omega_{x_1}$. This concludes the proof of (ii) for these models.

Consider models with $y_4 > 0$ or $y_4 = \infty$. It is proved in [19] that points ω with $y(\omega) = \infty$ are located as follows: one of them, say f_1 , is such that $\Im f_1 = 0$ and $\Re\omega_{x_4} \leq \Re f_1 \leq \Re\omega_{y_4} < \Re\omega_{x_1}$; the other, say f_2 , is symmetric to f_1 w.r.t. ω_{y_4} . Furthermore $\Im f_2 = 0$ and $\Re\omega_{y_4} \leq \Re f_2 \leq \Im\omega_{x_1}$ (in the limiting case $y_4 = \infty$, one has $f_1 = f_2 = \omega_{y_4}$). All other points where $y(\omega) = \infty$ are the shifts of f_1, f_2 by $m\omega_2 + p\omega_1$, for $m, p \in \mathbf{Z}$. Only points of type $f_1 + p\omega_1$ and $f_2 + p\omega_1$ with $p \in \mathbf{Z}$ can be in Δ . We have for them $\Re(f_i + p\omega_1) = \Re f_i \leq \Re\omega_{x_1}$, $i \in \{1, 2\}$. Then either $\Re(f_i + p\omega_1) < \Re\omega_{x_1}$ or $\Re(f_i + p\omega_1) = \Re\omega_{x_1} = \Re(\omega_{x_1} + p\omega_1)$. In the last case the observation $\Im(f_i + p\omega_1) = p\omega_1 = \Im(\omega_{x_1} + p\omega_1)$, $i \in \{1, 2\}$, leads to $f_i + p\omega_1 = \omega_{x_1} + p\omega_1$ and finishes the proof of (i) for this class of models. Points ω such that $y(\widehat{\xi}\omega) = \infty$ are $\widehat{\xi}(f_1 + m\omega_2 + p\omega_1)$ and $\widehat{\xi}(f_2 + m\omega_2 + p\omega_1)$, for $m, p \in \mathbf{Z}$. Only points $\widehat{\xi}(f_i + p\omega_1)$, $i \in \{1, 2\}$, can be in Δ . If $\Re(f_i + p\omega_1) < \Re\omega_{x_1}$, then $\Re\widehat{\xi}(f_i + p\omega_1) > \Re\omega_{x_1}$. If $f_i + p\omega_1 = \omega_{x_1} + p\omega_1$, then $\widehat{\xi}(f_i + p\omega_1) = \omega_{x_1} + (-p + 1)\omega_1$. This proves assertion (ii) for all these models.

It remains to prove (iii). Consider ω with $x(\omega) = \infty$ first for models with $x_4 < 0$. Then points of \mathbf{C} with $x(\omega) = \infty$ lie on the vertical lines $\{\omega \in \mathbf{C} : \Re\omega = \Re\omega_{x_4} + m\omega_2\}$, $m \in \mathbf{Z}$. Only those on the line $\{\omega \in \mathbf{C} : \Re\omega = \Re\omega_{x_4} + \omega_2\}$ can be in Δ , where we have $\Re\omega_{x_4} + \omega_2 > \Re\omega_{x_1}$.

Let us finally prove (iii) for models such that $x_4 > 0$ or $x_4 = \infty$. It has been proved in [19] that points ω with $x(\omega) = \infty$ are located as follows: one of them, say f_3 , is such that $\Im f_3 = 0$ and $\Re\omega_{x_1} < \Re\omega_{x_4} + \omega_2 \leq \Re f_3 < \Re\omega_{y_4} + \omega_2$; the other, f_4 , is symmetric to f_3 w.r.t. $\omega_{x_4} + \omega_2$ and $\Re\omega_{x_1} < \Re\omega_{y_2} \leq \Re f_4 \leq \Re\omega_{x_4} + \omega_2$ (in the limiting case $x_4 = \infty$, one has $f_3 = f_4 = \omega_{x_4} + \omega_2$). All other points with $x(\omega) = \infty$ are their shifts by $m\omega_2 + p\omega_1$, where $m, p \in \mathbf{Z}$. Only points $f_3 + p\omega_1$ and $f_4 + p\omega_1$ with $p \in \mathbf{Z}$ can be in Δ . We have $\Re(f_3 + p\omega_1) = \Re f_3 > \Re\omega_{x_1}$ and $\Re(f_4 + p\omega_1) = \Re f_4 > \Re\omega_{x_1}$, that finishes the proof of item (iii).

Let d be a pole of r_y in Δ . Since r_y is holomorphic in Δ_y , then $d \in \Delta_x$ with $|x(d)| < 1$ and $r_x(d) \neq \infty$. Furthermore by (25), at $\omega = d$,

$$(50) \quad r_y(\omega) = -r_x(\omega) + Q(0, 0)K(0, 0) + x(\omega)y(\omega).$$

Then necessarily $x(d)y(d) = \infty$, from where $y(d) = \infty$. By assertion (i) above, we have either $\Re d < \Re\omega_{x_1}$ or $d = \omega_{x_1} + p\omega_1$ with some $p \in \mathbf{Z}$. The last fact is impossible since for none of the models, $\lim_{x \rightarrow x_1} Y(x) = \infty$ and simultaneously $\lim_{x \rightarrow x_1} xY(x) = \infty$ (remember that x_1 is a branch point of $Y(x)$ with $|x_1| < 1$). Hence $\Re d < \Re\omega_{x_1}$. Consider now $\widehat{\xi}d$. We have $|x(\widehat{\xi}d)| = |x(d)| < 1$, so that $\widehat{\xi}d \in \Delta_x \subset \Delta$ and $\Re(\widehat{\xi}d) = \Re(-d + 2\omega_{x_2}) > \Re\omega_{x_1}$. Then again by (i), $y(\widehat{\xi}d) \neq \infty$, so that $x(d)y(\widehat{\xi}d) \neq \infty$. Then the function $f_y(\omega) = x(\omega)[y(\widehat{\xi}\omega) - y(\omega)]$ has a pole at $\omega = d$. Furthermore by (50), one has

$$(51) \quad r_y(\omega) = -r_x(\omega) + K(0, 0) - f_y(\omega) + x(\omega)y(\widehat{\xi}\omega),$$

from where (49) follows.

Let d be a pole of f_y in Δ with $\Re d < \Re\omega_{x_1}$. Then necessarily one of the three following equalities holds: $y(d) = \infty$, $y(\widehat{\xi}d) = \infty$, or $x(d) = \infty$. By (i), (ii) and (iii), $y(d) = \infty$, so

that $d \in \Delta_x$ and $r_x(d) \neq \infty$. Thus $|x(d)| = |x(\widehat{\xi}d)| < 1$, $\widehat{\xi}d \in \Delta_x \subset \Delta$ and $\Re(\widehat{\xi}d) > \Re\omega_{x_1}$. By (i) one has $y(\widehat{\xi}d) \neq \infty$, so that $x(d)y(\widehat{\xi}d) \neq \infty$. The point d being a pole of f_y , it follows that $x(d)y(d) = \infty$. By (50), d is a pole of r_y in Δ . Finally, (51) implies (49).

It follows from (51) that any pole d of f_y in Δ with $\Re d < \Re\omega_{x_1}$ (or equivalently, any pole d of r_y in Δ) is not a pole of $r_y + f_y$ in Δ . It was proved that any pole d' of f_y in Δ with $\Re d' > \Re\omega_{x_1}$ is not a pole of r_y in Δ . Then d' must be a pole of $r_y + f_y$ in Δ . Finally, let us show that f_y cannot have a pole d'' in Δ with $\Re d'' = \Re\omega_{x_1}$. In fact, for any pole d'' of f_y , one of the three equalities holds: $y(d'') = \infty$, $y(\widehat{\xi}d'') = \infty$, or $x(d'') = \infty$. Assumption $\Re d'' = \Re\omega_{x_1}$ combined with (i), (ii) and (iii) yields $d'' = \omega_{x_1} + p\omega_1$, with $p \in \mathbf{Z}$. In particular, $y(d'') = \infty$. Then x_1 should be a branch point of $Y(x)$ with $|x_1| < 1$, $\lim_{x \rightarrow x_1} Y(x) = \infty$ and $\lim_{x \rightarrow x_1} x(Y_0(x) - Y_1(x)) = \infty$, which is true for none of the models. Hence f_y has no poles d'' with $\Re d'' = \Re\omega_{x_1}$, and Lemma 5 is proved. \square

Our ultimate goal is to obtain explicit expressions for $r_x(\omega)$ and $r_y(\omega)$ from Theorem 4. The theorem hereafter is a key preliminary result: it gives expressions for the principal parts $R_{d,y}(\omega)$ of r_y in terms of the principal parts of the elliptic function f_y at its poles. The main tool for that are the meromorphic continuation formulæ (23) and (24).

In what follows, we extend our notation $R_{d,y}(\omega)$ of Theorem 4 for any $d \in \mathbf{C}$, as follows: $R_{d,y}(\omega)$ will be the principal part of r_y at d in the sense of Definition 3 if d is a pole of r_y , and $R_{d,y}(\omega) = 0$ if d is not a pole of r_y . Further, throughout, we shall denote by $[x]$ (resp. $\lceil x \rceil$) the lower integer part (resp. the upper integer part) of $x \in \mathbf{R}$.

Theorem 6. *Assume the assertion (33). Denote by $F_{f,y}(\omega)$ the principal part of $f_y(\omega)$ at a pole f .*

(i) *Let $d \in \mathbf{C}$ with $\Re d < \Re\omega_{y_1}$. Let*

$$\mathcal{N}_d^- = \{n \geq 0 : d + n\omega_3 \text{ is a pole of } f_y \text{ with } \Re\omega_{x_1} - k\omega_2 < \Re d + n\omega_3 < \Re\omega_{x_1}\}.$$

If d is a pole of r_y , then $\mathcal{N}_d^- \neq \emptyset$. Furthermore, for any $d \in \mathbf{C}$ with $\Re d < \Re\omega_{y_1}$ and $\mathcal{N}_d^- \neq \emptyset$, we have:

$$(52) \quad R_{d,y}(\omega) = \sum_{n \in \mathcal{N}_d^-} -(\lfloor n/\ell \rfloor + 1)F_{d+n\omega_3,y}(\omega + n\omega_3).$$

(ii) *Let $d \in \mathbf{C}$ with $\Re d > \Re\omega_{y_1}$. Let*

$$\mathcal{N}_d^+ = \{n \geq 1 : d - n\omega_3 \text{ is a pole of } f_y \text{ with } \Re\omega_{x_1} < \Re d - n\omega_3 < \Re\omega_{x_1} + k\omega_2\}.$$

If d is a pole of r_y , then $\mathcal{N}_d^+ \neq \emptyset$. Furthermore, for any $d \in \mathbf{C}$ with $\Re d > \Re\omega_{y_1}$ and $\mathcal{N}_d^+ \neq \emptyset$, we have:

$$(53) \quad R_{d,y}(\omega) = \sum_{n \in \mathcal{N}_d^+} (\lfloor (n-1)/\ell \rfloor + 1)F_{d-n\omega_3,y}(\omega - n\omega_3).$$

Proof. Let d be a pole of r_y with $\Re d < \Re\omega_{y_1}$. Let us first prove that $\mathcal{N}_d^- \neq \emptyset$. Assume first that in addition $d \in \Delta$. By Lemma 5, d is a pole of f_y in Δ and $\Re d < \Re\omega_{x_1}$, so that $\mathcal{N}_d^- \neq \emptyset$. Let now d be a pole of r_y with $\Re d < \Re\omega_{y_1}$, but $d \notin \Delta$. In what follows, we

shall denote the domains of \mathbf{C} on the left of Δ by D^- and on the right of Δ by D^+ . Then $d \in D^-$. By (24), we have that for any $\omega \in D^-$,

$$(54) \quad r_y(\omega) = r_y(\omega + n\omega_3) - f_y(\omega + (n-1)\omega_3) - f_y(\omega + (n-2)\omega_3) - \cdots - f_y(\omega),$$

where n is such that $\omega + n\omega_3 \in \Delta$, while $\omega + (n-1)\omega_3, \omega + (n-2)\omega_3, \dots, \omega \notin \Delta$. If $d \in D^-$ is a pole of r_y , then either $d + t\omega_3$ is a pole of f_y for some $t \in \{0, \dots, n-1\}$ (then of course $\Re(d + t\omega_3) < \Re\omega_{x_1}$ since $d + t\omega_3 \in D^-$), or $d + n\omega_3$ is a pole of r_y in Δ , that is (by Lemma 5) a pole of f_y in Δ with $\Re d + n\omega_3 < \Re\omega_{x_1}$. This proves that $\mathcal{N}_d^- \neq \emptyset$.

Let us note that by (33), for any ω pole of f_y , $\omega - \ell\omega_3 = \omega - k\omega_2$ is also a pole of f_y , since this function is ω_2 -periodic, and so is $\omega - m\ell\omega_3$, for any $m \geq 1$. Then (52) follows from Lemma 5, Equations (49) and (54).

Let d be a pole of r_y with $\Re d > \Re\omega_{y_1}$. Then $\Re d > \Re\omega_{y_1} > \Re\omega_{x_1}$ and, by Lemma 5, $d \notin \Delta$. Then $d \in D^+$. By (24), we have that for any $\omega \in D^+$,

$$(55) \quad r_y(\omega) = r_y(\omega - n\omega_3) + f_y(\omega - n\omega_3) + f_y(\omega - (n-1)\omega_3) + \cdots + f_y(\omega - \omega_3),$$

where n is chosen such that $\omega - n\omega_3 \in \Delta$, but $\omega - (n-1)\omega_3, \omega - (n-2)\omega_3, \dots, \omega \notin \Delta$. If $d \in D^+$ is a pole of r_y , then either $d - t\omega_3$ is a pole of f_y for some $t \in \{1, \dots, n-1\}$ (then of course $\Re d - t\omega_3 > \Re\omega_{x_1}$, since $d - t\omega_3 \in D^+$), or $d - n\omega_3$ is a pole of $f_y + r_y$. By Lemma 5, the last fact is possible if and only if $d - n\omega_3$ is a pole of f_y in Δ with $\Re(d - n\omega_3) > \Re\omega_{x_1}$. Hence $\mathcal{N}_d^+ \neq \emptyset$.

By (33), for any ω pole of f_y and any $m \geq 1$, $\omega + m\ell\omega_3 = \omega + k\omega_2$ is also a pole of f_y , due to ω_2 -periodicity of f_y . Then (53) follows from Lemma 5 and Equation (55). \square

We remark that a point d with $\Re d < \Re\omega_{y_1}$ (resp. $\Re d > \Re\omega_{y_1}$) and $\mathcal{N}_d^- \neq \emptyset$ (resp. $\mathcal{N}_d^+ \neq \emptyset$) is not necessarily a pole of r_y , since in the sum (54) (resp. (55)), terms producing poles of r_y can compensate each other. However, the formula (52) (resp. (53)) is valid for any such d ; if d is not a pole, then obviously the RHS in the latter equation equals 0.

5. REPRESENTATION OF THE GFs IN TERMS OF CONVERGENT SERIES

Using Theorem 4 and the analysis of poles and principal parts of r_y in Section 4, we now compute the GFs $r_x(\omega)$ and $r_y(\omega)$ independently of the construction of the parallelograms $(P_m)_{m \geq 1}$. In the next theorem the GFs are expressed in terms of the principal parts of function $f_y(\omega) = y(\omega)[x(-\omega + 2\omega_{y_2}) - x(\omega)]$ (resp. $f_x(\omega) = x(\omega)[y(-\omega + 2\omega_{x_2}) - y(\omega)]$) at their (at most six) poles in the parallelogram $\Pi_{0,0} - \omega_{x_1}$ (resp. $\Pi_{0,0} + \omega_{y_1}$). This is the main result of the present paper valid for any of 74 non-singular models of walks and any $z \in \mathcal{H}$.

Theorem 7. *Assume the assertion (33). Let the f_i^y 's be the poles of the function f_y , and let $F_{f_i^y, y}(\omega)$ be the principal parts of this function at them. We have*

$$(56) \quad r_y(\omega) - r_y(\omega_{y_2}) = \sum_{p=-\infty}^{\infty} \sum_{n=0}^{\infty} \sum_{s=0}^{k-1} \sum_{f_i^y \in \Pi_{0,0} - \omega_{x_1}} A_{s,p,n}^{f_i^y}(\omega),$$

where

$$(57) \quad \begin{aligned} A_{s,p,n}^{f_i^y}(\omega) &= -(\lfloor n/\ell \rfloor + 1)F_{f_i^y,y}(\omega + s\omega_2 + n\omega_3 + p\omega_1) \\ &\quad -(\lfloor n/\ell \rfloor + 1)F_{f_i^y,y}(-\omega + 2\omega_{y_2} + s\omega_2 + n\omega_3 + p\omega_1) \\ &\quad + 2(\lfloor n/\ell \rfloor + 1)F_{f_i^y,y}(\omega_{y_2} + s\omega_2 + n\omega_3 + p\omega_1). \end{aligned}$$

The series (56) of terms $(\sum_{f_i^y \in \Pi_{0,0}-\omega_{x_1}} A_{s,p,n}^{f_i^y}(\omega))$ is absolutely convergent.

Let the f_i^x 's be the poles of the function f_x , and let $F_{f_i^x,x}(\omega)$ be the principal parts of this function at them. We have

$$(58) \quad r_x(\omega) - r_x(\omega_{x_2}) = \sum_{p=-\infty}^{\infty} \sum_{n=0}^{\infty} \sum_{s=0}^{k-1} \sum_{f_i^y \in \Pi_{0,0}+\omega_{y_1}} B_{s,p,n}^{f_i^x}(\omega),$$

where

$$(59) \quad \begin{aligned} B_{s,p,n}^{f_i^x}(\omega) &= -(\lfloor n/\ell \rfloor + 1)F_{f_i^x,x}(\omega - s\omega_2 - n\omega_3 - p\omega_1) \\ &\quad -(\lfloor n/\ell \rfloor + 1)F_{f_i^x,x}(-\omega + 2\omega_{x_2} - s\omega_2 - n\omega_3 - p\omega_1) \\ &\quad + 2(\lfloor n/\ell \rfloor + 1)F_{f_i^x,x}(\omega_{x_2} - s\omega_2 - n\omega_3 - p\omega_1). \end{aligned}$$

The series (58) of terms $(\sum_{f_i^y \in \Pi_{0,0}+\omega_{y_1}} B_{s,p,n}^{f_i^x}(\omega))$ is absolutely convergent.

Let $\omega_0^x \in \Delta_x$ be such that $x(\omega_0^x) = 0$ and let $\omega_0^y \in \Delta_y$ be such that $y(\omega_0^y) = 0$. Then $r_x(\omega)$ and $r_y(\omega)$ can be found from (56) and (58) as follows:

$$(60) \quad r_y(\omega) = (r_x(\omega_x^0) - r_x(\omega_{x_2})) - (r_x(\omega) - r_x(\omega_{x_2})) + x(\omega)y(\omega),$$

$$(61) \quad r_x(\omega) = (r_y(\omega_y^0) - r_y(\omega_{y_2})) - (r_y(\omega) - r_y(\omega_{y_2})) + x(\omega)y(\omega).$$

Proof. Denote by

$$F_{f_i^y,y}(\omega) = \sum_{n=1}^t \frac{r_{i,n}}{(\omega - f_i^y)^n}$$

the principal part (see Definition 3) of f_y at the pole f_i^y . Since f_y is an elliptic function with periods ω_1, ω_2 , it follows that $\sum_{i=1}^6 r_{i,1} = 0$ (it is a classical property of elliptic functions that the sum of the residues in a fundamental parallelogram is zero, see (P10), Lemma 25 in Appendix B, where we have gathered all needed properties of elliptic functions). This fact implies the convergence of the series

$$\sum_{p=-\infty}^{\infty} \sum_{n=0}^{\infty} \sum_{s=0}^{k-1} \left| \sum_{f_i^y \in \Pi_{0,0}-\omega_{x_1}} A_{s,p,n}^{f_i^y}(\omega) \right|.$$

We use the construction of the parallelograms P_m 's in Section 3 bounded by $V_m^+, V_m^-, H_{p_m}^*$ and $H_{p_m}^{**}$. They are all centered in ω_{y_2} , so that $d \in P_m$ is equivalent to $\hat{\eta}d \in P_m$. Since $r_y(\omega) = r_y(\hat{\eta}\omega)$, then for any pole d of r_y , $R_{\hat{\eta}d,y}(\omega) = R_{d,y}(\hat{\eta}\omega) = R_{d,y}(-\omega + 2\omega_{y_2})$. For any $d \in P_m$ with $\Re d < \Re \omega_{y_2} = \Re \omega_{y_1}$, there corresponds $\hat{\eta}d \in P_m$ with $\Re \hat{\eta}d > \Re \omega_{y_1}$ and vice versa. Points with real part equal to $\Re \omega_{y_1}$ belong to Δ_y and therefore cannot be

poles of r_y . Then by Theorem 4,

$$\begin{aligned}
r_y(\omega) - r_y(\omega_{y_2}) &= \lim_{m \rightarrow \infty} \sum_{d_i \in P_m \text{ poles of } r_y} R_{d_i, y}(\omega) - R_{d_i, y}(\omega_{y_2}) \\
(62) \qquad &= \lim_{m \rightarrow \infty} \sum_{\substack{d_i \in P_m \text{ poles of } r_y \\ \Re d_i < \Re \omega_{y_1}}} [R_{d_i, y}(\omega) + R_{d_i, y}(-\omega + 2\omega_{y_2}) - 2R_{d_i, y}(\omega_{y_2})].
\end{aligned}$$

By Theorem 6 (i), for any d pole of r_y with $\Re d < \Re \omega_{y_1}$, we have $\mathcal{N}_d^- \neq \emptyset$. Then for any $\omega \in \mathbf{C}$,

$$(63) \qquad \sum_{\substack{d_i \in P_m \text{ poles of } r_y \\ \Re d_i < \Re \omega_{y_1}}} R_{d_i, y}(\omega) = \sum_{\substack{d_i \in P_m \\ \Re d_i < \Re \omega_{y_1} \text{ and } \mathcal{N}_d^- \neq \emptyset}} R_{d_i, y}(\omega),$$

where the sum in the RHS above is taken over all points of P_m (not only the poles) with $\Re d < \Re \omega_{y_1}$ and $\mathcal{N}_d^- \neq \emptyset$. For any d with $\Re d < \Re \omega_{y_1}$ and $\mathcal{N}_d^- \neq \emptyset$, there exist $n \geq 0$ and f_i^y pole of f_y with $\Re \omega_{x_1} - k\omega_2 < \Re f_i^y < \Re \omega_{x_1}$ such that $d + n\omega_3 = f_i^y$. If in addition $d \in P_m$, then $\Re d \geq \Re \omega^- - m\omega_3$, so that $n \leq m - \Re(\omega^- - f_i^y)/\omega_3$ and also $\Im(\omega^* - p_m\omega_1) \leq \Im d = \Im f_i^y \leq \Im(\omega^* + p_m\omega_1)$. On the other hand, for any f_i^y with real and imaginary parts bounded as above and for any non-negative integer $n \leq m - \Re(\omega^- - f_i^y)/\omega_3$, the point $f_i^y - n\omega_3$ belongs to P_m and it is such that $\mathcal{N}_d^- \neq \emptyset$. Therefore by (52) of Theorem 6, the quantity in the RHS of (63) equals

$$(64) \qquad \sum_{\substack{f_i^y: \Im(\omega^* - p_m\omega_1) \leq \Im f_i^y \leq \Im(\omega^* + p_m\omega_1) \\ \Re \omega_{x_1} - k\omega_2 < \Re f_i^y < \Re \omega_{x_1}}} \sum_{n=0}^{\lfloor m - \Re(\omega^- - f_i^y)/\omega_3 \rfloor} -(\lfloor n/\ell \rfloor + 1) F_{f_i^y, y}(\omega + n\omega_3).$$

For any f_i^y in the sum above, there exist unique $s \in \{0, 1, \dots, k-1\}$ and $p \in \mathbf{Z}$ such that $f_i^y = \bar{f}_i^y - s\omega_2 - p\omega_1$, where $\bar{f}_i^y \in \Pi_{0,0} - \omega_{x_1}$ is a pole of f_y . Furthermore, the bounds for $\Im f_i^y$ are equivalent to $-p_m + \Im(\bar{f}_i^y - \omega^*)/|\omega_1| \leq p \leq p_m + \Im(\bar{f}_i^y - \omega^*)/|\omega_1|$. We deduce that for any $\omega \in \mathbf{C}$, the sum in (64) equals

$$(65) \qquad \sum_{f_i^y \in \Pi_{0,0} - \omega_{x_1}} \sum_{s=0}^{k-1} \sum_{p=\lceil -p_m + \Im(\omega^* - f_i^y)/|\omega_1| \rceil}^{\lfloor p_m + \Im(\omega^* - f_i^y)/|\omega_1| \rfloor} \sum_{n=0}^{\lfloor m - \Re(\omega^- - f_i^y)/\omega_3 \rfloor} -(\lfloor n/\ell \rfloor + 1) F_{f_i^y, y}(\omega + n\omega_3 + s\omega_2 + p\omega_1).$$

By (57), (62), (63) and (65), we obtain

$$(66) \qquad r_y(\omega) - r_y(\omega_{y_2}) = \lim_{m \rightarrow \infty} \sum_{f_i^y \in \Pi_{0,0} - \omega_{x_1}} \sum_{s=0}^{k-1} \sum_{p=\lceil -p_m + \Im(\omega^* - f_i^y)/|\omega_1| \rceil}^{\lfloor p_m + \Im(\omega^* - f_i^y)/|\omega_1| \rfloor} \sum_{n=0}^{\lfloor m - \Re(\omega^- - f_i^y)/\omega_3 \rfloor} A_{s,n,p}^{f_i^y}(\omega).$$

To conclude the proof of (56), it remains to show that the difference

$$(67) \quad \sum_{f_i^y \in \Pi_{0,0} - \omega_{x_1}} \sum_{s=0}^{k-1} \sum_{p=\lceil -p_m + \Im(\omega^* - f_i^y)/|\omega_1| \rceil}^{\lfloor p_m + \Im(\omega^* - f_i^y)/|\omega_1| \rfloor} \sum_{n=0}^{\lfloor m - \Re(\omega^- - f_i^y)/\omega_3 \rfloor} A_{s,n,p}^{f_i^y}(\omega) \\ - \sum_{p=-p_m}^{p_m} \sum_{n=0}^m \sum_{s=0}^{k-1} \sum_{f_i^y \in \Pi_{0,0} - \omega_{x_1}} A_{s,n,p}^{f_i^y}(\omega)$$

converges to zero as $m \rightarrow \infty$. To that purpose, it is sufficient to prove that for any fixed $f_i^y \in \Pi_{0,0} - \omega_{x_1}$ and $s \in \{0, 1, \dots, k-1\}$, the limit of

$$(68) \quad \sum_{p=\lceil -p_m + \Im(\omega^* - f_i^y)/|\omega_1| \rceil}^{\lfloor p_m + \Im(\omega^* - f_i^y)/|\omega_1| \rfloor} \sum_{n=0}^{\lfloor m - \Re(\omega^- - f_i^y)/\omega_3 \rfloor} A_{s,n,p}^{f_i^y}(\omega) - \sum_{p=-p_m}^{p_m} \sum_{n=0}^m A_{s,n,p}^{f_i^y}(\omega)$$

is 0 as $m \rightarrow \infty$. Using the definition (57), it is easy to show that for any f_i^y and any $s = 0, 1, \dots, k-1$

$$(69) \quad \sum_{p=-\infty}^{\infty} |A_{s,n,p}^{f_i^y}(\omega)| = O(1/n), \quad n \rightarrow \infty, \quad \sum_{n=0}^{\infty} |A_{s,n,p}^{f_i^y}(\omega)| = O(1/p), \quad p \rightarrow \infty,$$

from where (68) follows. Hence (56) is proved.

The proof of (58) for r_x is completely analogous and is omitted. In order to show (60) and (61), we notice that $Q(0,0)K(0,0) = r_x(\omega_0^x) = r_y(\omega_0^y)$. This way, (60) and (61) are immediate corollaries of (25). \square

6. MISCELLANEOUS

6.1. More on the poles of f_y and principal parts at them. There are generally six poles of the function $f_y(\omega)$ in any parallelogram of periods ω_1, ω_2 on \mathbf{C} , see Section 3. We call f_i^y , $i \in \{1, \dots, 6\}$, the poles in $\Pi_{0,0} - \omega_{x_2}$. They can coincide for some models (see [19, Section 7]). We denote by

$$(70) \quad f_1^y = (x^*, \infty), \quad f_2^y = (x^{**}, \infty)$$

those where $y = \infty$, by

$$(71) \quad f_3^y = (\infty, y'), \quad f_4^y = (\infty, y'')$$

those where $x = \infty$, and by

$$(72) \quad f_5^y = (x^*, y^*) = \widehat{\xi} f_1^y - \omega_2, \quad f_6^y = (x^{**}, y^{**}) = \widehat{\xi} f_2^y - \omega_2$$

the remaining ones. The location of these poles heavily depends on the signs of the branch points x_4 and y_4 , see [19, Cases I, II et III in Section 7] or the proof of Lemma 5 above. The sign of x_4 and y_4 can be expressed in terms of the geometry of the step set \mathcal{S} , see [19, Remark 24].

In this section we are interested in the computation of the functions $F_{f_i^y, y}(\omega)$ determining our main result, that is the series (56) of Theorem 7. We first (Proposition 9 and its proof) compute the degree of $F_{f_i^y, y}(\omega)$ (i.e., the order of the pole f_i^y of the function $f_y(\omega)$). Then

(below Proposition 9) we explain how to concretely compute the functions $F_{f_i^y, y}(\omega)$. First of all, we need the following technical lemma:

Lemma 8. *On the fundamental rectangle $\omega_1[0, 1[+ \omega_2[0, 1[$, x has one double pole or two simple poles. It has one double pole if and only if*

$$(73) \quad \delta_{1,0} = \delta_{1,1} = 0 \quad \text{or} \quad \delta_{1,0} = \delta_{1,-1} = 0,$$

and two simple poles if and only if the two conditions in (73) are not satisfied.

Proof. The uniformization formulæ (12) show that on $\omega_1[0, 1[+ \omega_2[0, 1[$, x has one double pole (if and only if $x_4 = \infty$) or two simple poles (if and only if $x_4 \neq \infty$). Moreover, it is shown in [9, Lemma 2.3.8] that $x_4 = \infty$ if and only if $\delta_{1,0}^2 - 4\delta_{1,1}\delta_{1,-1} = 0$. Lemma 8 follows. \square

Proposition 9. *The order of the poles of f_y is 1, 2 or 3.*

Proof. We first notice that f_y defined in (36) can alternatively be written as

$$f_y(\omega) = \frac{x'(\omega)x(\omega)}{2a(x(\omega))},$$

where

$$a(x)/z = \delta_{1,1}x^2 + \delta_{0,1}x + \delta_{-1,1}.$$

To be complete, we have the following cases to consider (for the non-singular walks):

1. $a(x)/z = x^2$;
2. $a(x)/z = x$;
3. $a(x)/z = 1$;
4. $a(x)/z = x^2 + x$;
5. $a(x)/z = x^2 + 1$;
6. $a(x)/z = x + 1$;
7. $a(x)/z = x^2 + x + 1$.

In the cases 1 and 4, the poles are simple. Indeed, we have, respectively,

$$f_y(\omega) = \frac{1}{2z} \frac{x'(\omega)}{x(\omega)}, \quad f_y(\omega) = \frac{1}{2z} \frac{(x(\omega) + 1)'}{x(\omega) + 1}.$$

Case 1 in particular covers Kreweras' and Gessel's models.

In the cases 5 and 7, we have (with $\delta = 0$ and 1, respectively)

$$f_y(\omega) = \frac{1}{2z} \frac{x'(\omega)}{x(\omega) + \delta + 1/x(\omega)}.$$

If ω_0 is a simple pole of $x(\omega)$, then it is obviously also a simple pole of $f_y(\omega)$. If ω_0 is such that $x(\omega_0) + \delta + 1/x(\omega_0) = 0$, then it can be a zero of order one or two (remember that $x(\omega)$ is an elliptic function of order two). In both cases, an expansion in the expression of $f_y(\omega)$ above shows that f_y has a pole of order one at ω_0 .

In Case 2, $f_y(\omega) = x'(\omega)/(2z)$. Therefore, if x has one simple (resp. double) pole, then f_y has a double (resp. triple) pole. Both situations can happen: the simple walk (Figure 2) is such that the pole is simple, and the walk on the left on Figure 6 has a double pole.




FIGURE 6. Examples in the proof of Proposition 9

In Case 3, one has

$$f_y(\omega) = \frac{x'(\omega)x(\omega)}{2z}.$$

Therefore, if x has one simple (resp. double) pole, then f_y has a triple (resp. quintuple) pole. By definition of Case 3, one has $\delta_{1,1} = \delta_{0,1} = 0$. Further, if the pole of x is double, one of the two conditions in (73) must hold. This would imply that the walk is singular; this is a contradiction, since we do not consider these models here.

It remains to consider Case 6. For similar reasons as for Cases 5 and 7, if $x(\omega_0) + 1 = 0$ then ω_0 is a pole of order one of f_y . If x has a simple (resp. double) pole, then obviously f_y has a double (resp. triple) pole. The two situations can happen, as the second and third examples on Figure 6 illustrate. \square

We now explain in which extent it is possible to find explicit expressions for the functions $F_{f_i^y, y}(\omega)$. In fact, this heavily depends on the model under consideration. For instance, in the Cases 1 and 4, we do not need to simplify the uniformization (12): if ω_0 is a pole (resp. zero) of order p of x , then the residue of f_y at ω_0 is $-p/(2z)$ (resp. $p/(2z)$). For the other cases, one needs more information on (12). For example, in Case 2, we clearly need to know the principal parts of x at its poles in order to deduce the principal parts of f_y at its poles. Unfortunately, there is no general formula, and this should be done model by model.

6.2. Algebraicity of the GFs. Theorem 7 determines the GFs $Q(x, 0)$ and $Q(0, y)$ in terms of the principal parts of functions f_x and f_y at their poles in a parallelogram of periods. We would like now to identify the algebraic nature of these functions (question (ii) stated at the very beginning of this paper) in the same terms.

Given a pole $f_i^y \in \Pi_{0,0} - \omega_{x_1}$, consider the sum

$$\mathcal{F}_{f_i^y}^y(\omega) = F_{f_i^y, y}(\omega) + F_{f_i^y + \omega_3, y}(\omega + \omega_3) + \cdots + F_{f_i^y + (\ell-1)\omega_3, y}(\omega + (\ell-1)\omega_3).$$

(We recall that above, for $t \in \{0, \dots, \ell-1\}$, $F_{f_i^y + t\omega_3, y}$ is the principal part of f_y at $f_i^y + t\omega_3$. In particular, for t such that $f_i^y + t\omega_3$ is not a pole of f_y , the corresponding term $F_{f_i^y + t\omega_3, y}$ is zero.)

Proposition 10. *Under the assumption (33), the functions $x \mapsto Q(x, 0)$ and $y \mapsto Q(0, y)$ are algebraic functions with $2k$ branches if and only if*

$$(74) \quad \mathcal{F}_{f_i^y}^y(\omega) = 0, \quad \forall \omega \in \mathbf{C}, \quad \forall i \in \{1, \dots, 4\},$$

with the numbering of the poles (70)–(72) as above.

Proof. Assume first (74). Under (74), for all $i \in \{1, \dots, 4\}$, there exists $j \in \{1, \dots, 6\}$ (with an appropriate $F_{f_j^y}(\omega)$) such that $f_i^y + n\omega_3 = f_j^y + s\omega_2$ for some $n \in \{1, \dots, \ell - 1\}$ and $s \in \{0, \dots, k - 1\}$. Since

$$f_5^y = \widehat{\xi}\widehat{\eta}f_1^y = f_1^y - \omega_3, \quad f_6^y = \widehat{\xi}\widehat{\eta}f_2^y = f_2^y - \omega_3,$$

then under (74), the sum $\mathcal{F}_{f_i^y}^y(\omega)$ is also zero for $i \in \{5, 6\}$ and for all $\omega \in \mathbf{C}$. Therefore, under (74), the elliptic function (which in some sense is a half-orbit sum, see [6])

$$\mathcal{O}(\omega) = \sum_{n=0}^{\ell-1} f_y(\omega + n\omega_3)$$

has no poles in \mathbf{C} and thus, by a well-known property of elliptic functions (see Property (P2)) it is a constant: there exists $C \in \mathbf{C}$ such that $\mathcal{O}(\omega) = C$, for all $\omega \in \mathbf{C}$. We now prove that $C = 0$. One has in particular $\mathcal{O}(\omega_{y_2} + \omega_3) = \mathcal{O}(\omega_{y_2} - \ell\omega_3) = C$, and thus, by Equations (54) and (55),

$$\begin{aligned} r_y(\omega_{y_2} - \ell\omega_3) &= r_y(\omega_{y_2}) - \mathcal{O}(\omega_{y_2} - \ell\omega_3) = r_y(\omega_{y_2}) - C, \\ r_y(\omega_{y_2} + \ell\omega_3) &= r_y(\omega_{y_2}) + \mathcal{O}(\omega_{y_2} + \omega_3) = r_y(\omega_{y_2}) + C. \end{aligned}$$

Moreover, it follows from (26) that $r_y(\omega_{y_2} - \ell\omega_3) = r_y(\omega_{y_2} + \ell\omega_3)$, from where $C = 0$. Hence, under (74), we conclude that $\mathcal{O}(\omega) = 0$ for all $\omega \in \mathbf{C}$. It follows again from (54) and (55) that the function $r_y(\omega)$ is $\ell\omega_3 = k\omega_2$ -periodic, and then by (25), so is $r_x(\omega)$. By Property (P6), r_x and r_y are algebraic in $x(\omega)$ and $y(\omega)$ respectively. On the other hand, if (74) is not satisfied, then $\mathcal{O}(\omega)$ is not identically zero. It has been shown in [19, Section 6] that under this last condition, the functions $x \mapsto Q(x, 0)$ and $y \mapsto Q(0, y)$ are holonomic, but not algebraic. \square

It is worth noting that Condition (74), which is stated for f_y , is equivalent to its analogue for the function f_x . Further, it is immediate from Proposition 10 that under (74), the residues at the poles of r_x and r_y are bounded on \mathbf{C} . Moreover, under (74), the terms

$$A_{s,n,p}^{f_i^y}(\omega)$$

defined in (57) can be permuted and regrouped in their sum over

$$p \in \mathbf{Z}, \quad n \geq 0, \quad s \in \{0, \dots, k - 1\}, \quad f_i^y \in \Pi_{0,0} - \omega_{x_1}$$

in such a way that the coefficients in front of $F_{f_i^y, y}$ are bounded and this infinite sum stays equal to the one of (56). This condition can never hold true for models where $x_4 > 0$ or $y_4 > 0$ because of the position of the poles of f_y , see [19]. We do not know either for models with infinite group and such that $x_4 \in]-\infty, 0] \cup \{\infty\}$ and $y_4 \in]-\infty, 0] \cup \{\infty\}$, there exist values of $z \in \mathcal{H}$ such that (74) is satisfied. But among the 23 models with finite group, it is satisfied exactly for 4 models, and in particular for Kreweras' and Gessel's walks.

6.3. Conclusion and open problems. This article closes our study of the walks with small steps in the quarter plane. The study began with the finite group case [2, 5, 6, 10, 16, 18], then the interest moved to the infinite group case [7, 19, 23, 24, 25]. The present article presents an approach that works both for the two cases, and which is computationally efficient (see the example of Kreweras' model below, see also the forthcoming article [3] on Gessel's walks). Below we mention some open problems:

- (i) In this article, we find expressions as infinite series for $Q(x, 0; z)$ and $Q(0, y; z)$ for a dense set of values of z (namely, $z \in \mathcal{H}$). By continuity this must provide an expression for the GFs for all values of $z \in]0, 1/|\mathcal{S}|[$. It is an open problem to determine whether there exists an expression as an infinite series for the GFs when $z \in]0, 1/|\mathcal{S}| \setminus \mathcal{H}$.
- (ii) We know from [19] that (as functions of the variables x and y) the functions $Q(x, 0; z)$ and $Q(0, y; z)$ are non-holonomic for all $z \in]0, 1/|\mathcal{S}| \setminus \mathcal{H}$. It is an open problem to reprove this result, starting from the results (in particular Theorems 6 and 7) of this paper. In other words, is it possible to prove non-holonomy from holonomy?

Part 2. Example: GFs for Kreweras' model

In Sections 7, 8 and 9, we illustrate our computational results on the example of Kreweras' model. The step set \mathcal{S} of this model can be seen on Figure 2. We decompose this illustration into three steps. The first one (Section 7) consists in deriving from Theorems 6 and 7 an expression for the function $r_y(\omega) = zy(\omega)Q(0, y(\omega))$ in terms of Weierstrass ζ -functions, see Proposition 15. This expression is new up to our knowledge. The second step (Section 8) is to show how obtaining from this expression an algebraic function of the series of the excursions $Q(0, 0)$. Finally (Section 9), we explain how finding an algebraic expression for $Q(y, 0) = Q(0, y)$.

Proposition 11 ([2, 5, 6, 9, 13, 14, 17]). *Consider Kreweras' walks. Let W be the only power series in z satisfying $W = z(2 + W^3)$. Then*

$$(75) \quad Q(0, 0) = \frac{1}{2z}(W - W^4/4).$$

Further, one has

$$(76) \quad Q(y, 0) = Q(0, y) = \frac{1}{zy} \left(\frac{1}{2z} - \frac{1}{y} - \left(\frac{1}{W} - \frac{1}{y} \right) \sqrt{1 - yW^2} \right).$$

(Of course, (75) is an immediate consequence of (76), but in order to illustrate our approach, we shall first prove (75), and then (76).) There already exist many proofs of Proposition 11. The first one was proposed by Kreweras himself [17]. See also Flatto and Hahn [13] and Flatto [14], where methods using elliptic functions were used (see also [9, Section 4.6], where similar ideas are used). More recently, proofs were done in [5, 6], using an extension of the well-known kernel method. Finally, in [2], there is a proof based on a guessing-proving approach (which is also applied to Gessel's walks).

7. EXPRESSION OF THE GF COUNTING KREWERAS' WALKS IN TERMS OF
WEIERSTRASS ELLIPTIC FUNCTIONS

7.1. Uniformization in Kreweras' case. The function f_y defined in (36)—which appears in the meromorphic continuation procedure (24)—will be particularly important. To compute it, we need to simplify the coordinates of the uniformization $x(\omega), y(\omega)$ in (12). In the case of Kreweras, $x_4 = y_4 = \infty$, and one has

$$(77) \quad \begin{cases} x(\omega) &= \frac{\wp(\omega) - d''(0)/6}{d'''(0)}, \\ y(\omega) &= -\frac{b(x(\omega)) + \wp'(\omega)/(2d_3)}{2a(x(\omega))}, \end{cases}$$

with $d''(0)/2 = 1$, $d'''(0)/6 = -4z^2$, $b(x) = -x + z$ and $a(x) = zx^2$, see (9). We recall (Section 2) that in (77), $\wp(\omega)$ denotes the Weierstrass elliptic function with periods ω_1, ω_2 , see (10) and (11). We shall sometimes write, instead, $\wp(\omega; \omega_1, \omega_2)$. Finally, we recall that with ω_3 defined as in (13), in the case of Kreweras one has $\omega_3/\omega_2 = 2/3$ (this in particular implies that the group (4) has order 6). To apply our main results (Theorems 6 and 7), we need to know where the poles of f_y are located.

Lemma 12. *In the fundamental rectangle $\omega_1[0, 1[+ \omega_2[0, 1[$, the function f_y has poles at 0 , $\omega_2/3$ and $2\omega_2/3$. These poles are simple, with residues equal to $-1/z$, $1/(2z)$ and $1/(2z)$, respectively.*

Proof. Observe that with (36), (77), and with the equalities $\wp(-\omega) = \wp(\omega)$, $\wp'(-\omega) = -\wp'(\omega)$, we can write

$$(78) \quad f_y(\omega) = \frac{\wp'(\omega)}{2z(\wp(\omega) - 1/3)}.$$

The latter has a simple pole at 0 , with residue $-1/z$ (remember that in the neighborhood of 0 , $\wp(\omega) = 1/\omega^2 + O(\omega^2)$). It has also poles at points where $\wp(\omega) - 1/3 = 0$. For the model under consideration, one has $\wp(\omega_2/3) = 1/3$, see [25, Page 773]. Accordingly, $\wp(2\omega_2/3) = 1/3$ as well, and one concludes that $\omega_2/3$ and $2\omega_2/3$ are the two remaining poles of f_y on the fundamental rectangle. Making an expansion in (78), we deduce that the residues at $\omega_2/3$ and $2\omega_2/3$ are equal to $1/(2z)$. \square

Remark 13. The function f_y has a priori 6 poles (see Section 3). Lemma 12 shows that it may have strictly less than 6 poles. \square

The Weierstrass \wp -function with periods ω_1, ω_2 can alternatively be characterized by its periods ω_1, ω_2 (see (11)), or by its invariants g_2, g_3 , by the formula

$$(79) \quad \wp'(\omega)^2 = 4\wp(\omega)^3 - g_2\wp(\omega) - g_3.$$

Lemma 14. *For Kreweras' model, we have*

$$(80) \quad g_2 = 4/3 - 32z^3, \quad g_3 = -8/27 + 32z^3/3 - 64z^6.$$

Proof. The construction of the uniformization (see [9] or [25, Page 770], and (12) in this paper) is the following: with (77) one has $\wp(\omega) = g(x(\omega))$, where $g(x) = 1/3 - 4z^2x$. Then

$$g_2 = -4[g(x_1)g(x_2) + g(x_1)g(x_3) + g(x_2)g(x_3)], \quad g_3 = 4g(x_1)g(x_2)g(x_3).$$

One concludes by computing the branch points above (which are the roots of the polynomial (9)) in terms of z . \square

7.2. Expression of the GFs. Applying Theorem 7 to Kreweras' walks, we first obtain an expression of $r_y(\omega)$ in terms of special functions. In what follows, we denote by $\zeta_{1,2}$ and $\wp_{1,2}$ the Weierstrass ζ - and \wp -functions with periods $\omega_1, 2\omega_2$.

Proposition 15. *One has*

$$(81) \quad r_y(\omega) = c + \frac{1}{2z}\zeta_{1,2}(\omega + 2\omega_2/3) - \frac{1}{z}\zeta_{1,2}(\omega + \omega_2/3) + \frac{1}{z}\zeta_{1,2}(\omega) - \frac{1}{2z}\zeta_{1,2}(\omega - \omega_2/3), \quad \forall \omega \in \mathbf{C}.$$

Theorem 7 also gives the expression of the constant c in (81), but we shall compute it in Section 9 only.

Proof. There are several ways to deduce Proposition 15. A first one is to use Theorem 7, which gives an expression for r_y : one should group some terms of (56) together in order to deduce ζ -functions and to exploit (69).

The second one is to apply Theorem 6, which only concerns the poles of the GF (and not its expression), combined with the ellipticity of r_y proper to Kreweras' model. Namely, it is elementary to check Condition (74) of Proposition 10 using our analysis of poles of f_y above. Then by Proposition 10 (see also [19, Section 6]) r_y is algebraic with $2k = 4$ branches and hence elliptic with periods $\omega_1, 2\omega_2$. This remark allows us to study the poles of r_y only in the parallelogram $\omega_2[-3/2, 1/2[+ \omega_1[-1/2, 1/2[$. According to Theorem 6, the poles of r_y must satisfy $\mathcal{N}_d^- \neq \emptyset$, where (we recall that $\Re\omega_{x_1} = \omega_2/2$)

$$\mathcal{N}_d^- = \{n \geq 0 : d + n\omega_3 \text{ is a pole of } f_y \text{ with } \omega_2/2 - k\omega_2 < \Re d + n\omega_3 < \omega_2/2\}.$$

For Kreweras' model, one has $\omega_2/\omega_3 = 3/2$, so that $k = 2$, and the only poles d of f_y with $-3\omega_2/2 < \Re d + n\omega_3 < \omega_2/2$ and $|\Im d| \leq |\omega_1/2|$ are the points of the set

$$P = \{-4\omega_2/3, -\omega_2, -2\omega_2/3, -\omega_2/3, 0, \omega_2/3\}.$$

We thus have $\mathcal{N}_d^- = \{n \geq 0 : d + n\omega_3 \in P\}$, and it is obvious that the points of our parallelogram $\omega_2[-3/2, 1/2[+ \omega_1[-1/2, 1/2[$ such that $\mathcal{N}_d^- \neq \emptyset$ are among the points of P . Let us study closer each of them. We start with $d \in \{0, \omega_2/3\}$. We have $\mathcal{N}_d^- = \{0\}$, and according to Theorem 6 and Lemma 12, we find the following principal parts of r_y :

$$R_{\omega_2/3,y}(\omega) = -F_{\omega_2/3,y}(\omega) = \frac{-1/(2z)}{\omega - \omega_2/3},$$

$$R_{0,y}(\omega) = -F_{0,y}(\omega) = \frac{1/z}{\omega}.$$

Consider now the cases $d \in \{-2\omega_2/3, -\omega_2/3\}$, for which one has $\mathcal{N}_d^- = \{0, 1\}$. Then

$$\begin{aligned} R_{-\omega_2/3,y}(\omega) &= -F_{-\omega_2/3,y}(\omega) - F_{\omega_2/3,y}(\omega + 2\omega_2/3) = \frac{-1/z}{\omega + \omega_2/3}, \\ R_{-2\omega_2/3,y}(\omega) &= -F_{-2\omega_2/3,y}(\omega) - F_{0,y}(\omega + 2\omega_2/3) = \frac{1/(2z)}{\omega + 2\omega_2/3}. \end{aligned}$$

For the two remaining cases $d \in \{-4\omega_2/3, -\omega_2\}$, one has $\mathcal{N}_d^- = \{0, 1, 2\}$, and similar computations as above show that

$$R_{-\omega_2,y}(\omega) = R_{-4\omega_2/3,y}(\omega) = 0.$$

In other words, the points $-4\omega_2/3$ and $-\omega_2$ are removable singularities of the GF r_y .

To conclude, we notice that the elliptic function r_y with periods $\omega_1, 2\omega_2$, has in the fundamental parallelogram $\omega_2[-3/2, 1/2[+ \omega_1[-1/2, 1/2[$ four poles, with principal parts given above. It is immediate from the theory of elliptic functions (in particular Property (P5)) that Equation (81) holds. The proof is completed. \square

Remark 16. Starting from (81), we can recover (24). Introduce

$$k(\omega) = \frac{1}{2z}\zeta_{1,2}(\omega + \omega_2/3) - \frac{1}{z}\zeta_{1,2}(\omega) + \frac{1}{2z}\zeta_{1,2}(\omega - \omega_2/3).$$

An easy computation starting from (81) yields

$$r_y(\omega + \omega_3) - r_y(\omega) = k(\omega) + k(\omega + \omega_2).$$

Since $k(\omega)$ is $\omega_1, 2\omega_2$ elliptic (see Property (P4)), we deduce that $k(\omega) + k(\omega + \omega_2)$ is ω_1, ω_2 elliptic, hence $r_y(\omega + \omega_3) - r_y(\omega)$ is ω_1, ω_2 elliptic. We first show that $r_y(\omega + \omega_3) - r_y(\omega)$ has the same poles as $f_y(\omega)$ in $\omega_1[0, 1[+ \omega_2[0, 1[$. Since in $\omega_1[0, 1[+ \omega_2[0, 1[$, $\zeta_{1,2}$ has only one pole (which is of order 1, at 0 and with residue 1, see Property (P1)), we obtain that in $\omega_1[0, 1[+ \omega_2[0, 1[$, $r_y(\omega + \omega_3) - r_y(\omega)$ has only simple poles, at 0, $\omega_2/3$ and $2\omega_2/3$, with respective residues $-1/z$, $1/(2z)$ and $1/(2z)$. The same holds for f_y (Lemma 12). Thus, there exists a constant c such that

$$(82) \quad r_y(\omega + \omega_3) - r_y(\omega) = c + f_y(\omega), \quad \forall \omega \in \mathbf{C}.$$

Evaluating (82) at $\omega = \omega_2/2$ and using that $f_y(\omega_2/2) = 0$ (see Equation (78)), we obtain that $c = r_y(\omega_2/2 + \omega_3) - r_y(\omega_2/2) = k(\omega_2/2) + k(3\omega_2/2)$. Using the fact that $\zeta_{1,2}$ is odd and $2\omega_2$ -periodic, we obtain that $c = 0$, which yields (24). \square

8. FINDING AN ALGEBRAIC EXPRESSION FOR THE GF OF THE EXCURSIONS OF KREWERAS' WALKS

In this section we prove the first part of Proposition 11 (Equation (75)). Surprisingly, we shall do it without finding the value of the additive constant c in (81).

8.1. **Beginning of the proof.** One has from (25) that

$$r_x(\omega) + r_y(\omega) - K(0,0)Q(0,0) - x(\omega)y(\omega) = 0, \quad \forall \omega \in \mathbf{C}.$$

Setting $\omega_0^y = 2\omega_2/3$, one has $y(\omega_0^y) = 0$; further, it turns out that $x(\omega_0^y) = 0$ (see (12) and (77)). Since $K(0,0) = 0$ for Kreweras' model, one deduces that

$$\frac{r_x(\omega)}{x(\omega)} = y(\omega) - \frac{r_y(\omega) - r_y(\omega_0^y)}{x(\omega)} = y(\omega) - \frac{r_y(\omega) - r_y(\omega_0^y)}{\omega - \omega_0^y} \frac{\omega - \omega_0^y}{x(\omega) - x(\omega_0^y)}.$$

Then, if $\omega \rightarrow \omega_0^y$, one finds

$$zQ(0,0) = -\frac{r'_y(\omega_0^y)}{x'(\omega_0^y)}.$$

By (77), $x'(\omega_0^y) = \wp'(2\omega_2/3)/(d'''(0)/6)$. Since $\wp(2\omega_2/3) = 1/3$, using (79) and (80) leads to $\wp'(2\omega_2/3) = \pm 8z^3$. But $\wp'(\omega)$ is positive on $]\omega_2/2, \omega_2[$, so that finally $\wp'(2\omega_2/3) = 8z^3$. Since $d'''(0)/6 = -4z^2$, we reach the conclusion that $x'(\omega_0^y) = -2z$. Finally, one finds

$$(83) \quad Q(0,0) = \frac{1}{4z^3} [\wp_{1,2}(\omega_2/3) - \wp_{1,2}(4\omega_2/3) + 2(\wp_{1,2}(\omega_2) - \wp_{1,2}(2\omega_2/3))].$$

The difficulty now consists in transforming the above expression in an algebraic function of z . So far, we have already computed the values of $\wp(\omega)$ for some particular ω (e.g., we saw that $\wp(\omega_2/3) = \wp(2\omega_2/3) = 1/3$, and that $\wp(0) = \wp(\omega_2) = \infty$). On the other hand, we never computed $\wp_{1,2}(\omega)$ for some given ω . To do so, the simplest thing is to express $\wp_{1,2}$ (with periods $\omega_1, 2\omega_2$) in terms of \wp (with periods ω_1, ω_2). This is the aim of the next section.

8.2. **Intermezzo: transformation theory of elliptic functions.** Let $\wp_{1,2}$ be the Weierstrass elliptic function with periods $\omega_1, 2\omega_2$. Denote its invariants by $g_2^{1,2}$ and $g_3^{1,2}$. We also define $e_2^{1,2} = \wp_{1,2}(\omega_2)$, $e_1^{1,2} = \wp_{1,2}(\omega_1/2)$ and $e_{1+2}^{1,2} = \wp_{1,2}(\omega_1/2 + \omega_2)$. Note that the latter are the three solutions of $4X^3 - g_2^{1,2}X - g_3^{1,2} = 0$. To compute the latter quantities, it is convenient to introduce r , \tilde{r} and \hat{r} as the roots of

$$(84) \quad 4X^3 - g_2X + g_3 = 0,$$

with g_2, g_3 defined as in Lemma 14.⁶ We enumerate them in the following way: only one solution of (84) is a power series in z (see [27, Proposition 6.1.8]), we call it r . It admits the expansion

$$r = 2/3 - 8z^3 - 48z^6 - 640z^9 + O(z^{12}).$$

Lemma 17. *One has*

$$\begin{aligned} e_2^{1,2} &= r/2, \\ g_2^{1,2} &= 15r^2/4 - g_2/4, \\ g_3^{1,2} &= 11g_3/32 - 7rg_2/32. \end{aligned}$$

⁶It is worthwhile comparing these roots with $\wp(\omega_2/2)$, $\wp((\omega_2 + \omega_1)/2)$ and $\wp(\omega_1/2)$, which are solutions to $4X^3 - g_2X - g_3 = 0$.

Proof. Using the properties (P3) and (P8), one can write, for any $\omega \in \mathbf{C}$,

$$(85) \quad \wp(\omega) = \wp_{1,2}(\omega) + \wp_{1,2}(\omega + \omega_2) - \wp_{1,2}(\omega_2) = -2\wp_{1,2}(\omega_2) + \left(\frac{\wp'_{1,2}(\omega)}{\wp_{1,2}(\omega) - \wp_{1,2}(\omega_2)} \right)^2.$$

We then make an expansion of the LHS and the RHS of the above equation in the neighborhood of 0; we obtain

$$\begin{aligned} \frac{1}{\omega^2} + \frac{g_2}{20}\omega^2 + \frac{g_3}{28}\omega^4 + O(\omega^6) \\ = \frac{1}{\omega^2} + \left(3(e_2^{1,2})^2 - \frac{g_2^{1,2}}{5} \right) \omega^2 + \left(4(e_2^{1,2})^3 - \frac{g_2^{1,2}e_2^{1,2}}{2} - \frac{3g_3^{1,2}}{14} \right) \omega^4 + O(\omega^6). \end{aligned}$$

Identifying the expansions above, we obtain two equations for the three unknowns $e_2^{1,2}$, $g_2^{1,2}$ and $g_3^{1,2}$ (remember that g_2 and g_3 are known from Lemma 14). We add a third equation by noticing that $e_2^{1,2}$ is a root of $4X^3 - g_2^{1,2}X - g_3^{1,2} = 0$. We then have a (non-linear) system of three equations with three unknowns. Some computations finally lead to the expressions of $e_2^{1,2}$, $g_2^{1,2}$ and $g_3^{1,2}$ given in Lemma 17. \square

Remark 18. Contrary to g_2 and g_3 , $g_2^{1,2}$ and $g_3^{1,2}$ are not rational functions of z . However, they are algebraic (and so is $e_2^{1,2}$). \square

Thanks to Lemma 17, one can consider that the function $\wp_{1,2}$ is completely known, as its periods and its invariants are expressed in the variable z in an explicit way. The next result proposes an expression of $\wp_{1,2}$ in terms of \wp .

Lemma 19. *One has*

$$2\wp_{1,2}(\omega) = \wp(\omega) + e_2^{1,2} \pm \sqrt{(\wp(\omega) - e_2^{1,2})^2 + g_2^{1,2} - 12(e_2^{1,2})^2}, \quad \forall \omega \in \mathbf{C},$$

where $e_2^{1,2}$ and $g_2^{1,2}$ are as in Lemma 17.⁷

Proof. We start from Equation (85), which can be rewritten as

$$(86) \quad \wp(\omega) = -2e_2^{1,2} + \frac{(\wp_{1,2}(\omega) - e_1^{1,2})(\wp_{1,2}(\omega) - e_{1+2}^{1,2})}{\wp_{1,2}(\omega) - e_2^{1,2}}, \quad \forall \omega \in \mathbf{C}.$$

Further, we have the equalities

$$(87) \quad e_1^{1,2} + e_{1+2}^{1,2} + e_2^{1,2} = 0, \quad e_1^{1,2}e_{1+2}^{1,2} + e_1^{1,2}e_2^{1,2} + e_{1+2}^{1,2}e_2^{1,2} = -\frac{g_2^{1,2}}{4}.$$

To conclude, we solve Equation (86) as an equation (of the second order) in $\wp_{1,2}(\omega)$. Making use of the identities (87), we obtain Lemma 19. \square

⁷The sign \pm above depends on which half-parallelogram ω is located.

8.3. End of the proof. Using Lemma 19 and the fact that $\wp(\omega_2/3) = \wp(2\omega_2/3) = 1/3$, we obtain that

$$\begin{aligned}\wp_{1,2}(\omega_2/3) &= \frac{1}{2}(1/3 + r/2 + \sqrt{-2/9 - r/3 + r^2 + 8z^3}), \\ \wp_{1,2}(2\omega_2/3) = \wp_{1,2}(4\omega_2/3) &= \frac{1}{2}(1/3 + r/2 - \sqrt{-2/9 - r/3 + r^2 + 8z^3}).\end{aligned}$$

By (83), it follows that

$$(88) \quad Q(0,0) = \frac{1}{4z^3}(-1/3 + r/2 + 2\sqrt{-2/9 - r/3 + r^2 + 8z^3}).$$

The GF $Q(0,0)$ is now expressed as an algebraic function of z (in (83) it was not clear). To conclude, we notice that we can express r with W as follows (this can be achieved by computing the minimal polynomial of $W + W^4/4$ starting from the minimal polynomial of W):

$$r = \frac{2}{3} - 4z^2(W + W^4/4).$$

It is now obvious that (88) implies (75).

9. FINDING AN ALGEBRAIC EXPRESSION FOR THE GF OF KREWERAS' WALKS ENDING ON ONE AXIS

In what follows, we would like to express $r_y(\omega)$ in terms of $y(\omega)$ (to eventually find (76)). To that purpose we notice that the function $x(\omega)$ is simpler than $y(\omega)$, see (77). Further, due to [19, Equation (3.3)] and the symmetry of the model, we have $y(\omega) = x(\omega - \omega_3/2)$. In other words, it is equivalent to express $r_y(\omega)$ in terms of $y(\omega)$, or to express $r_y(\omega + \omega_3/2)$ in terms of $x(\omega)$. Since $\omega_3/2 = \omega_2/3$, (81) implies that

$$(89) \quad r_y(\omega + \omega_3/2) = c + \frac{1}{2z}\zeta_{1,2}(\omega - \omega_2) - \frac{1}{2z}\zeta_{1,2}(\omega - 2\omega_2) + \frac{1}{z}\zeta_{1,2}(\omega - 5\omega_2/3) - \frac{1}{z}\zeta_{1,2}(\omega - 4\omega_2/3).$$

We shall then structure the proof as follows: in Section 9.1 we find the constant c in (81) (or equivalently in (89)). In Section 9.2, for computational reasons, we replace $\zeta_{1,2}$ -functions by $\wp_{1,2}$ -functions. In Section 9.3 we are interested in (needed) technical lemmas. Finally, in Section 9.4 we give the proof of Equation (76) of Proposition 11.⁸

9.1. Finding the constant. To find the constant c in (81), we could use Theorem 7, which gives the exact expression for $r_y(\omega) - r_y(\omega_{y_2})$. It is also possible to use the following property of Kreweras' model: for ω such that $y(\omega) = 0$, we must have $r_y(\omega) = 0$. Indeed, remember that one has $r_y(\omega) = zy(\omega)Q(0, y(\omega))$. Since in this case $y(2\omega_2/3) = y(\omega_2) = 0$, we immediately find

$$(90) \quad c = \frac{1}{2z}(\zeta_{1,2}(\omega_2/3) - \zeta_{1,2}(4\omega_2/3)) + \frac{1}{z}(\zeta_{1,2}(\omega_2) - \zeta_{1,2}(2\omega_2/3)).$$

⁸In Section 8, we have shown how our theoretical results work for finding an expression for $Q(0,0)$ in the case of Kreweras. In Section 9 we now show how our methods also provide an expression for $Q(x,0)$ or $Q(0,y)$. For conciseness, we decided to state all necessary intermediate results to obtain such an expression, but not to prove all of them.

9.2. Replacing the Zeta-functions by Pe-functions. With (89) and (90), the function r_y is completely known. However, it would be more convenient to have $\wp_{1,2}$ -functions instead of $\zeta_{1,2}$ -functions in (89). To do so, we shall use (four times) the addition theorem (P3) in (81). Following this way, we obtain

$$(91) \quad \begin{aligned} r_y(\omega + \omega_3/2 - \omega_2/2) &= r_y(\omega + \omega_3/2 + 3\omega_2/2) \\ &= c + \frac{1}{2z}(\zeta_{1,2}(\omega + \omega_2/2) - \zeta_{1,2}(\omega - \omega_2/2)) + \frac{1}{z}(\zeta_{1,2}(\omega - \omega_2/6) - \zeta_{1,2}(\omega + \omega_2/6)) \\ &= c + \frac{\zeta_{1,2}(\omega_2/2)}{z} - \frac{\wp'_{1,2}(\omega_2/2)/(2z)}{\wp_{1,2}(\omega) - \wp_{1,2}(\omega_2/2)} - \frac{2\zeta_{1,2}(\omega_2/6)}{z} + \frac{\wp'_{1,2}(\omega_2/6)/z}{\wp_{1,2}(\omega) - \wp_{1,2}(\omega_2/6)}. \end{aligned}$$

9.3. Some technical computations. Equation (91) expresses $r_y(\omega + \omega_3/2)$ in terms of one single function, namely, $\wp_{1,2}(\omega + \omega_2/2)$. This is why we now need to compute $\wp_{1,2}(\omega + \omega_2/2)$ in terms of $x(\omega)$. This will be done in Lemma 21. But before stating and proving this result, we first express $x(\omega + \omega_2/2)$ in terms of $x(\omega)$.

Lemma 20. *One has*

$$x(\omega + \omega_2/2) = \sqrt{x_1} \frac{\sqrt{x_1}x(\omega) + 1}{x(\omega) - x_1}, \quad \forall \omega \in \mathbf{C}.$$

Proof. Introduce, as in the proof of Lemma 14, the function $g(x) = 1/3 - 4z^2x$. We have, for any $\omega \in \mathbf{C}$,

$$\wp(\omega + \omega_2/2) = e_2 + \frac{(e_2 - e_1)(e_2 - e_{1+2})}{\wp(\omega) - e_2} = g(x_1) + \frac{(g(x_1) - g(x_2))(g(x_1) - g(x_3))}{g(x(\omega)) - g(x_1)}.$$

Using the facts that $x_1 + x_2 + x_3 = 1/(4z^2)$, $x_1x_2x_3 = 1/4$ and $d(x_1) = 0$, and after some computations, we obtain that

$$\wp(\omega + \omega_2/2) = \frac{(1/3 - 4z^2x_1)x(\omega) + (-x_1/3 + 2z - 2z^2/x_1)}{3(x(\omega) - x_1)}, \quad \forall \omega \in \mathbf{C}.$$

Finally, using that $x(\omega + \omega_2/2) = g^{-1}(\wp(\omega + \omega_2/2))$, we obtain Lemma 20. \square

In Section 8, we introduced r to be the only power series solution to $4X^3 - g_2X + g_3 = 0$ (Equation (84)). The two other solutions of (84) are Puiseux series, and we set

$$\begin{aligned} \tilde{r} &= -1/3 + 4z^3 - 8z^{9/2} + 24z^6 - 84z^{15/2} + 320z^9 + O(z^{21/2}), \\ \hat{r} &= -1/3 + 4z^3 + 8z^{9/2} + 24z^6 + 84z^{15/2} + 320z^9 + O(z^{21/2}). \end{aligned}$$

Lemma 21. *One has*

$$(92) \quad \wp_{1,2}(\omega + \omega_2/2) = \frac{B_0 + B_1x(\omega) + \sqrt{A_0}\sqrt{1 - x(\omega)W^2}}{12(x(\omega) - x_1)}, \quad \forall \omega \in \mathbf{C},$$

where

$$(93) \quad B_0 = -(2x_1 + 24z^2\sqrt{x_1} + 3rx_1), \quad B_1 = 3(r - \tilde{r}), \quad A_0 = \frac{(B_0 + B_1x_1)^2}{1 - x_1W^2}.$$

Proof. The first step of the proof of (92) consists in evaluating at $\omega + \omega_2/2$ the expression in Lemma 19. Together with Lemma 20, this gives an expression of $\wp_{1,2}(\omega + \omega_2/2)$ in terms of $x(\omega)$. Some simplifications (that we do not write here) eventually lead to (92). \square

Corollary 22. *One has*

$$(94) \quad \wp_{1,2}(\omega_2/2) = \frac{B_1}{12}, \quad \wp_{1,2}(\omega_2/6) = \frac{\sqrt{A_0} - B_0}{12x_1}.$$

Proof. In the expression (92) of Lemma 21, evaluate at $\omega = 0$. Since $x(\omega)$ has a pole of order 2 at 0, the first part of Corollary 22 follows. We then evaluate (92) at $\omega = \omega_2/3$. Since $x(\omega_2/3) = 0$, we obtain the second part of Corollary 22. \square

Because of Equation (91), we also need to have an expression of $1/(\wp_{1,2}(\omega + \omega_2/2) - \wp_{1,2}(\omega_2/2))$ and $1/(\wp_{1,2}(\omega + \omega_2/2) - \wp_{1,2}(\omega_2/6))$ in terms of $x(\omega)$.

Lemma 23. *One has*

$$(95) \quad \frac{1}{\wp_{1,2}(\omega + \omega_2/2) - \wp_{1,2}(\omega_2/2)} = \frac{12}{A_0 W^2} (B_0 + B_1 x_1 - \sqrt{A_0} \sqrt{1 - x(\omega) W^2}), \quad \forall \omega \in \mathbf{C},$$

as well as

$$(96) \quad \frac{1}{\wp_{1,2}(\omega + \omega_2/2) - \wp_{1,2}(\omega_2/6)} = \frac{12x_1}{C_1 x(\omega)} (\sqrt{A_0} x_1 + (B_0 - \sqrt{A_0} + B_1 x_1) x(\omega) - x_1 \sqrt{A_0} \sqrt{1 - x(\omega) W^2}), \quad \forall \omega \in \mathbf{C},$$

with

$$(97) \quad C_1 = 2x_1 B_1 B_0 - 2B_1 \sqrt{A_0} x_1 - 2\sqrt{A_0} B_0 + x_1^2 B_1^2 + A_0 + B_0^2.$$

Proof. The proof of both (95) and (96) is immediate, and just uses Lemma 21 and Corollary 22. \square

9.4. Conclusion. Using the expression of (91) of $r_y(\omega + \omega_3/2 - \omega_2/2)$, we obtain that with c defined as in (90), one has

$$r_y(\omega + \omega_3/2) = c + \frac{\zeta_{1,2}(\omega_2/2)}{z} - \frac{\wp'_{1,2}(\omega_2/2)/(2z)}{\wp_{1,2}(\omega + \omega_2/2) - \wp_{1,2}(\omega_2/2)} - \frac{2\zeta_{1,2}(\omega_2/6)}{z} + \frac{\wp'_{1,2}(\omega_2/6)/z}{\wp_{1,2}(\omega + \omega_2/2) - \wp_{1,2}(\omega_2/6)}.$$

Using (95) and (96), we obtain that

$$(98) \quad r_y(\omega + \omega_3/2) = \alpha + \frac{\beta}{x(\omega)} + \left(\gamma + \frac{\delta}{x(\omega)} \right) \sqrt{1 - x(\omega) W^2},$$

where

$$(99) \quad \alpha = c + \frac{\zeta_{1,2}(\omega_2/2)}{z} - \frac{2\zeta_{1,2}(\omega_2/6)}{z} - \frac{6\phi'_{1,2}(\omega_2/2)(B_0 + B_1x_1)}{zA_0W^2} + \frac{12x_1\phi'_{1,2}(\omega_2/6)(B_0 - \sqrt{A_0} + B_1x_1)}{C_1z},$$

$$(100) \quad \beta = \frac{12x_1^2\sqrt{A_0}\phi'_{1,2}(\omega_2/6)}{C_1z},$$

$$(101) \quad \gamma = \frac{6\phi'_{1,2}(\omega_2/2)}{z\sqrt{A_0}W^2},$$

$$(102) \quad \delta = \frac{-12x_1^2\sqrt{A_0}\phi'_{1,2}(\omega_2/6)}{C_1z}.$$

Lemma 24. *One has $\alpha = 1/(2z)$, $\beta = -1$, $\gamma = -1/W$ and $\delta = 1$.*

Proof. The proof of Lemma 24 is long, and is based on arguments similar to those already used in other proofs of Sections 7–9. For brevity, we omit it. \square

With Equation (98), Lemma (24), and since $y(\omega) = x(\omega - \omega_3/2)$, we obtain Proposition 11. The proof is completed.

APPENDIX A. ANOTHER POSSIBLE APPROACH

In this section we discuss another approach which also could lead to representations for r_x and r_y on the universal covering. Contrary to the representations in terms of infinite series of meromorphic functions we obtain in Theorem 7, the expressions for the GFs we could obtain following this alternative approach would be directly in terms of elliptic functions. This other approach is based on methods proposed in [9, Chapter 4], and it could be applied in principle to any model of walks with non-singular set of increments \mathcal{S} and any $z \in \mathcal{H}$. The difficulty of obtaining in this way closed formulæ substantially depends on the values of \mathcal{S} and z . Moreover, this method does not produce explicit representations that can be unified for all models and all $z \in \mathcal{H}$, contrary to the one preferred in this paper.

We sketch this alternative approach below. It heavily relies on [9, Theorem 4.4.1]. This theorem states that if ω_2/ω_3 is rational, the function $r_x(\omega)$ can be written as

$$(103) \quad r_x(\omega) = w_1(\omega) + \tilde{\Phi}(\omega)\phi(\omega) + w(\omega)/s(\omega),$$

where $w_1(\omega)$ and $s(\omega)$ are rational functions in the variable $x(\omega)$, while $\phi(\omega)$ and $w(\omega)$ are algebraic in $x(\omega)$. Further, in [10, Lemma 2.1] it is shown that $\tilde{\Phi}(\omega)$ is holonomic (but not algebraic) in $x(\omega)$. Notice that [9, Theorem 4.4.1] is proved for $z = 1/|\mathcal{S}|$ only, but in [10] it is observed that this result also holds for $z \in]0, 1/|\mathcal{S}|[$.

Theorem 4.4.1 of [9] has already been used in [10] to give another proof, after [2, 6], that the 23 walks having a finite group of the walk have holonomic GFs. It has also been shown in [10] that for the five models with a finite group and a positive covariance

$$\sum_{(i,j) \in \mathcal{S}} ij - [\sum_{(i,j) \in \mathcal{S}} i][\sum_{(i,j) \in \mathcal{S}} j]$$

(including Kreweras' and Gessel's models), $\tilde{\Phi}(\omega)$ is identically zero. This implies that these models have algebraic GFs. Finally, [9, Theorem 4.4.1] was also applied in [19] to show that for the 51 non-singular walks with an infinite group (4), the GFs are holonomic for all $z \in \mathcal{H}$ (though the trivariate GF $Q(x, y; z)$ is not holonomic).

In these two examples of use, [9, Theorem 4.4.1] was used to answer to Problem (ii), stated at the very beginning of this article (i.e., to give some qualitative informations on the GFs). Later on, we realized that the same theorem could also be used to solve Problem (i), on an explicit expression for the GFs. The idea consists in using the two following facts simultaneously:

- With the exception of w , all the functions in (103) could be found (in particular, their poles could be computed) by using the same approach (via Galois theory) as in [9, Chapter 4]. As for w , its only known properties are that for all $\omega \in \mathbf{C}$,

$$(104) \quad w(\omega) = w(\omega + \omega_1),$$

$$(105) \quad w(\omega) = w(\omega_2 - \omega),$$

$$(106) \quad w(\omega) = w(\omega - \omega_3).$$

Equations (104), (105) and (106) are proved in [9, Lemma 4.3.3], [9, Theorem 4.3.1] and [9, Lemma 4.3.3], respectively. Below we shall give some consequences of these three equations, but before we pass to the second fact.

- For $z \in]0, 1[\setminus \mathcal{S}[$, $r_x(\omega)$ is a GF which is analytic on $\{\omega \in \mathbf{C} : |x(\omega)| < 1\}$, see (1). Therefore, it cannot have any singularities in the part of the fundamental parallelogram $\Pi_{0,0}$ where $|x(\omega)| < 1$.

For these reasons, the role of the function w is to compensate for the possible poles which appear because of w_1 , $\tilde{\Phi}\phi$ and $1/s$ in (103).

We now show how this compensation property of w can lead to eventually determine it. The function $w(\omega)$ is ω_1, ω_3 elliptic (see (104) and (106)). Further, with (105), we obtain that the function $v(\omega) = w(\omega + \omega_2/2)$ is even and ω_1, ω_3 elliptic. Using Property (P7), we reach the conclusion that there exists a rational function R such that $v(\omega) = R(\wp(\omega; \omega_1, \omega_3))$. This way, we obtain the existence of a rational function R such that

$$(107) \quad w(\omega) = R(\wp(\omega - \omega_2/2; \omega_1, \omega_3)), \quad \forall \omega \in \mathbf{C}.$$

Next, instead of finding an expression for r_x given in (103), we can be interested in the following weaker problem: finding a rational function R such that, with w defined as in (107), the function r_x has no pole in the domain $\{\omega \in \mathbf{C} : |x(\omega)| < 1\} \cap \Pi_{0,0}$. To be able to find R , we need to overcome three difficulties:

- (i) We first have to compare the domain $\{\omega \in \mathbf{C} : |x(\omega)| < 1\} \cap \Pi_{0,0}$ with a translated fundamental parallelogram $\omega_1[0, 1[+ \omega_3[0, 1[$. Indeed, since w is elliptic with periods ω_1, ω_3 , to determine it we need to know its poles in a translated parallelogram of $\omega_1[0, 1[+ \omega_3[0, 1[$.
- (ii) If, in Point (i), $\{\omega \in \mathbf{C} : |x(\omega)| < 1\} \cap \Pi_{0,0}$ contains a translated fundamental parallelogram $\omega_1[0, 1[+ \omega_3[0, 1[$, then we have to deduce from the expressions of w_1 ,

$\tilde{\Phi}\phi$ and $1/s$ the zeros and poles that w should have in a translated parallelogram of $\omega_1[0, 1[+ \omega_3[0, 1[$.

- (iii) Finally, we have to deduce from (ii) an expression for the rational function R and an expression for $\wp(\omega - \omega_2/3; \omega_1, \omega_3)$ in terms of $x(\omega)$.

Item (i) is difficult to prove—though purely technical. The difficulty of Item (ii) depends on the expression of the functions w_1 , $\tilde{\Phi}\phi$ and $1/s$ in (103). In the general case they are quite complicated, so that it is almost impossible to obtain an expression for R (this explains why we chose another approach in this article). However, in some particular cases, it may happen that these functions are simple to deal with, see just below for Kreweras' example. Item (iii) is doable, and similar computations are done for Kreweras' case (Sections 7–9).

To conclude Section A, we have a closer look at Kreweras' example. One could show that the following choices⁹ are suitable: $w_1(\omega) = -1/x(\omega)$, $\tilde{\Phi}(\omega)\phi(\omega) = 0$ and $1/s(\omega) = 1$. We could then show that the function R has the following simple form

$$R(X) = \frac{\wp'_{1,3}(\omega_2/6)/(2z)}{X - \wp_{1,3}(\omega_2/6)} + \beta.$$

To find β , we could use the same idea as in Theorem 7. Then it would remain to express $\wp(\omega - \omega_2/3; \omega_1, \omega_3)$ in terms of $x(\omega)$. For this we could use the same ideas and techniques as in Sections 7–9.

APPENDIX B. SOME PROPERTIES OF ELLIPTIC FUNCTIONS

In this appendix, we gather the results we used on Weierstrass \wp - and ζ -functions.

Lemma 25. *Let ζ and \wp be the Weierstrass functions with certain periods $\bar{\omega}, \hat{\omega}$.*

(P1) *We have the expansion*

$$\zeta(\omega) = \frac{1}{\omega} + \sum_{(\bar{n}, \hat{n}) \in \mathbf{Z}^2 \setminus \{(0,0)\}} \left(\frac{1}{\omega - (\bar{n}\bar{\omega} + \hat{n}\hat{\omega})} + \frac{1}{\bar{n}\bar{\omega} + \hat{n}\hat{\omega}} + \frac{\omega}{(\bar{n}\bar{\omega} + \hat{n}\hat{\omega})^2} \right), \quad \forall \omega \in \mathbf{C}.$$

As for the expansion of $\wp(\omega)$, it is given in (11). In particular, in the fundamental rectangle $\bar{\omega}[0, 1[+ \hat{\omega}[0, 1[$, ζ (resp. \wp) has a unique pole. It is of order 1 (resp. 2), at 0, and has residue 1 (resp. 0, and principal part $1/\omega^2$).

(P2) *An elliptic function with no poles in the fundamental rectangle $\bar{\omega}[0, 1[+ \hat{\omega}[0, 1[$ is constant.*

(P3) *We have the addition theorems*

$$\zeta(\omega + \tilde{\omega}) = \zeta(\omega) + \zeta(\tilde{\omega}) + \frac{1}{2} \frac{\wp'(\omega) - \wp'(\tilde{\omega})}{\wp(\omega) - \wp(\tilde{\omega})}, \quad \forall \omega, \tilde{\omega} \in \mathbf{C}.$$

and

$$\wp(\omega + \tilde{\omega}) = -\wp(\omega) - \wp(\tilde{\omega}) + \frac{1}{4} \left(\frac{\wp'(\omega) - \wp'(\tilde{\omega})}{\wp(\omega) - \wp(\tilde{\omega})} \right)^2, \quad \forall \omega, \tilde{\omega} \in \mathbf{C}.$$

⁹It is worth noting that there is no uniqueness of w_1 and $1/s$, as if we add some constants to the latter, we can subtract them in w .

(P4) For given $\tilde{\omega}_1, \dots, \tilde{\omega}_p \in \mathbf{C}$, define

$$(108) \quad f(\omega) = c + \sum_{1 \leq \ell \leq p} r_\ell \zeta(\omega - \tilde{\omega}_\ell), \quad \forall \omega \in \mathbf{C}.$$

The function f above is elliptic if and only if $\sum_{1 \leq \ell \leq p} r_\ell = 0$.

(P5) Let f be an elliptic function with periods $\bar{\omega}, \hat{\omega}$ such that in the fundamental rectangle $\bar{\omega}[0, 1[+ \hat{\omega}[0, 1[$, f has only poles of order 1, at $\tilde{\omega}_1, \dots, \tilde{\omega}_p$, with residues r_1, \dots, r_p , respectively. Then there exists a constant c such that (108) holds.

(P6) Let f and g be non-constant elliptic functions with the same periods $\bar{\omega}, \hat{\omega}$. Then there exists a non-zero polynomial P such that $P(f(\omega), g(\omega)) = 0$, for all $\omega \in \mathbf{C}$.

(P7) Let f be an even elliptic function with periods $\bar{\omega}, \hat{\omega}$, and let \wp be the Weierstrass elliptic function with periods $\bar{\omega}, \hat{\omega}$. Then there exists a rational function F such that $f(\omega) = F(\wp(\omega))$, for all $\omega \in \mathbf{C}$.

(P8) Let p be some positive integer. The Weierstrass elliptic function with periods $\bar{\omega}, \hat{\omega}/p$ can be written in terms of \wp as

$$\wp(\omega) + \sum_{1 \leq \ell \leq p-1} [\wp(\omega + \ell \hat{\omega}/p) - \wp(\ell \hat{\omega}/p)], \quad \forall \omega \in \mathbf{C}.$$

(P9) The function ζ is quasi-periodic, in the sense that

$$\zeta(\omega + \bar{\omega}) = \zeta(\omega) + 2\zeta(\bar{\omega}/2), \quad \zeta(\omega + \hat{\omega}) = \zeta(\omega) + 2\zeta(\hat{\omega}/2), \quad \forall \omega \in \mathbf{C}.$$

(P10) The sum of the residues of an elliptic function in a fundamental parallelogram is zero.

Proof. Property (P1) is proved in [28, Sections 20.2 and 20.4]. (P2) is known as Liouville's theorem, and can be found in [28, Section 20.1]. The addition theorems (P3) are stated in [26, Equations (5.5.3) and (5.5.16)]. For Properties (P4) and (P5), we refer to [28, Section 20.52]. (P6) is shown in [28, Section 20.54], and (P7) in [28, Section 20.51]. (P8) can be found in [28, Example 9 in page 456]. Finally, (P9) is proved in [28, Section 20.41], and (P10) is proved in [28, Section 20.12]. \square

ACKNOWLEDGMENTS

We would like to thank Alin Bostan for interesting discussions.

REFERENCES

- [1] Abramowitz, M., and Stegun, I.: *Handbook of mathematical functions with formulas, graphs, and mathematical tables*. National Bureau of Standards Applied Mathematics Series, Washington (1964)
- [2] Bostan, A., and Kauers, M.: The complete generating function for Gessel walks is algebraic. *Proc. Amer. Math. Soc.* **138** 3063–3078 (2010)
- [3] Bostan, A., Kurkova, I., and Raschel, K.: A human proof of Gessel's lattice path conjecture. *Preprint* (2013)
- [4] Bostan, A., Raschel, K., and Salvy, B.: Non-D-finite excursions in the quarter plane. *Preprint arXiv:1205.3300* (2012) *J. Combin. Theory Ser. A*, to appear
- [5] Bousquet-Mélou, M.: Walks in the quarter plane: Kreweras' algebraic model. *Ann. Appl. Probab.* **15** 1451–1491 (2005)

- [6] Bousquet-Mélou, M., and Mishna, M.: Walks with small steps in the quarter plane. *Contemp. Math.* **520** 1–40 (2010)
- [7] Bousquet-Mélou, M., and Petkovsek, M.: Walks confined in a quadrant are not always D-finite. *Theoret. Comput. Sci.* **307** 257–276 (2003)
- [8] Chabat, B.: *Introduction à l'analyse complexe. Tome 1*. Mir, Moscow (1990)
- [9] Fayolle, G., Iasnogorodski, R., and Malyshev, V.: *Random walks in the quarter-plane*. Springer-Verlag, Berlin (1999)
- [10] Fayolle, G., and Raschel, K.: On the holonomy or algebraicity of generating functions counting lattice walks in the quarter-plane. *Markov Process. Related Fields* **16** 485–496 (2010)
- [11] Fayolle, G., and Raschel, K.: Random walks in the quarter plane with zero drift: an explicit criterion for the finiteness of the associated group. *Markov Process. Related Fields* **17** 619–636 (2011)
- [12] Flajolet, P., and Sedgewick, R.: *Analytic combinatorics*. Cambridge University Press, Cambridge (2009)
- [13] Flatto, L., and Hahn, S.: Two parallel queues created by arrivals with two demands. I. *SIAM J. Appl. Math.* **44** 1041–1053 (1984)
- [14] Flatto, L.: Two parallel queues created by arrivals with two demands. II. *SIAM J. Appl. Math.* **45** 861–878 (1985)
- [15] Jones, G., and Singerman, D.: *Complex Functions*. Cambridge University Press, Cambridge (1987)
- [16] Kauers, M., Koutschan, C., and Zeilberger, D.: Proof of Ira Gessel's lattice path conjecture. *P. Natl. Acad. Sci. USA* **106** 11502–11505 (2009)
- [17] Kreweras, G.: Sur une classe de problèmes de dénombrement liés au treillis des partitions des entiers. *Cahiers du B.U.R.O.* **6** 5–105 (1965)
- [18] Kurkova, I., and Raschel, K.: Explicit expression for the generating function counting Gessel's walks. *Adv. in Appl. Math.* **47** 414–433 (2011)
- [19] Kurkova, I., and Raschel, K.: On the functions counting walks with small steps in the quarter plane. *Publ. Math. Inst. Hautes Études Sci.* **116** 69–114 (2012)
- [20] Malyshev, V.: *Random Walks, Wiener-Hopf Equations in the Quarter Plane, Galois Automorphisms. Lomonossov Moscow University Press* (1970) (in Russian)
- [21] Malyshev, V.: Positive random walks and Galois theory. *Uspehi Matem. Nauk* **26** 227–228 (1971)
- [22] Malyshev, V.: An analytical method in the theory of two-dimensional positive random walks. *Siberian Math. J.* **13** 1314–1329 (1972)
- [23] Melzer, S., and Mishna, M.: Singularity analysis via the iterated kernel method. *Preprint arXiv:1303.3236* (2013)
- [24] Mishna, M., and Rechnitzer, A.: Two non-holonomic lattice walks in the quarter plane. *Theor. Comput. Sci.* **410** 3616–3630 (2009)
- [25] Raschel, K.: Counting walks in a quadrant: a unified approach via boundary value problems. *J. Eur. Math. Soc.* **14** 749–777 (2012)
- [26] Sansone, G., and Gerretsen, J.: *Lectures on the theory of functions of a complex variable. I. Holomorphic functions*. P. Noordhoff, Groningen (1960)
- [27] Stanley, R.: *Enumerative Combinatorics*, vol. 2. Cambridge University Press, Cambridge (1999)
- [28] Watson, G., and Whittaker, E.: *A course of modern analysis*. Cambridge University Press, Cambridge (1962)

LABORATOIRE DE PROBABILITÉS ET MODÈLES ALÉATOIRES, UNIVERSITÉ PIERRE ET MARIE CURIE,
4 PLACE JUSSIEU, 75252 PARIS CEDEX 05, FRANCE

E-mail address: Irina.Kourkova@upmc.fr

CNRS & FÉDÉRATION DE RECHERCHE DENIS POISSON & LABORATOIRE DE MATHÉMATIQUES ET
PHYSIQUE THÉORIQUE, UNIVERSITÉ DE TOURS, PARC DE GRANDMONT, 37200 TOURS, FRANCE

E-mail address: Kilian.Raschel@lmpt.univ-tours.fr