



**HAL**  
open science

# Optimal continuous dependence estimates for fractional degenerate parabolic equations

Nathael Alibaud, Simone Cifani, Espen Jakobsen

► **To cite this version:**

Nathael Alibaud, Simone Cifani, Espen Jakobsen. Optimal continuous dependence estimates for fractional degenerate parabolic equations. 2013. hal-00841159v1

**HAL Id: hal-00841159**

**<https://hal.science/hal-00841159v1>**

Preprint submitted on 4 Jul 2013 (v1), last revised 5 Oct 2015 (v4)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# OPTIMAL CONTINUOUS DEPENDENCE ESTIMATES FOR FRACTIONAL DEGENERATE PARABOLIC EQUATIONS

NATHAËL ALIBAUD, SIMONE CIFANI, AND ESPEN R. JAKOBSEN

ABSTRACT. We obtain optimal continuous dependence estimates for weak entropy solutions of degenerate parabolic equations with nonlinear fractional diffusion. The diffusion term involves the fractional Laplace operator,  $\Delta^{\alpha/2}$  for  $\alpha \in (0, 2)$ , the generator of a pure jump Lévy process. Our results cover the dependence on the nonlinearities, and for the first time, also the explicit dependence on  $\alpha$ . The former estimate (dependence on nonlinearity) shows a clear dependence on  $\alpha$ , and it is stable in the limits  $\alpha \rightarrow 0$  and  $\alpha \rightarrow 2$ . In the limit  $\alpha \rightarrow 2$ ,  $\Delta^{\alpha/2}$  converges to the usual Laplacian, and we show rigorously that we recover the optimal continuous dependence result of [15] for local degenerate parabolic equations (thus providing an alternative proof).

## CONTENTS

1. Introduction	2
Notation	5
2. Preliminaries	5
3. The main results	6
4. Two general results from [3]	9
5. Continuous dependence in the supercritical case	10
6. Continuous dependence in the critical and subcritical cases	12
6.1. A technical lemma	12
6.2. Proof of Theorem 3.1	13
6.3. Proof of Theorem 3.8	21
7. Proof of Theorem 3.3	28
8. Optimal example	33
8.1. Optimality of Theorem 3.1	33
8.2. Optimality of Theorem 3.8	34
8.3. Proofs	34
Appendix A. Proof of some estimates used to prove Theorem 3.1	42
Appendix B. Some technical lemmas	43
Appendix C. The proof of Lemma 7.3	43
References	44

---

2010 *Mathematics Subject Classification.* 35R09, 35K65, 35L65, 35D30, 35B30.

*Key words and phrases.* Fractional/fractal conservation laws, nonlinear degenerate diffusions, fractional Laplacian, optimal continuous dependence estimates.

This research was supported by the Research Council of Norway (NFR) through the project "Integro-PDEs: Numerical methods, Analysis, and Applications to Finance", and by the "French ANR project CoToCoLa, no. ANR-11-JS01-006-01".

## 1. INTRODUCTION

In this paper we consider the following Cauchy problem:

$$(1.1) \quad \begin{cases} \partial_t u + \operatorname{div} f(u) + (-\Delta)^{\frac{\alpha}{2}} \varphi(u) = 0 & \text{in } Q_T := \mathbb{R}^d \times (0, T), \\ u(x, 0) = u_0(x), & \text{in } \mathbb{R}^d, \end{cases}$$

where  $T > 0$ ,  $u = u(x, t)$  is the unknown function,  $\operatorname{div}$  and  $\Delta$  denote divergence and Laplacian with respect to  $x$ , and  $(-\Delta)^{\frac{\alpha}{2}}$ ,  $\alpha \in (0, 2)$ , is the fractional Laplacian e.g. defined via the Fourier transform (in space) as  $(-\Delta)^{\frac{\alpha}{2}} \phi := \mathcal{F}^{-1}(|\cdot|^\alpha \mathcal{F}\phi)$ . Throughout the paper we assume that

$$(1.2) \quad u_0 \in L^\infty \cap L^1 \cap BV(\mathbb{R}^d),$$

$$(1.3) \quad f \in W_{\text{loc}}^{1,\infty}(\mathbb{R}; \mathbb{R}^d) \text{ with } f(0) = 0,$$

$$(1.4) \quad \varphi \in W_{\text{loc}}^{1,\infty}(\mathbb{R}) \text{ is nondecreasing with } \varphi(0) = 0.$$

*Remark 1.1.* Subtracting constants to  $f$  and  $\varphi$  if necessary, there is no loss of generality in assuming that  $f(0) = 0$  and  $\varphi(0) = 0$ .

Equation (1.1) could be seen as a nonlocal ‘‘interpolation’’ between the scalar conservation law

$$(1.5) \quad \partial_t u + \operatorname{div} f(u) + \varphi(u) = 0,$$

and the degenerate parabolic equation

$$(1.6) \quad \partial_t u + \operatorname{div} f(u) - \frac{\Delta}{4\pi^2} \varphi(u) = 0,$$

since formally  $-(-\Delta)^{\frac{\alpha}{2}}$  converges to the identity and  $\frac{\Delta}{4\pi^2}$  when  $\alpha$  converges to 0 and 2 respectively. Equation (1.1) is said to be supercritical if  $\alpha < 1$ , subcritical if  $\alpha > 1$ , and critical if  $\alpha = 1$ . It can be strongly degenerate (as can (1.6)), i.e.  $\varphi$  may vanish/degenerate on sets of positive measure. Equation (1.1) can therefore be of mixed hyperbolic parabolic type depending on the choice of  $\alpha$  and  $\varphi$ .

The fractional Laplacian is the generator of the symmetric  $\alpha$ -stable process, the most famous pure jump Lévy process. There is a large literature on Lévy processes, we refer to e.g. [35] for more details, and they are important in many modern applications. Being very selective, we mention radiation hydrodynamics [34, 37, 33], anomalous diffusion in semiconductor growth [38], over-driven gas detonations [14], mathematical finance [17], and flow in porous media [19]. In the mathematical community, interest in nonlinear nonlocal diffusions is in fact very recent, and only few results exist; cf. e.g. [6, 7, 10, 12, 3, 19] and the references therein.

The well-posedness of the Cauchy problem for (1.5) is well-known. The convection  $f$ -term can have a deregularizing effect so that shocks can be formed in finite time. In general, weak solutions are not unique and additional entropy conditions are needed to get uniqueness. The most famous uniqueness result relies on the notion of entropy solutions introduced in [29]. Much later, the adequate notion of entropy solutions for strongly degenerate second-order equations has been introduced in [9]. This paper focuses on some initial-boundary value problems on bounded domains. For equations on  $\mathbb{R}^d$  including (1.6), see e.g. [26]. For more references on conservation laws and degenerate parabolic equations, we refer the reader to the books [20, 18] or the references in [26].

At the same time, there has been a large interest in nonlocal versions of these equations. The first work seems to be [16] on nonlocal time fractional derivatives, cf. also [25]. The study of nonlocal diffusion terms has probably been initiated by [5]. Now, the well-posedness is quite well-understood in the nondegenerate linear

case where  $\varphi(u) = u$ . Smooth solutions exist and are unique for subcritical equations [5, 21], shocks could occur [4, 28] and weak solutions could be nonunique [2] for supercritical equations, entropy solutions exist and are always unique [1, 27]. Very recently, the well-posedness theory was extended in [12] to cover the full problem (1.1), even for strongly degenerate  $\varphi$ . For a more complete bibliography, we refer to [12] or also [3].

In this paper we investigate continuous dependence estimates for (1.1), i.e. we estimate explicitly the difference of two entropy solutions  $u$  and  $v$  in terms of the difference of their respective data  $(\alpha, u_0, f, \varphi)$  and  $(\beta, v_0, g, \psi)$ . In this setting, the first results appeared in [30] for scalar conservation laws via the Kruzkov doubling of variables technique; see also [8] and the references therein for a kinetic approach. Continuous dependence estimates for Equation (1.6) was established in [15]. These estimates are optimal and roughly speaking state that if  $u$  has the same data as  $v$  except for  $\varphi \neq \psi$ , then

$$(1.7) \quad \|u(\cdot, t) - v(\cdot, t)\|_{L^1} = O\left(\|\sqrt{\varphi'} - \sqrt{\psi'}\|_{\infty}\right),$$

where throughout this paper the  $L^{\infty}$ -norm is always taken over the range of  $u_0$ . This result has been generalized to anisotropic diffusions via the kinetic approach, see [11] for more details and references. To the best of our knowledge, the first continuous dependence estimate for nonlocal equations was given in [27]. It concerns the case of linear nondegenerate Lévy diffusions. In [3], the authors of the present paper extended this result to general nonlinear degenerate Lévy diffusions. In particular, the results of [3] apply also to (1.1). But, as it turns out, for the particular Lévy diffusion we are studying in this paper (the fractional Laplacian), they give optimal rates only in the supercritical case  $\alpha < 1$ .

In this paper we obtain optimal rates for all cases. To do so we restart the proofs from the beginning, by taking into account the homogeneity properties of the fractional Laplacian. The main ingredients are a new linearization argument à la Young measure theory/kinetic formulations, and for the linear case, a clever change of the (jump)  $z$ -variable in (2.1). This change of variable allow us to adapt ideas from viscosity solution theory for integro-PDEs developed e.g. in [31]. Let us also refer the reader to [36] for other applications of this change of variable in the context of viscosity solutions. Roughly speaking, we prove that

$$(1.8) \quad \|u(\cdot, t) - v(\cdot, t)\|_{L^1} = \begin{cases} O\left(\|(\varphi')^{\frac{1}{\alpha}} - (\psi')^{\frac{1}{\alpha}}\|_{\infty}\right), & \alpha > 1, \\ O\left(\|\varphi' \ln \varphi' - \psi' \ln \psi'\|_{\infty}\right), & \alpha = 1, \\ O\left(\|\varphi' - \psi'\|_{\infty}\right), & \alpha < 1. \end{cases}$$

Note well that just as in [3], our proofs do not need tools like chain rules or entropy defect measures. And even though these tools play a key role in the local second-order theory, the arguments here really seem to be less technical relying only on basic convex inequalities and integral calculus. Hence, it seems interesting to mention that we recover the optimal result (1.7) rigorously from (1.8) by passing to the limit as  $\alpha \rightarrow 2$ . Another remarkable feature is that a simple rescaling transforms the Kuznetsov type estimate (1.8) into the following time continuity estimate:

$$\|u(\cdot, t) - u(\cdot, s)\|_{L^1} = \begin{cases} O\left(|t^{\frac{1}{\alpha}} - s^{\frac{1}{\alpha}}|\right), & \alpha > 1, \\ O\left(|t \ln t - s \ln s|\right), & \alpha = 1, \\ O\left(|t - s|\right), & \alpha < 1. \end{cases}$$

This result is optimal and strictly better than earlier results in [13], see Remark 3.7. E.g. for positive times, we get Lipschitz regularity in time with values in  $L^1(\mathbb{R}^d)$ . This is a regularizing effect in time when  $\alpha \geq 1$  and  $u$  not more than  $BV$  initially.

In the second main contribution of this paper, we focus on the continuous dependence on the order of the fractional Laplacian. To the best of our knowledge, this is the first time that such a question is studied in detail. By general stability arguments, it is possible to show that the unique entropy solution  $u =: u^\alpha$  of (1.1) is continuous with respect to  $\alpha$  with values in  $L^1_{\text{loc}}$ . In this paper, we prove that it is in fact locally Lipschitz continuous by considering it as a function of  $\alpha \in (0, 2)$  into the Banach space  $C([0, T]; L^1)$ . It seems interesting to recall that the type of Equation (1.1) could change from parabolic when  $\alpha > 1$  to hyperbolic when  $\alpha < 1$ . As a consequence, quite different behaviors are observed in the  $\varphi$ - and  $t$ -continuity when  $\alpha$  crosses 1, cf. the continuous dependence estimates above. A natural question is thus whether such kind of phenomena arises in the  $\alpha$ -regularity? We prove that the answer is positive by carefully estimating the best Lipschitz constant of the function  $\alpha \rightarrow u^\alpha$  with respect to the position of  $\alpha$  and the other data. More precisely, for  $\lambda \in (0, 2)$  we define

$$\text{Lip}_\alpha(u; \lambda) := \limsup_{\alpha, \beta \rightarrow \lambda} \frac{\|u^\alpha - u^\beta\|_{C([0, T]; L^1)}}{|\alpha - \beta|},$$

and roughly speaking we prove that

$$\text{Lip}_\alpha(u; \lambda) = \begin{cases} O\left(M^{\frac{1}{\lambda}} |\ln M|\right), & \lambda > 1, \\ O\left(M \ln^2 M\right), & \lambda = 1, \\ O(M), & \lambda < 1, \end{cases}$$

for  $M := T \|\varphi'\|_\infty$ , and

$$\text{Lip}_\alpha(u; \lambda) = \begin{cases} O(|u_0|_{BV}), & \lambda > 1, \\ O\left(|u_0|_{BV} \ln^2 \frac{\|u_0\|_{L^1}}{|u_0|_{BV}}\right), & \lambda = 1, \\ O\left(\|u_0\|_{L^1}^{1-\lambda} |u_0|_{BV}^\lambda \ln \frac{\|u_0\|_{L^1}}{|u_0|_{BV}}\right), & \lambda < 1. \end{cases}$$

We also exhibit an example of an equation for which these estimates are optimal in the regimes where  $M$  is sufficiently small or  $\frac{\|u_0\|_{L^1}}{|u_0|_{BV}}$  is sufficiently large. Let us emphasize that here again the proofs rely on a clever change of (the jump) variable, so that the general results of [3] allow us to show the Lipschitz regularity in  $\alpha$  only in the supercritical case.

To conclude, note that even if we adapt some ideas from viscosity solution theory, the definition of relevant generalized solution and the mathematical arguments are very different from the ones in e.g. [31]. Moreover we obtain optimal results here, which was not the case in [31] or other papers treating viscosity solutions.

As far as open problems are concerned, a natural question is what is the best modulus of continuity of  $\alpha \rightarrow u^\alpha$ , at  $\alpha = 0$  and 2. At 0, this function could surprisingly be discontinuous with values in  $L^1$ ; but, this seems to be a problem of choosing the right norm, cf. Remark 3.11(2) for more details. For 2, the question relies on the rate of convergence of the entropy solutions of (1.1) toward (1.6) as  $\alpha \rightarrow 2$ . To the best of our knowledge, this problem is still open and rather difficult, at least in the nondegenerate cases where there could be some lack of regularity.

The rest of the paper is organized as follows. In Section 2, we recall the well-posedness theory for fractional degenerate parabolic equations. In Section 3, we state our main results: continuous dependence with respect to the nonlinearities and the order of the fractional Laplacian. In Section 4, we recall the general continuous dependence estimates of [3] along with a general Kuznetsov type of Lemma. Sections 5–7 are devoted to the proofs of our main results. In Section 8, we exhibit

an example of an equation for which we rigorously show that our estimates are optimal. Finally, there is an appendix containing technical lemmas and computations from the different proofs.

**Notation.** The symbols  $\nabla$  and  $\nabla^2$  denote the  $x$ -gradient and  $x$ -Hessian. The symbols  $\|\cdot\|$  and  $|\cdot|$  are used for norms and semi-norms, respectively. We denote by  $\mathcal{F}$  the Fourier transform in the  $x$ -variable  $\mathcal{F}\phi(\xi) = \int e^{-2i\pi x \cdot \xi} \phi(x) dx$ . The surface measure of the unit sphere of  $\mathbb{R}^d$  is denoted by  $S_d$ . The convex hull of  $\{a, b\} \subset \mathbb{R}$  is denoted by  $\text{co}\{a, b\} := (a \wedge b, a \vee b)$ .

## 2. PRELIMINARIES

In this section we recall some basic facts on the fractional Laplacian and fractional degenerate parabolic equations. We start by a Lévy–Khinchine type representation formula. For  $\alpha \in (0, 2)$  and all  $\phi \in C_c^\infty(\mathbb{R}^d)$ ,  $x \in \mathbb{R}^d$ , and  $r > 0$ ,

$$\begin{aligned} -(-\Delta)^{\frac{\alpha}{2}} \phi(x) &= G_d(\alpha) \int_{|z| < r} \frac{\phi(x+z) - \phi(x) - \nabla\phi(x) \cdot z}{|z|^{d+\alpha}} dz \\ &+ G_d(\alpha) \int_{|z| > r} \frac{\phi(x+z) - \phi(x)}{|z|^{d+\alpha}} dz \\ (2.1) \qquad \qquad \qquad &=: \mathcal{L}_r^\alpha[\phi](x) + \mathcal{L}^{\alpha,r}[\phi](x), \end{aligned}$$

where

$$(2.2) \qquad \qquad \qquad G_d(\alpha) := \frac{\alpha \Gamma\left(\frac{d+\alpha}{2}\right)}{2\pi^{\frac{d}{2}+\alpha} \Gamma\left(\frac{2-\alpha}{2}\right)}.$$

The result is standard, see e.g. [24, 22] and the references therein. Here are some properties on the coefficient that will be needed later:

$$(2.3) \qquad \begin{cases} G_d(\alpha) > 0 \text{ is smooth (and analytic) with respect to } \alpha \in (0, 2); \\ G_d(\alpha) \sim_{\alpha \rightarrow 0} S_d^{-1} \alpha \quad \text{and} \quad G_d(\alpha) \sim_{\alpha \rightarrow 2} \frac{d S_d^{-1}}{4\pi^2} (2 - \alpha), \end{cases}$$

where  $S_d$  is the surface measure of the unit sphere of  $\mathbb{R}^d$ .

We then proceed to define entropy solutions of (1.1). Let  $u$  be a function and consider the Kruzkov [29] entropies  $|u - k|$ ,  $k \in \mathbb{R}$ , and entropy fluxes

$$q_f(u, k) := \text{sgn}(u - k) (f(u) - f(k)) \in \mathbb{R}^d,$$

where throughout this paper we always consider the following everywhere representation of the sign function:

$$(2.4) \qquad \qquad \qquad \text{sgn}(u) := \begin{cases} \pm 1 & \text{if } \pm u > 0, \\ 0 & \text{if } u = 0. \end{cases}$$

By monotonicity (1.4) of  $\varphi$ ,

$$(2.5) \qquad \qquad \qquad \text{sgn}(u - k) (\varphi(u) - \varphi(k)) = |\varphi(u) - \varphi(k)|,$$

and then we formally deduce from (2.1) that

$$\text{sgn}(u - k) (-(-\Delta)^{\frac{\alpha}{2}} \varphi(u)) \leq \mathcal{L}_r^\alpha[|\varphi(u) - \varphi(k)|] + \text{sgn}(u - k) \mathcal{L}^{\alpha,r}[\varphi(u)].$$

This Kato type inequality is the starting point of the entropy formulation from [12].

**Definition 2.1** (Entropy solutions). *Let  $\alpha \in (0, 2)$ ,  $u_0 \in L^\infty \cap L^1(\mathbb{R}^d)$ , and (1.3)–(1.4) hold. We say that  $u \in L^\infty(Q_T) \cap L^\infty(0, T; L^1)$  is an entropy solution of (1.1) provided that for all  $k \in \mathbb{R}$ ,  $r > 0$ , and all nonnegative  $\phi \in C_c^\infty(\mathbb{R}^{d+1})$ ,*

$$(2.6) \quad \begin{aligned} & \int_{Q_T} |u - k| \partial_t \phi + q_f(u, k) \cdot \nabla \phi \, dx \, dt \\ & + \int_{Q_T} |\varphi(u) - \varphi(k)| \mathcal{L}_r^\alpha[\phi] + \operatorname{sgn}(u - k) \mathcal{L}^{\alpha, r}[\varphi(u)] \phi \, dx \, dt \\ & + \int_{\mathbb{R}^d} |u_0(x) - k| \phi(x, 0) \, dx \geq 0. \end{aligned}$$

*Remark 2.1.* Under our assumptions, the entropy solutions are continuous in time with values in  $L^1(\mathbb{R}^d)$  (cf. Theorem 2.2 below). Hence we get an equivalent definition if we take  $\phi \in C_c^\infty(\mathbb{R}^{d+1})$  and add the term  $-\int_{\mathbb{R}^d} |u(x, T) - k| \phi(x, T) \, dx$  to (2.6); see [12] for more details.

Here is the well-posedness result from [12].

**Theorem 2.2.** (Well-posedness) *Let  $\alpha \in (0, 2)$ ,  $u_0 \in L^\infty \cap L^1(\mathbb{R}^d)$ , and (1.3)–(1.4) hold. Then there exists a unique entropy solution  $u \in L^\infty(Q_T) \cap C([0, T]; L^1)$  of (1.1), satisfying*

$$(2.7) \quad \begin{cases} \operatorname{ess\,inf} u_0 \leq u \leq \operatorname{ess\,sup} u_0, \\ \|u\|_{C([0, T]; L^1)} \leq \|u_0\|_{L^1}, \\ |u|_{L^\infty(0, T; BV)} \leq |u_0|_{BV}. \end{cases}$$

*Moreover, if  $v$  is an entropy solution of (1.1) with  $v(\cdot, 0) = v_0(\cdot) \in L^\infty \cap L^1(\mathbb{R}^d)$ , then*

$$(2.8) \quad \|u - v\|_{C([0, T]; L^1)} \leq \|u_0 - v_0\|_{L^1}.$$

### 3. THE MAIN RESULTS

We state our main results in this section. They compare the entropy solution  $u$  of (1.1) to the entropy solution  $v$  of

$$(3.1) \quad \begin{cases} \partial_t v + \operatorname{div} g(v) + (-\Delta)^{\frac{\beta}{2}} \psi(v) = 0, \\ v(0, \cdot) = v_0(\cdot), \end{cases}$$

under the assumptions that

$$(3.2) \quad \begin{cases} \alpha, \beta \in (0, 2), \\ u_0 \in L^\infty \cap L^1 \cap BV(\mathbb{R}^d), v_0 \in L^\infty \cap L^1(\mathbb{R}^d), \\ f, g \in \left(W_{\operatorname{loc}}^{1, \infty}(\mathbb{R})\right)^d \text{ with } f(0) = 0 = g(0), \\ \varphi, \psi \in W_{\operatorname{loc}}^{1, \infty}(\mathbb{R}) \text{ are nondecreasing with } \varphi(0) = 0 = \psi(0). \end{cases}$$

From now on, we will use the shorthand notation

$$\begin{aligned} \|f' - g'\|_\infty &:= \operatorname{ess\,sup}_{I(u_0)} |f' - g'|, \\ \|\varphi' - \psi'\|_\infty &:= \operatorname{ess\,sup}_{I(u_0)} |\varphi' - \psi'|, \end{aligned}$$

where  $I(u_0) := (\operatorname{ess\,inf} u_0, \operatorname{ess\,sup} u_0)$ . We will also define

$$(3.3) \quad \mathbf{E}_i(u_0) := |u_0|_{BV} \left\{ 1 + \left( \ln \frac{\|u_0\|_{L^1}}{|u_0|_{BV}} \right)^i \right\} \mathbf{1}_{\frac{\|u_0\|_{L^1}}{|u_0|_{BV}} > 1},$$

with the convention that  $E_i(u_0) = 0$  if  $|u_0|_{BV} = 0$  ( $i = 1, 2$ ). Simple computations show that  $0 \leq E_i(u_0) \leq \|u_0\|_{L^1}$ ,

$$(3.4) \quad \frac{|u_0|_{BV}}{E_i(u_0)} = o(1) \quad \text{and} \quad \frac{E_i(u_0)}{\|u_0\|_{L^1}} = o(1) \quad \text{as} \quad \frac{\|u_0\|_{L^1}}{|u_0|_{BV}} \rightarrow +\infty.$$

Here is our first main result.

**Theorem 3.1.** (Continuous dependence on the nonlinearities) *Let (3.2) hold with  $\alpha = \beta$ , and let  $u$  and  $v$  be the entropy solutions of (1.1) and (3.1) respectively. Then for all  $T > 0$ ,*

$$(3.5) \quad \|u - v\|_{C([0,T];L^1)} \leq \|u_0 - v_0\|_{L^1} + T |u_0|_{BV} \|f' - g'\|_\infty + C \mathcal{E}_{T,\alpha,u_0}^{\varphi-\psi},$$

with  $C = C(d, \alpha)$  and

$$(3.6) \quad \mathcal{E}_{T,\alpha,u_0}^{\varphi-\psi} = \begin{cases} T^{\frac{1}{\alpha}} |u_0|_{BV} \|(\varphi')^{\frac{1}{\alpha}} - (\psi')^{\frac{1}{\alpha}}\|_\infty, & \alpha \in (1, 2), \\ T E_1(u_0) \|\varphi' - \psi'\|_\infty \\ \quad + T(1 + |\ln T|) |u_0|_{BV} \|\varphi' - \psi'\|_\infty \\ \quad + T |u_0|_{BV} \|\varphi' \ln \varphi' - \psi' \ln \psi'\|_\infty, & \alpha = 1, \\ T \|u_0\|_{L^1}^{1-\alpha} |u_0|_{BV}^\alpha \|\varphi' - \psi'\|_\infty, & \alpha \in (0, 1). \end{cases}$$

The proof of this result can be found in Sections 5 and 6.

*Remark 3.2.* We emphasize that this result is optimal with respect to the modulus in  $\varphi$ . In the regimes where  $T$  is sufficiently small or  $\frac{\|u_0\|_{L^1}}{|u_0|_{BV}}$  is sufficiently large, it is also optimal with respect to the dependence of  $T$  and  $u_0$ . See the example of Section 8 for more details. In particular, see Proposition 8.1 and Remark 8.2.

Note that our result is robust in the sense that the constant  $C = C(d, \alpha)$  in Theorem 3.1 has finite limits as  $\alpha \rightarrow 0$  or  $\alpha \rightarrow 2$ . This will be seen during the proof, cf. Remarks 5.1(1) and 6.2(1). Hence, we can recover the known continuous dependence results of the limiting cases  $\alpha = 0$  and  $\alpha = 2$  (cf. (1.7)), i.e. for Equations (1.5) and (1.6).

To show this we start by identifying the limits of the solutions  $u^\alpha$  of (1.1) as  $\alpha \rightarrow 0$  and  $\alpha \rightarrow 2$ .

**Theorem 3.3** (Limiting equations). *Let  $u_0 \in L^\infty \cap L^1(\mathbb{R}^d)$ , (1.3)–(1.4) hold, and for each  $\alpha \in (0, 2)$ , let  $u^\alpha$  denote the entropy solution of (1.1). Then  $u^\alpha$  converges in  $C([0, T]; L^1_{\text{loc}})$ , as  $\alpha \rightarrow 0$  (resp.  $\alpha \rightarrow 2$ ), to the unique entropy solution  $u \in L^\infty(Q_T) \cap C([0, T]; L^1)$  of (1.5) (resp. (1.6)) with initial condition  $u_0$ .*

Let us recall that under our assumptions there are unique entropy solutions of (1.5) and (1.6) with initial data  $u_0$ ; cf. [29, 9, 26]. The proof of Theorem 3.3 can be found in Section 7, as well as the definitions of entropy solutions of [29, 9, 26].

Now we prove that the estimates holds in the limits  $\alpha = 0$  and  $\alpha = 2$ .

**Corollary 3.4** (Limiting estimates). *Theorem 3.1 holds with  $\alpha \in [0, 2]$ .*

*Proof.* We only do the proof for  $\alpha = 2$ , the case  $\alpha = 0$  being similar. Let  $u$  and  $v$  denote the entropy solutions of (1.1) and (3.1) with  $\alpha = 2$  respectively. Moreover, for each  $\alpha \in (0, 2)$ , we denote by  $u^\alpha$  and  $v^\alpha$  the entropy solutions of (1.1) and (3.1) respectively, and  $\mathcal{E}(\alpha)$  the right hand side of (3.5). Then

$$u - v = (u - u^\alpha) + (u^\alpha - v^\alpha) + (v^\alpha - v),$$

and the triangle inequality and Theorems 3.1 and 3.3 imply that for all  $R > 0$ ,

$$\|(u - v)\mathbf{1}_{|x| < R}\|_{C([0,T];L^1)} \leq o(1) + \mathcal{E}(\alpha) + o(1)$$



as  $\alpha \rightarrow 2$  and  $R$  is fixed. By the monotone convergence theorem, Remark 6.2(1), and  $\alpha$ -continuity of  $\mathcal{E}_{T,\alpha,u_0}^{\varphi-\psi}$  at  $\alpha = 2$ , the result follows by first sending  $\alpha \rightarrow 2$  and then sending  $R \rightarrow +\infty$ .  $\square$

*Remark 3.5.* By our results for  $\alpha = 2$ , we get back the “modulus” of [15],

$$\mathcal{E}_{T,\alpha=2,u_0}^{\varphi-\psi} = T^{\frac{1}{2}} |u_0|_{BV} \|(\varphi')^{\frac{1}{2}} - (\psi')^{\frac{1}{2}}\|_{\infty}.$$

Our approach also gives an alternative and possibly less technical proof this result.

Optimal time regularity for (1.1) is another corollary of Theorem 3.3.

**Corollary 3.6** (Modulus of continuity in time). *Let  $\alpha \in [0, 2]$  and (1.2)–(1.4) hold. Let  $u$  be the entropy solution of (1.1). Then for all  $t, s \geq 0$ ,*

$$(3.7) \quad \|u(\cdot, t) - u(\cdot, s)\|_{L^1} \leq |u_0|_{BV} \|f'\|_{\infty} |t - s| + C \mathcal{E}_{\alpha,u_0,\varphi}^{t-s},$$

with  $C = C(d, \alpha)$ ,

$$\mathcal{E}_{\alpha,u_0,\varphi}^{t-s} = \begin{cases} |u_0|_{BV} \|(\varphi')^{\frac{1}{\alpha}}\|_{\infty} |t^{\frac{1}{\alpha}} - s^{\frac{1}{\alpha}}|, & \alpha \in (1, 2], \\ E_1(u_0) \|\varphi'\|_{\infty} |t - s| \\ \quad + |u_0|_{BV} \|\varphi'\|_{\infty} (1 + \|\ln \varphi'\|_{\infty}) |t - s| \\ \quad + |u_0|_{BV} \|\varphi'\|_{\infty} |t \ln t - s \ln s|, & \alpha = 1, \\ \|u_0\|_{L^1}^{1-\alpha} |u_0|_{BV}^{\alpha} \|\varphi'\|_{\infty} |t - s|, & \alpha \in [0, 1), \end{cases}$$

and where  $E_1(u_0)$  is defined in (3.3).

*Remark 3.7.* This result is optimal with respect to the modulus in time, and also with respect to the dependence of  $\varphi$  and  $u_0$  in the regimes where  $\varphi'$  is sufficiently small or the ratio  $\frac{\|u_0\|_{L^1}}{|u_0|_{BV}}$  is sufficiently large, cf. Remark 8.5. The result improves earlier results by the two last authors in [13] where the modulus was given as

$$\mathcal{E}_{\alpha,u_0,\varphi}^{t-s} = C(\alpha, u_0, \varphi) \begin{cases} |t - s|^{\frac{1}{\alpha}}, & \alpha > 1, \\ |t - s|(1 + |\ln |t - s||), & \alpha = 1, \\ |t - s|, & \alpha < 1. \end{cases}$$

The optimal new results give a strictly better modulus of continuity when  $\alpha \in [1, 2]$  at the initial time<sup>1</sup> and for positive times  $u \in W_{\text{loc}}^{1,\infty}((0, +\infty]; L^1)$ . The Lipschitz in time result is a regularizing effect when the solution is no more than BV initially.

*Proof.* We fix  $t, s > 0$  and introduce the rescaled solutions  $v(x, \tau) := u(x, t\tau)$  and  $w(x, \tau) := u(x, s\tau)$ . These are solutions of (1.1) with initial data  $u_0$ , new respective fluxes  $t f$  and  $s f$ , and new respective diffusion functions  $t \varphi$  and  $s \varphi$ . The result immediately follows from the preceding corollary applied at time  $\tau = 1$ .  $\square$

Next we consider the continuous dependence on  $\alpha$ . Given  $T > 0$  and  $\lambda \in (0, 2)$ , we define “the best Lipschitz constant” of  $\alpha \rightarrow u_{\alpha}$  at the position  $\alpha = \lambda$  as follows:

$$(3.8) \quad \text{Lip}_{\alpha}(u; \lambda) := \limsup_{\alpha, \beta \rightarrow \lambda} \frac{\|u^{\alpha} - u^{\beta}\|_{C([0,T]; L^1)}}{|\alpha - \beta|},$$

where  $u^{\alpha}$  denotes the unique entropy solution of (1.1).

---

<sup>1</sup>Simply take sequences  $t_n, s_n \rightarrow 0$  such that  $\frac{t_n}{s_n} \rightarrow 1$  and check that  $\lim_{n \rightarrow +\infty} \frac{|t_n^{\frac{1}{\alpha}} - s_n^{\frac{1}{\alpha}}|}{|t_n - s_n|^{\frac{1}{\alpha}}} = 0$  and  $\lim_{n \rightarrow +\infty} \frac{|t_n \ln t_n - s_n \ln s_n|}{|t_n - s_n| |\ln |t_n - s_n||} = 0$ .

**Theorem 3.8.** (Lipschitz continuity in  $\alpha$ ) *Let (1.2)–(1.4) hold. For all  $T > 0$  and  $\lambda \in (0, 2)$ ,*

$$(3.9) \quad \text{Lip}_\alpha(u; \lambda) \leq C \begin{cases} M^{\frac{1}{\lambda}} (1 + |\ln M|) |u_0|_{BV}, & \lambda \in (1, 2), \\ M E_2(u_0) + M (1 + \ln^2 M) |u_0|_{BV}, & \lambda = 1, \\ M \|u_0\|_{L^1}^{1-\lambda} |u_0|_{BV}^\lambda \left(1 + \left| \ln \frac{\|u_0\|_{L^1}}{|u_0|_{BV}} \right| \right), & \lambda \in (0, 1), \end{cases}$$

where  $C = C(d, \lambda)$ ,  $M := T \|\varphi'\|_\infty$  and  $E_2(u_0)$  is defined in (3.3). In particular, the function  $\alpha \in (0, 2) \rightarrow u^\alpha \in C([0, T]; L^1)$  is locally Lipschitz continuous.

The proof of Theorem 3.8 can be found in Sections 5 and 6.

*Remark 3.9.* This result is optimal with respect to the dependence of  $M$  and  $u_0$  in the regimes where  $M$  is sufficiently small or  $\frac{\|u_0\|_{L^1}}{|u_0|_{BV}}$  is sufficiently large. An example is given in Section 8, cf. Proposition 8.3 and Remark 8.4.

*Remark 3.10.* With Theorem 3.1, Corollary 3.6 and Theorem 3.8 in hands, we can easily get an explicit continuous dependence estimate of  $u$  with respect to the quintuplet  $(t, \alpha, u_0, f, \varphi)$  under (3.2).

*Remark 3.11.* (1) When  $\lambda$  approaches 0 or 2,  $C = C(d, \lambda)$  blows up. As explained below, we can not do better for  $\lambda = 0$  but this is not clear for  $\lambda = 2$ .

- (2) Concerning  $\lambda = 0$ , the function  $\alpha \rightarrow u^\alpha \in C([0, T]; L^1)$  is not even continuous at this point, even though this is always the case in  $C([0, T]; L^1_{\text{loc}})$ . A counter example is the initial value problem  $\partial_t u + (-\Delta)^{\frac{\alpha}{2}} u = 0$  and  $u(x, 0) = u_0(x)$  where  $\int u_0 \neq 0^2$ . Note that in this example the convergence holds in  $C([0, T]; L^p)$  for every  $p > 1$ .
- (3) Concerning  $\lambda = 2$ , the problem of finding the rate of convergence of the solutions of (1.1) to (1.6) is open.

#### 4. TWO GENERAL RESULTS FROM [3]

In this section we recall two key results developed in [3] for the more general case where the diffusion operator can be the generator of an arbitrary pure jump Lévy process. First we state the Kuznetsov type lemma of [3] that measures the  $L^1$ -distance between  $u$  and an arbitrary function  $v$ . From now on, let  $\epsilon$  and  $\nu$  be positive parameters and  $\phi^{\epsilon, \nu} \in C^\infty(\mathbb{R}^{2d+2})$  denote the test function

$$(4.1) \quad \phi^{\epsilon, \nu}(x, t, y, s) := \theta_\nu(t - s) \rho_\epsilon(x - y) := \frac{1}{\nu} \theta\left(\frac{t - s}{\nu}\right) \frac{1}{\epsilon^d} \rho\left(\frac{x - y}{\epsilon}\right),$$

where

$$(4.2) \quad \begin{cases} \theta \in C_c^\infty(\mathbb{R}), & \theta \geq 0, & \text{supp } \theta \subseteq [-1, 1], & \int \theta = 1, \\ \rho \in C_c^\infty(\mathbb{R}^d), & \rho \geq 0, & \text{and } \int \rho = 1. \end{cases}$$

We also let  $m_u(\nu)$  denote the modulus of continuity in time of  $u \in C([0, T]; L^1)$ .

**Lemma 4.1** (Kuznetsov type Lemma). *Let  $\alpha \in (0, 2)$ ,  $u_0 \in L^\infty \cap L^1 \cap BV(\mathbb{R}^d)$ , and let us assume (1.3)–(1.4). Let  $u$  be the entropy solution of (1.1) and let  $v \in$*

<sup>2</sup>By Fourier methods,  $u - v = \mathcal{F}^{-1} \left\{ (e^{-t|\cdot|^\alpha} - e^{-t}) \mathcal{F}u_0 \right\}$ ,  $v$  solution of (1.5) with  $v_0 = u_0$ . Then  $\|u(\cdot, t) - v(\cdot, t)\|_{L^1} \geq \left\| (e^{-t|\cdot|^\alpha} - e^{-t}) \mathcal{F}u_0(\cdot) \right\|_\infty \geq (1 - e^{-t}) \mathcal{F}u_0(0) \neq 0$ , for all  $\alpha$ .

$L^\infty(Q_T) \cap C([0, T]; L^1)$  be such that  $v(0, \cdot) = v_0(\cdot)$ . Then for all  $r, \epsilon > 0$  and  $T > \nu > 0$ ,

$$\begin{aligned}
(4.3) \quad & \|u(\cdot, T) - v(\cdot, T)\|_{L^1} \\
& \leq \|u_0 - v_0\|_{L^1} + C_\rho |u_0|_{BV} \epsilon + 2m_u(\nu) \vee m_v(\nu) \\
& \quad - \int_{Q_T^2} |v(x, t) - u(y, s)| \partial_t \phi^{\epsilon, \nu}(x, t, y, s) \, dw \\
& \quad - \int_{Q_T^2} q_f(v(x, t), u(y, s)) \cdot \nabla_x \phi^{\epsilon, \nu}(x, t, y, s) \, dw \\
& \quad + \int_{Q_T^2} |\varphi(v(x, t)) - \varphi(u(y, s))| \mathcal{L}_r^\alpha[\phi^{\epsilon, \nu}(x, t, \cdot, s)](y) \, dw \\
& \quad - \int_{Q_T^2} \operatorname{sgn}(v(x, t) - u(y, s)) \mathcal{L}^{\alpha, r}[\varphi(u(\cdot, s))](y) \phi^{\epsilon, \nu}(x, t, y, s) \, dw \\
& \quad + \int_{\mathbb{R}^d \times Q_T} |v(x, T) - u(y, s)| \phi^{\epsilon, \nu}(x, T, y, s) \, dx \, dy \, ds \\
& \quad - \int_{\mathbb{R}^d \times Q_T} |v_0(x) - u(y, s)| \phi^{\epsilon, \nu}(x, 0, y, s) \, dx \, dy \, ds
\end{aligned}$$

where  $dw := dx \, dt \, dy \, ds$  and the constant  $C_\rho$  only depends on  $\rho$ .

*Proof.* This is Lemma 3.1 of [3] with the particular diffusion operator (2.1).  $\square$

In the setting of this paper, the general continuous dependence estimates of [3] take the following form:

**Theorem 4.2.** *Let us assume (3.2) and let  $u$  and  $v$  be the respective entropy solutions of (1.1) and (3.1). Then for all  $T > 0$ , and all  $r > 0$ ,*

$$\|u - v\|_{C([0, T]; L^1)} \leq \|u_0 - v_0\|_{L^1} + T |u_0|_{BV} \|f' - g'\|_\infty + \mathcal{E}_{T, \alpha, \beta, u_0, \varphi}^{\alpha - \beta, \varphi - \psi}$$

with

$$(4.4) \quad \mathcal{E}_{T, \alpha, \beta, u_0, \varphi, r}^{\alpha - \beta, \varphi - \psi} = \begin{cases} T \int_{|z| > r} \|u_0(\cdot + z) - u_0(\cdot)\|_{L^1} \, d\mu_\alpha(z) \|\varphi' - \psi'\|_\infty \\ \quad + c_d \sqrt{T} |u_0|_{BV} \sqrt{\int_{|z| < r} |z|^2 \, d\mu_\alpha(z)} \|\varphi' - \psi'\|_\infty, & \alpha = \beta, \\ M \int_{|z| > r} \|u_0(\cdot + z) - u_0(\cdot)\|_{L^1} \, d|\mu_\alpha - \mu_\beta|(z) \\ \quad + c_d \sqrt{M} |u_0|_{BV} \sqrt{\int_{|z| < r} |z|^2 \, d|\mu_\alpha - \mu_\beta|(z)}, & \varphi = \psi, \end{cases}$$

where  $d\mu_\alpha(z) = \frac{G_d(\alpha)}{|z|^{d+\alpha}} \, dz$ ,  $M = T \|\varphi'\|_\infty$  and  $c_d = \sqrt{\frac{4d^2}{d+1}}$ .

*Proof.* This is Theorems 3.3 and 3.4 of [3] with the special choice of diffusion (2.1) and Lévy measure  $\frac{G_d(\alpha)}{|z|^{d+\alpha}} \, dz$ .  $\square$

## 5. CONTINUOUS DEPENDENCE IN THE SUPERCRITICAL CASE

In this section we use Theorem 4.2 to prove Theorems 3.1 and 3.8 for supercritical diffusions.

*Proof of Theorem 3.1 when  $\alpha < 1$ .* We use Estimate (4.4) with  $\beta = \alpha$ . Since  $\alpha < 1$ , the worst term  $\sqrt{\int_{|z| < r} |z|^2 \, d\mu_\alpha(z)} \|\varphi' - \psi'\|_\infty$  vanishes when  $r \rightarrow 0$ , and hence

$$\mathcal{E}_{T, \alpha, \beta, u_0, \varphi, r}^{\alpha - \beta, \varphi - \psi} \xrightarrow{r \rightarrow 0} I := T \int \|u_0(\cdot + z) - u_0(\cdot)\|_{L^1} \, d\mu_\alpha(z) \|\varphi' - \psi'\|_\infty.$$

To estimate this integral, we consider separately the domains  $|z| > \tilde{r}$  and  $|z| < \tilde{r}$  for arbitrary  $\tilde{r} > 0$ . In the second domain, we use the inequality  $\|u_0(\cdot+z) - u_0(\cdot)\|_{L^1} \leq |u_0|_{BV} |z|$ . A direct computation using the fact that  $\alpha < 1$ , then leads to

$$I \leq 2T \|u_0\|_{L^1} \|\varphi' - \psi'\|_\infty S_d \frac{G_d(\alpha)}{\alpha} \tilde{r}^{-\alpha} + T |u_0|_{BV} \|\varphi' - \psi'\|_\infty S_d \frac{G_d(\alpha)}{1-\alpha} \tilde{r}^{1-\alpha}$$

(where  $S_d$  is the surface measure of the unit sphere of  $\mathbb{R}^d$ ). We complete the proof by taking  $\tilde{r} = \|u_0\|_{L^1} |u_0|_{BV}^{-1}$ .  $\square$

- Remark 5.1.* (1) From the proof,  $C \leq S_d \left( \frac{2G_d(\alpha)}{\alpha} + \frac{G_d(\alpha)}{1-\alpha} \right)$  in (3.5) when  $\alpha < 1$ . By (2.3),  $\lim_{\alpha \rightarrow 0} C(d, \alpha)$  is finite and independent of  $d$ .
- (2) We also have  $C \leq S_d \left( \frac{2G_d(\alpha)}{\alpha} + \frac{G_d(\alpha)}{1-\alpha} \right)$  when  $\alpha < 1$  in (3.7), since we have seen that this estimate is a simple rewriting of the preceding one by rescaling the time variable.
- (3) When  $\alpha \geq 1$ , Theorem 4.2 is not sufficient to establish the optimal estimate. Indeed, since  $\int_{|z| < \tilde{r}} |z| d\mu_\alpha(z) = +\infty$ , we can not send  $r$  to zero and have to optimize with respect to both  $r$  and  $\tilde{r}$ . The results are modulus of continuity of the order  $\|\varphi' - \psi'\|_\infty^{\frac{1}{\alpha}}$  for  $\alpha > 1$  and  $\|\varphi' - \psi'\|_\infty |\ln \|\varphi' - \psi'\|_\infty|$  for  $\alpha = 1$ . These modulus are worse than those in (3.5).

*Proof of Theorem 3.8 when  $\lambda \in (0, 1)$ .* Given  $\alpha, \beta \in (0, 2)$ , we use Theorem 4.2 with  $u = u^\alpha$  and  $v = u^\beta$ , i.e. with  $(u_0, f, \varphi) = (v_0, g, \psi)$ . As in the preceding proof, we pass to the limit as  $r \rightarrow 0$  in (4.4) and we cut the remaining integral in two parts. We find that

$$(5.1) \quad \|u^\alpha - u^\beta\|_{C([0, T]; L^1)} \leq 2M \|u_0\|_{L^1} \underbrace{\int_{|z| > \tilde{r}} d|\mu_\alpha - \mu_\beta|(z)}_{=: J_1} + M |u_0|_{BV} \underbrace{\int_{|z| < \tilde{r}} |z| d|\mu_\alpha - \mu_\beta|(z)}_{=: J_2}.$$

In the rest of the proof we use the letter  $C$  to denote various constants  $C = C(d, \lambda)$ .

We have

$$(5.2) \quad \begin{aligned} J_1 &= \int_{|z| > \tilde{r}} |G_d(\alpha) |z|^{-d-\alpha} - G_d(\beta) |z|^{-d-\beta}| dz \\ &\leq |G_d(\alpha) - G_d(\beta)| \max_{\sigma=\alpha, \beta} \int_{|z| > \tilde{r}} \frac{dz}{|z|^{d+\sigma}} \\ &\quad + (G_d(\alpha) \vee G_d(\beta)) \underbrace{\int_{|z| > \tilde{r}} \left| |z|^{-d-\alpha} - |z|^{-d-\beta} \right| dz}_{=: \tilde{J}_1}, \end{aligned}$$

where  $\tilde{J}_1 \leq S_d \left| \frac{\tilde{r}^{-\alpha}}{\alpha} - \frac{\tilde{r}^{-\beta}}{\beta} \right| + 2S_d \left| \frac{1}{\alpha} - \frac{1}{\beta} \right| \mathbf{1}_{\tilde{r} < 1}$ . We have estimated  $\tilde{J}_1$  using the fact that  $|z|^{-d-\alpha} - |z|^{-d-\beta}$  has a sign both inside and outside the unit ball. By (2.3) and a simple passage to the limit under the integral sign,

$$\limsup_{\alpha, \beta \rightarrow \lambda} \frac{J_1}{|\alpha - \beta|} \leq \underbrace{C(\tilde{r}^{-\lambda} + \mathbf{1}_{\tilde{r} < 1})}_{\leq C \tilde{r}^{-\lambda}} + C \limsup_{\alpha, \beta \rightarrow \lambda} \frac{1}{|\alpha - \beta|} \underbrace{\left| \frac{\tilde{r}^{-\alpha}}{\alpha} - \frac{\tilde{r}^{-\beta}}{\beta} \right|}_{=: \tilde{\tilde{J}}_1}.$$

By the Taylor formula with integral remainder,

$$\tilde{\tilde{J}}_1 = \lim_{\alpha, \beta \rightarrow \lambda} \left| \int_0^1 \frac{\alpha_\tau \tilde{r}^{-\alpha_\tau} \ln \tilde{r} + \tilde{r}^{-\alpha_\tau}}{\alpha_\tau^2} d\tau \right| \leq C \tilde{r}^{-\lambda} (1 + |\ln \tilde{r}|),$$

where  $\alpha_\tau := \tau \alpha + (1 - \tau) \beta$ . We deduce the following estimate:

$$(5.3) \quad \limsup_{\alpha, \beta \rightarrow \lambda} \frac{J_1}{|\alpha - \beta|} \leq C \tilde{r}^{-\lambda} (1 + |\ln \tilde{r}|).$$

Let us notice that this estimate works for all  $\lambda \in (0, 2)$ . By similar arguments, we also have

$$\limsup_{\alpha, \beta \rightarrow \lambda} \frac{J_2}{|\alpha - \beta|} \leq C \tilde{r}^{1-\lambda} (1 + |\ln \tilde{r}|),$$

but this time we have to use that  $\lambda < 1$ . Inserting these inequalities into (5.1), we find that for all  $\tilde{r} > 0$ ,

$$\text{Lip}_\alpha(u; \lambda) \leq C M (1 + |\ln \tilde{r}|) (\|u_0\|_{L^1} \tilde{r}^{-\lambda} + |u_0|_{BV} \tilde{r}^{1-\lambda}).$$

To conclude we take  $\tilde{r} = \|u_0\|_{L^1} |u_0|_{BV}^{-1}$ .  $\square$

*Remark 5.2.* The preceding proof implies that  $\alpha \in (0, 1) \rightarrow u^\alpha \in C([0, T]; L^1)$  is locally Lipschitz continuous. But, for the same reason as in Remark 5.1(3), Theorem 4.2 does not allow us to show the local Lipschitz continuity on  $[1, 2)$ .

## 6. CONTINUOUS DEPENDENCE IN THE CRITICAL AND SUBCRITICAL CASES

Since we can not use Theorem 4.2 any more, we start from Lemma 4.1 and take advantage of the homogeneity of the fractional Laplacian. We thus use the Kruzkov type doubling of variables techniques introduced in [29] along with ideas from [30]; see also [16, 25, 1, 12] or [3] for results on nonlocal equations. We recall that the idea is to consider  $v$  to be a function of  $(x, t)$ ,  $u$  to be a function of  $(y, s)$ , and use the approximate unit  $\phi^{\epsilon, \nu}(x, t, y, s)$  in (4.1) as a test function. For brevity, we do not specify the variables of  $u, v$ , and  $\phi^{\epsilon, \nu}$  when the context is clear. Finally, we recall that  $dw = dx dt dy ds$ .

**6.1. A technical lemma.** In order to adapt the ideas of [30] to the nonlocal case, we need the following Kato type of inequality. The reader could skip this technical subsection at the first reading.

**Lemma 6.1.** *Let  $\alpha \in (0, 2)$ ,  $c, \tilde{c} \in \mathbb{R}$ ,  $\gamma, \tilde{\gamma} \in \mathbb{R}$  and  $I$  be a real interval with a positive lower bound. Let  $u, v \in L^1(Q_T)$ ,  $\varphi$  satisfy (1.4) and  $\phi^{\epsilon, \nu}$  be the test function in (4.1). Then*

$$\begin{aligned} \mathcal{E} &:= \int_{Q_T^2} \int_{|z| \in I} \text{sgn}(v(x, t) - u(y, s)) \\ &\quad \cdot \frac{\{\varphi(v(x + \tilde{c}|z|^{\tilde{\gamma}-1}z, t)) - \varphi(u(y + c|z|^{\gamma-1}z, s))\} - \{\varphi(v(x, t)) - \varphi(u(y, s))\}}{|z|^{d+\alpha}} \\ &\quad \cdot \phi^{\epsilon, \nu}(x, t, y, s) dz dw \\ &\leq \int_{Q_T^2} \int_{|z| \in I} |\varphi(v(x, t)) - \varphi(u(y, s))| \theta_\nu(t-s) \frac{\rho_\epsilon(x-y+h(z)) - \rho_\epsilon(x-y)}{|z|^{d+\alpha}} dz dw, \end{aligned}$$

with  $h(z) := (\tilde{c}|z|^{\tilde{\gamma}-1} - c|z|^{\gamma-1})z$ . In particular, if  $c = \tilde{c}$  and  $\gamma = \tilde{\gamma}$ , then  $\mathcal{E} \leq 0$ .

*Proof.* Note that  $\mathcal{E}$  is well-defined as ‘‘convolution-like integral of  $L^1$ -functions’’. Indeed,  $\phi^{\epsilon, \nu}(x, t, y, s) = \theta_\nu(t-s) \rho_\epsilon(x-y)$ , where  $\theta_\nu$  and  $\rho_\epsilon$  are approximate units,

so that by Fubini,

$$\begin{aligned} & \int_{Q_T^2} \int_{|z| \in I} \phi^{\epsilon, \nu} \\ & \cdot \left| \frac{\{\varphi(v(x + \tilde{c}|z|^{\tilde{\gamma}-1}z, t)) - \varphi(u(y + c|z|^{\gamma-1}z, s))\} - \{\varphi(v) - \varphi(u)\}}{|z|^{d+\alpha}} \right| dz dw \\ & \leq 2 (\|\varphi(u)\|_{L^1(Q_T)} + \|\varphi(v)\|_{L^1(Q_T)}) \int_{|z| \in I} \frac{dz}{|z|^{d+\alpha}} < +\infty, \end{aligned}$$

since  $u$  and  $v$  are  $L^\infty \cap L^1$ ,  $\varphi$  is  $W_{loc}^{1, \infty}$  with  $\varphi(0) = 0$ , and  $\inf I > 0$ .

Then by (2.5) and the nonnegativity of  $\phi^{\epsilon, \nu}$ ,

$$\begin{aligned} \mathcal{E} & \leq \int_{Q_T^2} \int_{|z| \in I} \phi^{\epsilon, \nu} \\ & \quad \frac{|\varphi(v(x + \tilde{c}|z|^{\tilde{\gamma}-1}z, t)) - \varphi(u(y + c|z|^{\gamma-1}z, s))| - |\varphi(v) - \varphi(u)|}{|z|^{d+\alpha}} dz dw \\ & = \int_{Q_T^2} \int_{|z| \in I} |\varphi(v) - \varphi(u)| \\ & \quad \underbrace{\{\phi^{\epsilon, \nu}(x + \tilde{c}|z|^{\tilde{\gamma}-1}z, t, y + c|z|^{\gamma-1}z, s) - \phi^{\epsilon, \nu}\}}_{=\theta_\nu(t-s)\{\rho_\epsilon(x-y+(\tilde{c}|z|^{\tilde{\gamma}-1}-c|z|^{\gamma-1})z) - \rho_\epsilon(x-y)\}} \frac{dz}{|z|^{d+\alpha}} dw; \end{aligned}$$

the last line has been obtained by splitting the integral in two pieces and using the change of variable  $(x + \tilde{c}|z|^{\tilde{\gamma}-1}z, t, y + c|z|^{\gamma-1}z, s, -z) \rightarrow (x, t, y, s, z)$ . The proof is complete.  $\square$

**6.2. Proof of Theorem 3.1.** During the proof we freeze the nonlinear diffusion functions and use a sort of linearization procedure. The techniques could look a little bit like the ones in Young measure theory and kinetic formulations [32, 8].

*Proof of Theorem 3.1.*

**1. Initial reduction.** We first reduce the proof to the case where

$$(6.1) \quad \begin{cases} v_0 = u_0, \\ \varphi' \text{ and } \psi' \text{ vanish outside } I(u_0) \text{ and take values in } (\underline{\Lambda}, \bar{\Lambda}), \end{cases}$$

with  $I(u_0) = (\text{ess inf } u_0, \text{ess sup } u_0)$  and for some  $\bar{\Lambda} \geq \underline{\Lambda} > 0$ . Let us justify that we can do this without loss of generality.

Since  $u$  takes its values in  $I(u_0)$  by (2.7), we can redefine  $\varphi$  to be constant outside this interval without changing the solutions of the initial value problem (1.1). Hence  $\bar{\Lambda}$  could be taken as a Lipschitz constant of  $\varphi$  on  $I(u_0)$ . In a similar way, we could also modify  $\psi$  outside  $I(u_0)$  if  $v_0 = u_0$ . The last assumption is no restriction. Indeed, by (2.8),

$$\|u - v\|_{C([0, T]; L^1)} \leq \|u - w\|_{C([0, T]; L^1)} + \underbrace{\|w - v\|_{C([0, T]; L^1)}}_{\leq \|u_0 - v_0\|_{L^1}}$$

for the entropy solution  $w$  of (3.1) with initial data  $u_0$ ; hence, (3.5) of Theorem 3.1 holds for  $u - v$  whenever it does for  $u - w$ . Finally, if  $\underline{\Lambda}$  does not exist, we can always consider sequences  $\varphi_n(\xi) := \varphi(\xi) + \frac{\xi}{n}$  and  $\psi_n(\xi) := \psi(\xi) + \frac{\xi}{n}$  for which it does. The associated entropy solutions  $u_n$  and  $v_n$  respectively converge to  $u$  and  $v$  in  $C([0, T]; L^1)$  by e.g. Theorem 4.2. Consequently, if we could prove (3.5) for  $u_n - v_n$ , it would follow for  $u - v$  by going to the limit.

In the rest of the proof we always assume (6.1).

**2. Applying Kuznetsov.** Let us use the entropy inequality (2.6) for  $v = v(x, t)$  with  $k = u(y, s)$  fixed and  $\phi(x, t) := \phi^{\epsilon, \nu}(x, t, y, s)$ . By Remark 2.1 and an integration of  $(y, s)$  over  $Q_T$ , we find that

$$\begin{aligned} & \int_{Q_T^2} |v - u| \partial_t \phi^{\epsilon, \nu} + q_g(v, u) \cdot \nabla_x \phi^{\epsilon, \nu} \, dw \\ & + \int_{Q_T^2} |\psi(v) - \psi(u)| \mathcal{L}_r^\alpha[\phi^{\epsilon, \nu}(\cdot, t, y, s)](x) \, dw \\ & + \int_{Q_T^2} \operatorname{sgn}(v - u) \mathcal{L}^{\alpha, r}[\psi(v(\cdot, t))](x) \phi^{\epsilon, \nu} \, dw \\ & - \int_{\mathbb{R}^d \times Q_T} |v(x, T) - u(y, t)| \phi^{\epsilon, \nu}(x, T, y, s) \, dx \, dy \, ds \\ & + \int_{\mathbb{R}^d \times Q_T} |v_0(x) - u(y, s)| \phi^{\epsilon, \nu}(x, 0, y, s) \, dx \, dy \, ds \geq 0. \end{aligned}$$

Inserting this inequality into the Kuznetsov inequality (4.3), we obtain for all  $r, \epsilon > 0$  and  $T > \nu > 0$ ,

$$\begin{aligned} (6.2) \quad & \|u(\cdot, T) - v(\cdot, T)\|_{L^1} \\ & \leq C(d) |u_0|_{BV} \epsilon + 2(m_u(\nu) \vee m_v(\nu)) \\ & \quad + \underbrace{\int_{Q_T^2} (q_g - q_f)(v, u) \cdot \nabla_x \phi^{\epsilon, \nu} \, dw}_{=:\mathcal{E}_1} \\ & \quad + \underbrace{\int_{Q_T^2} |\psi(v) - \psi(u)| \mathcal{L}_r^\alpha[\phi^{\epsilon, \nu}(\cdot, t, y, s)](x) + |\varphi(v) - \varphi(u)| \mathcal{L}_r^\alpha[\phi^{\epsilon, \nu}(x, t, \cdot, s)](y) \, dw}_{=:\mathcal{E}_2} \\ & \quad + \underbrace{\int_{Q_T^2} \operatorname{sgn}(v - u) (\mathcal{L}^{\alpha, r}[\psi(v(\cdot, t))](x) - \mathcal{L}^{\alpha, r}[\varphi(u(\cdot, s))](y)) \phi^{\epsilon, \nu} \, dw}_{=:\mathcal{E}_3} \end{aligned}$$

where  $C(d) = C_\rho$  from (4.3). During the proof,  $C(d)$  will denote various constant depending only on  $d$ .

**3. Estimates of  $\mathcal{E}_1$  and  $\mathcal{E}_2$ .** A standard estimate shows that

$$(6.3) \quad \mathcal{E}_1 \leq T |u_0|_{BV} \|f' - g'\|_\infty,$$

see e.g. [30, 18]. Let us estimate  $\mathcal{E}_2$ . By Taylor's formula,

$$\rho_\epsilon(x + z) - \rho_\epsilon(x) - \nabla \rho_\epsilon(x) \cdot z = \int_0^1 (1 - \tau) \nabla^2 \rho_\epsilon(x + \tau z) \cdot z^2 \, d\tau$$

for all  $x, z \in \mathbb{R}^d$ . Since  $\rho_\epsilon \in C_c^\infty(\mathbb{R}^d)$ , we infer that  $\mathcal{L}_r^\alpha[\rho_\epsilon] \in L^1(\mathbb{R}^d)$  with

$$\begin{aligned} \|\mathcal{L}_r^\alpha[\rho_\epsilon]\|_{L^1} & \leq G_d(\alpha) \int_{|z| < r} \int_0^1 (1 - \tau) |z|^{-d+2-\alpha} \int_{\mathbb{R}^d} |\nabla^2 \rho_\epsilon(x + \tau z)| \, dx \, d\tau \, dz \\ & = C(d, \alpha, \epsilon) r^{2-\alpha}. \end{aligned}$$

Moreover, by Definitions (2.1) and (4.1),

$$\mathcal{L}_r^\alpha[\phi^{\epsilon, \nu}(\cdot, t, y, s)](x) = \theta_\nu(t - s) \mathcal{L}_r^\alpha[\rho_\epsilon](x - y).$$

By Fubini and the convolution like structure of the integral, it follows that

$$\begin{aligned} & \int_{Q_T^2} |\psi(v(x, t)) - \psi(u(y, s))| \mathcal{L}_r^\alpha[\phi^{\epsilon, \nu}(\cdot, y, t, s)](x) \, dw \\ & \leq (\|\psi(v)\|_{L^1(Q_T)} + \|\psi(u)\|_{L^1(Q_T)}) C(d, \alpha, \epsilon) r^{2-\alpha}, \end{aligned}$$

since  $\int \theta_\nu = 1$ . In a similar way we can estimate the  $\varphi$ -integral and conclude that

$$(6.4) \quad \mathcal{E}_2 \leq C_\epsilon r^{2-\alpha}.$$

From now on  $C_\epsilon$  will denote various constants depending among other things on  $\epsilon$ , but not on  $r, \nu$ . For later use we note that  $\mathcal{E}_2 \rightarrow 0$  as  $r, \nu \rightarrow 0$  and  $\epsilon$  is fixed.

**4. Estimate of  $\mathcal{E}_3$  – the linear case.** We consider the case  $\varphi' \equiv a$  and  $\psi' \equiv b$  for  $a, b > 0$ . In this case

$$(6.5) \quad \begin{aligned} \mathcal{E}_3 &= G_d(\alpha) \int_{Q_T^2} \int_{|z|>r} \operatorname{sgn}(v-u) \phi^{\epsilon, \nu} \\ & \quad \cdot \frac{a(v(x+z, t) - v) - b(u(y+z, s) - u)}{|z|^{d+\alpha}} \, dz \, dw. \end{aligned}$$

By the change of variables  $z \rightarrow b^{\frac{1}{\alpha}} z$ , we see that

$$\begin{aligned} b \mathcal{L}^{\alpha, r}[v(\cdot, t)](x) &= G_d(\alpha) \int_{|z|>r} \frac{v(x+z, t) - v(x, t)}{|b^{-\frac{1}{\alpha}} z|^{d+\alpha}} b^{-\frac{d}{\alpha}} \, dz \\ &= G_d(\alpha) \int_{|z|>b^{-\frac{1}{\alpha}} r} \frac{v(x+b^{\frac{1}{\alpha}} z, t) - v(x, t)}{|z|^{d+\alpha}} \, dz, \end{aligned}$$

and similarly that

$$a \mathcal{L}^{\alpha, r}[u(\cdot, s)](y) = G_d(\alpha) \int_{|z|>a^{-\frac{1}{\alpha}} r} \frac{u(y+a^{\frac{1}{\alpha}} z, s) - u(y, s)}{|z|^{d+\alpha}} \, dz.$$

It follows that

$$(6.6) \quad \begin{aligned} \mathcal{E}_3 &= G_d(\alpha) \int_{Q_T^2} \int_{(a\nu b)^{-\frac{1}{\alpha}} r < |z| < (a\wedge b)^{-\frac{1}{\alpha}} r} \cdots \frac{dz}{|z|^{d+\alpha}} \\ & \quad + G_d(\alpha) \int_{Q_T^2} \int_{|z| > (a\wedge b)^{-\frac{1}{\alpha}} r} \operatorname{sgn}(v-u) \\ & \quad \cdot \frac{(v(x+b^{\frac{1}{\alpha}} z, t) - u(y+a^{\frac{1}{\alpha}} z, s)) - (v-u)}{|z|^{d+\alpha}} \phi^{\epsilon, \nu} \, dz \, dw \\ & =: \mathcal{E}_{3,1} + \mathcal{E}_{3,2}, \end{aligned}$$

where  $\mathcal{E}_{3,1}$  contains only the  $u$ -terms if  $a \geq b$ , or only the  $v$ -terms in the other case. In the  $u$ -case, e.g.,

$$\mathcal{E}_{3,1} = G_d(\alpha) \int_{Q_T^2} \int_{a^{-\frac{1}{\alpha}} r < |z| < b^{-\frac{1}{\alpha}} r} \operatorname{sgn}(u-v) \frac{u(y+a^{\frac{1}{\alpha}} z, s) - u}{|z|^{d+\alpha}} \phi^{\epsilon, \nu} \, dz \, dw.$$

The estimates for  $\mathcal{E}_{3,1}$  are similar in both cases, and we only detail the  $u$ -case. As in the proof of Lemma 6.1, we use that

$$\begin{aligned} & \operatorname{sgn}(u(y, s) - v(x, t)) \left( u(y+a^{\frac{1}{\alpha}} z, s) - u(y, s) \right) \\ & \leq \left| u(y+a^{\frac{1}{\alpha}} z, s) - v(x, t) \right| - |u(y, s) - v(x, t)|, \end{aligned}$$



to deduce that

$$\begin{aligned}
\mathcal{E}_{3,1} &\leq G_d(\alpha) \int_{Q_T^2} \int_{a^{-\frac{1}{\alpha}} r < |z| < b^{-\frac{1}{\alpha}} r} \frac{|u(y + a^{\frac{1}{\alpha}} z, s) - v(x, t)| - |u - v|}{|z|^{d+\alpha}} \phi^{\epsilon, \nu} dz dw \\
&= G_d(\alpha) \int_{Q_T^2} |u - v| \\
&\quad \cdot \int_{a^{-\frac{1}{\alpha}} r < |z| < b^{-\frac{1}{\alpha}} r} \underbrace{\left( \phi^{\epsilon, \nu}(x, t, y + a^{\frac{1}{\alpha}} z, s) - \phi^{\epsilon, \nu} \right)}_{=\theta_\nu(t-s)(\rho_\epsilon(x-y-a^{\frac{1}{\alpha}} z) - \rho_\epsilon(x-y))} |z|^{-d-\alpha} dz dw.
\end{aligned}$$

We continue as in the derivation of (6.4), and use a Taylor expansion with integral remainder of  $\rho_\epsilon$ . Since the first order term contains the factor

$$\int_{a^{-\frac{1}{\alpha}} r < |z| < b^{-\frac{1}{\alpha}} r} \frac{z}{|z|^{d+\alpha}} dz = 0,$$

we find an estimate similar to (6.4), namely

$$(6.7) \quad \mathcal{E}_{3,1} \leq C_\epsilon (\|u\|_{L^1(Q_T)} + \|v\|_{L^1(Q_T)}) r^{2-\alpha}.$$

We emphasize that  $C_\epsilon$  can be chosen to be independent of  $a$  and  $b$  by (6.1) (more precisely  $C_\epsilon = C(d, \alpha, \epsilon, \underline{\Delta}, \overline{\Delta})$ ); this will be important in the next step.

**5. Estimate of  $\mathcal{E}_{3,2}$ .** Note that  $a, b$  are arbitrary numbers such that (6.1) holds, i.e.  $\overline{\Delta} \geq a, b \geq \underline{\Delta}$ , and let  $r_2 \geq r_1 > 0$ . Since  $\underline{\Delta} > 0$  and  $r$  will be sent to 0, we assume without loss of generality that  $r_1 > \underline{\Delta}^{-\frac{1}{\alpha}} r$ . In particular,  $r_1 > (a \wedge b)^{-\frac{1}{\alpha}} r$ . Then

$$\begin{aligned}
\mathcal{E}_{3,2} &= \sum_{i=1}^3 G_d(\alpha) \int_{Q_T^2} \int_{|z| \in I_i} \operatorname{sgn}(v - u) \\
(6.8) \quad &\quad \cdot \frac{\left( v(x + b^{\frac{1}{\alpha}} z, t) - u(y + a^{\frac{1}{\alpha}} z, s) \right) - (v - u)}{|z|^{d+\alpha}} \phi^{\epsilon, \nu} dz dw \\
&=: \sum_{i=1}^3 \mathcal{E}_{3,2,i},
\end{aligned}$$

where  $I_1 = (r_2, +\infty)$ ,  $I_2 = (r_1, r_2)$  and  $I_3 = ((a \wedge b)^{-\frac{1}{\alpha}} r, r_1)$ .

By adding and subtracting  $\operatorname{sgn}(v - u) u(y + b^{\frac{1}{\alpha}} z, s)$  and using Lemma 6.1 with  $c = \tilde{c} = b^{\frac{1}{\alpha}}$  and  $\gamma = \tilde{\gamma} = 1$ , we find that

$$(6.9) \quad \mathcal{E}_{3,2,i} \leq G_d(\alpha) \int_{Q_T^2} \int_{|z| \in I_i} \operatorname{sgn}(v - u) \frac{u(y + b^{\frac{1}{\alpha}} z, s) - u(y + a^{\frac{1}{\alpha}} z, s)}{|z|^{d+\alpha}} \phi^{\epsilon, \nu} dz dw.$$

By the  $BV$ -regularity of  $u$ , we then immediately deduce that

$$\mathcal{E}_{3,2,2} \leq G_d(\alpha) |u|_{L^1(0,T;BV)} |a^{\frac{1}{\alpha}} - b^{\frac{1}{\alpha}}| \int_{r_1 < |z| < r_2} \frac{|z| dz}{|z|^{d+\alpha}}.$$

Moreover, going back to the original variables  $a^{\frac{1}{\alpha}} z \rightarrow z$  and  $b^{\frac{1}{\alpha}} z \rightarrow z$ , we find that

$$\begin{aligned}
&\int_{Q_T^2} \int_{|z| > r_2} \operatorname{sgn}(v - u) \frac{u(y + a^{\frac{1}{\alpha}} z, s)}{|z|^{d+\alpha}} \phi^{\epsilon, \nu} dz dw \\
&= a \int_{Q_T^2} \int_{|z| > a^{\frac{1}{\alpha}} r_2} \operatorname{sgn}(v - u) \frac{u(y + z, s)}{|z|^{d+\alpha}} \phi^{\epsilon, \nu} dz dw,
\end{aligned}$$

and a similar formula for the  $b$ -term. Hence we find that

$$\begin{aligned} \mathcal{E}_{3,2,1} &\leq G_d(\alpha) (b-a) \int_{Q_T^2} \int_{|z| > (a \vee b)^{\frac{1}{\alpha}} r_2} \operatorname{sgn}(v-u) \frac{u(y+z, s)}{|z|^{d+\alpha}} \phi^{\epsilon, \nu} dz dw \\ &\quad + G_d(\alpha) \operatorname{sgn}(a-b) (a \wedge b) \int_{Q_T^2} \int_{(a \wedge b)^{\frac{1}{\alpha}} r_2 < |z| < (a \vee b)^{\frac{1}{\alpha}} r_2} \dots, \end{aligned}$$

where the integrands are the same. Since  $\phi^{\epsilon, \nu}$  is an approximate unit,

$$\mathcal{E}_{3,2,1} \leq C(d) \frac{G_d(\alpha)}{\alpha} \|u\|_{L^1(Q_T)} \frac{|a-b|}{a \vee b} r_2^{-\alpha},$$

where  $C(d) = 2S_d$ .

It remains to estimate  $\mathcal{E}_{3,2,3}$  in (6.8). By Lemma 6.1, with  $c = a^{\frac{1}{\alpha}}$  and  $\tilde{c} = b^{\frac{1}{\alpha}}$ ,

$$(6.10) \quad \begin{aligned} \mathcal{E}_{3,2,3} &\leq G_d(\alpha) \int_{Q_T^2} \int_{(a \wedge b)^{-\frac{1}{\alpha}} r < |z| < r_1} |v-u| \theta_\nu(t-s) \\ &\quad \cdot \{\rho_\epsilon(x-y+h(z)) - \rho_\epsilon(x-y)\} |z|^{-d-\alpha} dz dw \end{aligned}$$

with  $h(z) := (b^{\frac{1}{\alpha}} - a^{\frac{1}{\alpha}})z$ . After a Taylor expansion of  $\rho_\epsilon$  with integral remainder, we find that

$$\begin{aligned} \mathcal{E}_{3,2,3} &\leq G_d(\alpha) \int_{Q_T^2} \int_{(a \wedge b)^{-\frac{1}{\alpha}} r < |z| < r_1} \int_0^1 (1-\tau) |v-u| \theta_\nu(t-s) |z|^{-d-\alpha} \\ &\quad \cdot \nabla^2 \rho_\epsilon(x-y+\tau h(z)) \cdot h(z)^2 d\tau dz dw. \end{aligned}$$

Remember that the integral of the first order term in  $z$  is zero by symmetry. By a standard argument,  $|v-u|$  is  $BV$  in  $y$  as composition of a  $BV$  with a Lipschitz function (cf. e.g. [8]). Hence, by an integration by parts with respect to  $y$ ,

$$\begin{aligned} \mathcal{E}_{3,2,3} &\leq G_d(\alpha) \int_0^T \int_{Q_T} \int_{(a \wedge b)^{-\frac{1}{\alpha}} r < |z| < r_1} \int_0^1 (1-\tau) \theta_\nu(t-s) |z|^{-d-\alpha} \\ &\quad \cdot \left\{ \int_{\mathbb{R}^d} \nabla \rho_\epsilon(x-y+\tau h(z)) \cdot h(z) h(z) \cdot d\nabla_y |v(x, t) - u(\cdot, s)|(y) \right\} \\ &\quad d\tau dz dx dt ds. \end{aligned}$$

We use the notation  $d\nabla_y |v(x, t) - u(\cdot, s)|(y)$  in case  $\nabla_y |v-u|$  is a measure. Then  $|\nabla_y |v-u|| \leq |\nabla u|$  in the sense of measures since  $y$  is the space variable of  $u$ . It follows that

$$\begin{aligned} \mathcal{E}_{3,2,3} &\leq G_d(\alpha) \int_0^T \int_{Q_T} \int_{|z| < r_1} \int_0^1 (1-\tau) \theta_\nu(t-s) |z|^{-d-\alpha} |h(z)|^2 \\ &\quad \cdot \left\{ \int_{\mathbb{R}^d} |\nabla \rho_\epsilon(x-y+\tau h(z))| d|\nabla u(\cdot, s)|(y) \right\} d\tau dz dx dt ds. \end{aligned}$$

By Fubini<sup>3</sup> we integrate with respect to  $(x, t)$  before  $(y, s)$ , and then we use that  $h(z) = (b^{\frac{1}{\alpha}} - a^{\frac{1}{\alpha}})z$  and  $\int |\nabla \rho_\epsilon| = \frac{1}{\epsilon} \int |\nabla \rho| = \frac{C(d)}{\epsilon}$  (by (4.1)), to see that

$$(6.11) \quad \begin{aligned} \mathcal{E}_{3,2,3} &\leq G_d(\alpha) \int_0^T \int_{|z| < r_1} \int_0^1 (1-\tau) |z|^{-d-\alpha} \\ &\quad \cdot |h(z)|^2 |u(\cdot, s)|_{BV} d\tau dz ds \int_{Q_T} \theta_\nu |\nabla \rho_\epsilon| dx dt \\ &\leq C(d) \frac{G_d(\alpha)}{2-\alpha} |u|_{L^1(0, T; BV)} (a^{\frac{1}{\alpha}} - b^{\frac{1}{\alpha}})^2 \frac{r_1^{2-\alpha}}{\epsilon}. \end{aligned}$$

<sup>3</sup>applied for fixed  $s$ , so that  $d|\nabla u(\cdot, s)|(y) dz dx dt$  is a tensor product of  $\sigma$ -finite measures!

**6.** *Estimate of  $\mathcal{E}_3$  – conclusion in the linear case.* By the estimates of part 4 and 5, (6.6), (6.8), ..., we can then conclude that

$$\begin{aligned}
(6.12) \quad \mathcal{E}_3 &\leq \mathcal{E}_{3,1} + \mathcal{E}_{3,2,1} + \mathcal{E}_{3,2,2} + \mathcal{E}_{3,2,3} \\
&\leq C_\epsilon (\|u\|_{L^1(Q_T)} + \|v\|_{L^1(Q_T)}) r^{2-\alpha} \\
&\quad + C(d) G_d(\alpha) \left\{ \frac{1}{\alpha} \|u\|_{L^1(Q_T)} \frac{|a-b|}{a \vee b} r_2^{-\alpha} \right. \\
&\quad + |u|_{L^1(0,T;BV)} |a^{\frac{1}{\alpha}} - b^{\frac{1}{\alpha}}| \int_{r_1 < |z| < r_2} \frac{|z| dz}{|z|^{d+\alpha}} \\
&\quad \left. + \frac{1}{2-\alpha} |u|_{L^1(0,T;BV)} (a^{\frac{1}{\alpha}} - b^{\frac{1}{\alpha}})^2 \frac{r_1^{2-\alpha}}{\epsilon} \right\},
\end{aligned}$$

for arbitrary  $r_2 \geq r_1 > \underline{\Delta}^{-\frac{1}{\alpha}} r$ . Note that the  $\frac{1}{a \vee b}$ -term has to be handled with care since it could be large in the general case when  $\varphi'$  and  $\psi'$  can be degenerate.

We conclude the estimate of  $\mathcal{E}_3$  by choosing the values of constants  $r_1$  and  $r_2$ . In the critical case where  $\alpha = 1$ , we take  $r_1 = T \wedge 1$  and  $r_2 = 1 \vee \frac{\|u_0\|_{L^1}}{(a \vee b) |u_0|_{BV}}$ . Notice that if  $|u_0|_{BV} = 0$ , then  $u_0 \equiv 0$  as constant integrable function, and (3.5) reduces to (2.7). In the sequel, we thus assume without loss of generality that  $|u_0|_{BV} \neq 0$ . Note then that  $+\infty > r_2 \geq r_1 = T \wedge 1 > \underline{\Delta}^{-\frac{1}{\alpha}} r$  for  $r$  small enough ( $r \rightarrow 0$  in the end). By easy computation and Lemma B.1 of the Appendix,

$$\begin{aligned}
|a-b| \int_{r_1 < |z| < r_2} \frac{|z| dz}{|z|^{d+1}} &= C |a-b| (\ln r_2 - \ln r_1) \\
&\leq C |a-b| \left\{ |\ln T| + \mathbf{1}_{\frac{\|u_0\|_{L^1}}{|u_0|_{BV}} > 1} \ln \frac{\|u_0\|_{L^1}}{|u_0|_{BV}} + (-\ln(a \vee b))^+ \right\} \\
&\leq C \left\{ \left(1 + |\ln T| + |u_0|_{BV}^{-1} E_1(u_0)\right) |a-b| + |a \ln a - b \ln b| \right\},
\end{aligned}$$

where  $C = C(d)$  and where  $E_1(u_0)$  is defined in (3.3). We finally deduce from (6.12) that, when  $\alpha = 1$ ,

$$\begin{aligned}
(6.13) \quad \mathcal{E}_3 &\leq C_\epsilon (\|u\|_{L^1(Q_T)} + \|v\|_{L^1(Q_T)}) r \\
&\quad + C(d) \left\{ \|u\|_{L^1(Q_T)} \frac{|u_0|_{BV}}{\|u_0\|_{L^1}} |a-b| \right. \\
&\quad + (1 + |\ln T| + |u_0|_{BV}^{-1} E_1(u_0)) |u|_{L^1(0,T;BV)} |a-b| \\
&\quad + |u|_{L^1(0,T;BV)} |a \ln a - b \ln b| \\
&\quad \left. + T |u|_{L^1(0,T;BV)} (a-b)^2 \frac{1}{\epsilon} \right\},
\end{aligned}$$

for all  $T \wedge 1 > \underline{\Delta}^{-1} r$ . To divide by  $\|u_0\|_{L^1}$ , we have assumed without loss of generality that we are not in the case where  $\|u_0\|_{L^1} = 0$ , for which (3.5) also reduces to (2.7).

When  $\alpha > 1$ , we simply choose  $r_2 = +\infty$  in (6.12) and we get

$$\begin{aligned}
(6.14) \quad \mathcal{E}_3 &\leq C_\epsilon (\|u\|_{L^1(Q_T)} + \|v\|_{L^1(Q_T)}) r^{2-\alpha} \\
&\quad + C(d) G_d(\alpha) \left\{ \frac{1}{\alpha-1} |u|_{L^1(0,T;BV)} |a^{\frac{1}{\alpha}} - b^{\frac{1}{\alpha}}| r_1^{1-\alpha} \right. \\
&\quad \left. + \frac{1}{2-\alpha} |u|_{L^1(0,T;BV)} (a^{\frac{1}{\alpha}} - b^{\frac{1}{\alpha}})^2 \frac{r_1^{2-\alpha}}{\epsilon} \right\},
\end{aligned}$$

for all  $r_1 > \underline{\Delta}^{-\frac{1}{\alpha}} r$ .

**7.** *Estimate of  $\mathcal{E}_3$  - the general case via linearization.* The idea is now to reduce to the linear case in step 4 by freezing the ‘‘diffusion coefficients’’  $\varphi'(\xi)$  and  $\psi'(\xi)$ . To

do so, we introduce the function

$$(6.15) \quad \chi_a^b(\xi) := \operatorname{sgn}(b - a) \mathbf{1}_{(a \wedge b, a \vee b)}(\xi),$$

for  $\xi, a, b \in \mathbb{R}$ . By (6.2) and the fundamental theorem of calculus, we then find that

$$(6.16) \quad \begin{aligned} \mathcal{E}_3 &= G_d(\alpha) \int_{Q_T^2} \int_{|z|>r} \operatorname{sgn}(v - u) \\ &\quad \cdot \frac{\int_{v(x,t)}^{v(x+z,t)} \psi'(\xi) \, d\xi - \int_{u(y,s)}^{u(y+z,s)} \varphi'(\xi) \, d\xi}{|z|^{d+\alpha}} \phi^{\epsilon,\nu} \, dz \, dw \\ &= G_d(\alpha) \int_{Q_T^2} \int_{|z|>r} \int \operatorname{sgn}(v - u) \\ &\quad \cdot \frac{\chi_{v(x,t)}^{v(x+z,t)}(\xi) \psi'(\xi) - \chi_{u(y,s)}^{u(y+z,s)}(\xi) \varphi'(\xi)}{|z|^{d+\alpha}} \phi^{\epsilon,\nu} \, d\xi \, dz \, dw. \end{aligned}$$

Let us notice that this integral is well-defined, since  $\int |\chi_a^b(\xi)| \, d\xi = |b - a|$  and,  $\varphi'$  and  $\psi'$  are assumed bounded by (6.1).

For each  $\delta > 0$ , we define a regularized version of  $\mathcal{E}_3$  as

$$(6.17) \quad \begin{aligned} \mathcal{E}_3(\delta) &:= G_d(\alpha) \int_{Q_T^2} \int_{|z|>r} \int \int \operatorname{sgn}(v - u) \\ &\quad \cdot \frac{\chi_{v(x,t)}^{v(x+z,t)}(\zeta) \psi'(\xi) - \chi_{u(y,s)}^{u(y+z,s)}(\zeta) \varphi'(\xi)}{|z|^{d+\alpha}} \phi^{\epsilon,\nu} \omega_\delta(\xi - \zeta) \, d\zeta \, d\xi \, dz \, dw, \end{aligned}$$

where the approximate unit  $\omega_\delta(\xi) := \frac{1}{\delta} \omega\left(\frac{\xi}{\delta}\right)$ , and

$$(6.18) \quad \omega \in C_b^\infty \cap L^1(\mathbb{R}), \quad \omega > 0, \quad \int \omega = 1.$$

For each  $\zeta, \xi \in \mathbb{R}$ , let  $\Omega_\xi(\zeta) := \int_{-\infty}^\zeta \omega_\delta(\xi - w) \, dw - \int_{-\infty}^0 \omega_\delta(\xi - w) \, dw$ , and note that

$$\int \chi_{v(x,t)}^{v(x+z,t)}(\zeta) \omega_\delta(\xi - \zeta) \, d\zeta = \int_{v(x,t)}^{v(x+z,t)} \Omega'_\xi(\zeta) \, d\zeta = \Omega_\xi(v(x+z,t)) - \Omega_\xi(v(x,t)).$$

Moreover,  $\operatorname{sgn}(v - u) = \operatorname{sgn}(\Omega_\xi(v) - \Omega_\xi(u))$  since  $\Omega_\xi(\cdot)$  is increasing, and since  $\Omega_\xi(\cdot)$  is smooth and vanishes at zero,  $\Omega_\xi(u)$  and  $\Omega_\xi(v)$  have similar boundedness, integrability, and regularity properties as  $u$  and  $v$ . It follows that

$$\begin{aligned} \mathcal{E}_3(\delta) &= G_d(\alpha) \int \int_{Q_T^2} \int_{|z|>r} \operatorname{sgn}(\Omega_\xi(v) - \Omega_\xi(u)) \phi^{\epsilon,\nu} \\ &\quad \cdot \frac{\psi'(\xi) (\Omega_\xi(v(x+z,t)) - \Omega_\xi(v)) - \varphi'(\xi) (\Omega_\xi(u(y+z,s)) - \Omega_\xi(u))}{|z|^{d+\alpha}} \, dz \, dw \, d\xi. \end{aligned}$$

This integrand has similar form and properties as the one in (6.5) for fixed  $\xi$ !

We continue in the critical case when  $\alpha = 1$ . We argue as in step 4 with  $a = \varphi'(\xi)$  and  $b = \psi'(\xi)$ . By (6.13) we get that for all  $T \wedge 1 > \underline{\Lambda}^{-1} r$ ,

$$\begin{aligned}
\mathcal{E}_3(\delta) &\leq \int C_\epsilon (\|\Omega_\xi(u)\|_{L^1(Q_T)} + \|\Omega_\xi(v)\|_{L^1(Q_T)}) r \, d\xi \\
&\quad + C(d) \int \left\{ \|\Omega_\xi(u)\|_{L^1(Q_T)} \frac{|u_0|_{BV}}{\|u_0\|_{L^1}} |\varphi'(\xi) - \psi'(\xi)| \right. \\
&\quad + (1 + |\ln T| + |u_0|_{BV}^{-1} E_1(u_0)) |\Omega_\xi(u)|_{L^1(0,T;BV)} |\varphi'(\xi) - \psi'(\xi)| \\
&\quad + |\Omega_\xi(u)|_{L^1(0,T;BV)} |\varphi'(\xi) \ln \varphi'(\xi) - \psi'(\xi) \ln \psi'(\xi)| \\
&\quad \left. + T |\Omega_\xi(u)|_{L^1(0,T;BV)} (\varphi'(\xi) - \psi'(\xi))^2 \frac{1}{\epsilon} \right\} d\xi \\
&\leq C_\epsilon r \int \|\Omega_\xi(u)\|_{L^1(Q_T)} + \|\Omega_\xi(v)\|_{L^1(Q_T)} \, d\xi \\
&\quad + C(d) \left\{ A \frac{|u_0|_{BV}}{\|u_0\|_{L^1}} \|\varphi' - \psi'\|_\infty \right. \\
&\quad + (1 + |\ln T| + |u_0|_{BV}^{-1} E_1(u_0)) B \|\varphi' - \psi'\|_\infty \\
&\quad + B \|\varphi' \ln \varphi' - \psi' \ln \psi'\|_\infty \\
&\quad \left. + T B \|\varphi' - \psi'\|_\infty^2 \frac{1}{\epsilon} \right\},
\end{aligned}$$

with  $A = \int \|\Omega_\xi(u)\|_{L^1(Q_T)} \, d\xi$ ,  $B = \int |\Omega_\xi(u)|_{L^1(0,T;BV)} \, d\xi$ , and

$$\|\varphi' - \psi'\|_\infty = \operatorname{ess\,sup}_{I(u_0)} |\varphi' - \psi'|.$$

The supremum above can be taken only on  $I(u_0)$ , since  $\varphi'$  and  $\psi'$  are assumed to vanish outside this interval by (6.1). Note also that  $C_\epsilon = C(d, \alpha, \epsilon, \underline{\Lambda}, \overline{\Lambda})$  can be chosen independent of  $\varphi'(\xi)$  and  $\psi'(\xi)$  as discussed below (6.7). A standard argument, see Appendix A, then reveals that

$$(6.19) \quad \int \|\Omega_\xi(u)\|_{L^1(Q_T)} \, d\xi = \|u\|_{L^1(Q_T)},$$

$$(6.20) \quad \int |\Omega_\xi(u)|_{L^1(0,T;BV)} \, d\xi = |u|_{L^1(0,T;BV)},$$

and hence that  $A \leq T \|u_0\|_{L^1}$  and  $B \leq T |u_0|_{BV}$  by (2.7).

By standard computations given in Appendix A,

$$(6.21) \quad \lim_{\delta \rightarrow 0} \mathcal{E}_3(\delta) = \mathcal{E}_3,$$

and it follows after going to the limit in the estimate above, that

$$\begin{aligned}
\mathcal{E}_3 &\leq C_\epsilon r \\
&\quad + C(d) \left\{ T E_1(u_0) \|\varphi' - \psi'\|_\infty \right. \\
(6.22) \quad &\quad + T (1 + |\ln T|) |u_0|_{BV} \|\varphi' - \psi'\|_\infty \\
&\quad + T |u_0|_{BV} \|\varphi' \ln \varphi' - \psi' \ln \psi'\|_\infty \\
&\quad \left. + T^2 |u_0|_{BV} \|\varphi' - \psi'\|_\infty^2 \frac{1}{\epsilon} \right\},
\end{aligned}$$

for all  $T \wedge 1 > \underline{\Lambda}^{-1} r$  when  $\alpha = 1$ .

When  $\alpha > 1$ , similar arguments using (6.14) show that for all  $r_1 > \underline{\Delta}^{-\frac{1}{\alpha}} r$ ,

$$(6.23) \quad \begin{aligned} \mathcal{E}_3 &\leq C_\epsilon r^{2-\alpha} \\ &+ C(d) \left\{ \frac{G_d(\alpha)}{\alpha-1} T |u_0|_{BV} \|(\varphi')^{\frac{1}{\alpha}} - (\psi')^{\frac{1}{\alpha}}\|_\infty r_1^{1-\alpha} \right. \\ &\left. + \frac{G_d(\alpha)}{2-\alpha} T |u_0|_{BV} \|(\varphi')^{\frac{1}{\alpha}} - (\psi')^{\frac{1}{\alpha}}\|_\infty^2 \frac{r_1^{2-\alpha}}{\epsilon} \right\}. \end{aligned}$$

**8. Conclusion.** We have to insert the estimates of the three preceding steps into (6.2). Let us begin by the case where  $\alpha = 1$ . By (6.3), (6.4) and (6.22),

$$\begin{aligned} \|u(\cdot, T) - v(\cdot, T)\|_{L^1} &\leq 2(m_u(\nu) \vee m_v(\nu)) + C_\epsilon r \\ &+ T |u_0|_{BV} \|f' - g'\|_\infty \\ &+ C(d) \left\{ |u_0|_{BV} \epsilon \right. \\ &+ T E_1(u_0) \|\varphi' - \psi'\|_\infty \\ &+ T (1 + |\ln T|) |u_0|_{BV} \|\varphi' - \psi'\|_\infty \\ &+ T |u_0|_{BV} \|\varphi' \ln \varphi' - \psi' \ln \psi'\|_\infty \\ &\left. + T^2 |u_0|_{BV} \|\varphi' - \psi'\|_\infty^2 \frac{1}{\epsilon} \right\}, \end{aligned}$$

for all  $r, \epsilon > 0$  and  $T > \nu > 0$  such that  $T \wedge 1 > \underline{\Delta}^{-1} r$ . We complete the proof by sending  $r$  and  $\nu$  to zero, and taking  $\epsilon = T \|\varphi' - \psi'\|_\infty$ .

When  $\alpha > 1$ , we find using (6.23) that

$$\begin{aligned} \|u(\cdot, T) - v(\cdot, T)\|_{L^1} &\leq 2(m_u(\nu) \vee m_v(\nu)) + C_\epsilon r^{2-\alpha} \\ &+ T |u_0|_{BV} \|f' - g'\|_\infty \\ &+ C(d) \left\{ |u_0|_{BV} \epsilon \right. \\ &+ \frac{G_d(\alpha)}{\alpha-1} T |u_0|_{BV} \|(\varphi')^{\frac{1}{\alpha}} - (\psi')^{\frac{1}{\alpha}}\|_\infty r_1^{1-\alpha} \\ &\left. + \frac{G_d(\alpha)}{2-\alpha} T |u_0|_{BV} \|(\varphi')^{\frac{1}{\alpha}} - (\psi')^{\frac{1}{\alpha}}\|_\infty^2 \frac{r_1^{2-\alpha}}{\epsilon} \right\}, \end{aligned}$$

for all  $r, \epsilon > 0$ ,  $T > \nu > 0$  and  $r_1 > \underline{\Delta}^{-\frac{1}{\alpha}} r$ . We conclude by choosing  $\epsilon = T^{\frac{1}{\alpha}} \|(\varphi')^{\frac{1}{\alpha}} - (\psi')^{\frac{1}{\alpha}}\|_\infty$  and  $r_1 = T^{\frac{1}{\alpha}}$ . The proof of Theorem 3.1 is complete.  $\square$

*Remark 6.2.* (1) From the proof, we find that  $C \leq C(d) \left(1 + \frac{G_d(\alpha)}{\alpha-1} + \frac{G_d(\alpha)}{2-\alpha}\right)$  in (3.5) when  $\alpha > 1$ . By (2.3),  $\lim_{\alpha \rightarrow 2} C(d, \alpha)$  is finite and only depends on  $d$ .

(2) In particular,  $C \leq C(d) \left(1 + \frac{G_d(\alpha)}{\alpha-1} + \frac{G_d(\alpha)}{2-\alpha}\right)$  when  $\alpha > 1$  also in (3.7).

**6.3. Proof of Theorem 3.8.** Here no linearization procedure is needed since  $\varphi = \psi$ . The new difficulty comes from the fact that the two Lévy measures are different. A key idea is to change variables to work with only one measure.

*Proof of Theorem 3.8.* We argue as in the preceding proof with  $u = u^\alpha$  and  $v = u^\beta$ , i.e.  $(u_0, f, \varphi) = (v_0, g, \psi)$ . To simplify references to similar computations, we still use the letters  $u$  and  $v$  for a while.

**1. Applying Kuznetsov, initial estimates.** As in step 1 in the proof of Theorem 3.1, we apply Lemma 4.1 and estimate the  $\mathcal{L}_r$ -terms. We obtain estimates similar to

(6.2), and (6.4), and conclude that for all  $\alpha, \beta \in (0, 2)$ ,  $r, \epsilon > 0$  and  $T > \nu > 0$ ,

$$(6.24) \quad \begin{aligned} & \|u(\cdot, T) - v(\cdot, T)\|_{L^1} \\ & \leq C(d) |u_0|_{BV} \epsilon + 2 (m_u(\nu) \vee m_v(\nu)) + C_\epsilon (r^{2-\alpha} + r^{2-\beta}) \\ & \quad + \underbrace{\int_{Q_T^2} \operatorname{sgn}(v - u) (\mathcal{L}^{\beta, r}[\varphi(v(\cdot, t))](x) - \mathcal{L}^{\alpha, r}[\varphi(u(\cdot, s))](y)) \phi^{\epsilon, \nu} dw}_{=: \mathcal{E}_3}. \end{aligned}$$

The new  $r^{2-\beta}$ -term comes from the new  $\mathcal{L}_r^\beta$ -term in the estimate corresponding to  $\mathcal{E}_2$ . Note that the terms in  $\mathcal{E}_3$  only involve one function  $\varphi$ , but different  $\alpha, \beta$ . Most of the remaining proof consists in estimating  $\mathcal{E}_3$ .

**2. Change of variables and first estimate of  $\mathcal{E}_3$ .** We perform several changes of variables to move the differences between  $\mathcal{L}^{\alpha, r}$  and  $\mathcal{L}^{\beta, r}$  from the Lévy measure to the  $z$ -translations. This is similar in spirit to what we did in the preceding proof to obtain (6.6). First we let  $\tilde{z} = |z|^{\gamma^{-1}-1} z$  ( $\gamma > 0$ ), and note that  $d\tilde{z} = \gamma^{-1} |z|^{d(\gamma^{-1}-1)} dz^d$  so that  $|z|^{-d-\beta} dz = \gamma |\tilde{z}|^{-d-\beta\gamma} d\tilde{z}$ . Take  $\gamma = \gamma_\beta := \sqrt{\frac{\alpha}{\beta}}$ , and check that  $-d - \beta\gamma = -d - \sqrt{\alpha\beta}$  and

$$\mathcal{L}^{\beta, r}[\varphi(v(\cdot, t))](x) = G_d(\beta) \gamma_\beta \int_{|z| > r^{\gamma_\beta^{-1}}} \frac{\varphi(v(x + |z|^{\gamma_\beta^{-1}} z, t)) - \varphi(v(x, t))}{|z|^{d+\sqrt{\alpha\beta}}} dz.$$

Then we use the change of variable  $z \rightarrow (G_d(\beta) \gamma_\beta)^{\frac{1}{\sqrt{\alpha\beta}}} z$  and get that

$$(6.25) \quad \mathcal{L}^{\beta, r}[\varphi(v(\cdot, t))](x) = \int_{|z| > r_\beta} \frac{\varphi(v(x + c_\beta |z|^{\gamma_\beta^{-1}} z, t)) - \varphi(v(x, t))}{|z|^{d+\sqrt{\alpha\beta}}} dz,$$

where  $c_\beta := (G_d(\beta) \gamma_\beta)^{\frac{1}{\beta}} > 0$  and  $r_\beta := (G_d(\beta) \gamma_\beta)^{-\frac{1}{\sqrt{\alpha\beta}}} r^{\gamma_\beta^{-1}} > 0$ . Similar computations for  $u$  show that

$$(6.26) \quad \mathcal{L}^{\alpha, r}[\varphi(u(\cdot, s))](y) = \int_{|z| > r_\alpha} \frac{\varphi(u(y + c_\alpha |z|^{\gamma_\alpha^{-1}} z, s)) - \varphi(u(y, s))}{|z|^{d+\sqrt{\alpha\beta}}} dz,$$

where  $\gamma_\alpha := \sqrt{\frac{\beta}{\alpha}}$ ,  $c_\alpha := (G_d(\alpha) \gamma_\alpha)^{\frac{1}{\alpha}}$  and  $r_\alpha := (G_d(\alpha) \gamma_\alpha)^{-\frac{1}{\sqrt{\alpha\beta}}} r^{\gamma_\alpha^{-1}}$ . Hence

$$\begin{aligned} \mathcal{E}_3 &= \int_{Q_T^2} \int_{r_\alpha \wedge r_\beta < |z| < r_\alpha \vee r_\beta} \dots \frac{dz}{|z|^{d+\sqrt{\alpha\beta}}} \\ & \quad + \int_{Q_T^2} \int_{|z| > r_\alpha \vee r_\beta} \operatorname{sgn}(v - u) \\ & \quad \cdot \frac{\{\varphi(v(x + c_\beta |z|^{\gamma_\beta^{-1}} z, t)) - \varphi(u(y + c_\alpha |z|^{\gamma_\alpha^{-1}} z, s))\} - \{\varphi(v) - \varphi(u)\}}{|z|^{d+\sqrt{\alpha\beta}}} \\ & \quad \cdot \phi^{\epsilon, \nu} dz dw \\ & =: \mathcal{E}_{3,1} + \mathcal{E}_{3,2}, \end{aligned}$$

where the integrand of  $\mathcal{E}_{3,1}$  only contains either  $u$ -terms or  $v$ -terms. As in the preceding proof, cf. (6.6) and (6.7), we find that

$$(6.27) \quad \mathcal{E}_{3,1} \leq C_\epsilon (r_\alpha \vee r_\beta)^{2-\sqrt{\alpha\beta}}.$$

<sup>4</sup>Indeed,  $d\tilde{z} = F(z) dz$  for  $F(z) = |\det(D(|z|^{\gamma^{-1}-1} z))|$  and

$$D(|z|^{\gamma^{-1}-1} z) = (\gamma^{-1} - 1) |z|^{\gamma^{-1}-3} z \otimes z + |z|^{\gamma^{-1}-1} \operatorname{Id}.$$

Hence  $F$  is positive,  $F(\lambda z) = |\lambda|^{d(\gamma^{-1}-1)} F(z)$  for all  $\lambda \in \mathbb{R}$ , and radial since

$$F(Re) = |\det((\gamma^{-1} - 1) Re (Re)^t + R R^t)| = |\det(R((\gamma^{-1} - 1) e e^t + \operatorname{Id}) R^t)| = \gamma^{-1},$$

for all orthogonal matrices  $R \in \mathbb{R}^{d \times d}$  and column vectors  $e$  of the canonical basis.

Most of the remaining proof consists in estimating  $\mathcal{E}_{3,2}$ . Before continuing, let us list the following properties that will be needed:

$$(6.28) \quad \forall \alpha \in (0, 2), \forall \beta \in (0, 2), \lim_{r \rightarrow 0} r_\alpha \vee r_\beta = 0;$$

$$(6.29) \quad \forall \lambda \in (0, 2), \begin{cases} \lim_{\alpha, \beta \rightarrow \lambda} \gamma_\alpha = \lim_{\alpha, \beta \rightarrow \lambda} \gamma_\beta = 1, \\ \lim_{\alpha, \beta \rightarrow \lambda} c_\alpha = \lim_{\alpha, \beta \rightarrow \lambda} c_\beta = G_d(\lambda)^{\frac{1}{\lambda}} > 0, \\ \lim_{\alpha, \beta \rightarrow \lambda} \frac{|\gamma_\alpha - \gamma_\beta|}{|\alpha - \beta|} = \frac{1}{\lambda}, \\ \exists C = C(d, \lambda), \lim_{\alpha, \beta \rightarrow \lambda} \frac{|c_\alpha - c_\beta|}{|\alpha - \beta|} = C. \end{cases}$$

These are immediate consequences of (2.3).

**3. First estimate of  $\mathcal{E}_{3,2}$ .** We introduce parameters  $r_2 \geq r_1 > 0$ . By (6.28),  $r_1 > r_\alpha \vee r_\beta$  for sufficiently small  $r$ . Let us define

$$(6.30) \quad \begin{aligned} \mathcal{E}_{3,2} &= \sum_{i=1}^3 \mathcal{E}_{3,2,i} := \sum_{i=1}^3 \int_{Q_T^2} \int_{|z| \in I_i} \operatorname{sgn}(v - u) \\ &\cdot \frac{\{\varphi(v(x + c_\beta |z|^{\gamma_\beta - 1} z, t)) - \varphi(u(y + c_\alpha |z|^{\gamma_\alpha - 1} z, s))\}}{|z|^{d + \sqrt{\alpha\beta}}} \\ &\cdot \phi^{\epsilon, \nu} dz dw \end{aligned}$$

for  $I_1 = (r_2, +\infty)$ ,  $I_2 = (r_1, r_2)$  and  $I_3 = (r_\alpha \vee r_\beta, r_1)$ . An application of Lemma 6.1 with  $c = \tilde{c} = c_\beta$  and  $\gamma = \tilde{\gamma} = \gamma_\beta$ , shows that

$$(6.31) \quad \begin{aligned} \mathcal{E}_{3,2,i} &\leq \int_{Q_T^2} \int_{|z| \in I_i} \operatorname{sgn}(v - u) \phi^{\epsilon, \nu} \\ &\cdot \frac{\varphi(u(y + c_\beta |z|^{\gamma_\beta - 1} z, s)) - \varphi(u(y + c_\alpha |z|^{\gamma_\alpha - 1} z, s))}{|z|^{d + \sqrt{\alpha\beta}}} dz dw. \end{aligned}$$

We now estimate these terms.

Let us begin with  $\mathcal{E}_{3,2,1}$ . Going back to the original variables,  $c_\alpha |z|^{\gamma_\alpha - 1} z \rightarrow z$ ,

$$\begin{aligned} &\int_{Q_T^2} \int_{|z| > r_2} \operatorname{sgn}(v - u) \frac{\varphi(u(y + c_\alpha |z|^{\gamma_\alpha - 1} z, s))}{|z|^{d + \sqrt{\alpha\beta}}} \phi^{\epsilon, \nu} dz dw \\ &= G_d(\alpha) \int_{Q_T^2} \int_{|z| > c_\alpha r_2^{\gamma_\alpha}} \operatorname{sgn}(v - u) \frac{\varphi(u(y + z, s))}{|z|^{d + \alpha}} \phi^{\epsilon, \nu} dz dw. \end{aligned}$$

Let us continue by assuming that  $c_\alpha r_2^{\gamma_\alpha} \geq c_\beta r_2^{\gamma_\beta}$ . By the above identity and a similar one for the  $\beta$ -term, we then find that

$$\begin{aligned} \mathcal{E}_{3,2,1} &\leq \int_{Q_T^2} \int_{|z| > c_\alpha r_2^{\gamma_\alpha}} \operatorname{sgn}(v - u) \varphi(u(y + z, s)) \phi^{\epsilon, \nu} \left( \frac{G_d(\beta)}{|z|^{d + \beta}} - \frac{G_d(\alpha)}{|z|^{d + \alpha}} \right) dz dw \\ &+ G_d(\beta) \int_{Q_T^2} \int_{c_\beta r_2^{\gamma_\beta} < |z| < c_\alpha r_2^{\gamma_\alpha}} \operatorname{sgn}(v - u) \varphi(u(y + z, s)) \phi^{\epsilon, \nu} \frac{dz dw}{|z|^{d + \beta}}. \end{aligned}$$

By (1.4) and (2.7),  $\|\varphi(u)\|_{L^1(Q_T)} \leq M \|u_0\|_{L^1}$  for  $M = T \operatorname{ess\,sup}_{I(u_0)} |\varphi'|$ , and then by Fubini,

$$(6.32) \quad \begin{aligned} \mathcal{E}_{3,2,1} &\leq M \|u_0\|_{L^1} \left\{ \int_{|z| > c_\alpha r_2^{\gamma_\alpha}} \left| \frac{G_d(\beta)}{|z|^{d + \beta}} - \frac{G_d(\alpha)}{|z|^{d + \alpha}} \right| dz + G_d(\beta) \int_{c_\beta r_2^{\gamma_\beta} < |z| < c_\alpha r_2^{\gamma_\alpha}} \frac{dz}{|z|^{d + \beta}} \right\}. \end{aligned}$$



Doing the same reasoning when  $c_\alpha r_2^{\gamma_\alpha} < c_\beta r_2^{\gamma_\beta}$  and taking the maximum, we finally get

$$(6.32) \quad \begin{aligned} \mathcal{E}_{3,2,1} &\leq M \|u_0\|_{L^1} \int_{|z| > (c_\alpha r_2^{\gamma_\alpha}) \wedge (c_\beta r_2^{\gamma_\beta})} \left| \frac{G_d(\beta)}{|z|^{d+\beta}} - \frac{G_d(\alpha)}{|z|^{d+\alpha}} \right| dz \\ &\quad + C(d) M \|u_0\|_{L^1} \max_{\sigma=\alpha,\beta} \int_{|z| \in \text{co}\{c_\alpha r_2^{\gamma_\alpha}, c_\beta r_2^{\gamma_\beta}\}} \frac{dz}{|z|^{d+\sigma}}, \end{aligned}$$

where  $C(d) = \max_{[0,2]} G_d$  is finite by (2.3) and from now on  $\text{co}\{a, b\}$  designs the interval  $(a \wedge b, a \vee b)$ .

Next, by (1.4) and (2.7),  $|\varphi(u)|_{L^1(0,T;BV)} \leq M |u_0|_{BV}$ . Hence by integrating first with respect to  $y$  in (6.31), we find that

$$(6.33) \quad \mathcal{E}_{3,2,2} \leq M |u_0|_{BV} \int_{r_1 < |z| < r_2} |c_\alpha |z|^{\gamma_\alpha} - c_\beta |z|^{\gamma_\beta}| \frac{dz}{|z|^{d+\sqrt{\alpha\beta}}}.$$

Finally, by Lemma 6.1

$$\begin{aligned} \mathcal{E}_{3,2,3} &\leq \int_{Q_T^2} \int_{r_\alpha \vee r_\beta < |z| < r_1} |\varphi(v) - \varphi(u)| \theta_\nu(t-s) \\ &\quad \cdot \{\rho_\epsilon(x-y+h(z)) - \rho_\epsilon(x-y)\} \frac{dz dw}{|z|^{d+\sqrt{\alpha\beta}}}, \end{aligned}$$

with  $h(z) := (c_\beta |z|^{\gamma_\beta-1} - c_\alpha |z|^{\gamma_\alpha-1}) z$ . This estimate is similar to (6.10), but with a new displacement, new functions  $\varphi(u)$  and  $\varphi(v)$ , and the new power  $\sqrt{\alpha\beta}$ . By arguing as before, we find that

$$\mathcal{E}_{3,2,3} \leq \frac{C(d)}{\epsilon} \int_0^T \int_{|z| < r_1} \int_0^1 (1-\tau) |z|^{-d-\sqrt{\alpha\beta}} |h(z)|^2 |\varphi(u(\cdot, s))|_{BV} d\tau dz ds,$$

instead of (6.11). Since  $|\varphi(u)|_{L^1(0,T;BV)} \leq M |u_0|_{BV}$ , we get that

$$(6.34) \quad \mathcal{E}_{3,2,3} \leq C(d) M |u_0|_{BV} \frac{1}{\epsilon} \int_{|z| < r_1} |c_\beta |z|^{\gamma_\beta} - c_\alpha |z|^{\gamma_\alpha}|^2 \frac{dz}{|z|^{d+\sqrt{\alpha\beta}}}.$$

**4. The general estimate.** Let us resume the preceding estimates. By (6.24), (6.27), (6.32), (6.33), (6.34) and the fact that  $\mathcal{E}_3 = \mathcal{E}_{3,1} + \mathcal{E}_{3,2,1} + \mathcal{E}_{3,2,2} + \mathcal{E}_{3,2,3}$ , we have proved that for all  $\alpha, \beta \in (0, 2)$ ,  $\epsilon > 0$ ,  $T > \nu > 0$ ,  $r_2 \geq r_1 > 0$  and  $r > 0$  small enough,

$$\begin{aligned} &\|u^\alpha(\cdot, T) - u^\beta(\cdot, T)\|_{L^1} \\ &\leq 2 (m_u(\nu) \vee m_v(\nu)) + C_\epsilon \left( r^{2-\alpha} + r^{2-\beta} + (r_\alpha \vee r_\beta)^{2-\sqrt{\alpha\beta}} \right) \\ &\quad + C(d) |u_0|_{BV} \epsilon \\ &\quad + M \|u_0\|_{L^1} \int_{|z| > (c_\alpha r_2^{\gamma_\alpha}) \wedge (c_\beta r_2^{\gamma_\beta})} \left| \frac{G_d(\beta)}{|z|^{d+\beta}} - \frac{G_d(\alpha)}{|z|^{d+\alpha}} \right| dz \\ &\quad + C(d) M \|u_0\|_{L^1} \max_{\sigma=\alpha,\beta} \int_{|z| \in \text{co}\{c_\alpha r_2^{\gamma_\alpha}, c_\beta r_2^{\gamma_\beta}\}} \frac{dz}{|z|^{d+\sigma}} \\ &\quad + M |u_0|_{BV} \int_{r_1 < |z| < r_2} |c_\alpha |z|^{\gamma_\alpha} - c_\beta |z|^{\gamma_\beta}| \frac{dz}{|z|^{d+\sqrt{\alpha\beta}}} \\ &\quad + C(d) M |u_0|_{BV} \frac{1}{\epsilon} \int_{|z| < r_1} |c_\beta |z|^{\gamma_\beta} - c_\alpha |z|^{\gamma_\alpha}|^2 \frac{dz}{|z|^{d+\sqrt{\alpha\beta}}}. \end{aligned}$$

Now, we pass to the limit as  $\nu, r \rightarrow 0$ , thanks to (6.28). Next, we replace the  $L^1$ -norm at time  $T$  by the  $C([0, T]; L^1)$ -norm, which can be done without loss of generality since  $t \|\varphi'\|_\infty \leq T \|\varphi'\|_\infty = M$ , for all  $t \leq T$ . Finally, we replace  $\epsilon$

by  $\epsilon|\alpha - \beta|$ , which can also be done since  $\epsilon$  is arbitrary. We deduce that for all  $\alpha, \beta \in (0, 2)$ ,  $T \geq 0$  and  $r_2 \geq r_1 > 0$ ,

$$\begin{aligned}
& \|u^\alpha - u^\beta\|_{C([0,T];L^1)} \\
& \leq C(d) \|u_0\|_{BV} \epsilon |\alpha - \beta| \\
& \quad + M \|u_0\|_{L^1} \underbrace{\int_{|z| > (c_\alpha r_2^{\gamma_\alpha}) \wedge (c_\beta r_2^{\gamma_\beta})} \left| \frac{G_d(\beta)}{|z|^{d+\beta}} - \frac{G_d(\alpha)}{|z|^{d+\alpha}} \right| dz}_{=: J_1} \\
(6.35) \quad & \quad + C(d) M \|u_0\|_{L^1} \underbrace{\max_{\sigma=\alpha,\beta} \int_{|z| \in \text{co}\{c_\alpha r_2^{\gamma_\alpha}, c_\beta r_2^{\gamma_\beta}\}} \frac{dz}{|z|^{d+\sigma}}}_{=: J_2} \\
& \quad + M \|u_0\|_{BV} \underbrace{\int_{r_1 < |z| < r_2} |c_\alpha |z|^{\gamma_\alpha} - c_\beta |z|^{\gamma_\beta}| \frac{dz}{|z|^{d+\sqrt{\alpha\beta}}}}_{=: J_3} \\
& \quad + \frac{C(d) M \|u_0\|_{BV}}{\epsilon} \underbrace{\frac{1}{|\alpha - \beta|} \int_{|z| < r_1} |c_\beta |z|^{\gamma_\beta} - c_\alpha |z|^{\gamma_\alpha}|^2 \frac{dz}{|z|^{d+\sqrt{\alpha\beta}}}}_{=: J_4}.
\end{aligned}$$

The rest of proof consists in estimating  $\limsup_{\alpha,\beta \rightarrow \lambda} \frac{J_i}{|\alpha - \beta|}$  ( $i = 1, \dots, 4$ ). Most of the upper semi-limits will be in fact true limits; but, since we will not need to care about this, we will always use the notation “lim sup” to simplify the reasoning. We will also use the letter  $C$  to denote various constants  $C = C(d, \lambda)$ .

**5. The case  $\lambda \in (1, 2)$ .** We first let  $r_2 \rightarrow +\infty$ . By (6.29),  $(c_\alpha r_2^{\gamma_\alpha}) \wedge (c_\beta r_2^{\gamma_\beta}) \rightarrow +\infty$  and we get at the limit

$$(6.36) \quad J_1 = J_2 = 0$$

and  $J_3 = \int_{|z| > r_1} |c_\alpha |z|^{-d-\sigma_\alpha} - c_\beta |z|^{-d-\sigma_\beta}| dz$ , with  $\sigma_\alpha := \sqrt{\alpha\beta} - \gamma_\alpha$  and  $\sigma_\beta := \sqrt{\alpha\beta} - \gamma_\beta$ .

Let us estimate  $J_3$ . We recognize a term of the same form than in (5.2) with the new “locally Lipschitz” coefficients  $c_\alpha, c_\beta$  and powers  $\sigma_\alpha, \sigma_\beta$ . Since  $\sigma_\alpha, \sigma_\beta \rightarrow \lambda - 1$  as  $\alpha, \beta \rightarrow \lambda$ , we will get an estimate of the form (6.37) below, instead of (5.3). Let us show how to adapt the reasoning for the reader’s convenience. We have

$$J_3 \leq |c_\alpha - c_\beta| \max_{\sigma=\sigma_\alpha, \sigma_\beta} \int_{|z| > r_1} \frac{dz}{|z|^{d+\sigma}} + \underbrace{(c_\alpha \vee c_\beta) \int_{|z| > r_1} \left| |z|^{-d-\sigma_\alpha} - |z|^{-d-\sigma_\beta} \right| dz}_{=: \tilde{J}_3},$$

where  $\tilde{J}_3 \leq S_d \left| \frac{r_1^{-\sigma_\alpha}}{\sigma_\alpha} - \frac{r_1^{-\sigma_\beta}}{\sigma_\beta} \right| + 2 S_d \left| \frac{1}{\sigma_\alpha} - \frac{1}{\sigma_\beta} \right| \mathbf{1}_{r_1 < 1}$ . By (6.29),

$$\limsup_{\alpha,\beta \rightarrow \lambda} \frac{J_3}{|\alpha - \beta|} \leq C \underbrace{(r_1^{1-\lambda} + \mathbf{1}_{r_1 < 1})}_{\leq C r_1^{1-\lambda} \text{ if } \lambda > 1} + C \underbrace{\limsup_{\alpha,\beta \rightarrow \lambda} \frac{1}{|\alpha - \beta|} \left| \frac{r_1^{-\sigma_\alpha}}{\sigma_\alpha} - \frac{r_1^{-\sigma_\beta}}{\sigma_\beta} \right|}_{=: \tilde{J}_3},$$

where a Taylor expansion with integral remainder shows that

$$\tilde{J}_3 = \limsup_{\alpha,\beta \rightarrow \lambda} \frac{|\sigma_\alpha - \sigma_\beta|}{|\alpha - \beta|} \left| \int_0^1 \frac{\sigma_\tau r_1^{-\sigma_\tau} \ln r_1 + r_1^{-\sigma_\tau}}{\sigma_\tau^2} d\tau \right| \leq C r_1^{1-\lambda} (1 + |\ln r_1|),$$

with  $\sigma_\tau := \tau \sigma_\alpha + (1 - \tau) \sigma_\beta$ . We deduce the following estimate:

$$(6.37) \quad \limsup_{\alpha,\beta \rightarrow \lambda} \frac{J_3}{|\alpha - \beta|} \leq C r_1^{1-\lambda} (1 + |\ln r_1|).$$

Let us notice that this estimate fails when  $\lambda = 1$ , because  $\sigma_\tau \rightarrow 0$  for all  $\tau$ , as  $\alpha, \beta \rightarrow 1$ .

Let us now estimate  $J_4$ . By adding and subtracting terms,

$$J_4 \leq \frac{1}{2} \sum_{\pm} \frac{|c_\alpha \mp c_\beta|^2}{|\alpha - \beta|} \underbrace{\int_{|z| < r_1} (|z|^{\gamma_\alpha} \pm |z|^{\gamma_\beta})^2 \frac{dz}{|z|^{d+\sqrt{\alpha\beta}}}}_{=: J_{4,\pm}}.$$

By expanding the squares and integrating,

$$J_{4,\pm} = S_d \left( \frac{r_1^{2\gamma_\alpha - \sqrt{\alpha\beta}}}{2\gamma_\alpha - \sqrt{\alpha\beta}} + \frac{r_1^{2\gamma_\beta - \sqrt{\alpha\beta}}}{2\gamma_\beta - \sqrt{\alpha\beta}} \pm 2 \frac{r_1^{\gamma_\alpha + \gamma_\beta - \sqrt{\alpha\beta}}}{\gamma_\alpha + \gamma_\beta - \sqrt{\alpha\beta}} \right).$$

By (6.29), the limit of  $J_{4,+}$  is easy to compute and we get

$$\limsup_{\alpha, \beta \rightarrow 1} \frac{J_4}{|\alpha - \beta|} \leq C r_1^{2-\lambda} + C \underbrace{\limsup_{\alpha, \beta \rightarrow 1} \frac{J_{4,-}}{(\alpha - \beta)^2}}_{=: \tilde{J}_{4,-}}.$$

We estimate  $\tilde{J}_{4,-}$  by multiplying and dividing by  $(\gamma_\alpha - \gamma_\beta)^2$  and changing the variables by  $a := \gamma_\alpha - \frac{\sqrt{\alpha\beta}}{2}$  and  $b := \gamma_\beta - \frac{\sqrt{\alpha\beta}}{2}$ . We get

$$\begin{aligned} \tilde{J}_{4,-} &= \limsup_{\alpha, \beta \rightarrow 1} \frac{(\gamma_\alpha - \gamma_\beta)^2}{|\alpha - \beta|^2} \\ &\cdot \limsup_{a, b \rightarrow c} \frac{1}{|a - b|^2} \left( \frac{r_1^{2a}}{2a} + \frac{r_1^{2b}}{2b} - \frac{2r_1^{a+b}}{a+b} \right), \end{aligned}$$

where  $c := 1 - \frac{\lambda}{2} > 0$  is the limit of  $a, b$  as  $\alpha, \beta \rightarrow \lambda$ . By (6.29) and the estimation of the last limit in Lemma B.2(ii) in appendix,

$$\tilde{J}_{4,-} \leq C r_1^{2-\lambda} (1 + \ln^2 r_1).$$

We conclude that

$$(6.38) \quad \limsup_{\alpha, \beta \rightarrow 1} \frac{J_4}{|\alpha - \beta|} \leq C r_1^{2-\lambda} (1 + \ln^2 r_1).$$

Note that this estimate works even if  $\lambda = 1$ .

We are now ready to conclude the proof and show (3.9) when  $\lambda \in (1, 2)$ . Recall that we estimate  $\text{Lip}_\alpha(u; \lambda)$  using (6.35) with  $r_2 = +\infty$ . The limsups of the terms on the right-hand side are estimated by (6.36), (6.37) and (6.38). We get for all  $\epsilon > 0$ ,  $T \geq 0$  and  $r_1 > 0$ ,

$$\text{Lip}_\alpha(u; \lambda) \leq C |u_0|_{BV} \left\{ \epsilon + M \left( r_1^{1-\lambda} (1 + |\ln r_1|) + \frac{r_1^{2-\lambda}}{\epsilon} (1 + \ln^2 r_1) \right) \right\}.$$

We complete the proof by taking  $\epsilon = M^{\frac{1}{\lambda}} (1 + |\ln M|)$  and  $r_1 = M^{\frac{1}{\lambda}}$ .

**6.** *The case  $\lambda = 1$ .* We have to estimate again  $J_i$  in (6.35) ( $i = 1, \dots, 4$ ). This time, we do not let  $r_2 \rightarrow +\infty$ .

For  $J_1$ , we recognize again a term of the form (5.2) and we argue in the same way to estimate it. The only difference is that the fixed cutting parameter  $\tilde{r}$  is replaced by a moving one  $(c_\alpha r_2^{\gamma_\alpha}) \wedge (c_\beta r_2^{\gamma_\beta})$ . But, by (6.29) it follows that  $\lim_{\alpha, \beta \rightarrow 1} (c_\alpha r_2^{\gamma_\alpha}) \wedge (c_\beta r_2^{\gamma_\beta}) = G_d(1) r_2$  with  $G_d(1) > 0$ , and we leave it to the reader to verify that this is sufficient to extend the proof of (5.3) to the current case. Now, this estimate becomes

$$(6.39) \quad \limsup_{\alpha, \beta \rightarrow 1} \frac{J_1}{|\alpha - \beta|} \leq C (G_d(1) r_2)^{-1} (1 + |\ln(G_d(1) r_2)|) = C r_2^{-1} (1 + |\ln r_2|).$$

For  $J_2$ , we use that

$$\begin{aligned} J_2 &= S_d \max_{\sigma=\alpha,\beta} \frac{1}{\sigma} |(c_\alpha r_2^{\gamma_\alpha})^{-\sigma} - (c_\beta r_2^{\gamma_\beta})^{-\sigma}| \\ &= S_d \max_{\sigma=\alpha,\beta} |c_\alpha r_2^{\gamma_\alpha} - c_\beta r_2^{\gamma_\beta}| \int_0^1 (\tau c_\alpha r_2^{\gamma_\alpha} + (1-\tau) c_\beta r_2^{\gamma_\beta})^{-\sigma-1} d\tau, \end{aligned}$$

thanks to a Taylor expansion. By (6.29) and a passage to the limit under the integral sign,

$$\limsup_{\alpha,\beta \rightarrow 1} \frac{J_2}{|\alpha - \beta|} = C r_2^{-2} \limsup_{\alpha,\beta \rightarrow 1} \frac{|c_\alpha r_2^{\gamma_\alpha} - c_\beta r_2^{\gamma_\beta}|}{|\alpha - \beta|}.$$

To estimate the last limit, we write

$$|c_\alpha r_2^{\gamma_\alpha} - c_\beta r_2^{\gamma_\beta}| \leq |c_\alpha - c_\beta| (r_2^{\gamma_\alpha} \vee r_2^{\gamma_\beta}) + (c_\alpha \vee c_\beta) |r_2^{\gamma_\alpha} - r_2^{\gamma_\beta}|,$$

where  $|r_2^{\gamma_\alpha} - r_2^{\gamma_\beta}| = |\gamma_\alpha - \gamma_\beta| |\ln r_2| \int_0^1 r_2^{\tau \gamma_\alpha + (1-\tau) \gamma_\beta} d\tau$ . Hence, again by (6.29),

$$(6.40) \quad \limsup_{\alpha,\beta \rightarrow 1} \frac{J_2}{|\alpha - \beta|} \leq C r_2^{-1} (1 + |\ln r_2|).$$

We have to do again the estimate of  $J_3$ , since the preceding one (6.37) fails.

$$\begin{aligned} J_3 &\leq |c_\alpha - c_\beta| \max_{\sigma=\alpha,\beta} \int_{r_1 < |z| < r_2} |z|^{\gamma_\sigma} \frac{dz}{|z|^{d+\sqrt{\alpha\beta}}} \\ &\quad + (c_\alpha \vee c_\beta) \underbrace{\int_{r_1 < |z| < r_2} \left| |z|^{\gamma_\alpha} - |z|^{\gamma_\beta} \right| \frac{dz}{|z|^{d+\sqrt{\alpha\beta}}}}_{=: \tilde{J}_3}, \end{aligned}$$

so that by (6.29) and a passage to the limit under the integral sign,

$$(6.41) \quad \limsup_{\alpha,\beta \rightarrow 1} \frac{J_3}{|\alpha - \beta|} \leq C (|\ln r_1| + |\ln r_2|) + C \limsup_{\alpha,\beta \rightarrow 1} \frac{\tilde{J}_3}{|\alpha - \beta|}.$$

To estimate  $\tilde{J}_3$ , we first assume that  $\alpha, \beta \neq 1$ , so that  $\gamma_\alpha - \sqrt{\alpha\beta} = (1-\alpha)\gamma_\alpha \neq 0$  and  $\gamma_\beta - \sqrt{\alpha\beta} \neq 0$ . Hence,  $\tilde{J}_3 = S_d (\int_{r_1}^1 \dots + \int_1^{r_2} \dots)$  in polar coordinates, and

$$\tilde{J}_3 \leq S_d \sum_{i=1,2} \left| \frac{r_i^{\gamma_\alpha - \sqrt{\alpha\beta}} - 1}{\gamma_\alpha - \sqrt{\alpha\beta}} - \frac{r_i^{\gamma_\beta - \sqrt{\alpha\beta}} - 1}{\gamma_\beta - \sqrt{\alpha\beta}} \right|.$$

By Lemma B.2(i) in the appendix,

$$\tilde{J}_3 \leq 2 S_d |\gamma_\alpha - \gamma_\beta| \max_{i=1,2} \max_{\sigma=\alpha,\beta} (1 \vee r_i^{\gamma_\sigma - \sqrt{\alpha\beta}}) \ln^2 r_i.$$

By sending  $\alpha$  or  $\beta \rightarrow 1$ , we see that this inequality holds also when  $\alpha$  or  $\beta = 1$ . Hence, by (6.29) and (6.41),

$$(6.42) \quad \limsup_{\alpha,\beta \rightarrow 1} \frac{J_3}{|\alpha - \beta|} \leq C (|\ln r_1| \vee \ln^2 r_1 + |\ln r_2| \vee \ln^2 r_2).$$

Finally, for  $J_4$ , we use (6.38) which is still valid and we are ready to show (3.9) in the critical case. By (6.35), (6.39), (6.40), (6.42) and (6.38), we have for all  $\epsilon > 0$ ,  $T \geq 0$  and  $r_2 \geq r_1 > 0$ ,

$$\begin{aligned} \text{Lip}_\alpha(u; 1) &\leq C |u_0|_{BV} \epsilon \\ &\quad + C M \|u_0\|_{L^1} r_2^{-1} (1 + |\ln r_2|) \\ &\quad + C M |u_0|_{BV} (|\ln r_1| \vee \ln^2 r_1 + |\ln r_2| \vee \ln^2 r_2) \\ &\quad + C M |u_0|_{BV} \frac{r_1}{\epsilon} (1 + \ln^2 r_1). \end{aligned}$$

We complete the proof by taking  $\epsilon = M(1 + |\ln M|)$ ,  $r_1 = M \wedge 1$ ,  $r_2 = 1 \vee \frac{\|u_0\|_{L^1}}{|u_0|_{BV}}$ , and noting that  $\|u_0\|_{L^1} \leq |u_0|_{BV}$  if  $r_2 = 1$ .  $\square$

## 7. PROOF OF THEOREM 3.3

This section is devoted to the proof of Theorem 3.3. Let us first recall the notions of entropy solutions of (1.5) and (1.6) introduced in [29, 9]. For (1.6), we use an equivalent definition introduced in [26].

**Definition 7.1** (Entropy solutions). *Let  $u_0 \in L^\infty \cap L^1(\mathbb{R}^d)$  and (1.3)–(1.4) hold. Let  $u \in L^\infty(Q_T) \cap L^\infty(0, T; L^1)$ .*

(1)  *$u$  is an entropy solution of (1.5) if, for all  $k \in \mathbb{R}$  and all nonnegative  $\phi \in C_c^\infty(\mathbb{R}^{d+1})$ ,*

$$\begin{aligned} & \int_{Q_T} |u - k| \partial_t \phi + q_f(u, k) \cdot \nabla \phi - \operatorname{sgn}(u - k) \varphi(u) \phi \, dx \, dt \\ & + \int_{\mathbb{R}^d} |u_0(x) - k| \phi(x, 0) \, dx \geq 0. \end{aligned}$$

(2)  *$u$  is an entropy solution of (1.6) if,*

(a)  $\varphi(u) \in L^2(0, T; H^1)$ ,

(b) *and for all  $k \in \mathbb{R}$  and all nonnegative  $\phi \in C_c^\infty(\mathbb{R}^{d+1})$ ,*

$$\begin{aligned} & \int_{Q_T} |u - k| \partial_t \phi + q_f(u, k) \cdot \nabla \phi + |\varphi(u) - \varphi(k)| \frac{\Delta}{4\pi^2} \phi \, dx \, dt \\ & + \int_{\mathbb{R}^d} |u_0(x) - k| \phi(x, 0) \, dx \geq 0. \end{aligned}$$

To prove Theorem 3.3, we need to establish some technical lemmas. Let us begin by a compactness result.

**Lemma 7.1.** *Let  $u_0 \in L^\infty \cap L^1(\mathbb{R}^d)$ , (1.3)–(1.4) hold, and for each  $\alpha \in (0, 2)$ , let  $u^\alpha$  be the unique entropy solution to (1.1). Then, there exist  $u, w \in L^\infty(Q_T) \cap C([0, T]; L^1)$  such that  $u = \lim_{\alpha \rightarrow 0} u^\alpha$  and  $w = \lim_{\alpha \rightarrow 2} u^\alpha$ , up to subsequences, in  $C([0, T]; L^1_{\text{loc}})$ .*

*Proof.* Let  $\alpha_m$  be a sequence converging to 2 and define  $E := \{u^{\alpha_m}\}_m$ . We will show that  $E$  is relatively compact in  $C([0, T]; L^1_{\text{loc}})$ . First we take a sequence  $\{u_0^n\}_n \subset L^\infty \cap L^1 \cap BV(\mathbb{R}^d)$  that converges to  $u_0$  in  $L^1(\mathbb{R}^d)$ , and let  $E_n$  denote the family  $\{u_n^{\alpha_m}\}_m$  of entropy solutions to (1.1) with  $\alpha = \alpha_m$  and  $u_0^n$  as initial data. We begin by showing that  $E_n$  is relatively compact in  $C([0, T]; L^1_{\text{loc}})$ .

The family  $E_n$  is bounded and equicontinuous in  $C([0, T]; L^1)$  by (2.7), Corollary 3.6, and Remark 6.2(2). For each  $t \in [0, T]$ ,  $\{u_n^{\alpha_m}(\cdot, t)\}_m$  is relatively compact in  $L^1_{\text{loc}}(\mathbb{R}^d)$  by the  $BV$ -bound in (2.7) and Helly's theorem. By the Arzela-Ascoli theorem it then easily follows that  $E_n$  is relatively compact in  $C([0, T]; L^1_{\text{loc}})$  for any  $n \in \mathbb{N}$ .

The relative compactness of  $E$ , and thus the existence of  $w \in C([0, T]; L^1_{\text{loc}})$ , is now a consequence of the  $L^1$ -contraction principle since

$$(7.1) \quad \sup_{m \in \mathbb{N}} \|u^{\alpha_m} - u_n^{\alpha_m}\|_{C([0, T]; L^1)} \leq \|u_0 - u_0^n\|_{L^1} \rightarrow 0 \quad \text{as } n \rightarrow +\infty.$$

Taking a.e. converging subsequences and using the a priori estimate (2.7), we infer that  $w \in L^\infty(Q_T)$ . To prove that  $w \in C([0, T]; L^1)$ , we observe that  $E$  is equicontinuous in  $C([0, T]; L^1)$  by the triangle inequality, the convergence estimate

(7.1), and the equicontinuity of  $E_n$ . Hence, for any  $R > 0$ ,  $m \in \mathbb{N}$ , and  $t, s \in [0, T]$ ,

$$\begin{aligned} & \| (w(\cdot, t) - w(\cdot, s)) \mathbf{1}_{|x| < R} \|_{L^1} \\ & \leq \| u^{\alpha_m}(\cdot, t) - u^{\alpha_m}(\cdot, s) \|_{L^1} \\ & \quad + \| (w(\cdot, t) - u^{\alpha_m}(\cdot, t)) \mathbf{1}_{|x| < R} \|_{L^1} + \| (u^{\alpha_m}(\cdot, s) - w(\cdot, s)) \mathbf{1}_{|x| < R} \|_{L^1} \\ & \leq o(1) + 2 \| (w - u^{\alpha_m}) \mathbf{1}_{|x| < R} \|_{C([0, T]; L^1)}, \end{aligned}$$

where  $o(1) \rightarrow 0$  as  $|t - s| \rightarrow 0$  uniformly in  $R$  and  $m$ . We then conclude that

$$\| (w(\cdot, t) - w(\cdot, s)) \|_{L^1} \leq o(1) \quad \text{as} \quad |t - s| \rightarrow 0$$

by first sending  $m \rightarrow +\infty$  and then  $R \rightarrow +\infty$  using Fatou's lemma.  $\square$

Let us now verify that these limits satisfy the entropy inequalities of the preceding definition.

**Lemma 7.2.** *Under the assumptions of Lemma 7.1,  $u$  and  $w$  satisfy the entropy inequalities of Definition 7.1 (1) and (2b) respectively.*

In the proof we need the following lemma:

**Lemma 7.3.** *A function  $u$  is an entropy solution of (1.1) (cf. Definition 2.1) if and only if for all convex  $\eta \in C^1(\mathbb{R})$ , all  $r > 0$  and all nonnegative  $\phi \in C_c^\infty(\mathbb{R}^{d+1})$ ,*

$$\begin{aligned} & \int_{Q_T} \eta(u) \partial_t \phi + q_f^\eta(u) \cdot \nabla \phi \, dx \, dt \\ (7.2) \quad & + \int_{Q_T} q_\varphi^\eta(u) \mathcal{L}_r^\alpha[\phi] + \eta'(u) \mathcal{L}^{\alpha, r}[\varphi(u)] \phi \, dx \, dt \\ & + \int_{\mathbb{R}^d} \eta(u_0(x)) \phi(x, 0) \, dx \geq 0, \end{aligned}$$

where  $q_g^\eta(u) := \int_0^u \eta'(\tau) g'(\tau) \, d\tau$  for all  $u \in \mathbb{R}$  and  $g = f, \varphi$ .

This result is well-known for (local) conservation laws, see e.g. [23, p. 27]. Because of the presence of the discontinuous sign-function in the Kruzkov formulation (2.6), any proof will be more technical than in the local case and we therefore provide one in Appendix C.

*Proof of Lemma 7.3.*

**1. Entropy inequalities for  $w$ .** Using the definition of  $\mathcal{L}_r$  and  $\mathcal{L}^r$  in (2.1), we send  $r \rightarrow +\infty$  in the entropy inequality (2.6) and find that

$$\begin{aligned} & \int_{Q_T} |u^\alpha - k| \partial_t \phi + q_f(u^\alpha, k) \cdot \nabla \phi - |\varphi(u^\alpha) - \varphi(k)| (-\Delta)^{\frac{\alpha}{2}} \phi \, dx \, dt \\ (7.3) \quad & + \int_{\mathbb{R}^d} |u_0(x) - k| \phi(x, 0) \, dx \geq 0. \end{aligned}$$

Since  $(-\Delta)^{\frac{\alpha}{2}} \phi = \mathcal{F}(|\cdot|^\alpha \mathcal{F}\phi)$  and  $-\frac{\Delta}{4\pi^2} \phi = \mathcal{F}(|\cdot|^2 \mathcal{F}\phi)$ , by Plancherel

$$(7.4) \quad -(-\Delta)^{\frac{\alpha}{2}} \phi \rightarrow \frac{\Delta}{4\pi^2} \phi \quad \text{in} \quad L^2(\mathbb{R}^{d+1}) \quad \text{as} \quad \alpha \rightarrow 2.$$

To get the entropy inequalities of Definition 7.1(2b), we must pass to the limit in (7.3). This is straight forward for the local terms due to Lemma 7.1 and (2.7).

For the nonlocal term, we first observe that

$$\begin{aligned}
& - \int_{Q_T} |\varphi(u^\alpha) - \varphi(k)| (-\Delta)^{\frac{\alpha}{2}} \phi \, dx \, dt \\
& = \int_{Q_T} \underbrace{\left\{ |\varphi(u^\alpha) - \varphi(k)| - |\varphi(k)| \right\}}_{=:q(u^\alpha)} \left\{ \Delta\phi - \Delta\phi - (-\Delta)^{\frac{\alpha}{2}} \phi \right\} \, dx \, dt \\
& \leq \int_{Q_T} q(u^\alpha) \Delta\phi \, dx \, dt + \|q(u^\alpha)\|_{L^2(Q_T)} \|\Delta\phi + (-\Delta)^{\frac{\alpha}{2}} \phi\|_{L^2(Q_T)}.
\end{aligned}$$

By the  $C([0, T]; L^1_{\text{loc}})$ -convergence of  $u^\alpha$ , the first term converges as  $\alpha \rightarrow 2$  to

$$\int_{Q_T} |\varphi(u^\alpha) - \varphi(k)| \Delta\phi \, dx \, dt.$$

By (7.4), the second term tends to zero since  $\|q(u^\alpha)\|_{L^2(Q_T)}$  is bounded independently of  $\alpha$ . The boundedness follows from (2.7) and an  $(L^1, L^\infty)$ -interpolation argument since  $q \in W^{1, \infty}_{\text{loc}}(\mathbb{R})$  and  $q(0) = 0$ . This completes the proof for  $w$ .

**2. Entropy inequalities for  $u$ .** Let us fix  $r > 0$  for the duration of this proof and start from the entropy inequalities (7.2), written for convex and  $C^1$ -entropies  $\eta$ .

There is again no difficulty to pass to the limit as  $\alpha \rightarrow 0$  in the local terms of (7.2). For the first nonlocal term, we use that  $\mathcal{L}_r^\alpha[\phi] \rightarrow 0$  uniformly on  $\mathbb{R}^{d+1}$ . This is readily seen from (2.1) and (2.3). Let us also notice that  $q_\varphi^\eta$ , defined just below (7.2), satisfies  $q_\varphi^\eta \in W^{1, \infty}_{\text{loc}}(\mathbb{R})$  and  $q_\varphi^\eta(0) = 0$ . Hence

$$\int_{Q_T} q_\varphi^\eta(u^\alpha) \mathcal{L}_r^\alpha[\phi] \, dx \, dt \rightarrow 0,$$

since  $q_\varphi^\eta(u^\alpha)$  is bounded in  $C([0, T]; L^1)$ . For the remaining nonlocal term, we split the integral and get

$$\begin{aligned}
& \int_{Q_T} \eta'(u^\alpha) \mathcal{L}^{\alpha, r}[\varphi(u^\alpha)] \phi \, dx \, dt \\
& \leq - \underbrace{G_d(\alpha) \int_{|z|>r} \frac{dz}{|z|^{d+\alpha}}}_{=:I} \int_{Q_T} \eta'(u^\alpha) \varphi(u^\alpha) \phi \, dx \, dt \\
& \quad + C \underbrace{\frac{G_d(\alpha)}{r^{d+\alpha}}}_{=:J} \|\varphi(u^\alpha)\|_{C([0, T]; L^1)} \|\phi\|_{L^1(Q_T)},
\end{aligned}$$

where  $C$  is an  $L^\infty$ -bound on  $\eta'(u^\alpha)$ . Let us recall that  $C$  can be chosen independent on  $\alpha$  by (2.7). Notice that for all fixed  $r$ ,  $\lim_{\alpha \rightarrow 0} I = 1$  and  $\lim_{\alpha \rightarrow 0} J = 0$  by (2.3). Moreover, taking another subsequence if necessary, we can assume that  $u^\alpha \rightarrow u$  a.e.. Since  $\eta'$  is continuous, we can pass to the limit as  $\alpha \rightarrow 0$  in the inequality above, thanks to (2.7) and the dominated convergence theorem.

The limit in (7.2) then implies that

$$\int_{Q_T} \eta(u) \partial_t \phi + q_f^\eta(u) \cdot \nabla \phi - \eta'(u) \varphi(u) \phi \, dx \, dt + \int_{\mathbb{R}^d} \eta(u_0(x)) \phi(x, 0) \, dx \geq 0,$$

for all convex  $C^1$ -entropies  $\eta$  and fluxes  $q_f^\eta(u) = \int_0^u \eta'(\tau) f'(\tau) \, d\tau$ . It is then classical to get the desired Kruzkov entropy inequalities of Definition 7.1(1) from these inequalities, see e.g. the if part of the proof in Appendix C.  $\square$

To prove that  $w$  satisfies the item (2a), we need to derive an  $H^{\frac{\alpha}{2}}$ -estimate on  $u^\alpha$ . In the sequel,  $H^{\frac{\alpha}{2}}(\mathbb{R}^d)$  denotes the subspace of  $u \in L^2(\mathbb{R}^d)$  such that the semi-norm

$$|u|_{H^{\frac{\alpha}{2}}} := \left( \frac{G_d(\alpha)}{2} \iint_{\mathbb{R}^{2d}} \frac{(u(x) - u(y))^2}{|x - y|^{d+\alpha}} dx dy \right)^2 < +\infty.$$

The  $H^{\frac{\alpha}{2}}$ -norm is defined as  $\|u\|_{H^{\frac{\alpha}{2}}} := \sqrt{\|u\|_{L^2}^2 + |u|_{H^{\frac{\alpha}{2}}}^2}$ .

**Lemma 7.4.** *Let  $\alpha \in (0, 2)$ ,  $u_0 \in L^\infty \cap L^1(\mathbb{R}^d)$ , (1.3)–(1.4) hold, and  $u^\alpha$  be the unique entropy solution to (1.1). Then, for all  $T > 0$ ,*

$$\int_{\mathbb{R}^d} \Phi(u^\alpha(x, T)) dx + |\varphi(u^\alpha)|_{L^2(0, T; H^{\frac{\alpha}{2}})} \leq \int_{\mathbb{R}^d} \Phi(u_0(x)) dx,$$

where  $\Phi(u) := \int_0^u \varphi(\tau) d\tau$  is nonnegative, convex, and 0 at 0.

*Proof.* Take  $\eta = \Phi$  in (7.2) where  $\Phi(u) = \int_0^u \varphi(\tau) d\tau$  is  $C^1$ , nonnegative and convex, since  $\varphi$  is nondecreasing and  $\varphi(0) = 0$ . Using also Lemma 7.3 and the continuity of  $u^\alpha$  in time with values in  $L^1(\mathbb{R}^d)$ , as in Remark 2.1, we find that for all  $\phi \in C_c^\infty(\mathbb{R}^{d+1})$ ,

$$(7.5) \quad \begin{aligned} & \int_{Q_T} \Phi(u^\alpha) \partial_t \phi + q_f^\Phi(u^\alpha) \cdot \nabla \phi dx dt \\ & + \int_{Q_T} q_\varphi^\Phi(u^\alpha) \mathcal{L}_r^\alpha[\phi] + \varphi(u^\alpha) \mathcal{L}^{\alpha, r}[\varphi(u^\alpha)] \phi dx dt \\ & + \int_{\mathbb{R}^d} \Phi(u_0(x)) \phi(x, 0) dx \geq \int_{\mathbb{R}^d} \Phi(u^\alpha(x, T)) \phi(x, T) dx. \end{aligned}$$

Then take  $\phi(x, t) = \gamma_R(x)$ , where  $R > 0$  and  $\gamma_R$  is an approximation of  $\mathbf{1}_{|x| < R}$  such that  $\gamma_R \in C_c^\infty(\mathbb{R}^d)$ ,  $\{\gamma_R\}_{R > 0}$  is bounded in  $W^{2, \infty}(\mathbb{R}^d)$ ,  $\gamma_R \rightarrow 1$  in  $W_{\text{loc}}^{2, \infty}(\mathbb{R}^d)$  as  $R \rightarrow +\infty$ . It is obvious that the  $\nabla$ - and  $\mathcal{L}_r^\alpha$ -terms in (7.5) vanish as  $R \rightarrow +\infty$ , since  $q_g^\Phi \in W_{\text{loc}}^{1, \infty}(\mathbb{R})$  and  $q_g^\Phi(0) = 0$  for  $g = f, \varphi$ . For the  $\mathcal{L}^{\alpha, r}$ -term, a standard computation<sup>5</sup> shows that

$$(7.6) \quad - \int_{\mathbb{R}^d} u \mathcal{L}^{\alpha, r}[v] dx = \frac{G_d(\alpha)}{2} \iint_{|x-y|>r} \frac{(u(x) - u(y))(v(x) - v(y))}{|x - y|^{d+\alpha}} dx dy,$$

---

5

$$\begin{aligned} & - 2 \iint_{|z|>r} \frac{u(x+z) - u(x)}{|z|^{d+\alpha}} v(x) dz dx \\ & = \iint_{|x-y|>r} u(x) v(x) \frac{dx dy}{|x-y|^{d+\alpha}} + \iint_{|x-y|>r} u(y) v(y) \frac{dx dy}{|x-y|^{d+\alpha}} \\ & \quad - \iint_{|x-y|>r} u(x) v(y) \frac{dx dy}{|x-y|^{d+\alpha}} - \iint_{|x-y|>r} u(y) v(x) \frac{dx dy}{|x-y|^{d+\alpha}} \\ & = \iint_{|x-y|>r} \frac{(u(x) - u(y))(v(x) - v(y))}{|x-y|^{d+\alpha}} dx dy. \end{aligned}$$



for all  $u, v \in L^2(\mathbb{R}^d)$  and  $r > 0$ . Hence, by the dominated convergence theorem,

$$\begin{aligned} & \int_{Q_T} \varphi(u) \mathcal{L}^{\alpha, r}[\varphi(u)] \gamma_R \, dx \, dt \\ &= \frac{G_d(\alpha)}{2} \int_0^T \iint_{|x-y|>r} (\varphi(u(x, t)) - \varphi(u(y, t))) \\ & \quad \cdot (\varphi(u(x, t)) \gamma_R(x) - \varphi(u(y, t)) \gamma_R(y)) \frac{dx \, dy}{|x-y|^{d+\alpha}} \, dt \\ & \rightarrow \frac{G_d(\alpha)}{2} \int_0^T \iint_{|x-y|>r} \frac{(\varphi(u(x, t)) - \varphi(u(y, t)))^2}{|x-y|^{d+\alpha}} \, dx \, dy \, dt \end{aligned}$$

as  $R \rightarrow +\infty$ . We then find in the limit that

$$\begin{aligned} & \int_{\mathbb{R}^d} \Phi(u^\alpha(x, T)) \, dx \\ &+ \frac{G_d(\alpha)}{2} \int_0^T \iint_{|x-y|>r} \frac{(\varphi(u(x, t)) - \varphi(u(y, t)))^2}{|x-y|^{d+\alpha}} \, dx \, dy \, dt \\ &\leq \int_{\mathbb{R}^d} \Phi(u_0(x)) \, dx. \end{aligned}$$

The proof is complete by sending  $r \rightarrow 0$  and using the monotone convergence theorem.  $\square$

From this energy type of estimate, we have the following result:

**Lemma 7.5.** *Under the assumptions of Lemma 7.1,  $\varphi(w) \in L^2(0, T; H^1)$ .*

*Proof.* Recall first that by (2.7) and a  $(L^1, L^\infty)$ -interpolation argument,  $\{u^\alpha\}_{\alpha \in (0, 2)}$  is bounded in  $L^2(0, T; L^2)$ . Using in addition the preceding lemma, we find a constant  $C$  such that for all  $\alpha \in (0, 2)$ ,

$$\|u^\alpha\|_{L^2(0, T; H^{\frac{\alpha}{2}})} \leq C.$$

Using (2.1) and (7.6), it is classical that

$$\|u^\alpha\|_{L^2(0, T; H^{\frac{\alpha}{2}})}^2 = \int_{Q_T} (1 + |\xi|^\alpha) |\mathcal{F}(u^\alpha(\cdot, t))(\xi)|^2 \, d\xi \, dt.$$

Now we use the following inequalities: for all  $1 \leq \beta \leq \alpha < 2$  and all  $\xi \in \mathbb{R}^d$ ,

$$(1 + |\xi|^\beta) \leq (1 + |\xi|)^\beta \leq (1 + |\xi|)^\alpha \leq 2^{\alpha-1} (1 + |\xi|^\alpha).$$

We deduce that

$$\int_{Q_T} (1 + |\xi|^\beta) |\mathcal{F}(u^\alpha(\cdot, t))(\xi)|^2 \, d\xi \, dt \leq 2^{\alpha-1} C.$$

Going back to the integral formula (again via (2.1) and (7.6)),

$$\|u^\alpha\|_{L^2(0, T; L^2)}^2 + \frac{G_d(\beta)}{2} \int_0^T \iint_{\mathbb{R}^{2d}} \frac{(u^\alpha(x, t) - u^\alpha(y, t))^2}{|x-y|^{d+\beta}} \, dx \, dy \, dt \leq 2^{\alpha-1} C.$$

By Fatou's lemma, applied for  $\alpha \rightarrow 2$  with fixed  $\beta$ ,

$$\|w\|_{L^2(0, T; L^2)}^2 + \frac{G_d(\beta)}{2} \int_0^T \iint_{\mathbb{R}^{2d}} \frac{(w(x, t) - w(y, t))^2}{|x-y|^{d+\beta}} \, dx \, dy \, dt \leq 2C.$$

Finally, Fatou's lemma applied to the Fourier formula shows that

$$\begin{aligned} 2C &\geq \liminf_{\beta \rightarrow 2} \int_{Q_T} (1 + |\xi|^\beta) |\mathcal{F}(w(\cdot, t))(\xi)|^2 \, d\xi \, dt \\ &\geq \int_{Q_T} (1 + |\xi|^2) |\mathcal{F}(w(\cdot, t))(\xi)|^2 \, d\xi \, dt. \end{aligned}$$

The proof is complete.  $\square$

We end by the proof of Theorem 3.3.

*Proof of Theorem 3.3.* Let  $u, w \in L^\infty(Q_T) \cap C([0, T]; L^1)$  be defined in Lemma 7.1. By previous lemmas, they are entropy solutions of (1.5) and (1.6), respectively. By uniqueness (cf. [29, 9, 26]), the whole sequences converge and the proof is complete.  $\square$

## 8. OPTIMAL EXAMPLE

In this last section, we give an example for which Theorems 3.1 and 3.8 are optimal. Let us consider, for every  $\alpha \in [0, 2]$  and  $\gamma, a > 0$ ,

$$(8.1) \quad \begin{cases} \partial_t u + a(-\Delta)^{\frac{\alpha}{2}} u = 0, \\ u(x, 0) = \gamma \mathbf{1}_Q(\gamma^{-1} x), \end{cases}$$

where  $Q := [-1, 1]^d$ . This is (1.1) with  $u_0$  as above,  $f \equiv 0$  and  $\varphi' \equiv a$ . Notice that

$$(8.2) \quad \begin{cases} \|u_0\|_{L^1} = 2^d \gamma^{d+1}, \\ |u_0|_{BV} = d 2^d \gamma^d, \\ E_i(u_0) = d 2^d \gamma^d (1 + \ln^i \frac{2}{d}) \mathbf{1}_{\gamma > d}, \quad (i = 1, 2), \end{cases}$$

where  $E_i(u_0)$  is defined in (3.3).

**8.1. Optimality of Theorem 3.1.** Let us fix  $\alpha \in [0, 2]$  and let us use the notation  $u =: u_a$ . Given  $T > 0$  and other parameters  $b, c > 0$ , we define

$$\omega_{a-b} := \begin{cases} |a^{\frac{1}{\alpha}} - b^{\frac{1}{\alpha}}|, & \alpha > 1, \\ |a \ln a - b \ln b|, & \alpha = 1, \\ |a - b|, & \alpha < 1, \end{cases}$$

$$\sigma_T := \begin{cases} T^{\frac{1}{\alpha}}, & \alpha > 1, \\ T |\ln T|, & \alpha = 1, \\ T, & \alpha < 1, \end{cases}$$

$$\sigma_\gamma := \begin{cases} \gamma^d, & \alpha > 1, \\ \gamma^d \ln \gamma, & \alpha = 1, \\ \gamma^{d+1-\alpha}, & \alpha < 1. \end{cases}$$

We also introduce the ‘‘smallest Lipschitz constant’’ of  $a \rightarrow u_a$  at  $a = c$ :

$$\text{Lip}_\varphi(u; c) := \limsup_{a, b \rightarrow c} \frac{\|u_a - u_b\|_{C([0, T]; L^1)}}{|a - b|}.$$

Theorem 3.1 and (8.2) imply that the function  $a \geq 0 \rightarrow u_a \in C([0, T]; L^1)$  is continuous at  $a = 0$  and locally Lipschitz continuous for  $a > 0$  with for all  $c > 0$ ,

$$\begin{aligned} \|u_a - u_b\|_{C([0, T]; L^1)} &= O(\omega_{a-b}) \quad \text{as } a, b \rightarrow 0, \\ \text{Lip}_\varphi(u; c) &= O(\sigma_T) \quad \text{as } T \rightarrow 0, \\ \text{Lip}_\varphi(u; c) &= O(\sigma_\gamma) \quad \text{as } \gamma \rightarrow +\infty, \end{aligned}$$

while all the respective remaining parameters are fixed. The result below states that these estimates are optimal.

**Proposition 8.1.** *Let  $\alpha \in [0, 2]$  and  $c > 0$ .*

- (i) *For all  $T, \gamma > 0$ ,  $\liminf_{a, b \rightarrow 0} \frac{\|u_a - u_b\|_{C([0, T]; L^1)}}{\omega_{a-b}} > 0$ .*
- (ii) *For all  $\gamma > 0$ ,  $\liminf_{T \rightarrow 0} \frac{\text{Lip}_\varphi(u; c)}{\sigma_T} > 0$ .*

(iii) For all  $T > 0$ ,  $\liminf_{\gamma \rightarrow +\infty} \frac{\text{Lip}_\alpha(u; c)}{\sigma_\gamma} > 0$ .

*Remark 8.2.* (1) This result shows that the modulus of continuity in  $\varphi - \psi$  derived in (3.6) is optimal. It also shows that the  $T$ - and  $u_0$ -dependencies of this modulus are optimal in the limits  $T \rightarrow 0$  or  $\frac{\|u_0\|_{L^1}}{|u_0|_{BV}} \rightarrow +\infty$  (recall that  $\frac{\|u_0\|_{L^1}}{|u_0|_{BV}} \sim \gamma$  by (8.2)).

(2) The quantities  $E_1(u_0)$  and  $\|u_0\|_{L^1}^{1-\alpha} |u_0|_{BV}^\alpha$  in the optimal estimate (3.6) are always  $O(\|u_0\|_{L^1} + |u_0|_{BV})$ . But as  $\frac{\|u_0\|_{L^1}}{|u_0|_{BV}} \rightarrow +\infty$  (see (3.4) and (3.3)), we have the better estimate  $(\|u_0\|_{L^1} + |u_0|_{BV}) o(1)$ .

**8.2. Optimality of Theorem 3.8.** Let us now use the notation  $u =: u^\alpha$  to emphasize the dependence on  $\alpha$ . Given  $\lambda \in (0, 2)$ , we define

$$\tilde{\sigma}_M := \begin{cases} M^{\frac{1}{\lambda}} |\ln M|, & \lambda > 1, \\ M \ln^2 M, & \lambda = 1, \\ M, & \lambda < 1, \end{cases}$$

$$\tilde{\sigma}_\gamma := \begin{cases} \gamma^d, & \lambda > 1, \\ \gamma^d \ln^2 \gamma, & \lambda = 1, \\ \gamma^{d+1-\lambda} \ln \gamma, & \lambda < 1, \end{cases}$$

where  $M := Ta$ . We also consider the ‘‘smallest Lipschitz constant’’ of  $\alpha \rightarrow u^\alpha$  at  $\alpha = \lambda$  defined in (3.8). Then, Theorem 3.8 and (8.2) imply that for all  $\lambda \in (0, 2)$ ,

$$\text{Lip}_\alpha(u; \lambda) = O(\tilde{\sigma}_M) \quad \text{as } M \rightarrow 0,$$

$$\text{Lip}_\alpha(u; \lambda) = O(\tilde{\sigma}_\gamma) \quad \text{as } \gamma \rightarrow +\infty,$$

while all the respective remaining parameters are fixed. The result below states that these estimates are optimal.

**Proposition 8.3.** *Let  $T, a > 0$ ,  $M = Ta$ , and  $\lambda \in (0, 2)$ . There exist  $M_0, \gamma_0 > 0$  such that:*

- (i) For all  $\gamma_0 \geq \gamma > 0$ ,  $\liminf_{M \rightarrow 0} \frac{\text{Lip}_\alpha(u; \lambda)}{\tilde{\sigma}_M} > 0$ .  
(ii) For all  $M_0 \geq M > 0$ ,  $\liminf_{\gamma \rightarrow +\infty} \frac{\text{Lip}_\alpha(u; \lambda)}{\tilde{\sigma}_\gamma} > 0$ .

*Remark 8.4.* (1) This result shows that the  $M$ - and  $u_0$ -dependencies in (3.9) are optimal at the limits  $M = T \|\varphi'\|_\infty \rightarrow 0$  or  $\frac{\|u_0\|_{L^1}}{|u_0|_{BV}} \rightarrow +\infty$ .

(2) Here again the quantities  $\|u_0\|_{L^1}^{1-\alpha} |u_0|_{BV}^\alpha \ln \frac{\|u_0\|_{L^1}}{|u_0|_{BV}}$  and  $E_2(u_0)$  in (3.9) are always  $O(\|u_0\|_{L^1} + |u_0|_{BV})$ , but as  $\frac{\|u_0\|_{L^1}}{|u_0|_{BV}} \rightarrow +\infty$ , we have the better estimate  $(\|u_0\|_{L^1} + |u_0|_{BV}) o(1)$ .

### 8.3. Proofs.

*Proof of Proposition 8.1.* Let us prove each items in order.

**1. Item (i).** Let us first assume that  $T = \gamma = 1$ . The general case will follow from a rescaling argument given at the end of the proof. Let us define

$$(8.3) \quad \mathcal{E}_Q := \int_Q u_a(x, 1) dx - \int_Q u_b(x, 1) dx.$$

Since  $\|u_a - u_b\|_{C([0,1]; L^1)} \geq \|u_a(\cdot, 1) - u_b(\cdot, 1)\|_{L^1} \geq |\mathcal{E}_Q|$ , it suffices to show that  $\liminf_{a, b \rightarrow 0} \frac{|\mathcal{E}_Q|}{\omega_{a-b}} > 0$ . It is well-known that  $u_a(x, t) = \mathcal{F}^{-1}(e^{-ta} |\cdot|^\alpha) * \mathbf{1}_Q(x)$

and  $u_b(x, t) = \mathcal{F}^{-1}(e^{-tb|\cdot|^\alpha}) * \mathbf{1}_Q(x)$ . A short computation shows that

$$\begin{aligned}
 \mathcal{E}_Q &= \int \mathcal{F}^{-1}(e^{-a|\cdot|^\alpha} - e^{-b|\cdot|^\alpha}) (\mathbf{1}_Q * \mathbf{1}_Q) dx \\
 (8.4) \quad &= \int (e^{-a|\xi|^\alpha} - e^{-b|\xi|^\alpha}) (\mathcal{F}\mathbf{1}_Q)^2 d\xi \\
 &= \int \int_0^1 (b-a) |\xi|^\alpha e^{-(\tau a + (1-\tau)b)|\xi|^\alpha} (\mathcal{F}\mathbf{1}_Q)^2 d\tau d\xi,
 \end{aligned}$$

where  $\mathcal{F}\mathbf{1}_Q(\xi) = \prod_{i=1}^d \frac{\sin(2\pi\xi_i)}{\pi\xi_i}$ ,  $\xi = (\xi_1, \dots, \xi_d)$ . We now give separate arguments for the cases  $\alpha < 0$ ,  $\alpha = 1$ , and  $\alpha > 0$ .

**a.** *The case  $\alpha < 1$ .* The proof is obvious since  $0 < \int |\xi|^\alpha (\mathcal{F}\mathbf{1}_Q)^2 d\xi < +\infty$ .

**b.** *The case  $\alpha > 1$ .* Note that  $|\xi|^\alpha \leq d^{(\alpha-1)} \sum_{i=1}^d |\xi_i|^\alpha$ . Hence, by (8.4),

$$|\mathcal{E}_Q| \geq I_{a,b} \int \int_0^1 |a-b| |\xi_1|^\alpha e^{-d^{\alpha-1}(\tau a + (1-\tau)b)|\xi_1|^\alpha} \frac{\sin^2(2\pi\xi_1)}{\xi_1^2} d\tau d\xi_1$$

where  $I_{a,b} = \frac{1}{\pi^2} \prod_{i=2}^d \int e^{-d^{\alpha-1}(a \vee b)|\xi_i|^\alpha} \frac{\sin^2(2\pi\xi_i)}{\pi^2 \xi_i^2} d\xi_i$ . Since  $e^{-d^{\alpha-1}(a \vee b)|\xi_i|^\alpha} \rightarrow 1$  as  $a, b \rightarrow 0$ ,

$$I_{a,b} \geq C_0 := \frac{1}{2\pi^2} \prod_{i=2}^d \int \frac{\sin^2(2\pi\xi_i)}{\pi^2 \xi_i^2} d\xi_i > 0,$$

for all  $a, b > 0$  sufficiently small. Hence, assuming e.g. that  $a > b$ , we get

$$\begin{aligned}
 |\mathcal{E}_Q| &\geq C_0 \underbrace{\int a |\xi_1|^{\alpha-2} e^{-d^{\alpha-1}a|\xi_1|^\alpha} \sin^2(2\pi\xi_1) d\xi_1}_{=: I_a} \\
 (8.5) \quad &- C_0 \int b |\xi_1|^{\alpha-2} e^{-d^{\alpha-1}b|\xi_1|^\alpha} \sin^2(2\pi\xi_1) d\xi_1.
 \end{aligned}$$

Before continuing, notice that this estimate is valid for  $\alpha = 1$ ; this is will be useful later. Let us continue the case  $\alpha > 1$  by changing variables,

$$I_a = a^{\frac{1}{\alpha}} \int |\xi_1|^{\alpha-2} e^{-d^{\alpha-1}|\xi_1|^\alpha} \sin^2(2\pi a^{-\frac{1}{\alpha}} \xi_1) d\xi_1.$$

Doing the same for the  $b$ -integral and adding and subtracting term,

$$\begin{aligned}
 (8.6) \quad |\mathcal{E}_Q| &\geq C_0 (a^{\frac{1}{\alpha}} - b^{\frac{1}{\alpha}}) \int |\xi_1|^{\alpha-2} e^{-d^{\alpha-1}|\xi_1|^\alpha} \sin^2(2\pi a^{-\frac{1}{\alpha}} \xi_1) d\xi_1 \\
 &+ C_0 b^{\frac{1}{\alpha}} \int |\xi_1|^{\alpha-2} e^{-d^{\alpha-1}|\xi_1|^\alpha} \left\{ \sin^2(2\pi a^{-\frac{1}{\alpha}} \xi_1) - \sin^2(2\pi b^{-\frac{1}{\alpha}} \xi_1) \right\} d\xi_1 \\
 &=: C_0 (a^{\frac{1}{\alpha}} - b^{\frac{1}{\alpha}}) I_1 + C_0 b^{\frac{1}{\alpha}} I_2.
 \end{aligned}$$

By a Taylor expansion and an integration by parts,

$$\begin{aligned}
 |I_2| &\leq 2\pi (a^{\frac{1}{\alpha}} - b^{\frac{1}{\alpha}}) \left| \int_0^1 \int a_{\alpha,\tau}^{-2} \underbrace{|\xi_1|^{\alpha-2} \xi_1 e^{-d^{\alpha-1}|\xi_1|^\alpha}}_{=: f(\xi_1)} \right. \\
 &\quad \cdot \underbrace{2 \sin(2\pi a_{\alpha,\tau}^{-1} \xi_1) \cos(2\pi a_{\alpha,\tau}^{-1} \xi_1)}_{=: \sin(4\pi a_{\alpha,\tau}^{-1} \xi_1)} d\xi_1 d\tau \left. \right| \\
 &\leq \frac{1}{2} b^{-\frac{1}{\alpha}} (a^{\frac{1}{\alpha}} - b^{\frac{1}{\alpha}}) \left| \int_0^1 \int f'(\xi_1) \cos(4\pi a_{\alpha,\tau}^{-1} \xi_1) d\xi_1 d\tau \right|,
 \end{aligned}$$

where  $a_{\alpha,\tau} := \tau a^{\frac{1}{\alpha}} + (1-\tau)b^{\frac{1}{\alpha}}$  and  $f'$  is integrable when  $\alpha > 1$ . We deduce that  $C_0 b^{\frac{1}{\alpha}} I_2 = (a^{\frac{1}{\alpha}} - b^{\frac{1}{\alpha}}) o(1)$  as  $a, b \rightarrow 0$ , since for fixed  $\tau$ ,  $\cos(4\pi a_{\alpha,\tau}^{-1} \cdot)$  converges to its zero mean value in  $L^\infty$ -weak- $\star$ . By a similar argument  $\sin^2(2\pi a^{-\frac{1}{\alpha}} \cdot)$  also weakly- $\star$  converges to its positive mean value  $m$  and hence

$$\lim_{a,b \rightarrow 0} I_1 = m \int |\xi_1|^{\alpha-2} e^{-d^{\alpha-1} |\xi_1|^\alpha} d\xi_1 > 0.$$

We thus conclude the result from (8.6).

**c.** *The case  $\alpha = 1$ .* We restart from (8.5) assuming again that  $a > b$ ,  $a, b$  small. This time we cut  $I_a$  into three pieces.

$$I_a = \int_{1 < |\xi_1| < a^{-1}} \dots + \int_{|\xi_1| < 1} \dots + \int_{|\xi_1| > a^{-1}} \dots$$

We do the same for the  $b$ -integral and we get

$$\begin{aligned} |\mathcal{E}_Q| &\geq C_0 \int_{1 < |\xi_1| < a^{-1}} a |\xi_1|^{-1} e^{-a|\xi_1|} \sin^2(2\pi \xi_1) d\xi_1 \\ &\quad - C_0 \int_{1 < |\xi_1| < b^{-1}} b |\xi_1|^{-1} e^{-b|\xi_1|} \sin^2(2\pi \xi_1) d\xi_1 \\ &\quad + C_0 \left( \int_{|\xi_1| < 1} \dots - \int_{|\xi_1| < 1} \dots \right) + C_0 \left( \int_{|\xi_1| > a^{-1}} \dots - \int_{|\xi_1| > b^{-1}} \dots \right). \end{aligned}$$

The last two terms are  $O(a-b) = (b \ln b - a \ln a) o(1)$  as  $a, b \rightarrow 0$ . To show this, we follow line by line the arguments of **a** and **b** respectively, noting that all integrals are well-defined because of the new domains of integration. Let now  $I$  denote the remaining term. Recalling that  $a > b$ ,

$$\begin{aligned} I &= C_0 \int_{1 < |\xi_1| < a^{-1}} |\xi_1|^{-1} (a e^{-a|\xi_1|} - b e^{-b|\xi_1|}) \sin^2(2\pi \xi_1) d\xi_1 \\ &\quad - C_0 \int_{a^{-1} < |\xi_1| < b^{-1}} b |\xi_1|^{-1} e^{-b|\xi_1|} \sin^2(2\pi \xi_1) d\xi_1 \\ &=: I_1 + I_2. \end{aligned}$$

Note that

$$\begin{aligned} |I_2| &\leq C_0 \int_{a^{-1} < |\xi_1| < b^{-1}} b |\xi_1|^{-1} d\xi_1 \\ &= 2C_0 b (\ln a - \ln b) \leq 2C_0 (a - b) = (b \ln b - a \ln a) o(1) \end{aligned}$$

as  $a, b \rightarrow 0$ . Hence it remains to show that  $\liminf_{a,b \rightarrow 0} \frac{I_1}{b \ln b - a \ln a} > 0$ . Since

$$\begin{aligned} a e^{-a|\xi_1|} - b e^{-b|\xi_1|} &= (a-b) \int_0^1 \{1 - (\tau a + (1-\tau)b) |\xi_1|\} e^{-(\tau a + (1-\tau)b) |\xi_1|} d\tau \\ &\geq \begin{cases} \frac{e^{-1}}{2} (a-b) & \text{for all } |\xi_1| \leq \frac{a^{-1}}{2}, \\ 0 & \text{for all } |\xi_1| \leq a^{-1}, \end{cases} \end{aligned}$$

we find that

$$\begin{aligned} I_1 &\geq C_0 \frac{e^{-1}}{2} (a-b) \int_{1 < |\xi_1| < \frac{a^{-1}}{2}} |\xi_1|^{-1} \sin^2(2\pi \xi_1) d\xi_1 \\ &\geq C_0 \frac{e^{-1}}{2} (a-b) \int_{1 < |\xi_1| < \frac{9}{8}} |\xi_1|^{-1} \sin^2(2\pi \xi_1) d\xi_1 \\ &\quad + \frac{C_0 e^{-1}}{8} (a-b) \int_{\frac{9}{8} < |\xi_1| < \frac{a^{-1}}{2}} |\xi_1|^{-1} d\xi_1. \end{aligned}$$

To get the last line, we have used that since  $\sin^2(2\pi \cdot) \geq \frac{1}{4}$  on  $E := [\frac{1}{8}, \frac{3}{8}] + \frac{1}{2}\mathbb{Z}$  and  $\mathbb{R} \setminus E = E + \frac{1}{4}$ ,

$$(8.7) \quad \begin{aligned} \int_{\frac{9}{8} < |\xi_1| < \frac{a-1}{2}} |\xi_1|^{-1} \sin^2(2\pi \xi_1) d\xi_1 &= \int_{\frac{9}{8} < |\xi_1|} \underbrace{|\xi_1|^{-1} \mathbf{1}_{|\xi_1| < \frac{a-1}{2}}}_{=: g(|\xi_1|)} \sin^2(2\pi \xi_1) d\xi_1 \\ &\geq \frac{1}{4} \int_{E \cap \{\frac{9}{8} < |\xi_1|\}} g(|\xi_1|) d\xi_1 \\ &\geq \frac{1}{8} \int_{\frac{9}{8} < |\xi_1|} g(|\xi_1|) d\xi_1, \end{aligned}$$

by translation and since  $g$  is nonincreasing. It follows that

$$I_1 \geq \tilde{C}_0 (b-a) \ln a + O(a-b) \geq \tilde{C}_0 (b \ln b - a \ln a) + (b \ln b - a \ln a) o(1)$$

as  $a, b \rightarrow 0$ , where  $\tilde{C}_0 = \frac{C_0}{8e}$ . Here we have used that  $b \ln a \geq b \ln b$ , and since  $b \ln b - a \ln a > 0$  for small  $a > b > 0$ , the proof of (i) is complete under the assumption that  $T = \gamma = 1$ .

For general  $T, \gamma > 0$  fixed, the result follows from rescaling. Let  $w(x, t) = \gamma^{-1} u(\gamma x, Tt)$  and note that

$$\begin{cases} w_t + T \gamma^{-\alpha} a (-\Delta)^{\frac{\alpha}{2}} w = 0, \\ w(x, 0) = \mathbf{1}_Q(x). \end{cases}$$

Set  $\mu := T \gamma^{-\alpha}$  and  $w =: w_{\mu a}$  to emphasize the dependence on the new “nonlinearity”  $\mu a$ . Then by the results of the  $T = \gamma = 1$  case above,

$$\liminf_{a, b \rightarrow 0} \frac{\|w_{\mu a} - w_{\mu b}\|_{C([0,1]; L^1)}}{\omega_{\mu a - \mu b}} > 0,$$

where  $\omega_{\dots}$  is defined on page 33. By a simple change of variables,

$$\|u_a - u_b\|_{C([0,T]; L^1)} = \gamma^{d+1} \|w_{\mu a} - w_{\mu b}\|_{C([0,1]; L^1)},$$

and since  $\omega_{\mu a - \mu b} \sim \omega_{a-b}$  as  $a, b \rightarrow 0$  ( $\mu$  is fixed!), (i) holds for any  $T, \gamma > 0$ .

**2. Item (ii).** Let us adapt the preceding arguments. We only give the proof for the case  $c = 1$  and  $\gamma = 1$ , noting that the general result then easily follows from the rescaling  $w(x, t) = \gamma^{-1} u(\gamma x, \gamma^\alpha c^{-1} t)$ . We have

$$(8.8) \quad \begin{aligned} \text{Lip}_\varphi(u; 1) &\geq \lim_{a, b \rightarrow 1} \left| \frac{\int_Q u_a(x, T) dx - \int_Q u_b(x, T) dx}{a - b} \right| \\ &= \int T |\xi|^\alpha e^{-T|\xi|^\alpha} (\mathcal{F}\mathbf{1}_Q)^2 d\xi, \end{aligned}$$

thanks to (8.4) written for time  $T$ . Arguing as before, there exists  $C_0 > 0$  such that for all sufficiently small  $T$ ,

$$\text{Lip}_\varphi(u; 1) \geq C_0 \underbrace{\int T |\xi_1|^\alpha e^{-d^{\alpha-1} T |\xi_1|^\alpha} \frac{\sin^2(2\pi \xi_1)}{\xi_1^2} d\xi_1}_{=: I}.$$

It remains to prove that  $\liminf_{T \rightarrow 0} \frac{I}{\sigma_T} > 0$ . The case  $\alpha < 1$  follows from a direct passage to the limit, and the case  $\alpha > 1$  follows, as before, from the change of variable  $T^{\frac{1}{\alpha}} \xi_1 \rightarrow \xi_1$  and the  $L^\infty$ -weak- $\star$  convergence of  $\sin^2(2\pi T^{-\frac{1}{\alpha}} \cdot)$ . For the  $\alpha = 1$  case, we again split  $I$  into three parts,

$$I = \int_{1 < |\xi_1| < T^{-1}} \cdots + \int_{|\xi_1| < 1} \cdots + \int_{|\xi_1| > T^{-1}} \cdots$$

As in case (i), the two last terms are  $O(T) = T |\ln T| o(1)$  as  $T \rightarrow 0$ , and the remaining integral can be bounded below as in (8.7) by

$$\tilde{C}_0 \int_{\frac{9}{8} < |\xi_1| < T^{-1}} T |\xi_1|^{-1} d\xi_1 + O(T) \geq \tilde{C}_0 T |\ln T| + T |\ln T| o(1) \quad \text{as } T \rightarrow 0,$$

where  $\tilde{C}_0 > 0$  is another constant independent of  $T$  small enough. The proof is complete.

**3. Item (iii).** We assume that  $c = T = 1$ , and note that the general case follows by an easy rescaling argument using  $w(x, t) = u(c^{\frac{1}{\alpha}} T^{\frac{1}{\alpha}} x, Tt)$ . We start as in the preceding case, considering this time integrals on  $\gamma Q$  in (8.3). As in (8.4), we find that

$$\begin{aligned} \mathcal{E}_{\gamma Q} &= \int_{\gamma Q} u_a(x, 1) dx - \int_{\gamma Q} u_b(x, 1) dx \\ &= \gamma \int \int_0^1 (b-a) |\xi|^\alpha e^{-(\tau a + (a-\tau b)) |\xi|^\alpha} (\mathcal{F}\mathbf{1}_{\gamma Q})^2 d\tau d\xi, \end{aligned}$$

and hence

$$\begin{aligned} \text{Lip}_\varphi(u; 1) &\geq \lim_{a, b \rightarrow 1} \left| \frac{\mathcal{E}_{\gamma Q}}{a-b} \right| \\ &= \gamma \int |\xi|^\alpha e^{-|\xi|^\alpha} (\mathcal{F}\mathbf{1}_{\gamma Q})^2 d\xi \\ &= \gamma^{2d+1} \int |\xi|^\alpha e^{-|\xi|^\alpha} (\mathcal{F}\mathbf{1}_Q(\gamma \xi))^2 d\xi. \end{aligned}$$

After changing variables  $\gamma \xi \rightarrow \xi$ , we then get that

$$\text{Lip}_\varphi(u; 1) \geq \gamma^{d+1} \int \gamma^{-\alpha} |\xi|^\alpha e^{-\gamma^{-\alpha} |\xi|^\alpha} (\mathcal{F}\mathbf{1}_{\gamma Q})^2 d\xi.$$

This is the same expression as in (8.8) with  $\gamma^{-\alpha}$  in place of  $T$ . Note that

$$\gamma^{d+1} \sigma_T |_{T=\gamma^{-\alpha}} = \sigma_\gamma$$

according to the definitions of  $\sigma_T$  and  $\sigma_\gamma$  on page 33, and hence by the proof of (ii) we have that  $\liminf_{\gamma \rightarrow +\infty} \frac{\text{Lip}_\varphi(u; 1)}{\sigma_\gamma} > 0$ . The proof of (iii) is complete.  $\square$

*Remark 8.5.* In the proof of Corollary 3.6, a rescaling in time transformed the continuous dependence estimate (3.5) into the time continuity estimate (3.7). Hence, we leave it to the reader to verify that the same rescaling allow us to prove that (8.1) is also an example showing the optimality of Corollary 3.6.

*Proof of Proposition 8.3.* We adapt the arguments of the proof of Proposition 8.1(i).

**1. Item (i).** To avoid confusion with the proof of (ii) below, we denote the fixed parameter  $\gamma$  by  $\tilde{\gamma}$ . We consider the new difference

$$\mathcal{E}_Q := \int_{\tilde{\gamma} Q} u^\alpha(x, T) dx - \int_{\tilde{\gamma} Q} u^\beta(x, T) dx$$

with moving powers  $\alpha, \beta \in (0, 2)$  and time  $T$ . We let  $M = T a$  and argue as in (8.4) to see that

$$\begin{aligned} \mathcal{E}_Q &= \tilde{\gamma}^{2d+1} \int (e^{-M |\xi|^\alpha} - e^{-M |\xi|^\beta}) (\mathcal{F}\mathbf{1}_Q)^2 (\tilde{\gamma} \xi) d\xi \\ &= \tilde{\gamma}^{2d+1} \int \int_0^1 (\beta - \alpha) (\ln |\xi|) M |\xi|^{\tau \alpha + (1-\tau) \beta} e^{-M |\xi|^{\tau \alpha + (1-\tau) \beta}} (\mathcal{F}\mathbf{1}_Q)^2 (\tilde{\gamma} \xi) d\tau d\xi, \end{aligned}$$

so that

$$(8.9) \quad \text{Lip}_\alpha(u; \lambda) \geq \tilde{\gamma}^{2d+1} \left| \underbrace{\int (\ln |\xi|) M |\xi|^\lambda e^{-M |\xi|^\lambda} (\mathcal{F}\mathbf{1}_Q)^2 (\tilde{\gamma} \xi) d\xi}_{=: I} \right|.$$

To complete the proof, we must show that  $\liminf_{M \rightarrow 0} \frac{|I|}{\tilde{\sigma}_M} > 0$ .

**a. The case  $\lambda < 1$ .** Now  $\lim_{M \rightarrow 0} \frac{I}{M} = \int (\ln |\xi|) |\xi|^\lambda (\mathcal{F}\mathbf{1}_Q)^2 (\tilde{\gamma} \xi) d\xi =: I_{\tilde{\gamma}}$ . Since  $\mathcal{F}\mathbf{1}_Q(0) \neq 0$ ,  $\lim_{\tilde{\gamma} \rightarrow 0} I_{\tilde{\gamma}} = +\infty$ , and we see that (i) holds for  $\tilde{\gamma}$  small enough.

In the other two cases we split  $I$  in two,  $I = \int_{|\xi| < 1} \dots + \int_{|\xi| > 1} \dots$ . The first integral is of order  $O(M) = \tilde{\sigma}_M o(1)$  as  $M \rightarrow 0$ , by a direct passage to the limit. Arguing as in the preceding proof, the last integral can be bounded from below by

$$\int_{|\xi_1| > 1} (\ln |\xi_1|) M |\xi_1|^{\lambda-2} e^{-d^{\lambda-1} M |\xi_1|^\lambda} \sin^2(2\pi \tilde{\gamma} \xi_1) d\xi_1 =: J,$$

up to some positive multiplicative constant  $C_0$  independent of  $M$  small enough. Note that  $C_0$  will also depend on  $\tilde{\gamma} > 0$  which is constant in this proof. Hence it suffices to show that  $\liminf_{M \rightarrow 0} \frac{J}{\tilde{\sigma}_M} > 0$ .

**b. The case  $\lambda > 1$ .** By the change of variables  $M^{\frac{1}{\lambda}} \xi_1 \rightarrow \xi_1$ ,

$$\begin{aligned} J &= M^{\frac{1}{\lambda}} \int_{|\xi_1| > M^{\frac{1}{\lambda}}} (\ln |\xi_1|) |\xi_1|^{\lambda-2} e^{-d^{\lambda-1} |\xi_1|^\lambda} \sin^2(2\pi M^{-\frac{1}{\lambda}} \tilde{\gamma} \xi_1) d\xi_1 \\ &\quad - \lambda^{-1} M^{\frac{1}{\lambda}} \ln(M) \int_{|\xi_1| > M^{\frac{1}{\lambda}}} |\xi_1|^{\lambda-2} e^{-d^{\lambda-1} |\xi_1|^\lambda} \sin^2(2\pi M^{-\frac{1}{\lambda}} \tilde{\gamma} \xi_1) d\xi_1. \end{aligned}$$

It is clear that the first term is  $O(M^{\frac{1}{\lambda}}) = M^{\frac{1}{\lambda}} |\ln M| o(1)$  as  $M \rightarrow 0$ , and that the second one has the expected behavior due to  $L^\infty$ -weak- $\star$  convergence arguments.

**c. The case  $\lambda = 1$ .** We write  $J = \int_{|\xi_1| > M^{-1}} \dots + \int_{1 < |\xi_1| < M^{-1}} \dots$ . The first term is  $O(M |\ln M|) = M (\ln^2 M) o(1)$  as  $M \rightarrow 0$  by the change of variables argument of the  $\lambda > 1$  case. For the remaining term, we argue as in (8.7), using this time that  $\sin^2(2\pi \tilde{\gamma} \cdot)$  is bounded below by  $\frac{1}{4}$  on  $\tilde{\gamma}^{-1} E$ . Taking  $N$  so large that the new function  $g$  (defined below) is nonincreasing on  $(\frac{4N+1}{8} \tilde{\gamma}^{-1}, +\infty)$ , we get a lower bound of the form

$$\begin{aligned} &\int_{1 < |\xi_1| < M^{-1}} (\ln |\xi_1|) M |\xi_1|^{-1} e^{-M |\xi_1|} \sin^2(2\pi \tilde{\gamma} \xi_1) d\xi_1 \\ &\geq e^{-1} M \int_{1 < |\xi_1| < M^{-1}} \underbrace{(\ln |\xi_1|) |\xi_1|^{-1}}_{=: g(|\xi_1|)} \sin^2(2\pi \tilde{\gamma} \xi_1) d\xi_1 \\ &= e^{-1} M \int_{\frac{4N+1}{8} \tilde{\gamma}^{-1} < |\xi_1| < M^{-1}} \dots + e^{-1} M \int_{1 < |\xi_1| < \frac{4N+1}{8} \tilde{\gamma}^{-1}} \dots \\ &\geq \frac{e^{-1}}{8} M \int_{\frac{4N+1}{8} \tilde{\gamma}^{-1} < |\xi_1| < M^{-1}} |\xi_1|^{-1} \ln |\xi_1| d\xi_1 + O(M) \\ &= \frac{e^{-1}}{4} M \ln^2 M + M (\ln^2 M) o(1) \quad \text{as } M \rightarrow +\infty. \end{aligned}$$

The proof of (i) is now complete.



**2.** *Item (ii).* To avoid confusion with the preceding proof, we denote the fixed parameter  $M = T a$  by  $\tilde{M}$ . Then, by (8.9),

$$(8.10) \quad \text{Lip}_\alpha(u; \lambda) \geq \tilde{M} \underbrace{\left| \gamma^{2d+1} \int (\ln |\xi|) |\xi|^\lambda e^{-\tilde{M} |\xi|^\lambda} (\mathcal{F}\mathbf{1}_Q)^2(\gamma \xi) d\xi \right|}_{=: I}$$

and it suffices to show that  $\liminf_{\gamma \rightarrow +\infty} \frac{|I|}{\sigma_\gamma} > 0$ .

**a.** *The case  $\lambda > 1$ .* Since  $\ln |\xi|$  has different signs inside and outside the unit ball, we split the integral  $I$  in two,

$$I = \int_{|\xi| < 1} \dots + \int_{|\xi| > 1} \dots =: I_1 + I_2.$$

By the inequality  $|\ln |\xi|| |\xi|^\lambda \leq d^{\lambda-1} \sum_{i=1}^d |\ln |\xi_i|| |\xi_i|^\lambda$  for  $|\xi| < 1$  and the change of variables  $\gamma \xi_j \rightarrow \xi_j$  for  $j \neq i$ , we find that

$$\begin{aligned} |I_1| &\leq d^{\lambda-1} \gamma^{2d+1} \sum_{i=1}^d \int_{|\xi| < 1} |\ln |\xi_i|| |\xi_i|^\lambda (\mathcal{F}\mathbf{1}_Q)^2(\gamma \xi) d\xi \\ &\leq d^{\lambda-1} \gamma^d \sum_{i=1}^d \left\{ \int_{|\xi_i| < 1} \frac{|\ln |\xi_i|| |\xi_i|^\lambda}{\pi^2 \xi_i^2} d\xi_i \prod_{j \neq i} \int \frac{\sin^2(2\pi \xi_j)}{\pi^2 \xi_j^2} d\xi_j \right\}. \end{aligned}$$

Here we also have used that  $\sin^2(2\pi \gamma \xi_i) \leq 1$ . It follows that  $\limsup_{\gamma \rightarrow +\infty} \frac{|I_1|}{\gamma^d} \leq C(d, \lambda)$ , a constant that does not depend on  $\tilde{M}$ .

Let us now see that  $I_2$  is the dominant term provided that the fixed parameter  $\tilde{M}$  is chosen sufficiently small. We have

$$I_2 \geq \gamma^{2d+1} \int (\ln |\xi_1|) |\xi_1|^\lambda e^{-\tilde{M} |\xi_1|^\lambda} (\mathcal{F}\mathbf{1}_Q)^2(\gamma \xi) d\xi,$$

and then, letting  $\xi_\gamma := (\xi_1, \gamma^{-1} \xi_2, \dots, \gamma^{-1} \xi_d)$  and changing variables  $\gamma \xi_i \rightarrow \xi_i$  for  $i \neq 1$ , we find that

$$\begin{aligned} I_2 &\geq \gamma^d \int \left\{ (\ln |\xi_1|) |\xi_1|^{\lambda-2} \int e^{-\tilde{M} |\xi_\gamma|^\lambda} \frac{1}{\pi^2} \prod_{i=2}^d \frac{\sin^2(2\pi \xi_i)}{\pi^2 \xi_i^2} d\xi_2 \dots d\xi_d \right\} \\ &\quad \cdot \sin^2(2\pi \gamma \xi_1) d\xi_1. \end{aligned}$$

By  $L^\infty$ -weak- $\star$  convergence arguments,  $\liminf_{\gamma \rightarrow +\infty} \frac{I_2}{\gamma^d} \geq \tilde{m} I_{\tilde{M}}$ , where

$$\tilde{m} := \frac{1}{\pi^2} \int_0^1 \sin^2(2\pi \xi_1) d\xi_1 \prod_{i=2}^d \int \frac{\sin^2(2\pi \xi_i)}{\pi^2 \xi_i^2} d\xi_i > 0$$

and  $I_{\tilde{M}} := \int (\ln |\xi_1|) |\xi_1|^{\lambda-2} e^{-\tilde{M} |\xi_1|^\lambda} d\xi_1$ . Since  $\lim_{\tilde{M} \rightarrow 0} I_{\tilde{M}} = +\infty$ , it suffices to fix  $\tilde{M} > 0$  small to get (ii) in the  $\lambda > 1$  case.

**b.** *The case  $\lambda < 1$ .* We restart from (8.10), change the variables  $\gamma \xi \rightarrow \xi$ , and pass to the limit as  $\gamma \rightarrow +\infty$ . The result follows.

**c.** *The case  $\lambda = 1$ .* Let us rewrite  $I$  in (8.10) as

$$I = \underbrace{\gamma^{2d+1} \int_{\gamma^{-1} < |\xi| < 1} \dots}_{=: -J_1} + \underbrace{\gamma^{2d+1} \int_{|\xi| > 1} \dots}_{=: J_2} + \gamma^{2d+1} \int_{|\xi| < \gamma^{-1}} \dots$$

By the arguments of the  $\lambda < 1$  case, the last integral is of order  $O(\gamma^d \ln \gamma) = \gamma^d (\ln^2 \gamma) o(1)$  as  $\gamma \rightarrow +\infty$ . For  $J_2$ , we use that

$$\left| \int_{|\xi|>1} \dots \right| \leq \int_{|\xi_1|, \dots, |\xi_d| > \frac{1}{\sqrt{d}}} |\dots| \leq \sum_{i=1}^d \int_{\xi: |\xi_i| > \frac{1}{\sqrt{d}}} |\dots| = d \int_{\xi: |\xi_1| > \frac{1}{\sqrt{d}}} |\dots|$$

by symmetry of the  $I$ -integrand (cf. (8.10)). We then bound  $|\xi| |\ln |\xi|| e^{-\tilde{M}|\xi|}$  by some constant  $C$ , change the variables  $\gamma \xi_i \rightarrow \xi_i$  for  $i \neq 1$ , get

$$|J_2| \leq d C \gamma^d \int_{|\xi_1| > \frac{1}{\sqrt{d}}} \frac{d\xi_1}{\pi^2 \xi_1^2} \prod_{i=2}^d \int \frac{\sin^2(2\pi \xi_i)}{\pi^2 \xi_i^2} d\xi_i$$

and conclude that  $J_2 = O(\gamma^d) = \gamma^d (\ln^2 \gamma) o(1)$  as  $\gamma \rightarrow +\infty$ .

Since  $J_1 > 0$ , it remains to show that  $\liminf_{\gamma \rightarrow +\infty} \frac{J_1}{\gamma^d \ln^2 \gamma} > 0$ . It will be convenient to use the notation  $\hat{\xi} := (\xi_2, \dots, \xi_d)$ . By the change of variables  $\gamma \xi \rightarrow \xi$  and the inequality  $|\xi| \xi_1^{-2} \geq |\xi|^{-1}$ ,

$$J_1 \geq -\gamma^d \int_{1 < |\xi| < \gamma} \ln(\gamma^{-1} |\xi|) |\xi|^{-1} f(\hat{\xi}) \sin^2(2\pi \xi_1) d\xi,$$

where  $f(\hat{\xi}) := \frac{e^{-\tilde{M}}}{\pi^2} \prod_{i=2}^d \frac{\sin^2(2\pi \xi_i)}{\pi^2 \xi_i^2}$ . Let us restrict to the domain of integration

$$A := \left\{ (\xi_1, \hat{\xi}) \text{ s.t. } 1 < |\xi_1| < \frac{1}{2} \sqrt{\gamma^2 - |\hat{\xi}|^2} \text{ and } \epsilon^2 < |\hat{\xi}| < \epsilon \right\},$$

where  $\epsilon > 0$  is fixed and so small that  $A \subset \{1 < |\xi| < \gamma\}$  and  $C_0 := \min_A f(\hat{\xi}) > 0$ . Then,

$$J_1 \geq -C_0 \gamma^d \int_{\epsilon^2 < |\hat{\xi}| < \epsilon} \int_{1 < |\xi_1| < \frac{1}{2} \sqrt{\gamma^2 - |\hat{\xi}|^2}} \ln(\gamma^{-1} |\xi|) |\xi|^{-1} \sin^2(2\pi \xi_1) d\xi_1 d\hat{\xi}.$$

Arguing as in (8.7),

$$\begin{aligned} J_1 &\geq -C_0 \gamma^d \int_{\epsilon^2 < |\hat{\xi}| < \epsilon} \int_{1 < |\xi_1| < \frac{9}{8}} \ln(\gamma^{-1} |\xi|) |\xi|^{-1} \sin^2(2\pi \xi_1) d\xi_1 d\hat{\xi} \\ &\quad - \tilde{C}_0 \gamma^d \int_{\epsilon^2 < |\hat{\xi}| < \epsilon} \int_{\frac{9}{8} < |\xi_1| < \frac{1}{2} \sqrt{\gamma^2 - |\hat{\xi}|^2}} \ln(\gamma^{-1} |\xi|) |\xi|^{-1} d\xi_1 d\hat{\xi} \\ &=: J_{1,1} + J_{1,2}, \end{aligned}$$

where  $\tilde{C}_0 = \frac{C_0}{8}$ . Since  $\ln(\gamma^{-1} |\xi|) = \ln |\xi| - \ln \gamma$ , it is easy to see that  $J_{1,1} = O(\gamma^d \ln \gamma) = \gamma^d (\ln^2 \gamma) o(1)$  as  $\gamma \rightarrow +\infty$ . Let us verify that  $\liminf_{\gamma \rightarrow +\infty} \frac{J_{1,2}}{\gamma^d \ln^2 \gamma} > 0$ . If  $\gamma$  is large enough, then for all  $\xi \in A$ ,  $\gamma^{-1} |\xi| \leq \gamma^{-1} (|\xi_1| + |\hat{\xi}|) < 1$ . We can then use that

$$-\ln(\gamma^{-1} |\xi|) |\xi|^{-1} \geq -\ln(\gamma^{-1} (|\xi_1| + |\hat{\xi}|)) (|\xi_1| + |\hat{\xi}|)^{-1},$$

and by integrating the right-hand side, we get

$$\begin{aligned} J_{1,2} &\geq \tilde{C}_0 \gamma^d \int_{\epsilon^2 < |\hat{\xi}| < \epsilon} \ln^2 \left( \gamma^{-1} \left( \frac{9}{8} + |\hat{\xi}| \right) \right) d\hat{\xi} \\ &\quad - \tilde{C}_0 \gamma^d \int_{\epsilon^2 < |\hat{\xi}| < \epsilon} \ln^2 \left( \frac{1}{2} \sqrt{1 - \gamma^{-2} |\hat{\xi}|^2} + \gamma^{-1} |\hat{\xi}| \right) d\hat{\xi} \\ &\geq \tilde{C}_0 \gamma^d \ln^2 \gamma + \gamma^d (\ln^2 \gamma) o(1) \quad \text{as } \gamma \rightarrow +\infty. \end{aligned}$$

Here  $\tilde{C}_0$  is another positive constant independent of  $\gamma$  large enough. The proof is complete.  $\square$

## APPENDIX A. PROOF OF SOME ESTIMATES USED TO PROVE THEOREM 3.1

*Proof of (6.19) and (6.20).* Recall that  $\Omega_\xi(\cdot)$  is defined on page 19 and  $\chi_a^b$  in (6.15). For  $A$ , we use that

$$\begin{aligned} \int \|\Omega_\xi(u)\|_{L^1(Q_T)} d\xi &= \int \int_{Q_T} \left| \int \chi_0^{u(x,t)}(\zeta) \omega_\delta(\xi - \zeta) d\zeta \right| dx dt d\xi \\ &= \int \int_{Q_T} \left| \chi_0^{u(x,t)}(\zeta) \right| \int \omega_\delta(\xi - \zeta) d\xi dx dt d\xi = \|u\|_{L^1(Q_T)}. \end{aligned}$$

For  $B$ , we consider  $\{u_n\}_n \subset C([0, T]; W^{1,1})$  converging to  $u$  in  $C([0, T]; L^1)$  and such that  $\int_{Q_T} |\nabla u_n| \rightarrow |u|_{L^1(0, T; BV)}$ . Then

$$\begin{aligned} &\int \int_{Q_T} |\nabla \Omega_\xi(u_n)| dx dt d\xi \\ &= \int \int_{Q_T} \omega_\delta(\xi - u_n(x, t)) |\nabla u_n(x, t)| dx dt d\xi = \int_{Q_T} |\nabla u_n|, \end{aligned}$$

so that

$$\begin{aligned} \int |\Omega_\xi(u)|_{L^1(0, T; BV)} d\xi &\leq \int \int_0^T \left\{ \liminf_{n \rightarrow +\infty} \int_{\mathbb{R}^d} |\nabla \Omega_\xi(u_n)| dx \right\} dt d\xi \\ &\leq \lim_{n \rightarrow +\infty} \int \int_{Q_T} |\nabla \Omega_\xi(u_n)| dx dt d\xi = |u|_{L^1(0, T; BV)}, \end{aligned}$$

due to the lower semi-continuity of the  $BV$ -semi-norm with respect to the  $L^1$ -norm and to Fatou's lemma <sup>(6)</sup>.  $\square$

*Proof of (6.21).* Recall that  $\mathcal{E}_3$  and  $\mathcal{E}_3(\delta)$  are defined in (6.2) and (6.17), respectively. See also the original assumption (3.2) of the theorem, and the simplifying assumption (6.1). First we define

$$F_v(x, t, y, s, z, \xi) := \operatorname{sgn}(v - u) \frac{\chi_{v(x,t)}^{v(x+z,t)}(\xi)}{|z|^{d+\alpha}} \phi^{\epsilon, \nu}.$$

Let us recall that  $F_v$  is integrable on  $Q_T^2 \times \{|z| > r\} \times \mathbb{R}$  since  $\int |\chi_a^b(\xi)| d\xi = |b - a|$ . Hence, by Fubini the function

$$G_v(\xi) := \int_{Q_T^2} \int_{\{|z| > r\}} F_v(x, t, y, s, z, \xi) dz dw$$

is integrable with respect to  $\xi \in \mathbb{R}$ . But, by (6.16), (6.17),

$$\mathcal{E}_3 = G_d(\alpha) \int G_v(\xi) \psi'(\xi) - G_u(\xi) \varphi'(\xi) d\xi$$

and  $\mathcal{E}_3(\delta) = G_d(\alpha) \int \psi'(\xi) G_v * \omega_\delta(\xi) - \varphi'(\xi) G_u * \omega_\delta(\xi) d\xi$ , where  $*$  is the convolution product in  $\mathbb{R}$ . Since  $\omega_\delta$  is an approximate unit, the convolution products inside the integral respectively converge to  $G_v$  and  $G_u$  in  $L^1(\mathbb{R})$  as  $\delta \rightarrow 0$ . Using in addition that  $\varphi'$  and  $\psi'$  are bounded by (6.1),  $\lim_{\delta \rightarrow 0} \mathcal{E}_3(\delta) = \mathcal{E}_3$ .  $\square$

<sup>6</sup>For the reverse inequality, we use that, at all fixed time and for all  $\Phi \in C_c^1(\mathbb{R}^d, \mathbb{R}^d)$  such that  $|\Phi| \leq 1$ ,

$$\begin{aligned} \int |\Omega_\xi(u)|_{BV} d\xi &\geq \int \int_{\mathbb{R}^d} \Omega_\xi(u) \operatorname{div} \Phi dx d\xi \\ &= \int \int_{\mathbb{R}^d} \int \chi_0^u(\zeta) \omega_\delta(\xi - \zeta) \operatorname{div} \Phi d\zeta dx d\xi = \int_{\mathbb{R}^d} u \operatorname{div} \Phi dx, \end{aligned}$$

and next we take the supremum with respect to  $\Phi$ .

## APPENDIX B. SOME TECHNICAL LEMMAS

**Lemma B.1.** For all  $a, b > 0$ ,  $|a - b| (-\ln(a \vee b))^+ \leq |a - b| + |a \ln a - b \ln b|$ .

*Proof.* We assume without loss of generality that  $a \vee b = a$  and that  $a \leq e^{-1}$  (the result is trivial otherwise). Then

$$|a \ln(a) - b \ln(b)| = - \int_b^a (1 + \ln(\tau)) d\tau = -|a - b| - \int_b^a \ln(\tau) d\tau,$$

since  $1 + \ln(\tau)$  is negative, and hence

$$|a - b| + |a \ln(a) - b \ln(b)| \geq -|a - b| \ln(a),$$

since the logarithm is nondecreasing. This completes the proof.  $\square$

**Lemma B.2.** For all  $x > 0$ ,  $a, b \neq 0$  and  $c > 0$ ,

- (i)  $\left| \frac{x^a - 1}{a} - \frac{x^b - 1}{b} \right| \leq |a - b| (1 \vee x^a \vee x^b) \ln^2 x$ ,
- (ii)  $\lim_{a, b \rightarrow c} \frac{1}{(a-b)^2} \left| \frac{x^{2a}}{2a} + \frac{x^{2b}}{2b} - \frac{2x^{a+b}}{a+b} \right| \leq C x^{2c} (1 + \ln^2 x)$ ,

where  $C = C(c)$ .

*Proof.* (i) Let  $f(a) = \frac{x^a - 1}{a}$ . Observe that  $(\ln x) \int_0^1 x^{\tau a} d\tau = f(a)$  by a Taylor expansion of  $x^a$  at  $a = 0$ . It then follows that by differentiating under the integral sign that  $f'(a) = \ln^2 x \int_0^1 \tau x^{\tau a} d\tau$  and

$$\begin{aligned} f(a) - f(b) &= (a - b) \int_0^1 f'(\tau a + (1 - \tau)b) d\tau \\ &= (a - b) (\ln x)^2 \int_0^1 \int_0^1 \tilde{\tau} x^{\tilde{\tau}(\tau a + (1 - \tau)b)} d\tilde{\tau} d\tau. \end{aligned}$$

Since  $x^{\tilde{\tau}(\tau a + (1 - \tau)b)} \leq 1 \vee x^{a \vee b} \vee x^{a \wedge b}$ , we find that

$$|f(a) - f(b)| \leq |a - b| (\ln x)^2 (1 \vee x^a \vee x^b),$$

and the proof of (i) is complete.

(ii) Note that

$$\begin{aligned} \frac{x^{2a}}{2a} + \frac{x^{2b}}{2b} - \frac{2x^{a+b}}{a+b} &= \frac{b(a+b)x^{2a} + a(a+b)x^{2b} - 4abx^{a+b}}{2ab(a+b)} \\ &= \frac{(x^a - x^b)^2}{2(a+b)} + \frac{(bx^a - ax^b)^2}{2ab(a+b)}. \end{aligned}$$

The first term satisfies  $|x^a - x^b| \leq |a - b| (x^a \vee x^b) |\ln x|$ . Moreover, by adding and subtracting terms and the inequality  $\frac{(A+B)^2}{2} \leq A^2 + B^2$ , we find that the second term is bounded by

$$(bx^a - ax^b)^2 \leq \frac{1}{2} (a - b)^2 (x^a + x^b)^2 + \frac{1}{2} (a + b)^2 (x^a - x^b)^2.$$

The now proof follows from these two inequalities.  $\square$

## APPENDIX C. THE PROOF OF LEMMA 7.3

*Proof.* The if part follows by approximating the Kruzkov entropy  $|u - k|$  by a family of smooth convex entropies  $\eta_n(u) = \sqrt{(u - k)^2 + n^{-2}} - n^{-1}$ . The functions  $\eta_n(u)$  and  $\eta'_n(u)$  are locally uniformly bounded and converge pointwise to  $|u - k|$  and the everywhere representative of its weak derivative given by (2.4). Since  $u$  is bounded and  $\eta_n(u)$  satisfies (7.2), we use the dominated convergence theorem to pass to the limit and find that  $|u - k|$  satisfies (2.6).

To prove the only if part, we note that we may approximate (locally uniformly) any  $C^1$ -convex entropy  $\eta$  by a family of piecewise linear functions  $\tilde{\eta}_n$  of the form

$$(C.1) \quad \tilde{\eta}(u) = a + b u + \sum_{i=1}^m c_i |u - k_i|$$

where  $a, b, k_i \in \mathbb{R}$ ,  $c_i \geq 0$ , and  $m \in \mathbb{N}$ . See e.g. [23, p. 27] for a proof. We need to refine this construction to ensure everywhere convergence of the derivatives  $\tilde{\eta}'_n$ . Consider the everywhere defined representative of  $\tilde{\eta}'$  defined by

$$(C.2) \quad \tilde{\eta}'(u) = b + \sum_{i=1}^m c_i \operatorname{sgn}(u - k_i),$$

where the  $\operatorname{sgn}$ -function is everywhere defined by (2.4). Since  $\eta'$  is continuous, it can be approximated uniformly on compact sets by piecewise constant functions of the form (C.2). Take such a sequence  $\{\tilde{\eta}'_n\}_n$  that converges on  $|u| \leq \|u_0\|_{L^\infty}$ , and redefine  $\{\tilde{\eta}_n\}_n$  to be the primitives such that  $\tilde{\eta}_n(0) = \eta(0)$ , i.e. functions of the form (C.1). It follows that both  $\tilde{\eta}_n$  and  $\tilde{\eta}'_n$  converge pointwise everywhere and uniformly towards  $\eta$  and  $\eta'$  on  $|u| \leq \|u_0\|_{L^\infty}$ .

Next we note that the left hand side of the entropy inequality (7.2) is linear with respect to  $\eta$ , that (7.2) holds with  $\eta(u) = a + b u$  since  $u$  is a weak solution of (1.1), and that (7.2) holds with  $\eta(u) = c_i |u - k_i|$  by the Kruzkov inequality (2.6) since  $c_i \geq 0$ . The reader may then check that (7.2) also holds with  $\eta = \tilde{\eta}$  and the everywhere representative of  $\tilde{\eta}'$  given by (C.2).

Since  $u$  is bounded,  $\tilde{\eta}_n(u)$  and  $\tilde{\eta}'_n(u)$  will be uniformly bounded, and we may use the dominated convergence theorem to pass to the limit in (7.2) with  $\eta = \tilde{\eta}_n$  to find that (7.2) holds also for the  $\eta$  in the limit. The proof is complete.  $\square$

## REFERENCES

- [1] N. Alibaud. Entropy formulation for fractal conservation laws. *J. Evol. Equ.* 7(1):145-175, 2007.
- [2] N. Alibaud and B. Andreianov. Non-uniqueness of weak solutions for the fractal Burgers equation. *Ann. Inst. H. Poincaré Anal. Non Linéaire* 27(4):997-1016, 2010.
- [3] N. Alibaud, S. Cifani and E. R. Jakobsen. Continuous dependence estimates for nonlinear fractional convection-diffusion equations. *SIAM Journal on Mathematical Analysis* 44(2): 603-632, 2012.
- [4] N. Alibaud, J. Droniou and J. Vovelle. Occurrence and non-appearance of shocks in fractal Burgers equation. *Journal of Hyperbolic Differential Equations* 4(3):479-499, 2007.
- [5] P. Biler, T. Funaki and W. Woźniński. Fractal Burgers Equations. *J. Diff. Equations* 148:9-46, 1998.
- [6] P. Biler, G. Karch and C. Imbert. Fractal porous medium equation. *C. R. Acad. Sci. Paris, Ser. I* 349 (2011) 641-645.
- [7] P. Biler, G. Karch and R. Monneau. A nonlinear diffusion of dislocation density and self-similar solutions. *Commun. Math. Physics* 294:145-168, 2010.
- [8] F. Bouchut and B. Perthame. Kruzkov's estimates for scalar conservation laws revisited. *Trans. Amer. Math. Soc.* 7:2847-2870, 1998.
- [9] J. Carrillo. Entropy Solutions for nonlinear Degenerate Problems. *Arch. Ration. Mech. Anal.* 147:269-361, 1999.
- [10] L. Caffarelli and J.-L. Vásquez. Nonlinear porous medium flow with fractional potential pressure. *Arch. Ration. Mech. Anal.* 202(2): 537-565, 2011.
- [11] G. Chen and K. H. Karlsen.  $L^1$ -framework for continuous dependence and error estimates for quasilinear anisotropic degenerate parabolic equations. *Trans. Amer. Math. Soc.* 358(3):937-963 (electronic), 2006.
- [12] S. Cifani and E. R. Jakobsen. Entropy formulation for degenerate fractional order convection-diffusion equations. *Ann. Inst. H. Poincaré Anal. Non Linéaire* 28(3):413-441, 2011.
- [13] S. Cifani and E. R. Jakobsen. On numerical methods and error estimates for degenerate fractional convection-diffusion equations. *Submitted for publication*, 2012.
- [14] P. Clavin. Instabilities and nonlinear patterns of overdriven detonations in gases. *Nonlinear PDE's in Condensed Matter and Reactive Flows. Kluwer*, 49-97, 2002.

- [15] B. Cockburn and G. Gripenberg. Continuous Dependence on the nonlinearities of Solutions of Degenerate Parabolic Equations. *J. Diff. Equations* 151:231–251, 1999.
- [16] B. Cockburn, G. Gripenberg and S.-O. Londen. On convergence of entropy solutions to a single conservation law. *J. Diff. Equations* 128(1):206–251, 1996.
- [17] R. Cont and P. Tankov. Financial modelling with jump processes. *Chapman & Hall/CRC Financial Mathematics Series*, Chapman & Hall/CRC, Boca Raton (FL), 2004.
- [18] C. Dafermos. *Hyperbolic Conservation Laws in Continuum Physics*. Springer, 2005.
- [19] A. de Pablo, F. Quiros, A. Rodriguez and J. L. Vázquez. A fractional porous medium equation. *Advances in Mathematics* 226(2):1378–1409, 2011.
- [20] E. DiBenedetto. *Degenerate parabolic equations*. Springer-Verlag, New-York, 1993.
- [21] J. Droniou, T. Gallouët and J. Vovelle. Global solution and smoothing effect for a nonlocal regularization of a hyperbolic equation. *J. Evol. Equ.* 4(3):479–499, 2003.
- [22] J. Droniou and C. Imbert. Fractal first order partial differential equations. *Arch. Ration. Mech. Anal.* 182(2):299–331, 2006.
- [23] H. Holden and N. H. Risebro. Front Tracking for Hyperbolic Conservation Laws. *Applied Mathematical Sciences*, 152, Springer, 2007.
- [24] C. Imbert. A non-local regularization of first order Hamilton-Jacobi equations. *J. Diff. Equations* 211:214–246, 2005.
- [25] V. G. Jakubowski and P. Wittbold. On a nonlinear elliptic/parabolic integro-differential equation with  $L^1$ -data. *J. Differential Equations* 197(2):427–445, 2003.
- [26] K. H. Karlsen and N. H. Risebro. On the uniqueness and stability of entropy solutions of nonlinear degenerate parabolic equations with rough coefficients. *Discrete Contin. Dyn. Syst.*, 9(5):1081–1104, 2003.
- [27] K. H. Karlsen and S. Ulusoy. Stability of entropy solutions for Lévy mixed hyperbolic parabolic equations. *Electronic Journal of Differential Equations*, 2011(116):1–23, 2011.
- [28] A. Kiselev, F. Nazarov and R. Shterenberg. Blow up and regularity for fractal Burgers equation. *Dynamics of PDE* 5(3):211–240, 2008.
- [29] S. N. Kruzkov. First order quasilinear equations with several independent variables. *Math. Sb. (N.S.)* 81(123):228–255, 1970.
- [30] N. N. Kuznetsov. Accuracy of some approximate methods for computing the weak solutions of a first-order quasi-linear equation. *USSR. Comput. Math. Phys.* 16:105–119, 1976.
- [31] E. R. Jakobsen and K. H. Karlsen. Continuous dependence estimates for viscosity solutions of integro-PDEs. *J. Differential Equations* 212(2): 278–318, 2005.
- [32] P.-L. Lions, B. Perthame and E. Tadmor. A kinetic formulation of multidimensional scalar conservation laws and related local equations. *J. Amer. Math. Soc.* 7:169–191, 1994.
- [33] C. Rohde and W.-A. Yong. The nonrelativistic limit in radiation hydrodynamics. I. Weak entropy solutions for a model problem. *J. Differential Equations* 234(1):91–109, 2007.
- [34] P. Rosenau. Extending hydrodynamics via the regularization of the Chapman-Enskog expansion. *Phys. Rev. A* 40:7193–7196, 1989.
- [35] K. Sato. Lévy processes and infinitely divisible distributions. *Cambridge Studies in Advanced Mathematics*, 68, Cambridge University Press, 1999.
- [36] A. Sayah, Equations d’Hamilton-Jacobi du premier ordre avec termes intégral-différentiels, Parties I et II, *Comm P.D.E.* 16: 1057–1093, 1991.
- [37] S. Schochet and E. Tadmor. Regularized Chapman-Enskog expansion for scalar conservation laws. *Arch. Rational Mech. Anal.* 119:95–107, 1992.
- [38] W. Woyczyński. Lévy processes in the physical sciences. *Lévy processes*, 241–266, Birkhäuser, Boston, 2001.

(Nathael Alibaud) UMR CNRS 6623, UNIVERSITÉ DE FRANCHE-COMTÉ, 16 ROUTE DE GRAY, 25 030 BESANÇON CEDEX, FRANCE AND DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, PRINCE OF SONGKLA UNIVERSITY, HAT YAI, SONGKHLA, THAILAND, 90112

E-mail address: [nathael.alibaud@ens2m.fr](mailto:nathael.alibaud@ens2m.fr)

(Simone Cifani) DEPARTMENT OF MATHEMATICS, NORWEGIAN UNIVERSITY OF SCIENCE AND TECHNOLOGY (NTNU), N-7491 TRONDHEIM, NORWAY

E-mail address: [simone.cifani@math.ntnu.no](mailto:simone.cifani@math.ntnu.no)

(Espen R. Jakobsen) DEPARTMENT OF MATHEMATICS, NORWEGIAN UNIVERSITY OF SCIENCE AND TECHNOLOGY (NTNU), N-7491 TRONDHEIM, NORWAY

E-mail address: [erj@math.ntnu.no](mailto:erj@math.ntnu.no)

URL: <http://www.math.ntnu.no/~erj/>