



**HAL**  
open science

## Bang-bang property for time optimal control of semilinear heat equation

Kim Dang Phung, Lijuan Wang, Can Zhang

► **To cite this version:**

Kim Dang Phung, Lijuan Wang, Can Zhang. Bang-bang property for time optimal control of semilinear heat equation. *Annales de l'Institut Henri Poincaré C, Analyse non linéaire*, 2014, pp.477-499. 10.1016/j.anihpc.2013.04.005 . hal-00845777

**HAL Id: hal-00845777**

**<https://hal.science/hal-00845777v1>**

Submitted on 17 Jul 2013

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Bang-bang property for time optimal control of semilinear heat equation

Kim Dang Phung<sup>\*</sup> , Lijuan Wang<sup>†</sup> and Can Zhang<sup>‡</sup>

## Abstract

This paper studies the bang-bang property for time optimal controls governed by semilinear heat equation in a bounded domain with control acting locally in a subset. Also, we present the null controllability cost for semilinear heat equation and an observability estimate from a positive measurable set in time for the linear heat equation with potential.

**Keywords.** semilinear heat equation; time optimal control; bang-bang property; observability estimate from measurable sets.

## 1 Introduction and main result

This paper continues the investigations carried out in [14]. Our main result deals with the bang-bang property for time optimal controls governed by semilinear heat equations with control acting locally. We complete the result in [14] in two directions: the nonlinearity of the equation; the geometry on which the equation takes place.

Let  $\Omega$  be a bounded connected open set of  $\mathbb{R}^n$ ,  $n \geq 1$ , with boundary  $\partial\Omega$  of class  $C^2$ . Let  $\omega$  be an open and non-empty subset of  $\Omega$  and denote  $1_\omega$  for the characteristic function of a set in the place where  $\cdot$  stays. Let  $y_0 \in L^2(\Omega)$  and  $v \in L^\infty(0, +\infty; L^2(\Omega))$ . Consider the following semilinear heat equation with initial data  $y_0$  and external force  $v$ :

$$\begin{cases} \partial_t y - \Delta y + f(y) = 1_\omega v & \text{in } \Omega \times (0, +\infty) , \\ y = 0 & \text{on } \partial\Omega \times (0, +\infty) , \\ y(\cdot, 0) = y_0 & \text{in } \Omega . \end{cases}$$

---

<sup>\*</sup>Université d'Orléans, Laboratoire MAPMO, CNRS UMR 7349, Fédération Denis Poisson, FR CNRS 2964, Bâtiment de Mathématiques, B.P. 6759, 45067 Orléans Cedex 2, France. (kim.dang\_phung@yahoo.fr).

<sup>†</sup>School of Mathematics and Statistics, Wuhan University, Wuhan 430072, China. (ljwang.math@whu.edu.cn). This work was partially supported by the National Science Foundation of China under grant 10971158 and 11161130003.

<sup>‡</sup>School of Mathematics and Statistics, Wuhan University, Wuhan 430072, China (canzhang@whu.edu.cn).

Existence and uniqueness of the solution  $y$  is ensured with the following assumptions:  $f : \mathbb{R} \rightarrow \mathbb{R}$  is globally Lipschitz and satisfies the “good-sign” condition  $f(s) s \geq 0$  for all  $s \in \mathbb{R}$  (and consequently,  $f(0) = 0$ ). In such case, for any  $T > 0$ , the solution  $y$  is in  $C([0, T]; L^2(\Omega))$  and the above equation holds in the sense of distributions in  $\Omega \times (0, T)$ .

Our motivation is a null control problem for semilinear heat equations which means that our goal consists to find  $v \in L^\infty(0, +\infty; L^2(\Omega))$  such that  $y(\cdot, T) = 0$  in  $\Omega$ .

The first natural null control problem solved in the literature is the following.

*Question 1:* what are the assumptions on  $f$  in order that the property

$$\left\{ \begin{array}{l} \forall y_0 \in L^2(\Omega), \forall T > 0, \exists M > 0, \exists v \in L^\infty(0, +\infty; L^2(\Omega)), \\ \text{such that } y(\cdot, T) = 0 \text{ in } \Omega \text{ and } \|v\|_{L^\infty(0, +\infty; L^2(\Omega))} \leq M \end{array} \right.$$

holds. Notice that the existence of a null control  $v$  gives the one of the bound  $M$ . This property is intensively studied in the literature (see e.g. [2],[8],[7]) and is called null controllability for semilinear heat equation. It holds for any nonlinear terms which are locally Lipschitz and slightly superlinear. Precisely, it is enough for  $f$  to satisfy  $f(0) = 0$  and

$$\lim_{|s| \rightarrow \infty} \frac{|f(s)|}{|s| \ln^{3/2}(1 + |s|)} = 0 .$$

In particular, if we assume that  $f$  is globally Lipschitz with  $f(0) = 0$ , then null controllability for the corresponding semilinear heat equation holds.

However, we can formulate another type of null control problem as follows.

*Question 2:* what are the assumptions on  $f$  in order that the property

$$\left\{ \begin{array}{l} \forall y_0 \in L^2(\Omega), \forall M > 0, \exists T > 0, \exists v \in L^\infty(0, +\infty; L^2(\Omega)), \\ \text{such that } y(\cdot, T) = 0 \text{ in } \Omega \text{ and } \|v\|_{L^\infty(0, +\infty; L^2(\Omega))} \leq M \end{array} \right.$$

holds. In this article, we will prove the existence of  $T$  and  $v$  under the assumption that  $f$  is globally Lipschitz and satisfies the “good-sign” condition. Once existence of a couple  $(y, v)$  is established for  $y_0 \in L^2(\Omega) \setminus \{0\}$  and  $M > 0$  given, via suitable assumption on  $f$ , we introduce the following admissible set of controls

$$\mathcal{V}_M = \left\{ v \in L^\infty(0, +\infty; L^2(\Omega)); \|v\|_{L^\infty(0, +\infty; L^2(\Omega))} \leq M \text{ and the solution } y \text{ corresponding to } v \text{ satisfies } y(\cdot, T) = 0 \text{ in } \Omega \text{ for some } T > 0 \right\} .$$

Among all the control functions  $v \in \mathcal{V}_M$ , we select the infimum of all such time:

$$T^* = \inf \{T; v \in \mathcal{V}_M\} ,$$

i.e., the minimal time needed to drive the system to rest with control functions in  $\mathcal{V}_M$ .

A control  $v^*$  such that the corresponding solution  $y$  satisfies  $y(\cdot, T^*) = 0$  in  $\Omega$  is called time optimal control. In this article, we shall prove the existence of a time

optimal control  $v^*$  under the assumption that  $f$  is globally Lipschitz and satisfies the “good-sign” condition.

Now, we are able to state our main result.

*Theorem 1 .- Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be a globally Lipschitz function satisfying  $f(s) s \geq 0$  for all  $s \in \mathbb{R}$ . Then for any  $y_0 \in L^2(\Omega) \setminus \{0\}$  and any  $M > 0$ , any time optimal control  $v^*$  satisfies the bang-bang property:  $\|v^*(\cdot, t)\|_{L^2(\Omega)} = M$  for a.e.  $t \in (0, T^*)$ .*

Clearly, bang-bang property is of high importance in optimal control theory as mentioned in [6] and [11]. In particular, the bang-bang property for certain time optimal controls governed by parabolic equations can be provided by making use of Pontryagin’s maximum principle (see [9],[10],[17]). Another approach to get bang-bang property for linear heat equation consists to follow a strategy based on null controllability with control functions acting on measurable set in time variable as in [12] and [16]. Recently, the authors in [1] established an observability inequality for the linear heat equation, where the observation is a subset of positive measure in space and time. And from which they obtained another kind of bang-bang property of time optimal problem for the linear heat equation with bounded controls in space and time. Naturally, the extension of this strategy for nonlinear parabolic equations requires a fixed point argument and an observability inequality for heat equations with space and time-dependent potentials.

This paper is organized as follows. Section 2 is devoted to the null controllability for semilinear heat equation with control functions acting on  $\omega \times E$  where  $|E| > 0$ . We present (see Theorem 2) and prove an estimate of the cost of the control functions when  $f$  is globally Lipschitz. Before giving the proof of Theorem 2, we recall the linear case and the observability estimate needed (see Theorem 4). In section 3, applying Theorem 2 in a very special case, we prove the existence for admissible control (see Theorem 5) when  $f$  is globally Lipschitz and satisfies the “good-sign” condition. Next we deduce the existence of time optimal (see Theorem 6). The proof of our main result, Theorem 1, concerning the bang-bang property for time optimal controls governed by semilinear heat equation with local control is given in section 4. Finally, in section 5, we prove the observability estimate of Theorem 4.

## 2 Null controllability for semilinear heat equation

The goal of this section is to present the null controllability for semilinear heat equation with control functions acting on  $\omega \times E$  where  $|E| > 0$ . A particular attention is given on the cost estimate.

*Theorem 2 .- Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be a globally Lipschitz function. Let  $0 \leq T_0 < T_1 < T_2$  and  $E \subset (T_1, T_2)$  with  $|E| > 0$ . Then for any  $\phi \in C([T_0, T_2], L^2(\Omega))$  and any  $w_0 \in L^2(\Omega)$ , there are a constant  $\kappa > 0$  and a function  $v_1 \in L^\infty(0, +\infty; L^2(\Omega))$  such that*

$$\|v_1\|_{L^\infty(0, +\infty; L^2(\Omega))} \leq \kappa \|w_0\|_{L^2(\Omega)}$$

and the solution  $w = w(x, t)$  of

$$\begin{cases} \partial_t w - \Delta w + f(\phi + w) - f(\phi) = 1_{|\omega \times E} v_1 & \text{in } \Omega \times (T_0, T_2) , \\ w = 0 & \text{on } \partial\Omega \times (T_0, T_2) , \\ w(\cdot, T_0) = w_0 & \text{in } \Omega , \end{cases}$$

satisfies  $w(\cdot, T_2) = 0$  in  $L^2(\Omega)$ . Further,

$$\kappa = e^{\tilde{K}} e^{c(T_1 - T_0)} e^{K(1+c+c(T_2 - T_1))} .$$

Here,  $c = c(f)$ ,  $K = K(\Omega, \omega) > 1$  and  $\tilde{K} = \tilde{K}(\Omega, \omega, E)$  are positive constants which do not depend on  $T_0$ .

*Remark 1* .- When  $E = (T_1, T_2)$ , then  $\kappa = e^{c(T_1 - T_0)} e^{K\left(1 + \frac{1}{T_2 - T_1} + c + c(T_2 - T_1)\right)}$ .

## 2.1 Linear case

In this section, we treat the case  $f(\phi + w) = aw + f(\phi)$  that is the linear heat equation with potential.

*Theorem 3* .- Let  $0 \leq T_0 < T_1 < T_2$  and  $E \subset (T_1, T_2)$  with  $|E| > 0$ . Let  $a \in L^\infty(\Omega \times (T_0, T_2))$ . Then for any  $z_0 \in L^2(\Omega)$ , there is a function  $v_0 \in L^\infty(\Omega \times (0, +\infty))$  such that the solution  $z = z(x, t)$  of

$$\begin{cases} \partial_t z - \Delta z + az = 1_{|\omega \times E} v_0 & \text{in } \Omega \times (T_0, T_2) , \\ z = 0 & \text{on } \partial\Omega \times (T_0, T_2) , \\ z(\cdot, T_0) = z_0 & \text{in } \Omega , \end{cases}$$

satisfies  $z(\cdot, T_2) = 0$  in  $L^2(\Omega)$ . Further,

$$\begin{aligned} \|v_0\|_{L^\infty(\Omega \times (0, +\infty))} &\leq e^{\tilde{K}} e^{(T_1 - T_0)\|a\|_{L^\infty(\Omega \times (T_0, T_1))}} \\ &\quad \times e^{K\left(1 + (T_2 - T_1)\|a\|_{L^\infty(\Omega \times (T_1, T_2))} + \|a\|_{L^\infty(\Omega \times (T_1, T_2))}^2\right)} \|z_0\|_{L^2(\Omega)} . \end{aligned}$$

Here,  $K = K(\Omega, \omega) > 1$  and  $\tilde{K} = \tilde{K}(\Omega, \omega, E)$  are positive constants which do not depend on  $T_0$ .

*Remark 2* .- When  $E = (T_1, T_2)$ , then  $\tilde{K} = K \frac{1}{T_2 - T_1}$ .

*Proof* .- We divide its proof into three steps. In the first step, we start to solve

$$\begin{cases} \partial_t z - \Delta z + az = 0 & \text{in } \Omega \times (T_0, T_1) , \\ z = 0 & \text{on } \partial\Omega \times (T_0, T_1) , \\ z(\cdot, T_0) = z_0 & \text{in } \Omega . \end{cases}$$

Therefore,  $z(\cdot, T_1) \in L^2(\Omega)$  and it is well-known that

$$\|z(\cdot, T_1)\|_{L^2(\Omega)} \leq e^{(T_1 - T_0)\|a\|_{L^\infty(\Omega \times (T_0, T_1))}} \|z_0\|_{L^2(\Omega)} .$$

The second step consists on establishing the existence of a function  $v \in L^\infty(\Omega \times (0, +\infty))$  such that the solution  $\tilde{z} = \tilde{z}(x, t)$  of

$$\begin{cases} \partial_t \tilde{z} - \Delta \tilde{z} + a\tilde{z} = 1_{|\omega \times E} v & \text{in } \Omega \times (T_1, T_2) , \\ \tilde{z} = 0 & \text{on } \partial\Omega \times (T_1, T_2) , \\ \tilde{z}(\cdot, T_1) = z(\cdot, T_1) & \text{in } \Omega , \end{cases}$$

satisfies  $\tilde{z}(\cdot, T_2) = 0$  in  $L^2(\Omega)$ . Further,

$$\|v\|_{L^\infty(\Omega \times (T_1, T_2))} \leq e^{\tilde{K}} e^{K(1+(T_2-T_1)\|a\|_{L^\infty(\Omega \times (T_1, T_2))} + \|a\|_{L^\infty(\Omega \times (T_1, T_2))}^{2/3})} \|z(\cdot, T_1)\|_{L^2(\Omega)} .$$

Here,  $K = K(\Omega, \omega) > 1$  and  $\tilde{K} = \tilde{K}(\Omega, \omega, E)$  are positive constants which do not depend on  $T_0$ . Finally, in the last step, we choose

$$v_0(\cdot, t) = \begin{cases} 0 & \text{if } t \in (0, T_1) \cup [T_2, +\infty) , \\ v(\cdot, t) & \text{if } t \in [T_1, T_2) . \end{cases}$$

Since

$$\|v_0\|_{L^\infty(\Omega \times (0, +\infty))} = \|v\|_{L^\infty(\Omega \times (T_1, T_2))} ,$$

the desired result holds. It is standard to get the existence of the above function  $v$  from an observability estimate. More precisely, we apply the following result. Its proof is provided in Section 5.

*Theorem 4 .- Observability estimate .-* Let  $\omega$  be an open and non-empty subset of  $\Omega$ . Let  $T > 0$  and  $E$  be a subset of positive measure in  $(0, T)$ . Then there are two constants  $K = K(\Omega, \omega)$  and  $\tilde{K} = \tilde{K}(\Omega, \omega, E) > 0$  such that for any  $a = a(x, t) \in L^\infty(\Omega \times (0, T))$  and any  $\varphi_0 \in L^2(\Omega)$ , the solution  $\varphi = \varphi(x, t)$  of

$$\begin{cases} \partial_t \varphi - \Delta \varphi + a\varphi = 0 & \text{in } \Omega \times (0, T) , \\ \varphi = 0 & \text{on } \partial\Omega \times (0, T) , \\ \varphi(\cdot, 0) = \varphi_0 & \text{in } \Omega , \end{cases}$$

satisfies

$$\|\varphi(\cdot, T)\|_{L^2(\Omega)} \leq e^{\tilde{K}} e^{K(1+T\|a\|_{L^\infty(\Omega \times (0, T))} + \|a\|_{L^\infty(\Omega \times (0, T))}^{2/3})} \int_{\omega \times E} |\varphi(x, t)| dx dt .$$

*Remark 3 .-* This is a refined observability estimate. When  $E = (0, T)$ , then the observability constant becomes

$$e^{K(1+\frac{1}{T}+T\|a\|_{L^\infty(\Omega \times (0, T))} + \|a\|_{L^\infty(\Omega \times (0, T))}^{2/3})} .$$

This is in accordance with the work of [4]. When  $E$  is a positive measurable set with 0 its Lebesgue point, then the observability constant becomes, for some  $\ell_1 \in E \cap (0, T)$ ,

$$e^{K(1+\frac{1}{\ell_1}+\ell_1\|a\|_{L^\infty(\Omega \times (0, T))} + \|a\|_{L^\infty(\Omega \times (0, T))}^{2/3})} .$$

## 2.2 Nonlinear case with Kakutani's fixed point

In this section, we prove Theorem 2. Let  $0 \leq T_0 < T_1 < T_2$  and  $E \subset (T_1, T_2)$  with  $|E| > 0$ . Let  $\phi \in C([T_0, T_2], L^2(\Omega))$  and  $w_0 \in L^2(\Omega)$ .

By a classical density argument, we may assume that  $f \in C^1$ . We shall use the Kakutani's fixed point theorem to prove the result. First, define for any  $(x, t) \in \Omega \times (T_0, T_2)$ ,

$$a(x, t, r) = \begin{cases} \frac{f(\phi(x, t) + r) - f(\phi(x, t))}{r} & \text{if } r \neq 0, \\ f'(\phi(x, t)) & \text{if } r = 0. \end{cases}$$

And consider

$$\mathcal{K} = \{\xi \in L^2(\Omega \times (T_0, T_2)); \|\xi\|_{L^2(T_0, T_2; H_0^1(\Omega)) \cap H^1(T_0, T_2; H^{-1}(\Omega))} \leq \widehat{\kappa}\}$$

where  $\widehat{\kappa} > 0$  will be determined later. Since  $f : \mathbb{R} \rightarrow \mathbb{R}$  is a globally Lipschitz function, we have that for a.e.  $(x, t) \in \Omega \times (T_0, T_2)$  and any  $r \in \mathbb{R}$

$$|a(x, t, r)| \leq L(f)$$

where  $L(f) > 0$  is the Lipschitz constant of the function  $f$ .

Next, using the fact  $L^\infty(\Omega \times (0, +\infty)) \subset L^\infty(0, +\infty; L^2(\Omega))$ , we know by Theorem 3 that for any  $\xi \in L^2(\Omega \times (T_0, T_2))$ , there are a function  $v_0 \in L^\infty(0, +\infty; L^2(\Omega))$  and a corresponding solution  $z = z(x, t)$  of

$$\begin{cases} \partial_t z - \Delta z + a(\cdot, \cdot, \xi(\cdot, \cdot)) z = 1_{|\omega \times E} v_0 & \text{in } \Omega \times (T_0, T_2), \\ z = 0 & \text{on } \partial\Omega \times (T_0, T_2), \\ z(\cdot, T_0) = w_0 & \text{in } \Omega, \end{cases} \quad (2.2.1)$$

such that

$$z(\cdot, T_2) = 0 \text{ in } L^2(\Omega) \quad (2.2.2)$$

and

$$\|v_0\|_{L^\infty(0, +\infty; L^2(\Omega))} \leq \widehat{K} \|w_0\|_{L^2(\Omega)}. \quad (2.2.3)$$

Here and throughout the proof of Theorem 2,

$$\widehat{K} = e^{\widetilde{K} e^{(T_1 - T_0)L(f)}} e^{K(1 + (T_2 - T_1)L(f) + L(f)^{2/3})}$$

where  $K = K(\Omega, \omega) > 1$  and  $\widetilde{K} = \widetilde{K}(\Omega, \omega, E)$  are positive constants which do not depend on  $T_0$ . Therefore, we can define the map

$$\begin{aligned} \Lambda : \mathcal{K} &\rightarrow L^2(\Omega \times (T_0, T_2)) \\ \xi &\mapsto z \end{aligned}$$

where (2.2.1)-(2.2.2)-(2.2.3) holds.

Now, we check that Kakutani's fixed point theorem is applicable. For convenience, let us state this result (see e.g. [3]).

Theorem (Kakutani's fixed point) .- Let  $\mathcal{Z}$  be a Banach space and  $\Pi$  be a nonempty convex compact subset of  $\mathcal{Z}$ . Let  $\Lambda : \Pi \rightarrow \mathcal{Z}$  be a set-valued mapping satisfying the following assumptions:

- i)  $\Lambda(\xi)$  is a nonempty convex set of  $\mathcal{Z}$  for every  $\xi \in \Pi$ .
- ii)  $\Lambda(\Pi) \subset \Pi$ .
- iii)  $\Lambda : \Pi \rightarrow \mathcal{Z}$  is upper semicontinuous in  $\mathcal{Z}$ .

Then  $\Lambda$  possesses a fixed point in the set  $\Pi$ .

Here  $\mathcal{Z} = L^2(\Omega \times (T_0, T_2))$  and  $\Pi = \mathcal{K}$  with an adequate choice of  $\widehat{\kappa}$  given below. Clearly,  $\mathcal{K}$  is a nonempty convex compact set in  $L^2(\Omega \times (T_0, T_2))$ . Further, from the above arguments,  $\Lambda(\xi)$  is a nonempty convex set in  $L^2(\Omega \times (T_0, T_2))$ . Thus *i*) holds.

Let us prove that *ii*) holds with an adequate choice of  $\widehat{\kappa}$ . By a standard energy method, using the fact that  $|a| \leq L(f)$  and (2.2.1)-(2.2.2)-(2.2.3), there exists  $C > 0$  such that

$$\|z\|_{C([T_0, T_2]; L^2(\Omega))}^2 + \int_{T_0}^{T_2} \|z(\cdot, t)\|_{H_0^1(\Omega)}^2 dt \leq C \|w_0\|_{L^2(\Omega)}^2 .$$

Combining the latter with the fact that  $|a| \leq L(f)$ , we deduce that the solution  $z$  satisfies

$$\|z\|_{L^2(T_0, T_2; H_0^1(\Omega)) \cap H^1(T_0, T_2; H^{-1}(\Omega))} \leq C \|w_0\|_{L^2(\Omega)} ,$$

for some  $C = C(\Omega, \omega, E, T_2, L(f))$  which is a positive constant which does not depend on  $T_0$ . Hence, if we take  $\widehat{\kappa}$  as follows

$$\widehat{\kappa} = C \|w_0\|_{L^2(\Omega)}$$

then  $\Lambda(\mathcal{K}) \subset \mathcal{K}$ .

Let us finally prove the upper semicontinuity of  $\Lambda : \mathcal{K} \rightarrow L^2(\Omega \times (T_0, T_2))$ . We need to prove that if  $\xi_m \in \mathcal{K} \rightarrow \xi$  strongly in  $L^2(\Omega \times (T_0, T_2))$  and if  $p_m \in \Lambda(\xi_m) \rightarrow p$  strongly in  $L^2(\Omega \times (T_0, T_2))$ , then  $p \in \Lambda(\xi)$ . To this end, firstly, we claim that there exists a subsequence of  $(m)_{m \geq 1}$ , still denoted in the same manner, such that

$$a(\cdot, \cdot, \xi_m(\cdot, \cdot))p_m \rightarrow a(\cdot, \cdot, \xi(\cdot, \cdot))p \text{ strongly in } L^2(\Omega \times (T_0, T_2)) . \quad (2.2.4)$$

Indeed, since  $\xi_m \rightarrow \xi$  strongly in  $L^2(\Omega \times (T_0, T_2))$ , we have that there exists a subsequence of  $(m)_{m \geq 1}$ , still denoted by itself, such that

$$\xi_m(x, t) \rightarrow \xi(x, t) \text{ for a.e. } (x, t) \in \Omega \times (T_0, T_2) .$$

On one hand, for  $(x, t)$  with  $\xi(x, t) \neq 0$ , by the above, there exists a positive integer  $m_0$  depending on  $(x, t)$  such that

$$\xi_m(x, t) \neq 0 \quad \forall m \geq m_0 ,$$

which implies by the definition of  $a$ ,

$$a(x, t, \xi_m(x, t)) \rightarrow a(x, t, \xi(x, t)) \quad \text{as } m \rightarrow +\infty . \quad (2.2.5)$$



On the other hand, for any  $(x, t)$  such that  $\xi(x, t) = 0$ , by the definition of  $a$ , we have that  $a(x, t, \xi(x, t)) = f'(\phi(x, t))$ . Since

$$a(x, t, \xi_m(x, t)) = \begin{cases} \frac{f(\phi(x, t) + \xi_m(x, t)) - f(\phi(x, t))}{\xi_m(x, t)} & \text{if } \xi_m(x, t) \neq 0, \\ f'(\phi(x, t)) & \text{if } \xi_m(x, t) = 0, \end{cases}$$

it gives

$$a(x, t, \xi_m(x, t)) \rightarrow a(x, t, \xi(x, t)) \quad \text{as } m \rightarrow +\infty.$$

This, combined with (2.2.5), implies

$$a(x, t, \xi_m(x, t)) \rightarrow a(x, t, \xi(x, t)) \quad \text{for a.e. } (x, t) \in \Omega \times (T_0, T_2).$$

From the latter, the fact that  $|a| \leq L(f)$  and the Lebesgue's dominated convergence theorem it follows that

$$\begin{aligned} & \|a(\cdot, \cdot, \xi_m(\cdot, \cdot))p_m - a(\cdot, \cdot, \xi(\cdot, \cdot))p\|_{L^2(\Omega \times (T_0, T_2))}^2 \\ & \leq 2\|a(\cdot, \cdot, \xi_m(\cdot, \cdot))(p_m - p)\|_{L^2(\Omega \times (T_0, T_2))}^2 + 2\|(a(\cdot, \cdot, \xi_m(\cdot, \cdot)) - a(\cdot, \cdot, \xi(\cdot, \cdot)))p\|_{L^2(\Omega \times (T_0, T_2))}^2 \\ & \leq 2L(f)^2\|p_m - p\|_{L^2(\Omega \times (T_0, T_2))}^2 + 2\|(a(\cdot, \cdot, \xi_m(\cdot, \cdot)) - a(\cdot, \cdot, \xi(\cdot, \cdot)))p\|_{L^2(\Omega \times (T_0, T_2))}^2 \\ & \rightarrow 0. \end{aligned}$$

This completes the proof of (2.2.4). Secondly, since  $p_m \in \Lambda(\xi_m)$ , there exists  $(v_m)_{m \geq 1}$  satisfying

$$\begin{cases} (p_m)_t - \Delta p_m + a(\cdot, \cdot, \xi_m(\cdot, \cdot))p_m = 1_{|\omega \times E} v_m & \text{in } \Omega \times (T_0, T_2), \\ p_m = 0 & \text{on } \partial\Omega \times (T_0, T_2), \\ p_m(\cdot, T_0) = w_0 & \text{in } \Omega, \\ p_m(\cdot, T_2) = 0 & \text{in } \Omega, \end{cases} \quad (2.2.6)$$

$$\begin{aligned} & \|p_m\|_{L^2(T_0, T_2; H_0^1(\Omega)) \cap H^1(T_0, T_2; H^{-1}(\Omega))} \\ & + \|p_m\|_{L^2(T_1, T_2; H^2(\Omega) \cap H_0^1(\Omega)) \cap H^1(T_1, T_2; L^2(\Omega))} \leq C, \end{aligned}$$

where  $C > 0$  is a constant independent on  $m$ , and

$$\|v_m\|_{L^\infty(0, +\infty; L^2(\Omega))} \leq \widehat{K} \|w_0\|_{L^2(\Omega)}. \quad (2.2.7)$$

Thus, we deduce the existence of  $v$  and subsequences  $(v_{m'})_{m' \geq 1}$  and  $(p_{m'})_{m' \geq 1}$  such that

$$v_{m'} \rightarrow v \text{ weakly star in } L^\infty(0, +\infty; L^2(\Omega)), \quad (2.2.8)$$

$$p_{m'} \rightarrow p \text{ weakly in } L^2(T_0, T_2; H_0^1(\Omega)) \cap H^1(T_0, T_2; H^{-1}(\Omega)), \quad (2.2.9)$$

$$p_{m'}(\cdot, T_2) \rightarrow p(\cdot, T_2) \text{ strongly in } L^2(\Omega). \quad (2.2.10)$$

Finally, passing to the limit for  $m' \rightarrow +\infty$  in (2.2.6) and (2.2.7), by (2.2.4) and (2.2.8)-(2.2.9)-(2.2.10), we obtain that  $p \in \Lambda(\xi)$ .

By the Kakutani's fixed point theorem, we conclude that there exists  $w \in \mathcal{K}$  with an adequate choice of  $\widehat{\kappa}$  such that  $w \in \Lambda(w)$ , i.e., there is a control  $v_1 \in L^\infty(0, +\infty; L^2(\Omega))$  satisfying

$$\|v_1\|_{L^\infty(0, +\infty; L^2(\Omega))} \leq \widehat{K} \|w_0\|_{L^2(\Omega)},$$

with the same  $\widehat{K}$  given at (2.2.3), and the corresponding solution  $w = w(x, t)$  solves

$$\begin{cases} \partial_t w - \Delta w + a(\cdot, \cdot, w(\cdot, \cdot)) w = 1_{|\omega \times E} v_1 & \text{in } \Omega \times (T_0, T_2) , \\ w = 0 & \text{on } \partial\Omega \times (T_0, T_2) , \\ w(\cdot, T_0) = w_0 & \text{in } \Omega , \\ w(\cdot, T_2) = 0 & \text{in } \Omega . \end{cases}$$

Since for any  $(x, t) \in \Omega \times (T_0, T_2)$ ,

$$a(x, t, w(x, t))w(x, t) = f(\phi(x, t) + w(x, t)) - f(\phi(x, t)) ,$$

we finally get

$$\begin{cases} \partial_t w - \Delta w + f(\phi + w) - f(\phi) = 1_{|\omega \times E} v_1 & \text{in } \Omega \times (T_0, T_2) , \\ w = 0 & \text{on } \partial\Omega \times (T_0, T_2) , \\ w(\cdot, T_0) = w_0 & \text{in } \Omega , \\ w(\cdot, T_2) = 0 & \text{in } \Omega . \end{cases}$$

This completes the proof of Theorem 2.

### 3 Existence of time optimal control

In this section, we start to prove the existence of admissible controls (see e.g. [15]). In other words we prove that

*Theorem 5 .-* Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be a globally Lipschitz function satisfying  $f(s) s \geq 0$  for all  $s \in \mathbb{R}$ . Then for any  $y_0 \in L^2(\Omega) \setminus \{0\}$  and any  $M > 0$ , there are a time  $T > 0$  and an admissible control  $v \in L^\infty(0, +\infty; L^2(\Omega))$  such that  $\|v\|_{L^\infty(0, +\infty; L^2(\Omega))} \leq M$  and the solution  $y$  corresponding to  $v$  satisfies  $y(\cdot, T) = 0$  in  $\Omega$ .

Proof .- We divide its proof into many steps.

Step 1 .- We consider the following equation

$$\begin{cases} \partial_t y - \Delta y + f(y) = 0 & \text{in } \Omega \times (0, T_0) , \\ y = 0 & \text{on } \partial\Omega \times (0, T_0) , \\ y(\cdot, 0) = y_0 & \text{in } \Omega , \end{cases}$$

where  $T_0 > 0$  will be determined later. By a standard energy method, using the fact that  $f(s) s \geq 0$ , we have

$$\|y(\cdot, T_0)\|_{L^2(\Omega)} \leq e^{-\lambda_1 T_0} \|y_0\|_{L^2(\Omega)} ,$$

where  $\lambda_1 > 0$  is the first eigenvalue of  $-\Delta$  with Dirichlet boundary condition.

Step 2 .- We apply Theorem 2 with  $T_1 = T_0 + 1$ ,  $T_2 = T_0 + 2$ ,  $E = (T_1, T_2)$ ,  $\phi = 0$  and  $w_0 = y(\cdot, T_0)$  in order that there are a constant  $\kappa > 0$  and a function  $\tilde{v} \in L^\infty(0, +\infty; L^2(\Omega))$  such that the solution  $w = w(x, t)$  of

$$\begin{cases} \partial_t w - \Delta w + f(w) = 1_{|\omega \times E} \tilde{v} & \text{in } \Omega \times (T_0, T_0 + 2) , \\ w = 0 & \text{on } \partial\Omega \times (T_0, T_0 + 2) , \\ w(\cdot, T_0) = y(\cdot, T_0) & \text{in } \Omega , \end{cases}$$

satisfies  $w(\cdot, T_0 + 2) = 0$  in  $L^2(\Omega)$ . Further,

$$\|\tilde{v}\|_{L^\infty(0, +\infty; L^2(\Omega))} \leq \kappa \|y(\cdot, T_0)\|_{L^2(\Omega)} ,$$

and  $\kappa$  does not depend on  $T_0$ .

Step 3 .- We can easily check that the function

$$v(\cdot, t) = \begin{cases} 0 & \text{if } t \in (0, T_0 + 1] \cup [T_0 + 2, +\infty) , \\ \tilde{v}(\cdot, t) & \text{if } t \in (T_0 + 1, T_0 + 2) , \end{cases}$$

is an admissible control with  $T = T_0 + 2$  when  $T_0 > 0$  is taken such that

$$T_0 = \frac{1}{\lambda_1} \ln \left( 1 + \frac{\kappa \|y_0\|_{L^2(\Omega)}}{M} \right)$$

in order that

$$\|v\|_{L^\infty(0, +\infty; L^2(\Omega))} = \|\tilde{v}\|_{L^\infty(T_0+1, T_0+2; L^2(\Omega))} \leq \kappa e^{-\lambda_1 T_0} \|y_0\|_{L^2(\Omega)} \leq M .$$

This completes the proof of Theorem 5.

Now, we establish the existence of time optimal controls (see e.g. [15]). In other words, we shall prove that

*Theorem 6 .- Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be a globally Lipschitz function satisfying  $f(s) s \geq 0$  for all  $s \in \mathbb{R}$ . Then for any  $y_0 \in L^2(\Omega) \setminus \{0\}$  and any  $M > 0$ , there is a time optimal control  $v^* \in L^\infty(0, +\infty; L^2(\Omega))$  such that  $\|v^*\|_{L^\infty(0, +\infty; L^2(\Omega))} \leq M$  and the solution  $y$  corresponding to  $v^*$  satisfies  $y(\cdot, T^*) = 0$  in  $\Omega$  where  $T^* = \inf\{T; v \in \mathcal{V}_M\}$ .*

Proof .- By Theorem 5 and the definition of  $T^*$ ,  $0 \leq T^* < \bar{T}$  for some  $\bar{T} > 0$ . Therefore, there exist sequences  $(T_m)_{m \geq 1}$  of positive real number and  $(v_m)_{m \geq 1}$  of function in  $L^\infty(0, +\infty; L^2(\Omega))$  such that  $T^* = \lim_{m \rightarrow \infty} T_m$ ,  $\|v_m\|_{L^\infty(0, +\infty; L^2(\Omega))} \leq M$  and the solution  $y_m = y_m(x, t)$  corresponding to  $v_m$  satisfies

$$\begin{cases} \partial_t y_m - \Delta y_m + f(y_m) = 1_{|\omega} v_m & \text{in } \Omega \times (0, \bar{T}) , \\ y_m = 0 & \text{on } \partial\Omega \times (0, \bar{T}) , \\ y_m(\cdot, 0) = y_0 & \text{in } \Omega , \\ y_m(\cdot, T_m) = 0 & \text{in } \Omega . \end{cases}$$

We have by a standard energy method, using the bound  $M$  on  $v_m$  and the “good-sign” condition on  $f$ ,

$$\|y_m\|_{C([0,\bar{T}];L^2(\Omega))}^2 + \int_0^{\bar{T}} \|y_m(\cdot, t)\|_{H_0^1(\Omega)}^2 dt \leq C .$$

Here and throughout the proof,  $C$  denotes a generic constant independent of  $m$ . Since  $f$  is globally Lipschitz and  $f(0) = 0$ , the above inequality implies

$$\|f(y_m)\|_{L^2(\Omega \times (0,\bar{T}))} = \|f(y_m) - f(0)\|_{L^2(\Omega \times (0,\bar{T}))} \leq C .$$

Therefore, from the boundeness of  $-f(y_m) + 1_{|\omega} v_m$ , the sequence  $(y_m)_{m \geq 1}$  is bounded in  $H^1(0, \bar{T}; H^{-1}(\Omega))$ .

Now, we deduce the existence of  $v^* \in L^\infty(0, +\infty; L^2(\Omega))$  and subsequences  $(v_{m'})_{m' \geq 1}$  and  $(y_{m'})_{m' \geq 1}$  such that

$$v_{m'} \rightarrow v^* \text{ weakly star in } L^\infty(0, +\infty; L^2(\Omega)) \text{ with } \|v^*\|_{L^\infty(0, +\infty; L^2(\Omega))} \leq M ,$$

$$\begin{aligned} y_{m'} &\rightarrow y^* \text{ weakly in } L^2(0, \bar{T}; H_0^1(\Omega)) \cap H^1(0, \bar{T}; H^{-1}(\Omega)) , \\ &\text{strongly in } C([0, \bar{T}]; L^2(\Omega)) . \end{aligned}$$

Further,

$$\begin{cases} \partial_t y^* - \Delta y^* + f(y^*) = 1_{|\omega} v^* & \text{in } \Omega \times (0, \bar{T}) , \\ y^* = 0 & \text{on } \partial\Omega \times (0, \bar{T}) , \\ y^*(\cdot, 0) = y_0 & \text{in } \Omega , \end{cases}$$

and

$$\begin{aligned} \|y^*(\cdot, T^*)\|_{L^2(\Omega)} &\leq \|y^*(\cdot, T^*) - y^*(\cdot, T_{m'})\|_{L^2(\Omega)} + \|y^*(\cdot, T_{m'}) - y_{m'}(\cdot, T_{m'})\|_{L^2(\Omega)} \\ &\rightarrow 0 \text{ when } m' \rightarrow \infty . \end{aligned}$$

This gives  $y^*(\cdot, T^*) = 0$  in  $\Omega$  and consequently,  $v^*$  is a time optimal control. This completes the proof.

## 4 Bang-bang property for time optimal control (proof of Theorem 1)

We want to prove that if  $v^*$  is a time optimal control corresponding to the optimal time  $T^* = \inf\{T; v \in \mathcal{V}_M\}$ , then  $\|v^*(\cdot, t)\|_{L^2(\Omega)} = M$  for a.e.  $t \in (0, T^*)$ . To prove this, we work by contradiction. Suppose that there are  $\varepsilon \in (0, M)$  and a positive measurable subset  $E^* \subset (0, T^*)$  such that

$$\|v^*(\cdot, t)\|_{L^2(\Omega)} \leq M - \varepsilon \quad \forall t \in E^*$$

and the solution  $y^* = y^*(x, t)$  corresponding to  $v^*$  satisfies

$$\begin{cases} \partial_t y^* - \Delta y^* + f(y^*) = 1_{|\omega} v^* & \text{in } \Omega \times (0, T^*) , \\ y^* = 0 & \text{on } \partial\Omega \times (0, T^*) , \\ y^*(\cdot, 0) = y_0 & \text{in } \Omega , \\ y^*(\cdot, T^*) = 0 & \text{in } \Omega . \end{cases}$$

We claim that there exist a real number  $\delta \in (0, T^*)$  and a couple  $(y, v)$  such that

$$\begin{cases} \partial_t y - \Delta y + f(y) = 1_{|\omega} v & \text{in } \Omega \times (0, T^* - \delta) , \\ y = 0 & \text{on } \partial\Omega \times (0, T^* - \delta) , \\ y(\cdot, 0) = y_0 & \text{in } \Omega , \\ y(\cdot, T^* - \delta) = 0 & \text{in } \Omega , \end{cases}$$

and  $v \in L^\infty(0, +\infty; L^2(\Omega))$  with  $\|v\|_{L^\infty(0, +\infty; L^2(\Omega))} \leq M$ . This is clearly a contradiction with the time optimal assumption  $T^* = \inf\{T; v \in \mathcal{V}_M\}$ .

Now, we prove our claim. We divide its proof into many steps.

Step 1 .-  $T^* > 0$  and  $0 < |E^*| \leq T^*$  being given, let  $\delta_0 = |E^*|/2$  and denote

$$E = E^* \cap (\delta_0, T^*).$$

Then

$$|E| > 0 .$$

Indeed,  $|E^* \cap (\delta_0, T^*)| \geq |E^*| - \delta_0 \geq |E^*|/2$ .

Step 2 .- We apply Theorem 2 with  $0 < T_0 < T_1 < T_2$ ,  $E \subset (T_1, T_2)$  with  $|E| > 0$  and  $\phi = y^*$ , in order that there are a constant  $\kappa > 0$  and a function  $v_1 \in L^\infty(0, +\infty; L^2(\Omega))$  such that

$$\begin{cases} \partial_t w - \Delta w + f(y^* + w) - f(y^*) = 1_{|\omega \times E} v_1 & \text{in } \Omega \times (T_0, T_2) , \\ w = 0 & \text{on } \partial\Omega \times (T_0, T_2) , \\ w(\cdot, T_0) = w_0 & \text{in } \Omega , \\ w(\cdot, T_2) = 0 & \text{in } \Omega , \\ \|v_1\|_{L^\infty(0, +\infty; L^2(\Omega))} \leq \kappa \|w_0\|_{L^2(\Omega)} & , \end{cases}$$

and further  $\kappa$  does not depend on  $T_0$ .

Step 3 . We apply step 2 with  $T_0 = \delta$ ,  $T_1 = \delta_0$ ,  $T_2 = T^*$ ,  $w_0 = y_0 - y^*(\cdot, \delta)$ , in order that  $z = y^* + w$  solves

$$\begin{cases} \partial_t z - \Delta z + f(z) = 1_{|\omega} (v^* + 1_{|E} v_1) & \text{in } \Omega \times (\delta, T^*) , \\ z = 0 & \text{on } \partial\Omega \times (\delta, T^*) , \\ z(\cdot, \delta) = y_0 & \text{in } \Omega , \\ z(\cdot, T^*) = 0 & \text{in } \Omega . \end{cases}$$

Denote  $v_2 = v^* + 1_{|E} v_1$ . On one hand, if  $t \in (0, +\infty) \setminus E$ , then  $\|v_2(\cdot, t)\|_{L^2(\Omega)} = \|v^*(\cdot, t)\|_{L^2(\Omega)} \leq M$ . On the other hand, if  $t \in E$ , then

$$\begin{aligned} \|v_2(\cdot, t)\|_{L^2(\Omega)} &\leq \|v^*(\cdot, t)\|_{L^2(\Omega)} + \|v_1(\cdot, t)\|_{L^2(\Omega)} \\ &\leq M - \varepsilon + \kappa \|y^*(\cdot, 0) - y^*(\cdot, \delta)\|_{L^2(\Omega)} . \end{aligned}$$

Now, we choose  $\delta$  sufficiently closed to 0 in order that

$$\|y^*(\cdot, 0) - y^*(\cdot, \delta)\|_{L^2(\Omega)} \leq \varepsilon/\kappa .$$

This is possible because  $y^* \in C([0, T^*], L^2(\Omega))$ . Consequently,  $\|v_2(\cdot, t)\|_{L^2(\Omega)} \leq M$  for a.e.  $t \in (0, +\infty)$ .

Step 4 .- Let  $v(\cdot, t) = v_2(\cdot, t + \delta)$ . Then  $v \in L^\infty(0, +\infty; L^2(\Omega))$  and further we can check that  $\|v(\cdot, t)\|_{L^2(\Omega)} \leq M$  for a.e.  $t \in (0, +\infty)$ . Let  $y(x, t) = z(x, t + \delta)$ . Then it solves

$$\begin{cases} \partial_t y - \Delta y + f(y) = 1_{|\omega} v & \text{in } \Omega \times (0, T^* - \delta), \\ y = 0 & \text{on } \partial\Omega \times (0, T^* - \delta), \\ y(\cdot, 0) = y_0 & \text{in } \Omega, \\ y(\cdot, T^* - \delta) = 0 & \text{in } \Omega. \end{cases}$$

This is the desired claim.

## 5 The heat equation with potential (proof of Theorem 4)

The proof of Theorem 4 is based on many lemmas. From now,  $\varphi$  denotes the solution of

$$\begin{cases} \partial_t \varphi - \Delta \varphi + a\varphi = 0 & \text{in } \Omega \times (0, T), \\ \varphi = 0 & \text{on } \partial\Omega \times (0, T), \\ \varphi(\cdot, 0) = \varphi_0 & \text{in } \Omega, \end{cases}$$

where  $a = a(x, t) \in L^\infty(\Omega \times (0, T))$ . We also denote  $\|a\|_\infty = \|a\|_{L^\infty(\Omega \times (0, T))}$ .

*Lemma 1 .- For any  $\varphi_0 \in L^2(\Omega)$ , the solution  $\varphi$  satisfies the two following estimates for any  $t \in (0, T]$ ,*

$$\int_{\Omega} |\varphi(x, t)|^2 dx \leq e^{2t\|a\|_\infty} \int_{\Omega} |\varphi_0(x)|^2 dx \text{ and } \int_{\Omega} |\nabla \varphi(x, t)|^2 dx \leq \frac{e^{3t\|a\|_\infty}}{t} \int_{\Omega} |\varphi_0(x)|^2 dx .$$

This result is deduced by energy estimate and is standard. Its proof is omitted here.

Let  $x_0 \in \Omega$ . Denote  $B_R = B(x_0, R)$  the ball of center  $x_0$  and radius  $R$ .

*Lemma 2 .- Let  $R_0 > 0$  and  $\lambda > 0$ . Introduce for  $t \in [0, T]$  and  $x_0 \in \Omega$ ,*

$$G_\lambda(x, t) = \frac{1}{(T - t + \lambda)^{n/2}} e^{-\frac{|x-x_0|^2}{4(T-t+\lambda)}} .$$

*Define for  $u \in H^1(0, T; L^2(\Omega \cap B_{R_0})) \cap L^2(0, T; H^2 \cap H_0^1(\Omega \cap B_{R_0}))$  and  $t \in (0, T]$ ,*

$$N_\lambda(t) = \frac{\int_{\Omega \cap B_{R_0}} |\nabla u(x, t)|^2 G_\lambda(x, t) dx}{\int_{\Omega \cap B_{R_0}} |u(x, t)|^2 G_\lambda(x, t) dx}, \text{ whenever } \int_{\Omega \cap B_{R_0}} |u(x, t)|^2 dx \neq 0 .$$

*The following two properties hold.*

i)

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \int_{\Omega \cap B_{R_0}} |u(x, t)|^2 G_\lambda(x, t) dx + \int_{\Omega \cap B_{R_0}} |\nabla u(x, t)|^2 G_\lambda(x, t) dx \\ &= \int_{\Omega \cap B_{R_0}} u(x, t) (\partial_t - \Delta) u(x, t) G_\lambda(x, t) dx . \end{aligned} \quad (5.1)$$

ii) When  $\Omega \cap B_{R_0}$  is star-shaped with respect to  $x_0$ ,

$$\frac{d}{dt} N_\lambda(t) \leq \frac{1}{T-t+\lambda} N_\lambda(t) + \frac{\int_{\Omega \cap B_{R_0}} |(\partial_t - \Delta) u(x, t)|^2 G_\lambda(x, t) dx}{\int_{\Omega \cap B_{R_0}} |u(x, t)|^2 G_\lambda(x, t) dx} . \quad (5.2)$$

Proof. The identity follows from some direct computations. The proof of the second one is the same as that in [13, pp. 1240-1245] or [5, Lemma 2].

*Lemma 3 .-* Let  $R > 0$  and  $\delta \in (0, 1]$ . Then there are two constants  $C_1, C_2 > 0$ , only dependent on  $(R, \delta)$  such that for any  $\varphi_0 \in L^2(\Omega)$  with  $\varphi_0 \neq 0$ , the quantity

$$h_0 = \frac{C_1}{\ln \left( (1 + C_2) \left( e^{1 + \frac{2C_1}{T} + 3T\|a\|_\infty + \|a\|_\infty^{2/3}} \right) \frac{\int_{\Omega} |\varphi_0(x)|^2 dx}{\int_{\Omega \cap B_R} |\varphi(x, T)|^2 dx} \right)} \quad (5.3)$$

has the following two properties.

i)

$$0 < \left( 1 + \frac{2C_1}{T} + T\|a\|_\infty + \|a\|_\infty^{2/3} \right) h_0 < C_1 . \quad (5.4)$$

ii) For any  $t \in [T - h_0, T]$ , it holds

$$e^{3T\|a\|_\infty} \int_{\Omega} |\varphi_0(x)|^2 dx \leq e^{1 + C_3 \frac{1}{h_0}} \int_{\Omega \cap B_{(1+\delta)R}} |\varphi(x, t)|^2 dx \quad (5.5)$$

for some  $C_3 > C_1$  only dependent on  $(R, \delta)$ .

*Remark 4 .-* By the strong unique continuation property for parabolic equations with zero Dirichlet boundary condition, it is impossible to have  $\int_{\Omega \cap B_R} |\varphi(x, T)|^2 dx = 0$  if  $\varphi_0 \in L^2(\Omega)$  with  $\varphi_0 \neq 0$ .

*Remark 5 .-* From (5.4), we have  $h_0 < T/2$  and therefore  $T/2 < T - h_0 < T$ . Here, (5.5) says that for any  $t$  sufficiently closed to  $T$ , the following Hölder interpolation estimate holds.

$$\begin{aligned} \int_{\Omega \cap B_R} |\varphi(x, T)|^2 dx &\leq \left( (1 + C_2) e^{1 + \frac{2C_1}{T} + \|a\|_\infty^{2/3}} \right) \left( e^{3T\|a\|_\infty} \int_{\Omega} |\varphi_0(x)|^2 dx \right)^{\frac{C_3 - C_1}{C_3}} \\ &\quad \times \left( e \int_{\Omega \cap B_{(1+\delta)R}} |\varphi(x, t)|^2 dx \right)^{\frac{C_1}{C_3}} . \end{aligned}$$

This can be compared with [5, Lemma 1].

Proof .- The property (5.4) is clearly true because the following inequality

$$\int_{\Omega \cap B_R} |\varphi(\cdot, T)|^2 dx \leq e^{2T\|a\|_\infty} \int_{\Omega} |\varphi_0|^2 dx$$

holds by Lemma 1. We prove (5.5) as follows. Let  $h > 0$ ,  $\rho(x) = |x - x_0|^2$ ,  $\chi \in C_0^\infty(B(x_0, (1 + \delta)R))$  be such that  $0 \leq \chi \leq 1$ ,  $\chi = 1$  on  $\{x; |x - x_0| \leq (1 + 3\delta/4)R\}$ . We multiply the equation  $\partial_t \varphi - \Delta \varphi + a\varphi = 0$  by  $e^{-\rho/h} \chi^2 \varphi$  and integrate over  $\Omega \cap B_{(1+\delta)R}$ . We get

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \int_{\Omega \cap B_{(1+\delta)R}} e^{-\rho/h} |\chi \varphi|^2 dx + \int_{\Omega \cap B_{(1+\delta)R}} \nabla \varphi \nabla (e^{-\rho/h} \chi^2 \varphi) dx \\ &= - \int_{\Omega \cap B_{(1+\delta)R}} a e^{-\rho/h} |\chi \varphi|^2 dx . \end{aligned}$$

But,  $\nabla (e^{-\rho/h} \chi^2 \varphi) = \frac{-1}{h} \nabla \rho e^{-\rho/h} \chi^2 \varphi + 2e^{-\rho/h} \chi \nabla \chi \varphi + e^{-\rho/h} \chi^2 \nabla \varphi$ . Therefore,

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \int_{\Omega \cap B_{(1+\delta)R}} e^{-\rho/h} |\chi \varphi|^2 dx + \int_{\Omega \cap B_{(1+\delta)R}} e^{-\rho/h} |\chi \nabla \varphi|^2 dx \\ & \leq \int_{\Omega \cap B_{(1+\delta)R}} (e^{-\rho/(2h)} |\chi \nabla \varphi|) \left( \frac{2}{h} |x - x_0| e^{-\rho/(2h)} \chi |\varphi| + 2 |\nabla \chi| e^{-\rho/(2h)} |\varphi| \right) dx \\ & \quad + \|a\|_\infty \int_{\Omega \cap B_{(1+\delta)R}} e^{-\rho/h} |\chi \varphi|^2 dx \end{aligned}$$

which gives by Cauchy-Schwarz inequality

$$\begin{aligned} & \frac{d}{dt} \int_{\Omega \cap B_{(1+\delta)R}} e^{-\rho/h} |\chi \varphi|^2 dx \\ & \leq \left( \frac{4((1+\delta)R)^2}{h^2} + 2\|a\|_\infty \right) \int_{\Omega \cap B_{(1+\delta)R}} e^{-\rho/h} |\chi \varphi|^2 dx \\ & \quad + 4 \int_{\Omega \cap \{x; (1+3\delta/4)R \leq \sqrt{\rho(x)} \leq (1+\delta)R\}} |\nabla \chi|^2 e^{-\rho/h} |\varphi|^2 dx . \end{aligned}$$

Thus,

$$\begin{aligned} & \frac{d}{dt} \left( e^{-\left(\frac{4((1+\delta)R)^2}{h^2} + 2\|a\|_\infty\right)t} \int_{\Omega \cap B_{(1+\delta)R}} e^{-\rho/h} |\chi \varphi|^2 dx \right) \\ & \leq 4 \|\nabla \chi\|_{L^\infty}^2 e^{-\left(\frac{4((1+\delta)R)^2}{h^2} + 2\|a\|_\infty\right)t} e^{-\frac{((1+3\delta/4)R)^2}{h}} e^{2t\|a\|_\infty} \int_{\Omega} |\varphi_0|^2 dx \end{aligned}$$

which gives by integration between  $t$  and  $T$ ,

$$\begin{aligned} & \int_{\Omega \cap B_{(1+\delta)R}} e^{-\rho/h} |\chi \varphi(\cdot, T)|^2 dx \\ & \leq e^{\left(\frac{4((1+\delta)R)^2}{h^2} + 2\|a\|_\infty\right)(T-t)} \int_{\Omega \cap B_{(1+\delta)R}} e^{-\rho/h} |\chi \varphi(\cdot, t)|^2 dx \\ & \quad + e^{\left(\frac{4((1+\delta)R)^2}{h^2} + 2\|a\|_\infty\right)T} \int_t^T e^{-\frac{4((1+\delta)R)^2}{h^2}s} ds 4 \|\nabla \chi\|_{L^\infty}^2 e^{-\frac{((1+3\delta/4)R)^2}{h}} \int_{\Omega} |\varphi_0|^2 dx . \end{aligned}$$



Therefore,

$$\begin{aligned} & \int_{\Omega \cap B_{(1+\delta)R}} e^{-\rho/h} |\chi \varphi(\cdot, T)|^2 dx \\ & \leq e^{\frac{c_1 R^2}{h^2}(T-t)} e^{2T\|a\|_\infty} \int_{\Omega \cap B_{(1+\delta)R}} e^{-\rho/h} |\chi \varphi(\cdot, t)|^2 dx \\ & \quad + e^{\frac{c_1 R^2}{h^2}(T-t)} (T-t) 4 \|\nabla \chi\|_{L^\infty}^2 e^{2T\|a\|_\infty} e^{-\frac{c_2 R^2}{h}} \int_{\Omega} |\varphi_0|^2 dx, \end{aligned}$$

with  $c_1 = 4(1+\delta)^2$  and  $c_2 = (1+3\delta/4)^2$ . Set  $c_3 = (1+\delta/2)^2$ . In particular,  $1 < c_3 < c_2$ . Recall that  $t \leq T$ . Now suppose that the positive real number  $h$  is such that

$$0 < T - \frac{c_2 - c_3}{c_1} h \leq t,$$

then  $\frac{c_1}{h^2}(T-t) \leq \frac{c_2 - c_3}{h}$  and

$$\begin{aligned} \int_{\Omega \cap B_{(1+\delta)R}} e^{-\rho/h} |\chi \varphi(\cdot, T)|^2 dx & \leq e^{\frac{(c_2 - c_3)R^2}{h}} e^{2T\|a\|_\infty} \int_{\Omega \cap B_{(1+\delta)R}} e^{-\rho/h} |\chi \varphi(\cdot, t)|^2 dx \\ & \quad + 4 \|\nabla \chi\|_{L^\infty}^2 e^{2T\|a\|_\infty} \frac{c_2 - c_3}{c_1} h e^{-\frac{c_3 R^2}{h}} \int_{\Omega} |\varphi_0|^2 dx. \end{aligned}$$

Since  $\chi = 1$  on  $\{x; |x - x_0| \leq R\}$ , the above estimate yields

$$\begin{aligned} \int_{\Omega \cap B_R} |\varphi(x, T)|^2 dx & \leq e^{\frac{(c_2 - c_3 + 1)R^2}{h}} e^{2T\|a\|_\infty} \int_{\Omega \cap B_{(1+\delta)R}} |\varphi(x, t)|^2 dx \\ & \quad + 4e^{2T\|a\|_\infty} \|\nabla \chi\|_{L^\infty}^2 \frac{c_2 - c_3}{c_1} h e^{-\frac{(c_3 - 1)R^2}{h}} \int_{\Omega} |\varphi_0(x)|^2 dx, \end{aligned} \quad (5.6)$$

whenever  $0 < T - \frac{c_2 - c_3}{c_1} h \leq t$  and  $t \leq T$ . Recall that  $h_0 < T$  from (5.4). Now, choose  $h \in \left(0, \frac{c_1}{c_2 - c_3} T\right)$  as follows.

$$h = \frac{c_1}{c_2 - c_3} h_0 = \frac{c_1}{c_2 - c_3} \frac{C_1}{\ln \left( \left( e^{1 + \frac{2C_1}{T}} \right) \frac{(1+C_2)e^{3T\|a\|_\infty + \|a\|_\infty^{2/3}} \int_{\Omega} |\varphi_0(x)|^2 dx}{\int_{\Omega \cap B_R} |\varphi(x, T)|^2 dx} \right)}$$

with  $C_1 = \frac{(c_2 - c_3)(c_3 - 1)R^2}{c_1}$  and  $C_2 = 4 \|\nabla \chi\|_{L^\infty}^2 C_1$ , in order that for any  $T - \frac{c_2 - c_3}{c_1} h \leq t \leq T$ ,

$$\begin{aligned} & 4e^{2T\|a\|_\infty} \|\nabla \chi\|_{L^\infty}^2 \frac{c_2 - c_3}{c_1} h e^{-\frac{(c_3 - 1)R^2}{h}} \int_{\Omega} |\varphi_0(x)|^2 dx \\ & = e^{2T\|a\|_\infty} C_2 \frac{h_0}{C_1} e^{-\frac{(c_3 - 1)R^2}{h}} \int_{\Omega} |\varphi_0(x)|^2 dx \\ & \leq \frac{h_0}{C_1} e^{-\frac{(c_3 - 1)R^2}{h}} (1 + C_2) e^{3T\|a\|_\infty + \|a\|_\infty^{2/3}} \int_{\Omega} |\varphi_0|^2 dx \\ & \leq e^{-\frac{(c_3 - 1)R^2}{h}} \frac{1}{e^{1 + \frac{2C_1}{T}}} e^{\frac{(c_3 - 1)R^2}{h}} \int_{\Omega \cap B_R} |\varphi(x, T)|^2 dx \\ & \leq \frac{1}{e} \int_{\Omega \cap B_R} |\varphi(x, T)|^2 dx \end{aligned} \quad (5.7)$$

where we have used in the third line the fact that  $h_0 \leq C_1$  from (5.4). The definition of  $h$  along with (5.3) was applied in the fourth line. Since  $t \in [T - h_0, T]$ , it yields that  $0 < T - \frac{c_2 - c_3}{c_1} h \leq t \leq T$  and further, combining (5.6) and (5.7) we have

$$\int_{\Omega \cap B_R} |\varphi(x, T)|^2 dx \leq e^{\frac{(c_2 - c_3 + 1)R^2}{h}} e^{2T\|a\|_\infty} \int_{\Omega \cap B_{(1+\delta)R}} |\varphi(x, t)|^2 dx + \frac{1}{e} \int_{\Omega \cap B_R} |\varphi(x, T)|^2 dx$$

which gives

$$\left(1 - \frac{1}{e}\right) \int_{\Omega \cap B_R} |\varphi(x, T)|^2 dx \leq e^{\frac{(c_2 - c_3 + 1)(c_2 - c_3)R^2}{c_1} \frac{1}{h_0}} e^{2T\|a\|_\infty} \int_{\Omega \cap B_{(1+\delta)R}} |\varphi(x, t)|^2 dx .$$

On the other hand, by the definition of  $h_0$  and the fact that  $T\|a\|_\infty h_0 \leq C_1$  from (5.4),

$$\begin{aligned} e^{3T\|a\|_\infty} \int_{\Omega} |\varphi_0(x)|^2 dx &\leq e^{3T\|a\|_\infty} e^{C_1 \frac{1}{h_0}} \int_{\Omega \cap B_R} |\varphi(x, T)|^2 dx . \\ &\leq e^{4C_1 \frac{1}{h_0}} \int_{\Omega \cap B_R} |\varphi(x, T)|^2 dx . \end{aligned}$$

We conclude that

$$\left(1 - \frac{1}{e}\right) e^{3T\|a\|_\infty} \int_{\Omega} |\varphi_0(x)|^2 dx \leq e^{C_3 \frac{1}{h_0}} \int_{\Omega \cap B_{(1+\delta)R}} |\varphi(x, t)|^2 dx$$

with  $C_3 = \frac{(c_2 - c_3 + 1)(c_2 - c_3)R^2}{c_1} + 6C_1$ . This completes the proof.

*Lemma 4 .-* Let  $0 < r < R$ . Suppose that  $B_r \subset \Omega$  and  $\Omega \cap B_{(1+2\delta)R}$  is star-shaped with respect to  $x_0$  for some  $\delta \in (0, 1]$ . Then there are  $C_1, C_2 > 0$  and  $\beta \in (0, 1)$  such that for any  $T > 0$  and  $\varphi_0 \in L^2(\Omega)$ ,

$$\begin{aligned} &\int_{\Omega \cap B_R} |\varphi(x, T)|^2 dx \\ &\leq \left( (1 + C_2) e^{1 + \frac{2C_1}{T} + 3T\|a\|_\infty + \|a\|_\infty^2/3} \int_{\Omega} |\varphi_0(x)|^2 dx \right)^\beta \left( 2 \int_{B_r} |\varphi(x, T)|^2 dx \right)^{1-\beta} . \end{aligned}$$

Here  $C_1, C_2 > 0$  are only dependent on  $(R, \delta)$ .  $\beta$  only depends on  $(n, r, R, \delta)$ .

*Proof .-* There is no loss of generality in assuming that  $\varphi_0 \neq 0$ . Let  $0 < r < R$  and  $R_0 = (1 + 2\delta)R$ . Let  $\chi \in C_0^\infty(B_{R_0})$ ,  $0 \leq \chi \leq 1$ ,  $\chi = 1$  on  $\{x; |x - x_0| \leq (1 + 3\delta/2)R\}$ . We will apply Lemma 2 with  $u = \chi\varphi$ . First,  $(\partial_t - \Delta)u = -au - 2\nabla\chi\nabla\varphi - \Delta\chi\varphi$ . Next, define  $g = -2\nabla\chi\nabla\varphi - \Delta\chi\varphi$ .

Step 1.- Notice that  $g$  is supported on  $\{x; (1 + 3\delta/2)R \leq |x - x_0| \leq R_0\}$ . Recall the fact that  $\chi = 1$  on  $\{x; |x - x_0| \leq (1 + \delta)R\}$ . Then there is  $C = C(R, \delta) > 0$  such that we have

$$\frac{\int_{\Omega \cap B_{R_0}} u(x, t) g(x, t) G_\lambda(x, t) dx}{\int_{\Omega \cap B_{R_0}} |u(x, t)|^2 G_\lambda(x, t) dx} \leq \frac{C(1 + t^{-1/2}) e^{3t\|a\|_\infty} \int_{\Omega} |\varphi_0(x)|^2 dx}{\int_{\Omega \cap B_{(1+\delta)R}} |\varphi(x, t)|^2 dx} e^{-\frac{C_4}{T-t+\lambda}}$$

and

$$\int_t^T \frac{\int_{\Omega \cap B_{R_0}} |g(x, s)|^2 G_\lambda(x, s) dx}{\int_{\Omega \cap B_{R_0}} |u(x, s)|^2 G_\lambda(x, s) dx} ds \leq \int_t^T \frac{C(1+s^{-1}) e^{3s\|a\|_\infty} \int_{\Omega} |\varphi_0(x)|^2 dx}{\int_{\Omega \cap B_{(1+\delta)R}} |\varphi(x, s)|^2 dx} e^{-\frac{C_4}{T-s+\lambda}} ds$$

with  $C_4 = -\frac{((1+\delta)R)^2}{4} + \frac{((1+3\delta/2)R)^2}{4} > 0$ . Then we have the existence of  $c = c(R, \delta) > 0$  such that for any  $t \in [T - h_0, T)$ ,

$$\frac{\int_{\Omega \cap B_{R_0}} u(x, t) g(x, t) G_\lambda(x, t) dx}{\int_{\Omega \cap B_{R_0}} |u(x, t)|^2 G_\lambda(x, t) dx} \leq c \left(1 + \frac{1}{\sqrt{T}}\right) e^{C_3 \frac{1}{h_0}} e^{-\frac{C_4}{T-t+\lambda}}$$

and

$$\int_t^T \frac{\int_{\Omega \cap B_{R_0}} |g(x, s)|^2 G_\lambda(x, s) dx}{\int_{\Omega \cap B_{R_0}} |u(x, s)|^2 G_\lambda(x, s) dx} ds \leq c e^{C_3 \frac{1}{h_0}} e^{-\frac{C_4}{T-t+\lambda}}$$

by using (5.4) that gives the two inequalities  $h_0 < C_1$ ,  $T/2 < T - h_0 \leq t < T$  and Lemma 3 saying that

$$\frac{e^{3T\|a\|_\infty} \int_{\Omega} |\varphi_0(x)|^2 dx}{\int_{\Omega \cap B_{(1+\delta)R}} |\varphi(x, t)|^2 dx} \leq e^{1+C_3 \frac{1}{h_0}} \text{ if } T - h_0 \leq t < T.$$

Step 2.- Now, our plan is to bound  $\lambda N_\lambda(T)$ . We apply Lemma 2 as follows. First of all, by (5.2)

$$\frac{d}{dt} N_\lambda(t) \leq \frac{1}{T-t+\lambda} N_\lambda(t) + \frac{\int_{\Omega \cap B_{R_0}} |(-au + g)(x, t)|^2 G_\lambda(x, t) dx}{\int_{\Omega \cap B_{R_0}} |u(x, t)|^2 G_\lambda(x, t) dx}$$

becomes

$$\frac{d}{dt} [(T-t+\lambda) N_\lambda(t)] \leq (T-t+\lambda) \frac{\int_{\Omega \cap B_{R_0}} |(-au + g)(x, t)|^2 G_\lambda(x, t) dx}{\int_{\Omega \cap B_{R_0}} |u(x, t)|^2 G_\lambda(x, t) dx}.$$

Thus, it holds

$$\frac{d}{dt} [(T-t+\lambda) N_\lambda(t)] \leq 2(T-t+\lambda) \left( \|a\|_\infty^2 + \frac{\int_{\Omega \cap B_{R_0}} |g(x, t)|^2 G_\lambda(x, t) dx}{\int_{\Omega \cap B_{R_0}} |u(x, t)|^2 G_\lambda(x, t) dx} \right)$$

which gives

$$\begin{aligned} \lambda N_\lambda(T) &\leq (T-t+\lambda) N_\lambda(t) + 2 \|a\|_\infty^2 \int_t^T (T-s+\lambda) ds \\ &\quad + 2 \int_t^T (T-s+\lambda) \frac{\int_{\Omega \cap B_{R_0}} |g(x,s)|^2 G_\lambda(x,s) dx}{\int_{\Omega \cap B_{R_0}} |u(x,s)|^2 G_\lambda(x,s) dx} ds. \end{aligned}$$

Therefore, for any  $0 < T - \varepsilon \leq t < T$  (where  $\varepsilon \in (0, h_0]$  will be determined later)

$$\frac{1}{\varepsilon + \lambda} \lambda N_\lambda(T) \leq N_\lambda(t) + 2\varepsilon \|a\|_\infty^2 + 2 \int_t^T \frac{\int_{\Omega \cap B_{R_0}} |g(x,s)|^2 G_\lambda(x,s) dx}{\int_{\Omega \cap B_{R_0}} |u(x,s)|^2 G_\lambda(x,s) dx} ds. \quad (5.8)$$

Secondly, by (5.1),

$$\begin{aligned} &\frac{1}{2} \frac{d}{dt} \int_{\Omega \cap B_{R_0}} |u(x,t)|^2 G_\lambda(x,t) dx + \int_{\Omega \cap B_{R_0}} |\nabla u(x,t)|^2 G_\lambda(x,t) dx \\ &= \int_{\Omega \cap B_{R_0}} u(x,t) (-au + g)(x,t) G_\lambda(x,t) dx \end{aligned}$$

becomes

$$\begin{aligned} &\frac{1}{2} \frac{d}{dt} \int_{\Omega \cap B_{R_0}} |u(x,t)|^2 G_\lambda(x,t) dx + N_\lambda(t) \int_{\Omega \cap B_{R_0}} |u(x,t)|^2 G_\lambda(x,t) dx \\ &= - \int_{\Omega \cap B_{R_0}} a(x,t) |u(x,t)|^2 G_\lambda(x,t) dx \\ &\quad + \frac{\int_{\Omega \cap B_{R_0}} u(x,t) g(x,t) G_\lambda(x,t) dx}{\int_{\Omega \cap B_{R_0}} |u(x,t)|^2 G_\lambda(x,t) dx} \int_{\Omega \cap B_{R_0}} |u(x,t)|^2 G_\lambda(x,t) dx. \end{aligned} \quad (5.9)$$

Therefore, combining (5.8) and (5.9), we obtain that for any  $0 < T - \varepsilon \leq t < T$

$$\begin{aligned} &\frac{1}{2} \frac{d}{dt} \int_{\Omega \cap B_{R_0}} |u(x,t)|^2 G_\lambda(x,t) dx + \frac{1}{\varepsilon + \lambda} \lambda N_\lambda(T) \int_{\Omega \cap B_{R_0}} |u(x,t)|^2 G_\lambda(x,t) dx \\ &\leq (\|a\|_\infty + 2\varepsilon \|a\|_\infty^2) \int_{\Omega \cap B_{R_0}} |u(x,t)|^2 G_\lambda(x,t) dx + \int_{\Omega \cap B_{R_0}} |u(x,t)|^2 G_\lambda(x,t) dx \\ &\quad \times \left( \frac{\int_{\Omega \cap B_{R_0}} u(x,t) g(x,t) G_\lambda(x,t) dx}{\int_{\Omega \cap B_{R_0}} |u(x,t)|^2 G_\lambda(x,t) dx} + 2 \int_t^T \frac{\int_{\Omega \cap B_{R_0}} |g(x,s)|^2 G_\lambda(x,s) dx}{\int_{\Omega \cap B_{R_0}} |u(x,s)|^2 G_\lambda(x,s) dx} ds \right). \end{aligned}$$

Now, define, for any  $\varepsilon \in (0, h_0]$ ,

$$Q_{h_0, \varepsilon, \lambda} = c \left( 3 + \frac{1}{\sqrt{T}} \right) e^{(C_3 + C_4) \frac{1}{h_0}} e^{-\frac{C_4}{\varepsilon + \lambda}} \quad (5.10)$$

given from the fact that using step 1,

$$\begin{aligned} & \frac{\int_{\Omega \cap B_{R_0}} u(x, t) g(x, t) G_\lambda(x, t) dx}{\int_{\Omega \cap B_{R_0}} |u(x, t)|^2 G_\lambda(x, t) dx} + 2 \int_t^T \frac{\int_{\Omega \cap B_{R_0}} |g(x, s)|^2 G_\lambda(x, s) dx}{\int_{\Omega \cap B_{R_0}} |u(x, s)|^2 G_\lambda(x, s) dx} ds \\ & \leq Q_{h_0, \varepsilon, \lambda} \text{ for any } 0 < T - \varepsilon \leq t < T \text{ with } \varepsilon \in (0, h_0]. \end{aligned}$$

Then, it holds

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \int_{\Omega \cap B_{R_0}} |u(x, t)|^2 G_\lambda(x, t) dx \\ & \leq - \left( \frac{1}{\varepsilon + \lambda} \lambda N_\lambda(T) - \|a\|_\infty - 2\varepsilon \|a\|_\infty^2 - Q_{h_0, \varepsilon, \lambda} \right) \int_{\Omega \cap B_{R_0}} |u(x, t)|^2 G_\lambda(x, t) dx \end{aligned}$$

which implies

$$\frac{d}{dt} \left( e^{2 \left( \frac{1}{\varepsilon + \lambda} \lambda N_\lambda(T) - \|a\|_\infty - 2\varepsilon \|a\|_\infty^2 - Q_{h_0, \varepsilon, \lambda} \right) t} \int_{\Omega \cap B_{R_0}} |u(x, t)|^2 G_\lambda(x, t) dx \right) \leq 0$$

for  $0 < T - \varepsilon \leq t$ . Integrating over  $(T - \varepsilon, T - \varepsilon/2)$ , we get

$$\begin{aligned} & e^{\frac{\varepsilon}{\varepsilon + \lambda} \lambda N_\lambda(T)} \int_{\Omega \cap B_{R_0}} |u(x, T - \varepsilon/2)|^2 G_\lambda(x, T - \varepsilon/2) dx \\ & \leq e^{\varepsilon \|a\|_\infty + 2\varepsilon^2 \|a\|_\infty^2} e^{\varepsilon Q_{h_0, \varepsilon, \lambda}} \int_{\Omega \cap B_{R_0}} |u(x, T - \varepsilon)|^2 G_\lambda(x, T - \varepsilon) dx \end{aligned}$$

that is

$$\begin{aligned} & e^{\frac{\varepsilon}{\varepsilon + \lambda} \lambda N_\lambda(T)} \int_{\Omega \cap B_{R_0}} |u(x, T - \varepsilon/2)|^2 \frac{1}{(\varepsilon/2 + \lambda)^{n/2}} e^{-\frac{|x-x_0|^2}{4(\varepsilon/2 + \lambda)}} dx \\ & \leq e^{\varepsilon \|a\|_\infty + 2\varepsilon^2 \|a\|_\infty^2} e^{\varepsilon Q_{h_0, \varepsilon, \lambda}} \int_{\Omega \cap B_{R_0}} |u(x, T - \varepsilon)|^2 \frac{1}{(\varepsilon + \lambda)^{n/2}} e^{-\frac{|x-x_0|^2}{4(\varepsilon + \lambda)}} dx. \end{aligned}$$

Thus,

$$e^{\frac{\varepsilon}{\varepsilon + \lambda} \lambda N_\lambda(T)} \leq e^{\varepsilon \|a\|_\infty + 2\varepsilon^2 \|a\|_\infty^2} e^{\varepsilon Q_{h_0, \varepsilon, \lambda}} \frac{\int_{\Omega \cap B_{R_0}} |u(x, T - \varepsilon)|^2 e^{-\frac{|x-x_0|^2}{4(\varepsilon + \lambda)}} dx}{\int_{\Omega \cap B_{R_0}} |u(x, T - \varepsilon/2)|^2 e^{-\frac{|x-x_0|^2}{4(\varepsilon/2 + \lambda)}} dx}.$$

Now, since  $\frac{\varepsilon}{2} \in (0, h_0]$ ,

$$\begin{aligned} & \frac{\int_{\Omega \cap B_{R_0}} |u(x, T - \varepsilon)|^2 e^{-\frac{|x-x_0|^2}{4(\varepsilon + \lambda)}} dx}{\int_{\Omega \cap B_{R_0}} |u(x, T - \varepsilon/2)|^2 e^{-\frac{|x-x_0|^2}{4(\varepsilon/2 + \lambda)}} dx} \leq \frac{e^{\frac{((1+\delta)R)^2}{2\varepsilon}} e^{3T\|a\|_\infty} \int_{\Omega} |\varphi_0(x)|^2 dx}{\int_{\Omega \cap B_{(1+\delta)R}} |\varphi(x, T - \varepsilon/2)|^2 dx} \\ & \leq e^{\frac{((1+\delta)R)^2}{2\varepsilon}} e^{1+C_3 \frac{1}{h_0}}. \end{aligned}$$

Indeed, by Lemma 3, we know that  $\frac{e^{3T\|a\|_\infty} \int_{\Omega} |\varphi_0(x)|^2 dx}{\int_{\Omega \cap B_{(1+\delta)R}} |\varphi(x,t)|^2 dx} \leq e^{1+C_3 \frac{1}{h_0}}$  if  $T - h_0 \leq t < T$ .  
Therefore, for any  $\varepsilon \in (0, h_0]$ ,

$$\begin{aligned} \lambda N_\lambda(T) &\leq \frac{\varepsilon + \lambda}{\varepsilon} \ln \left( e^{\varepsilon \|a\|_\infty + 2\varepsilon^2 \|a\|_\infty^2} e^{\varepsilon Q_{h_0, \varepsilon, \lambda}} e^{\frac{((1+\delta)R)^2}{2\varepsilon}} e^{1+C_3 \frac{1}{h_0}} \right) \\ &\leq \frac{\varepsilon + \lambda}{\varepsilon} \left( \frac{((1+\delta)R)^2}{2\varepsilon} + \varepsilon \|a\|_\infty + 2\varepsilon^2 \|a\|_\infty^2 + \varepsilon Q_{h_0, \varepsilon, \lambda} + 1 + C_3 \frac{1}{h_0} \right). \end{aligned} \quad (5.11)$$

Step 3 .- Now, we choose  $\lambda = \mu\varepsilon$  with  $\mu \in (0, 1)$  which will be determined later and

$$\varepsilon = \frac{C_4}{2(C_3 + C_4)} h_0$$

in order that  $Q_{h_0, \varepsilon, \lambda}$  given by (5.10) satisfies the following bound

$$Q_{h_0, \varepsilon, \lambda} \leq c \left( 3 + \frac{1}{\sqrt{T}} \right) e^{\frac{C_3 + C_4}{h_0} (1 - \frac{2}{1+\mu})} = c \left( 3 + \frac{1}{\sqrt{T}} \right) e^{\frac{C_3 + C_4}{h_0} (\frac{\mu-1}{1+\mu})} \leq c \left( 3 + \frac{1}{\sqrt{T}} \right)$$

and further, using the fact that  $\varepsilon \leq h_0$ , (5.11) becomes

$$\begin{aligned} \lambda N_\lambda(T) &\leq 2 \left( 1 + h_0 \|a\|_\infty + 2h_0^2 \|a\|_\infty^2 + (C_3 + C_4) \left( 1 + \frac{(1+\delta)^2 R^2}{C_4} \right) \frac{1}{h_0} + c \left( 3 + \frac{1}{\sqrt{T}} \right) h_0 \right). \end{aligned}$$

Next, we deduce that

$$\begin{aligned} \varepsilon \lambda N_\lambda(T) &\leq 2 \left( \varepsilon + \varepsilon h_0 \|a\|_\infty + 2\varepsilon h_0^2 \|a\|_\infty^2 + \varepsilon (C_3 + C_4) \left( 1 + \frac{(1+\delta)^2 R^2}{C_4} \right) \frac{1}{h_0} + \varepsilon c \left( 3 + \frac{1}{\sqrt{T}} \right) h_0 \right) \\ &\leq 2 \left( h_0 + h_0 T \|a\|_\infty + 2h_0^3 \|a\|_\infty^2 + \frac{1}{2} (C_4 + (1+\delta)^2 R^2) + c \left( 3h_0^2 + \sqrt{\frac{h_0}{T}} h_0^{3/2} \right) \right) \\ &\leq 2 \left( 2C_1 + 2C_1^3 + \frac{1}{2} (C_4 + (1+\delta)^2 R^2) + c \left( 3C_1^2 + C_1^{3/2} \right) \right) \end{aligned}$$

where in the last line, we used the following four inequalities  $h_0 < C_1$ ,  $h_0 < T$ ,  $h_0 T \|a\|_\infty < C_1$  and  $h_0^3 \|a\|_\infty^2 < C_1^3$  obtained in (5.4) of Lemma 3. Therefore, we conclude from the above bound of  $\varepsilon \lambda N_\lambda(T)$  that

$$\begin{aligned} \frac{16\lambda}{r^2} \left( \frac{n}{4} + \lambda N_\lambda(T) \right) &\leq \frac{16}{r^2} \mu \left( \frac{n}{4} C_1 + \varepsilon \lambda N_\lambda(T) \right) \\ &\leq \mu (1 + C_0) \end{aligned} \quad (5.12)$$

for some  $C_0 > 0$  only depending on  $(n, r, R, \delta)$ .

Step 4 .- Now, we are able to bound  $\int_{\Omega \cap B_{R_0}} |u(x, T)|^2 e^{-\frac{|x-x_0|^2}{4\lambda}} dx$  as follows. Since  $\Omega \cap B_{R_0}$  is star-shaped with respect to  $x_0$ , we have (see, for example, [13, pp. 1238] or [14, Lemma 2.5]- [5, Lemma 3]),

$$\begin{aligned} &\frac{1}{16\lambda} \int_{\Omega \cap B_{R_0}} |x - x_0|^2 |u(x, T)|^2 e^{-\frac{|x-x_0|^2}{4\lambda}} dx \\ &\leq \frac{n}{4} \int_{\Omega \cap B_{R_0}} |u(x, T)|^2 e^{-\frac{|x-x_0|^2}{4\lambda}} dx + \lambda \int_{\Omega \cap B_{R_0}} |\nabla u(x, T)|^2 e^{-\frac{|x-x_0|^2}{4\lambda}} dx \end{aligned}$$

which implies

$$\begin{aligned}
& \int_{\Omega \cap B_{R_0}} |u(x, T)|^2 e^{-\frac{|x-x_0|^2}{4\lambda}} dx \\
& \leq \int_{B_r} |u(x, T)|^2 e^{-\frac{|x-x_0|^2}{4\lambda}} dx + \int_{(\Omega \cap B_{R_0}) \setminus B_r} \frac{|x-x_0|^2}{r^2} |u(x, T)|^2 e^{-\frac{|x-x_0|^2}{4\lambda}} dx \\
& \leq \int_{B_r} |u(x, T)|^2 e^{-\frac{|x-x_0|^2}{4\lambda}} dx \\
& \quad + \frac{1}{r^2} \left[ 4\lambda n \int_{\Omega \cap B_{R_0}} |u(x, T)|^2 e^{-\frac{|x-x_0|^2}{4\lambda}} dx + \int_{\Omega \cap B_{R_0}} 16\lambda^2 |\nabla u(x, T)|^2 e^{-\frac{|x-x_0|^2}{4\lambda}} dx \right] \\
& \leq \int_{B_r} |\varphi(x, T)|^2 e^{-\frac{|x-x_0|^2}{4\lambda}} dx + \frac{16\lambda}{r^2} \left[ \frac{n}{4} + \lambda N_\lambda(T) \right] \int_{\Omega \cap B_{R_0}} |u(x, T)|^2 e^{-\frac{|x-x_0|^2}{4\lambda}} dx,
\end{aligned}$$

where in the last line we used the definition of  $N_\lambda(T)$  and the fact that  $u = \varphi$  in  $B_r$ . Combining the above inequality and (5.12), we deduce that

$$\begin{aligned}
\int_{\Omega \cap B_{R_0}} |u(x, T)|^2 e^{-\frac{|x-x_0|^2}{4\lambda}} dx & \leq \int_{B_r} |\varphi(x, T)|^2 e^{-\frac{|x-x_0|^2}{4\lambda}} dx \\
& \quad + \mu(1 + C_0) \int_{\Omega \cap B_{R_0}} |u(x, T)|^2 e^{-\frac{|x-x_0|^2}{4\lambda}} dx. \tag{5.13}
\end{aligned}$$

Step 5 .- Now, we choose  $\mu \in (0, 1)$  as follows.

$$\mu = \frac{1}{2} \frac{1}{(1 + C_0)}.$$

Then,  $\lambda = \mu\varepsilon = \mu \frac{C_4}{2(C_3+C_4)} h_0 = \frac{1}{4} \frac{C_4}{(1+C_0)(C_3+C_4)} h_0$  and by using the definition of  $h_0$ , we have

$$\begin{aligned}
& \int_{\Omega \cap B_R} |\varphi(x, T)|^2 dx \\
& \leq e^{\frac{R^2}{4\lambda}} \int_{\Omega \cap B_{R_0}} |u(x, T)|^2 e^{-\frac{|x-x_0|^2}{4\lambda}} dx \\
& \leq 2e^{\frac{R^2}{4\lambda}} \int_{B_r} |\varphi(x, T)|^2 e^{-\frac{|x-x_0|^2}{4\lambda}} dx \quad \text{by (5.13),} \\
& \leq 2e^{\frac{(1+C_0)(C_3+C_4)R^2}{C_4} \frac{1}{h_0}} \int_{B_r} |\varphi(x, T)|^2 dx \\
& \leq 2 \left( (1 + C_2) \frac{e^{1 + \frac{2C_1}{T} + e^{3T}\|a\|_\infty + \|a\|_\infty^2/3} \int_{\Omega} |\varphi_0(x)|^2 dx}{\int_{\Omega \cap B_R} |\varphi(x, T)|^2 dx} \right)^{\frac{(1+C_0)(C_3+C_4)R^2}{C_1 C_4}} \int_{B_r} |\varphi(x, T)|^2 dx.
\end{aligned}$$

We conclude that the desired estimate of Lemma 4 holds with  $\beta = \frac{(1+C_0)(C_3+C_4)R^2}{C_1 C_4 + (1+C_0)(C_3+C_4)R^2} \in (0, 1)$ . This completes the proof of Lemma 4.

*Lemma 5 .-* Let  $\tilde{\omega}$  be a non-empty open set of  $\Omega$ . Then there are  $C = C(\tilde{\omega}, \Omega) > 0$  and  $\tilde{\beta} = \tilde{\beta}(\tilde{\omega}, \Omega) \in (0, 1)$  such that for any  $T > 0$  and  $\varphi_0 \in L^2(\Omega)$ ,

$$\int_{\Omega} |\varphi(x, T)|^2 dx \leq e^{C(1+\frac{1}{T}+T\|a\|_{\infty}+\|a\|_{\infty}^{2/3})} \left( \int_{\Omega} |\varphi_0(x)|^2 dx \right)^{\tilde{\beta}} \left( \int_{\tilde{\omega}} |\varphi(x, T)|^2 dx \right)^{1-\tilde{\beta}}.$$

*Proof .-* Firstly, by Lemma 4 and constructing a sequence of balls chained along a curve, we claim that, for any compact sets  $\Theta_1$  and  $\Theta_2$  with non-empty interior in  $\Omega$ , there are constants  $C = C(\Theta_1, \Theta_2, \Omega) > 0$  and  $\alpha_1 = \alpha_1(\Theta_1, \Theta_2, \Omega) \in (0, 1)$  such that

$$\int_{\Theta_1} |\varphi(x, T)|^2 dx \leq e^{C(1+\frac{1}{T}+T\|a\|_{\infty}+\|a\|_{\infty}^{2/3})} \left( \int_{\Omega} |\varphi_0(x)|^2 dx \right)^{\alpha_1} \left( \int_{\Theta_2} |\varphi(x, T)|^2 dx \right)^{1-\alpha_1}. \quad (5.14)$$

Indeed, since  $\Theta_1$  is a compact set in  $\Omega$ , there are  $R > 0$  and finitely many points  $x_1, \dots, x_M$  such that  $\Theta_1 \subset \bigcup_{i=1, \dots, M} B(x_i, R)$  and  $B(x_i, 3R) \subset \Omega$ . Next, for each  $i \in \{1, \dots, M\}$ , we choose  $\rho \in (0, R)$  and finitely many points  $q_0, \dots, q_m$  with the following properties:

$$\begin{cases} x_i = q_m \\ \Theta_2 \supset B(q_0, \rho) \\ B(q_{j+1}, \rho/2) \subset B(q_j, \rho) \quad \forall j = 0, \dots, m-1 \\ B(q_j, 3\rho) \subset \Omega \quad \forall j = 0, \dots, m. \end{cases}$$

Thanks to Lemma 4, there exist  $\sigma, \sigma_1, \alpha_1 \in (0, 1)$ , such that

$$\begin{aligned} & \int_{B(x_i, R)} |\varphi(x, T)|^2 dx \\ & \leq e^{C(1+\frac{1}{T}+T\|a\|_{\infty}+\|a\|_{\infty}^{2/3})} \left( \int_{\Omega} |\varphi_0(x)|^2 dx \right)^{\sigma} \left( \int_{B(x_i, \rho/2)} |\varphi(x, T)|^2 dx \right)^{1-\sigma} \\ & = e^{C(1+\frac{1}{T}+T\|a\|_{\infty}+\|a\|_{\infty}^{2/3})} \left( \int_{\Omega} |\varphi_0(x)|^2 dx \right)^{\sigma} \left( \int_{B(q_m, \rho/2)} |\varphi(x, T)|^2 dx \right)^{1-\sigma} \\ & \leq e^{C(1+\frac{1}{T}+T\|a\|_{\infty}+\|a\|_{\infty}^{2/3})} \left( \int_{\Omega} |\varphi_0(x)|^2 dx \right)^{\sigma} \left( \int_{B(q_{m-1}, \rho)} |\varphi(x, T)|^2 dx \right)^{1-\sigma} \\ & \leq e^{C(1+\frac{1}{T}+T\|a\|_{\infty}+\|a\|_{\infty}^{2/3})} \left( \int_{\Omega} |\varphi_0(x)|^2 dx \right)^{\sigma} \\ & \quad \times \left( e^{C(1+\frac{1}{T}+T\|a\|_{\infty}+\|a\|_{\infty}^{2/3})} \left( \int_{\Omega} |\varphi_0(x)|^2 dx \right)^{\sigma_1} \left( \int_{B(q_{m-1}, \rho/2)} |\varphi(x, T)|^2 dx \right)^{1-\sigma_1} \right)^{1-\sigma} \\ & \leq \dots \\ & \leq e^{C(1+\frac{1}{T}+T\|a\|_{\infty}+\|a\|_{\infty}^{2/3})} \left( \int_{\Omega} |\varphi_0(x)|^2 dx \right)^{\alpha_1} \left( \int_{B(q_0, \rho)} |\varphi(x, T)|^2 dx \right)^{1-\alpha_1}, \end{aligned}$$

where  $C > 0$  may change value from line to line. This implies the desired inequality (5.14).

Secondly, since  $\Omega$  is bounded with a  $C^2$  boundary, there is a finite set of triplet  $(q_j, R_j, \delta_j) \in \Omega \times \mathbb{R}_+^* \times (0, 1]$ ,  $j = 1, \dots, m$ , such that

$$\partial\Omega \subset \bigcup_{j=1, \dots, m} B(q_j, (1+2\delta_j)R_j)$$



and  $\Omega \cap B(q_j, (1 + 2\delta_j) R_j)$  is star-shaped with center  $q_j$  for some  $\delta_j$ . Then we apply Lemma 4 with  $\Omega \cap B(q_j, (1 + 2\delta_j) R_j)$  for  $j = 1, \dots, m$ , and the same arguments as above to get that, when  $\vartheta$  is a neighborhood of  $\partial\Omega$  and  $\Theta_3$  is a compact set with non-empty interior in  $\Omega$ , there are constants  $C = C(\vartheta, \Theta_3, \Omega) > 0$  and  $\alpha_2 = \alpha_2(\vartheta, \Theta_3, \Omega) \in (0, 1)$  such that

$$\int_{\vartheta} |\varphi(x, T)|^2 dx \leq e^{C(1 + \frac{1}{T} + T\|a\|_{\infty} + \|a\|_{\infty}^{2/3})} \left( \int_{\Omega} |\varphi_0|^2 dx \right)^{\alpha_2} \left( \int_{\Theta_3} |\varphi(x, T)|^2 dx \right)^{1 - \alpha_2}.$$

Finally, we derive the desired estimate from the previous two statements with  $\Omega \subset (\vartheta \cup \Theta_1)$  and  $(\Theta_2 \cup \Theta_3) \subset \tilde{\omega}$ . This completes the proof.

Now, we are able to present the proof of the observability estimate of Theorem 4.

Proof of Theorem 4 .- We start with the following interpolation estimate deduced by Lemma 5 and the Young inequality. For any  $0 \leq t_1 < t_2 \leq T$ ,

$$\|\varphi(\cdot, t_2)\|_{L^2(\Omega)} \leq \frac{K_1}{\varepsilon^{\alpha}} e^{\frac{K_2}{t_2 - t_1}} \|\varphi(\cdot, t_2)\|_{L^2(\tilde{\omega})} + \varepsilon \|\varphi(\cdot, t_1)\|_{L^2(\Omega)} \quad \forall \varepsilon > 0.$$

Here,  $\tilde{\omega} \Subset \omega \subset \Omega$ ,  $K_1 = e^{\frac{C}{2(1-\beta)}(1+T\|a\|_{\infty} + \|a\|_{\infty}^{2/3})}$  and  $K_2 = \frac{C}{2(1-\beta)}$  in Lemma 5,  $\alpha = \frac{\tilde{\beta}}{1-\tilde{\beta}}$  in Lemma 5. By Nash inequality and Poincaré inequality,

$$\|\varphi(\cdot, t_2)\|_{L^2(\tilde{\omega})} \leq \frac{K_3}{\delta^{n/2}} \|\varphi(\cdot, t_2)\|_{L^1(\omega)} + \delta \|\nabla \varphi(\cdot, t_2)\|_{L^2(\Omega)} \quad \forall \delta > 0.$$

Here  $K_3 > 0$  only depends on  $(\tilde{\omega}, \omega, \Omega)$ . By Lemma 1, we know that

$$\|\nabla \varphi(\cdot, t_2)\|_{L^2(\Omega)} \leq \frac{K_4}{(t_2 - t_1)^{1/2}} \|\varphi(\cdot, t_1)\|_{L^2(\Omega)}$$

with  $K_4 = e^{2T\|a\|_{\infty}}$ . Therefore, from the above three estimate with  $\frac{K_1}{\varepsilon^{\alpha}} e^{\frac{K_2}{t_2 - t_1}} \delta \frac{K_4}{(t_2 - t_1)^{1/2}} = \varepsilon$ , we get

$$\begin{aligned} \|\varphi(\cdot, t_2)\|_{L^2(\Omega)} &\leq \frac{K_1}{\varepsilon^{\alpha}} e^{\frac{K_2}{t_2 - t_1}} \left( \frac{K_3}{\delta^{n/2}} \|\varphi(\cdot, t_2)\|_{L^1(\omega)} + \delta \frac{K_4}{(t_2 - t_1)^{1/2}} \|\varphi(\cdot, t_1)\|_{L^2(\Omega)} \right) \\ &\quad + \varepsilon \|\varphi(\cdot, t_1)\|_{L^2(\Omega)} \\ &\leq \frac{K_1 K_3}{\varepsilon^{\alpha}} e^{\frac{K_2}{t_2 - t_1}} \left( \frac{1}{\varepsilon^{\alpha+1}} e^{\frac{K_2}{t_2 - t_1}} \frac{K_1 K_4}{(t_2 - t_1)^{1/2}} \right)^{n/2} \|\varphi(\cdot, t_2)\|_{L^1(\omega)} + 2\varepsilon \|\varphi(\cdot, t_1)\|_{L^2(\Omega)} \\ &\leq \frac{K_1 K_3}{\varepsilon^{\alpha + (\alpha+1)n/2}} e^{\left(\frac{3n}{4} + 1\right) \frac{K_2}{t_2 - t_1}} \left( \frac{K_1 K_4}{\sqrt{K_2}} \right)^{n/2} \|\varphi(\cdot, t_2)\|_{L^1(\omega)} + 2\varepsilon \|\varphi(\cdot, t_1)\|_{L^2(\Omega)} \\ &\leq \frac{K_5}{(2\varepsilon)^{\gamma}} e^{\frac{K_6}{t_2 - t_1}} \|\varphi(\cdot, t_2)\|_{L^1(\omega)} + 2\varepsilon \|\varphi(\cdot, t_1)\|_{L^2(\Omega)} \quad \forall \varepsilon > 0, \end{aligned}$$

denoting  $\gamma = \alpha \left(1 + \frac{n}{2}\right) + \frac{n}{2}$ ,  $K_5 = 2^{\alpha + (\alpha+1)n/2} K_1 K_3 \left(\frac{K_1 K_4}{\sqrt{K_2}}\right)^{n/2}$  and  $K_6 = \left(\frac{3n}{4} + 1\right) K_2$ .

On another hand, let  $E$  be a subset of positive measure in  $(0, T)$ . Let  $\ell$  be a density point of  $E$ . Using [14, Proposition 2.1], for each  $\tau > 1$ , there exists  $\ell_1 \in (\ell, T)$ , depending on  $\tau$  and  $E$ , such that the sequence  $\{\ell_m\}_{m \geq 1}$ , given by

$$\ell_{m+1} = \ell + \frac{1}{\tau^m} (\ell_1 - \ell),$$

satisfies

$$\ell_m - \ell_{m+1} \leq 3 |E \cap (\ell_{m+1}, \ell_m)| .$$

Next, let  $0 < \ell_{m+2} < \ell_{m+1} \leq t < \ell_m < \ell_1 < T$ . We apply the above interpolation inequality to get

$$\|\varphi(\cdot, t)\|_{L^2(\Omega)} \leq \frac{K_5}{\varepsilon^\gamma} e^{\frac{K_6}{t-\ell_{m+2}}} \|\varphi(\cdot, t)\|_{L^1(\omega)} + \varepsilon \|\varphi(\cdot, \ell_{m+2})\|_{L^2(\Omega)} \quad \forall \varepsilon > 0 .$$

Recall that by Lemma 1

$$\|\varphi(\cdot, \ell_m)\|_{L^2(\Omega)} \leq K_4 \|\varphi(\cdot, t)\|_{L^2(\Omega)} .$$

Therefore,

$$\|\varphi(\cdot, \ell_m)\|_{L^2(\Omega)} \leq K_4 \left( \frac{K_5}{\varepsilon^\gamma} e^{\frac{K_6}{t-\ell_{m+2}}} \|\varphi(\cdot, t)\|_{L^1(\omega)} + \varepsilon \|\varphi(\cdot, \ell_{m+2})\|_{L^2(\Omega)} \right) \quad \forall \varepsilon > 0 .$$

Finally, with  $K_7 = (K_4)^{1+\gamma} K_5$ ,

$$\|\varphi(\cdot, \ell_m)\|_{L^2(\Omega)} \leq \frac{K_7}{\varepsilon^\gamma} e^{\frac{K_6}{t-\ell_{m+2}}} \|\varphi(\cdot, t)\|_{L^1(\omega)} + \varepsilon \|\varphi(\cdot, \ell_{m+2})\|_{L^2(\Omega)} \quad \forall \varepsilon > 0 .$$

Integrating it over  $t \in E \cap (\ell_{m+1}, \ell_m)$ , it yields that

$$\begin{aligned} |E \cap (\ell_{m+1}, \ell_m)| \|\varphi(\cdot, \ell_m)\|_{L^2(\Omega)} &\leq \frac{K_7}{\varepsilon^\gamma} e^{\frac{K_6}{\ell_{m+1}-\ell_{m+2}}} \int_{\ell_{m+1}}^{\ell_m} 1_E \|\varphi(\cdot, t)\|_{L^1(\omega)} dt \\ &\quad + \varepsilon |E \cap (\ell_{m+1}, \ell_m)| \|\varphi(\cdot, \ell_{m+2})\|_{L^2(\Omega)} \quad \forall \varepsilon > 0 . \end{aligned}$$

That is, using the fact that  $\ell_m - \ell_{m+1} = \frac{1}{\tau^m} (\tau - 1) (\ell_1 - \ell)$ ,

$$\begin{aligned} \|\varphi(\cdot, \ell_m)\|_{L^2(\Omega)} &\leq \frac{1}{|E \cap (\ell_{m+1}, \ell_m)|} \frac{K_7}{\varepsilon^\gamma} e^{K_6 \left[ \frac{1}{\ell_1 - \ell} \frac{\tau^{m+1}}{\tau - 1} \right]} \int_{\ell_{m+1}}^{\ell_m} 1_E \|\varphi(\cdot, t)\|_{L^1(\omega)} dt \\ &\quad + \varepsilon \|\varphi(\cdot, \ell_{m+2})\|_{L^2(\Omega)} \\ &\leq 3 \left[ \frac{1}{\ell_1 - \ell} \frac{\tau^m}{\tau - 1} \right] \frac{K_7}{\varepsilon^\gamma} e^{K_6 \left[ \frac{1}{\ell_1 - \ell} \frac{\tau^{m+1}}{\tau - 1} \right]} \int_{\ell_{m+1}}^{\ell_m} 1_E \|\varphi(\cdot, t)\|_{L^1(\omega)} dt \\ &\quad + \varepsilon \|\varphi(\cdot, \ell_{m+2})\|_{L^2(\Omega)} \quad \forall \varepsilon > 0 . \end{aligned}$$

Therefore,

$$\begin{aligned} \|\varphi(\cdot, \ell_m)\|_{L^2(\Omega)} &\leq \frac{1}{\varepsilon^\gamma} \frac{3 K_7}{\tau K_6} e^{2K_6 \left[ \frac{1}{\ell_1 - \ell} \frac{\tau^{m+1}}{\tau - 1} \right]} \int_{\ell_{m+1}}^{\ell_m} 1_E \|\varphi(\cdot, t)\|_{L^1(\omega)} dt \\ &\quad + \varepsilon \|\varphi(\cdot, \ell_{m+2})\|_{L^2(\Omega)} \quad \forall \varepsilon > 0 . \end{aligned}$$

Take  $d = 2K_6 \left[ \frac{1}{\ell_1 - \ell} \frac{1}{\tau(\tau - 1)} \right]$ . It guarantees that

$$\begin{aligned} &\varepsilon^\gamma e^{-d\tau^{m+2}} \|\varphi(\cdot, \ell_m)\|_{L^2(\Omega)} - \varepsilon^{1+\gamma} e^{-d\tau^{m+2}} \|\varphi(\cdot, \ell_{m+2})\|_{L^2(\Omega)} \\ &\leq \frac{3 K_7}{\tau K_6} \int_{\ell_{m+1}}^{\ell_m} 1_E \|\varphi(\cdot, t)\|_{L^1(\omega)} dt \quad \forall \varepsilon > 0 . \end{aligned}$$

Take  $\varepsilon = e^{-d\tau^{m+2}}$ , then

$$\begin{aligned} & e^{-(\gamma+1)d\tau^{m+2}} \|\varphi(\cdot, \ell_m)\|_{L^2(\Omega)} - e^{-(2+\gamma)d\tau^{m+2}} \|\varphi(\cdot, \ell_{m+2})\|_{L^2(\Omega)} \\ & \leq \frac{3}{\tau} \frac{K_7}{K_6} \int_{\ell_{m+1}}^{\ell_m} 1_E \|\varphi(\cdot, t)\|_{L^1(\omega)} dt . \end{aligned}$$

Take  $\tau = \sqrt{\frac{\gamma+2}{\gamma+1}}$ , then

$$\begin{aligned} & e^{-(2+\gamma)d\tau^m} \|\varphi(\cdot, \ell_m)\|_{L^2(\Omega)} - e^{-(2+\gamma)d\tau^{m+2}} \|\varphi(\cdot, \ell_{m+2})\|_{L^2(\Omega)} \\ & \leq \frac{3}{\tau} \frac{K_7}{K_6} \int_{\ell_{m+1}}^{\ell_m} 1_E \|\varphi(\cdot, t)\|_{L^1(\omega)} dt . \end{aligned}$$

Change  $m$  to  $2m'$  and sum the above from  $m' = 1$  to infinity give the desired result. Indeed,

$$\begin{aligned} & \frac{1}{K_4} e^{-(2+\gamma)d\tau^2} \|\varphi(\cdot, T)\|_{L^2(\Omega)} \\ & \leq e^{-(2+\gamma)d\tau^2} \|\varphi(\cdot, \ell_2)\|_{L^2(\Omega)} \\ & \leq \sum_{m'=1}^{+\infty} \left( e^{-(2+\gamma)d\tau^{2m'}} \|\varphi(\cdot, \ell_{2m'})\|_{L^2(\Omega)} - e^{-(2+\gamma)d\tau^{2m'+2}} \|\varphi(\cdot, \ell_{2m'+2})\|_{L^2(\Omega)} \right) \\ & \leq \frac{3}{\tau} \frac{K_7}{K_6} \sum_{m'=1}^{+\infty} \int_{\ell_{2m'+1}}^{\ell_{2m'}} 1_E \|\varphi(\cdot, t)\|_{L^1(\omega)} dt \\ & \leq \frac{3}{\tau} \frac{K_7}{K_6} \int_0^T 1_E \|\varphi(\cdot, t)\|_{L^1(\omega)} dt . \end{aligned}$$

This concludes the proof of Theorem 4.

*Remark 6* .- When  $E = (0, T)$ , then we can take the sequence  $\{\ell_m\}_{m \geq 1}$ , as follows

$$\ell_{m+1} = \frac{T}{\tau^m} ,$$

so that the observability constant becomes

$$e^{K\left(1+\frac{1}{T}+T\|a\|_\infty+\|a\|_\infty^{2/3}\right)} .$$

When  $E$  is a positive measurable set with 0 its Lebesgue point, then we can take the sequence  $\{\ell_m\}_{m \geq 1}$ , as follows

$$\ell_{m+1} = \frac{\ell_1}{\tau^m} ,$$

where the existence of  $\ell_1$  comes from [14, Proposition 2.1], so that the observability constant becomes

$$e^{K\left(1+\frac{1}{\ell_1}+\ell_1\|a\|_\infty+\|a\|_\infty^{2/3}\right)} .$$

## References

- [1] J. Apraiz, L. Escauriaza, G. Wang and C. Zhang, Observability inequalities and measurable sets, to appear in *J. Eur. Math. Soc.*
- [2] S. Anita and D. Tataru, Null controllability for the dissipative semilinear heat equation, *Appl. Math. Optim.*, 46(2002), 97-105.
- [3] V. Barbu, *Analysis and Control of Nonlinear Infinite Dimensional Systems*, Academic Press, Boston, 1993.
- [4] T. Duyckaerts, X. Zhang and E. Zuazua, On the optimality of the observability inequalities for parabolic and hyperbolic systems with potentials, *Annales de l'Institut Henri Poincaré, Analyse Non Linéaire*, 25(2008), 1-41.
- [5] L. Escauriaza, F.J. Fernandez and S. Vessella, Doubling properties of caloric functions, *Applicable analysis*, 85(2006), 205-223.
- [6] H.O. Fattorini, *Infinite Dimensional Linear Control Systems: The Time Optimal and Norm Optimal Problems*, North-Holland Mathematics Studies 201, Elsevier, Amsterdam, 2005.
- [7] E. Fernández-Cara and S. Guerrero, Global Carleman inequalities for parabolic systems and applications to controllability, *SIAM J. Control Optim.*, 45(2006), 1399-1446.
- [8] E. Fernández-Cara and E. Zuazua, Null and approximate controllability for weakly blowing up semilinear heat equations, *Annales de l'Institut Henri Poincaré, Analyse Non Linéaire*, 17(2000), 583-616.
- [9] K. Kunisch and L. Wang, Time optimal controls of the linear Fitzhugh-Nagumo equation with pointwise control constraints, *J. Math. Anal. Appl.*, 395(2012), 114-130.
- [10] K. Kunisch and L. Wang, Time optimal control of the heat equation with pointwise control constraints, to appear in *ESAIM:COCV*.
- [11] J.-L. Lions, *Optimal Control of Systems Governed by Partial Differential Equations*, Springer, Berlin, 1971.
- [12] V. Mizel and T. Seidman, An abstract bang-bang principle and time optimal boundary control of the heat equation, *SIAM Journal on Control and Optimization*, 35(1997), 1204-1216.
- [13] K.D. Phung and G. Wang, Quantitative unique continuation for the semilinear heat equation in a convex domain, *J. Funct. Anal.*, 259(2010), 1230-1247.
- [14] K.D. Phung and G. Wang, An observability estimate for parabolic equations from a measurable set in time and its applications, *J. Eur. Math. Soc.*, 15(2013), 681-703.

- [15] G. Wang, The existence of time optimal control of semilinear parabolic equations, System & Control Letters, 53(2004) 171-175.
- [16] G. Wang,  $L^\infty$ -null controllability for the heat equation and its consequences for the time optimal control problem, SIAM J. Control Optim., 47(2008), 1701-1720.
- [17] G. Wang and L. Wang, The Bang-Bang principle of time optimal controls for the heat equation with internal controls, Systems and Control Letters, 56(2007), 709-713.