



**HAL**  
open science

# Simplicial localization of homotopy algebras over a prop

Sinan Yalin

► **To cite this version:**

| Sinan Yalin. Simplicial localization of homotopy algebras over a prop. 2013. hal-00845807v1

**HAL Id: hal-00845807**

**<https://hal.science/hal-00845807v1>**

Preprint submitted on 17 Jul 2013 (v1), last revised 2 May 2014 (v2)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# SIMPLICIAL LOCALIZATION OF HOMOTOPY ALGEBRAS OVER A PROP

SINAN YALIN

**ABSTRACT.** We prove that a weak equivalence between two cofibrant (colored) props in chain complexes induces a Dwyer-Kan equivalence between the simplicial localizations of the associated categories of algebras. This homotopy invariance under base change implies that the homotopy category of homotopy algebras over a prop  $P$  does not depend on the choice of a cofibrant resolution of  $P$ , and gives thus a coherence to the notion of algebra up to homotopy in this setting. The result is established more generally for algebras in combinatorial monoidal dg categories.

*Keywords :* props, bialgebras category, simplicial localization,  $(\infty, 1)$ -category, homotopical algebra, homotopy invariance.

*AMS :* 18G55 ; 18D50 ; 18D10 ; 55U10.

## CONTENTS

1. Introduction	1
2. Recollections on props, algebras and homotopy	3
3. Simplicial localization of algebras over cofibrant props	5
3.1. Step 1. From simplicial categories to relative categories	6
3.2. Step 2. From relative categories to complete Segal spaces	7
3.3. Step 3. Comparison of classification diagrams	7
References	10

## 1. INTRODUCTION

Props are combinatorial devices introduced by MacLane in [21] in order to parametrize operations with multiple inputs and outputs. The usual categories of bialgebras, like the classical associative bialgebras, the Lie bialgebras, ..., which are defined by combining an algebra and a coalgebra structure constrained by a distribution relation (see [18] for many examples), can be modelled by props. The category of Frobenius bialgebras gives another example of a bialgebra structure modelled by a prop (of a more general form). Under suitable conditions on the ambient category, the category of props admits a model category structure [13]. Let  $P$  be a prop. We define a  $P$ -algebra up to homotopy, or homotopy  $P$ -algebra, as an algebra over a cofibrant resolution  $Q(P)$  of  $P$ . When going from a  $P$ -algebra structure to a  $Q(P)$ -algebra structure, the relations defining the  $P$ -algebra structure are relaxed and are only satisfied up to a (whole) set of (coherent) homotopies. This definition involves the choice of a cofibrant resolution, so a natural coherence request is to ask for the homotopy theory of homotopy algebras to not depend on this choice.

Let  $(C, W)$  be a pair of categories such that  $W$  is a subcategory of weak equivalences of  $C$  in the sense of [9]. Such a pair is called a relative category in [1]. To every relative category one can associate its simplicial localization [8], denoted by  $L(C, W)$ . It is a simplicial category which encodes the whole homotopy theory of  $(C, W)$ . The category of connected components  $\pi_0 L(C, W)$  gives the homotopy category  $C[W^{-1}]$ . Our main contribution in this paper is to prove an optimal homotopical invariance result for the homotopy theory of algebras over props, which provides a definitive answer to our coherence problem. We work with props in the category  $Ch_{\mathbb{K}}$  of chain complexes over a field  $\mathbb{K}$  (dg props for short). Let  $Ch_{\mathbb{K}}^P$  denote the category of algebras over a dg prop  $P$ , and  $wCh_{\mathbb{K}}^P$  its subcategory of morphisms of  $P$ -algebras which are quasi-isomorphisms in  $Ch_{\mathbb{K}}$ . The main result in this paper reads:

**Theorem 1.1.** *A weak equivalence  $P \xrightarrow{\sim} Q$  between cofibrant (colored) dg props induces a zigzag of Dwyer-Kan equivalences of simplicial localizations*

$$L(Ch_{\mathbb{K}}^Q, wCh_{\mathbb{K}}^Q) \cong L(Ch_{\mathbb{K}}^P, wCh_{\mathbb{K}}^P).$$

Thus the homotopy theory (both the primary and secondary information) of homotopy  $P$ -algebras does not depend on the choice of a cofibrant resolution of  $P$ . Theorem 1.1 implies in particular the existence of an equivalence of homotopy categories

$$Ch_{\mathbb{K}}^Q[(wCh_{\mathbb{K}}^Q)^{-1}] \cong Ch_{\mathbb{K}}^P[(wCh_{\mathbb{K}}^P)^{-1}].$$

We actually need the result of theorem 1.1 to establish this more basic relation. Indeed, other classical methods fail. For instance, since algebras over a cofibrant prop do not form a model category in general, their homotopy category is difficult to handle directly and we cannot use the machinery of Quillen equivalences. We combine simplicial localization techniques with the properties of  $\infty$ -categories, and elaborate on methods introduced in [26] about the invariance of classification spaces in the propic setting.

*Relation with the previous works.* Homotopy invariance for algebraic structures has a long history, going back to [7]. There are two kinds of homotopy invariance results that one wants to establish, giving rise to two parallel stories of results. The first sort consists in proving that homotopy algebraic structures can be transferred over homotopy equivalences in the ground category. This goes back to Boardman and Vogt [7] in the topological setting. It was established by numerous authors for different special cases of algebras, before the general result for homotopy algebras over dg operads by Markl [22]. Fresse finally proved it for props in monoidal model categories [13], a result extended by Johnson and Yau to the colored case [17].

The second sort is a homotopy invariance under base change, that is an invariance of the homotopy category of homotopy algebras, up to equivalence, under weak equivalences of cofibrant operads. It goes back to the preprint of Getzler and Jones [14] for dg operads and the thesis of Rezk [23] for simplicial operads. The general case of algebras over cofibrant operads in monoidal model categories has been treated by Berger and Moerdijk [3], then extended to the colored case [4]. It has been extended to colored props by Johnson and Yau [17] under the strong assumption of the existence of a free algebra functor. This covers the case of cartesian categories but exclude for instance chain complexes.

This paper brings a final answer for homotopy invariance under base change of algebras over cofibrant colored props in dg categories, with new methods different

from those of the preceding works (which rely on the existence of a model structure on algebras).

*Organization of the paper.*

Section 2 consists in brief recollections about symmetric monoidal model categories over a base category and props and algebras over a prop in this setting. Section 3 is the heart of this paper. It has been divided in three steps, each one using properties of  $(\infty, 1)$ -categories to finally reduce the problem to a comparison of classification diagrams. This comparison is proved by using a homotopy invariance result obtained in [26].

**Acknowledgements.** I would like to thank Denis-Charles Cisinski for decisive remarks on results of [26] which lead to this work.

## 2. RECOLLECTIONS ON PROPS, ALGEBRAS AND HOMOTOPY

We use the following relative version of the notion of a symmetric monoidal category:

**Definition 2.1.** Let  $\mathcal{C}$  be a symmetric monoidal category. A symmetric monoidal category over  $\mathcal{C}$  is a symmetric monoidal category  $(\mathcal{E}, \otimes, 1_{\mathcal{E}})$  endowed with an external tensor product  $\otimes : \mathcal{C} \times \mathcal{E} \rightarrow \mathcal{E}$  satisfying the following natural unit, associativity and symmetry constraints:

- (1)  $\forall X \in \mathcal{E}, 1_{\mathcal{C}} \otimes X \cong X$ ,
- (2)  $\forall X \in \mathcal{E}, \forall C, D \in \mathcal{C}, (C \otimes D) \otimes X \cong C \otimes (D \otimes X)$ ,
- (3)  $\forall C \in \mathcal{C}, \forall X, Y \in \mathcal{E}, C \otimes (X \otimes Y) \cong (C \otimes X) \otimes Y \cong X \otimes (C \otimes Y)$ .

We apply this definition to the case  $\mathcal{C} = Ch_{\mathbb{K}}$ , the category of chain complexes over a field  $\mathbb{K}$ . We assume that all small limits and small colimits exist in  $\mathcal{E}$ , and suppose moreover that the external tensor product of  $\mathcal{E}$  also preserves colimits in each variable. This last condition implies (under mild assumptions) the existence of an external hom bifunctor  $Hom_{\mathcal{E}}(-, -) : \mathcal{E}^{op} \times \mathcal{E} \rightarrow \mathcal{C}$  satisfying an adjunction relation

$$\forall C \in \mathcal{C}, \forall X, Y \in \mathcal{E}, Mor_{\mathcal{E}}(C \otimes X, Y) \cong Mor_{\mathcal{C}}(C, Hom_{\mathcal{E}}(X, Y))$$

(so  $\mathcal{E}$  is naturally an enriched category over  $\mathcal{C}$ ). We also assume that  $\mathcal{E}$  is equipped with such a hom-bifunctor in the sequel.

When  $\mathcal{C}$  is a symmetric monoidal category equipped with a model structure, we require the following compatibility axioms:

**Definition 2.2.** (1) A symmetric monoidal model category is a symmetric monoidal category  $\mathcal{C}$  endowed with a model category structure such that the following axioms hold:

**MM0.** The unit object  $1_{\mathcal{C}}$  of  $\mathcal{C}$  is cofibrant.

**MM1.** The pushout-product  $(i_*, j_*) : X \otimes T \oplus_{X \otimes Z} Y \otimes Z \rightarrow Y \otimes T$  of cofibrations  $i : X \rightarrow Y$  and  $j : Z \rightarrow T$  is a cofibration which is also acyclic as soon as  $i$  or  $j$  is so.

(2) Suppose that  $\mathcal{C}$  is a symmetric monoidal model category. A symmetric monoidal category  $\mathcal{E}$  over  $\mathcal{C}$  is a symmetric monoidal model category over  $\mathcal{C}$  if the axiom MM0 holds and the axiom MM1 holds for both the internal and external tensor products of  $\mathcal{E}$ .

Axiom MM0 ensures the existence of a unit for the monoidal structure of the homotopy category. Axiom MM1 provides the necessary assumptions to make the

tensor product a Quillen bifunctor. Let us mention that axiom MM0 is weakened in the usual definition of a monoidal model category. We refer the reader to [16] for more details about monoidal model categories. The category  $Ch_{\mathbb{K}}$  of chain complexes over a field  $\mathbb{K}$  is our main working example of symmetric monoidal model category.

A differential graded (dg)  $\Sigma$ -bobject is a sequence  $M = \{M(m, n)\}_{m, n \in \mathbb{N}}$  of chain complexes such that each  $M(m, n)$  is endowed with a left action of the symmetric group  $\Sigma_m$  and a right action of the symmetric group  $\Sigma_n$  commuting with the left one. We can see  $M(m, n)$  as a space of operations with  $m$  inputs and  $n$  outputs, and the action of the symmetric groups as permutations of the inputs and the outputs. We call  $(m, n)$  the biarity of such an operation. Composing operations of two  $\Sigma$ -bobjects  $M$  and  $N$  amounts to considering 2-levelled directed graphs (with no loops) with the first level indexed by operations of  $M$  and the second level by operations of  $N$ . Vertical composition by grafting and horizontal composition by concatenation allows one to define props.

**Definition 2.3.** A dg prop is a symmetric monoidal category  $P$ , enriched over  $Ch_{\mathbb{K}}$ , with  $\mathbb{N}$  as object set and the tensor product given by  $m \otimes n = m + n$  on objects.

Equivalently, a prop is a  $\Sigma$ -bobject equipped with horizontal products

$$\circ_h : P(m_1, n_1) \otimes P(m_2, n_2) \rightarrow P(m_1 + m_2, n_1 + n_2),$$

vertical composition products

$$\circ_v : P(k, n) \otimes P(m, k) \rightarrow P(m, n)$$

and units  $1 \rightarrow P(n, n)$  corresponding to identity morphisms of the objects  $n \in \mathbb{N}$  in  $P$ . These operations satisfy relations coming from the axioms of symmetric monoidal categories. We refer the reader to Enriquez and Etingof [11] for an explicit description of props in the context of modules over a ring. We denote by  $\mathcal{P}$  the category of props.

There exists a free prop functor  $\mathcal{P}$  fitting an adjunction

$$\mathcal{F} : \Sigma \rightleftarrows \mathcal{P} : U$$

with the forgetful functor  $U$  between the category of props and the category of  $\Sigma$ -bobjects. There is an explicit construction of the free prop for which we refer the reader to [13].

For a prop  $P$  in  $Ch_{\mathbb{K}}$ , we can define the notion of  $P$ -algebra in a symmetric monoidal category over  $Ch_{\mathbb{K}}$ :

**Definition 2.4.** Let  $\mathcal{E}$  be a symmetric monoidal category over  $Ch_{\mathbb{K}}$ .

(1) The endomorphism prop of  $X \in \mathcal{E}$  is given by  $End_X(m, n) = Hom_{\mathcal{E}}(X^{\otimes m}, X^{\otimes n})$  where  $Hom_{\mathcal{E}}(-, -)$  is the external hom bifunctor of  $\mathcal{E}$ .

(2) Let  $P$  be a prop in  $Ch_{\mathbb{K}}$ . A  $P$ -algebra in  $\mathcal{E}$  is an object  $X \in \mathcal{E}$  equipped with a prop morphism  $P \rightarrow End_X$ .

Now we are interested in doing homotopical algebra with props and their algebras. Concerning props, we have to distinguish two cases. In the case where  $\mathbb{K}$  is of characteristic zero, we can use the adjunction  $\mathcal{F} : \Sigma \rightleftarrows \mathcal{P} : U$  to transfer the cofibrantly generated model category structure on  $\Sigma$ -bobjects to the whole category of props (see [13], theorem 5.5). The weak equivalences and fibrations are the

componentwise quasi-isomorphisms and surjections, and the generating (acyclic) cofibrations are the images under the free prop functor of the generating (acyclic) cofibrations of the category of diagrams  $\Sigma$ , which we equip with the usual projective model structure (see [15], theorem 11.6.1).

In the positive characteristic case, this construction works well only with the subcategory of props with non-empty inputs (respectively, outputs), and does not give a full model structure, but only a semi-model structure on this subcategory (see [13]). A prop  $P$  has non-empty inputs if it satisfies

$$P(0, n) = \begin{cases} \mathbb{K}, & \text{if } n = 0, \\ 0 & \text{otherwise.} \end{cases}$$

We define in a symmetric way a prop with non-empty outputs. We refer the reader to theorem 4.9 of [13] for more details about this result. The semi-model structure is close enough to a full model structure to define the homotopy category and do similarly homotopical algebra. In the remaining part of the paper, we work in any of these settings. In the positive characteristic case, we tacitly assume that our props satisfy the non-empty inputs requirement.

Concerning algebras over a dg prop, the situation is far more involved than in the operadic case. In general, there is no model structure on such a category, there is even no limits or colimits (and no free algebra functor). However, there are other ways to recover information of a homotopical nature about algebras over a prop. In order to get the full homotopy theory of any category with weak equivalences, one has to consider its simplicial localization. This is the topic of the next section.

### 3. SIMPLICIAL LOCALIZATION OF ALGEBRAS OVER COFIBRANT PROPS

Let  $\mathcal{E}$  be a combinatorial symmetric monoidal model category over  $Ch_{\mathbb{K}}$  and  $(\mathcal{E}^c)$  its subcategory of cofibrant objects. A combinatorial category is a cofibrantly generated model category which is also locally presentable, see the appendix of [20] for the precise definition of this notion. Let  $(\mathcal{E}^c)^P$  denote the category of algebras over a dg prop  $P$ , and  $w(\mathcal{E}^c)^P$  its subcategory of morphisms of  $P$ -algebras which are weak equivalences in  $\mathcal{E}$ . The goal of this section is to prove the following result of homotopy invariance, which implies theorem 1.1 as a special case:

**Theorem 3.1.** *A weak equivalence  $P \xrightarrow{\sim} Q$  between two cofibrant dg (colored) props induces a zigzag of Dwyer-Kan equivalences between the associated simplicial localizations*

$$L^H((\mathcal{E}^c)^Q, w(\mathcal{E}^c)^Q) \xleftarrow{\sim} \dots \xrightarrow{\sim} L^H((\mathcal{E}^c)^P, w(\mathcal{E}^c)^P).$$

The proof is divided in three steps. Each step consists roughly in reducing the equivalence of theorem 3.1, by rewriting it in terms of different models of  $(\infty, 1)$ -categories until we abut to a comparison problem of bisimplicial sets. We solve this problem by using an improved version of the homotopy invariance theorem of classification spaces obtained in [26].

Here we use three models of  $(\infty, 1)$ -categories: complete Segal spaces ([24]), simplicial categories ([8], [9],[10], [5]) and relative categories ([1],[2]). We will not enter in the details of these theories but just recall informally, when it will be necessary, the definitions and properties we need.

**Set theoretic warning.** The construction of these models of  $(\infty, 1)$ -categories is a priori established in the setting of small categories, in order to avoid set theoretic

problems like simplicial proper classes instead of sets or categories which are not locally small. However in practice one often wants to apply these results to large categories. We adopt therefore Grothendieck axiom of universes to sort out this issue: for every set there exists a universe in which this set is an element. Thus there exists a universe  $U$  in which the categories  $(\mathcal{E}^c)^Q$  and  $(\mathcal{E}^c)^P$  are  $U$ -small.

**3.1. Step 1. From simplicial categories to relative categories.** A simplicial category is a category enriched over simplicial sets. We denote by  $SCat$  the category of simplicial categories. There exists functorial cosimplicial resolutions and simplicial resolutions in any model category ([10],[15]), so model categories provide examples of simplicial categories. One recovers the morphisms of the homotopy category between a cofibrant resolution and a fibrant resolution by taking the set of connected components of the corresponding simplicial mapping space. Another more general example is the simplicial localization developed by Dwyer and Kan [8]. Let  $(C, W)$  be a pair of categories such that  $W$  is a subcategory of  $C$  containing all the objects of  $W$ . We call  $W$  the category of weak equivalences of  $C$ . Such a pair is called a relative category in [1]. To any relative category Dwyer and Kan associates a simplicial category  $L(C, W)$  called its simplicial localization. They developed also another simplicial localization, the hammock localization  $L^H(C, W)$  [9]. The two are actually equivalent, and each one has its own advantages. The hammock localization has an explicit description of the simplicial mapping spaces. By taking the sets of connected components of the mapping spaces, we get  $\pi_0 L(C, W) \cong C[W^{-1}]$  where  $C[W^{-1}]$  is the localization of  $C$  with respect to  $W$  (the homotopy category of  $(C, W)$ ).

Let us define Dwyer-Kan equivalences:

**Definition 3.2.** Let  $C$  and  $D$  be two simplicial categories. A functor  $f : C \rightarrow D$  is a Dwyer-Kan equivalence if it induces weak equivalences of simplicial sets  $Map_C(X, Y) \xrightarrow{\sim} Map_D(fX, fY)$  for every  $X, Y \in C$  and an equivalence of categories  $\pi_0 C \xrightarrow{\sim} \pi_0 D$ .

In particular, every Quillen equivalence of model categories gives rise to a Dwyer-Kan equivalence of their simplicial localizations. According to [5], there exists a model category structure on the category of (small) simplicial categories with the Dwyer-Kan equivalences as weak equivalences. Since every simplicial category is Dwyer-Kan equivalent to the simplicial localization of a certain relative category, this model structure forms a homotopy theory of homotopy theories.

Let us denote by  $RelCat$  the category of relative categories. According to [1], there is an adjunction between the category of bisimplicial sets and the category of relative categories

$$K_\xi : sSets^{\Delta^{op}} \rightleftarrows RelCat : N_\xi$$

which lifts any Bousfield localization of the Reedy model structure of bisimplicial sets into a model structure on  $RelCat$ . In the particular case of the Bousfield localization defining the complete Segal spaces [24], one obtains a Quillen equivalent homotopy theory of the homotopy theories in  $RelCat$  [1]. We refer the reader to [1] for the definition of the functor  $N_\xi$ . The only property of  $N_\xi$  we will need is that it induces an equivalence of homotopy categories.

Recall that we want to get a zigzag of Dwyer-Kan equivalences

$$L^H((\mathcal{E}^c)^Q, w(\mathcal{E}^c)^Q) \xleftarrow{\sim} \dots \xrightarrow{\sim} L^H((\mathcal{E}^c)^P, w(\mathcal{E}^c)^P).$$

According to [2], a morphism of relative categories is a weak equivalence if and only if it induces a Dwyer-Kan equivalence of the associated simplicial localizations. The simplicial localization functor induces an equivalence of homotopy categories between  $Ho(RelCat)$  and  $Ho(SCat)$  [2]. So the existence of such a zigzag amounts to the existence of a zigzag of weak equivalences of  $RelCat$

$$((\mathcal{E}^c)^Q, w(\mathcal{E}^c)^Q) \xleftarrow{\cdot} \xrightarrow{\cdot} \dots \xrightarrow{\cdot} ((\mathcal{E}^c)^P, w(\mathcal{E}^c)^P).$$

**3.2. Step 2. From relative categories to complete Segal spaces.** Although simplicial categories are the most intuitive model for  $(\infty, 1)$ -categories, Dwyer-Kan equivalences are difficult to detect. A nice model with weak equivalences easier to handle has been developed by Rezk in [24], namely the category of complete Segal spaces. We denote this category by  $CSS$ . It has a model structure defined by a certain left Bousfield localization of the standard Reedy model structure on bisimplicial sets. The fibrant objects of  $CSS$  are precisely the complete Segal spaces. To each complete Segal space one can associate the homotopy theory of a certain relative category. The Reedy weak equivalences between two complete Segal spaces are precisely the Dwyer-Kan equivalences between their associated homotopy theories. An important instance of complete Segal space is the classification diagram of a model category, which will be defined in definition 3.3. Thus a Reedy weak equivalence of classification diagrams of model categories corresponds to an equivalence of their homotopy categories.

Recall that according to [1], the functor  $N_\xi$  induces an equivalence of homotopy categories

$$RN_\xi : Ho(RelCat) \xrightarrow{\sim} Ho(CSS).$$

Consequently, obtaining an isomorphism between  $((\mathcal{E}^c)^Q, w(\mathcal{E}^c)^Q)$  and  $((\mathcal{E}^c)^P, w(\mathcal{E}^c)^P)$  in  $Ho(RelCat)$  amounts to establishing the existence of an isomorphism of the form

$$RN_\xi((\mathcal{E}^c)^Q, w(\mathcal{E}^c)^Q) \cong RN_\xi((\mathcal{E}^c)^P, w(\mathcal{E}^c)^P)$$

in  $Ho(CSS)$  where  $R(-)$  stands for the right derived Quillen functor. Such an isomorphism corresponds to a zigzag of weak equivalences in  $CSS$ .

**3.3. Step 3. Comparison of classification diagrams.** Let us denote by  $sSets$  the category of simplicial sets.

**Definition 3.3.** Let  $(C, W)$  be a relative category. Its classification diagram, denoted by  $N(C, W)$ , is the bisimplicial set  $N(C, W) : \Delta^{op} \rightarrow sSets$  defined by

$$N(C, W)([n]) = \mathcal{N}we(C^{[n]})$$

where  $C^{[n]}$  is the category of diagrams over  $[n] = 0 \rightarrow \dots \rightarrow n$ , we write  $we(C^{[n]})$  for the subcategory defined by the (pointwise) weak-equivalences in this diagram category, and  $\mathcal{N}$  refers to the nerve functor.

The simplicial set  $N(C, W)([0]) = \mathcal{N}W$  is the classification space of  $(C, W)$ . This construction gives rise to a functor  $N : RelCat \rightarrow CSS$ .

According to lemma 5.4 of [1], there is a natural Reedy weak equivalence

$$N \xrightarrow{\sim} N_\xi$$

so the zigzag of weak equivalences between  $N_\xi((\mathcal{E}^c)^Q, w(\mathcal{E}^c)^Q)$  and  $N_\xi((\mathcal{E}^c)^P, w(\mathcal{E}^c)^P)$  can be obtained by comparing  $N((\mathcal{E}^c)^Q, w(\mathcal{E}^c)^Q)$  and  $N((\mathcal{E}^c)^P, w(\mathcal{E}^c)^P)$ . We have the following homotopy invariance result:



**Theorem 3.4.** *A weak equivalence of cofibrant dg props  $\varphi : P \xrightarrow{\sim} Q$  induces a Reedy weak equivalence of classification diagrams*

$$N((\mathcal{E}^c)^Q, w(\mathcal{E}^c)^Q) \xrightarrow{\sim} N((\mathcal{E}^c)^P, w(\mathcal{E}^c)^P).$$

Since Reedy equivalences are also weak equivalences in *CSS* by construction, we finally proved the existence of a zigzag of weak equivalences of *CSS*

$$N_\xi((\mathcal{E}^c)^Q, w(\mathcal{E}^c)^Q) \xleftarrow{\sim} N((\mathcal{E}^c)^Q, w(\mathcal{E}^c)^Q) \xrightarrow{\sim} N((\mathcal{E}^c)^P, w(\mathcal{E}^c)^P) \xrightarrow{\sim} N_\xi((\mathcal{E}^c)^P, w(\mathcal{E}^c)^P).$$

*Proof.* Let  $\mathcal{E}$  be a combinatorial symmetric monoidal model category over  $Ch_{\mathbb{K}}$ . Let  $I$  be a small category and  $\mathcal{E}^I$  the category of  $I$ -diagrams in  $\mathcal{E}$ . We equip  $\mathcal{E}^I$  with the standard injective model structure. We use the following result which holds for any category of diagrams as soon as we assume that  $\mathcal{E}$  is combinatorial (see loc. cit.):

**Proposition 3.5.** *(Lurie, proposition A.2.8.2 of [20]) Let us suppose that  $\mathcal{E}$  is combinatorial. Then the injective model structure on the category of diagrams  $\mathcal{E}^I$ , with point-wise weak-equivalences and cofibrations, is combinatorial (in particular, it is cofibrantly generated).*

In our setting, we also have the following proposition which we use in the proof of theorem 3.4:

**Proposition 3.6.** *The model category  $(\mathcal{E}^I)_{inj}$  is a symmetric monoidal model category over  $Ch_{\mathbb{K}}$ .*

*Proof.* The proof works in three steps. First we have to prove that the model category  $(\mathcal{E}^I)_{inj}$  is a symmetric monoidal model category for the pointwise tensor product. The pushout-product is a pointwise pushout-product since colimits are created pointwise, and (acyclic) cofibrations are the pointwise (acyclic) cofibrations so one has just to apply axiom MM1 of  $\mathcal{E}$ .

Then we claim that  $(\mathcal{E}^I)_{inj}$  is a symmetric monoidal model category over  $\mathcal{E}$  for the external tensor product defined for every  $E \in \mathcal{E}$  and  $F \in \mathcal{E}^I$  by

$$\forall i \in I, (E \otimes_e F)(i) = E \otimes F(i).$$

This follows from the following argument. Let  $f$  be a cofibration of  $\mathcal{E}$  and  $\phi$  be a cofibration of  $(\mathcal{E}^I)_{inj}$ . We form the pushout-product  $f \square \phi$ . By definition of the external tensor product, for every  $i \in I$  the map  $(f \square \phi)(i) = f \square \phi(i)$  is a pushout-product of  $f$  and  $\phi(i)$  in  $\mathcal{E}$ . We apply axiom MM1 in  $\mathcal{E}$  and use the fact that cofibrations are defined pointwise in  $(\mathcal{E}^I)_{inj}$ , so  $f \square \phi$  is a cofibration. If one of these two maps is acyclic then  $f \square \phi$  is acyclic by the same argument.

Finally we prove the following general property: let  $\mathcal{E}$  be a symmetric monoidal model category over  $\mathcal{C}$  and  $\mathcal{D}$  be a symmetric monoidal model category over  $\mathcal{E}$ . Then  $\mathcal{D}$  is a symmetric monoidal model category over  $\mathcal{C}$ .

We have a symmetric monoidal functor  $\eta_{\mathcal{E}} : \mathcal{C} \rightarrow \mathcal{E}$  defined by  $\eta_{\mathcal{E}}(C) = C \otimes_{\mathcal{C}}^{\mathcal{E}} 1_{\mathcal{E}}$  where  $\otimes_{\mathcal{C}}^{\mathcal{E}}$  is the external tensor product of  $\mathcal{E}$  over  $\mathcal{C}$  and  $1_{\mathcal{E}}$  is the unit of  $\mathcal{E}$ . According to axiom MM0 in  $\mathcal{E}$ , the unit  $1_{\mathcal{E}}$  is cofibrant, so by applying axiom MM1 for the external tensor product we see that  $\eta_{\mathcal{E}}$  preserves cofibrations and acyclic cofibrations.

Now, let  $\otimes_{\mathcal{E}}^{\mathcal{D}}$  denote the external tensor product of  $\mathcal{D}$  over  $\mathcal{E}$ . We can define an external tensor product of  $\mathcal{D}$  over  $\mathcal{C}$  by

$$C \otimes_{\mathcal{C}}^{\mathcal{D}} D = \eta_{\mathcal{E}}(C) \otimes_{\mathcal{E}}^{\mathcal{D}} D$$

where  $C \in \mathcal{C}$  and  $D \in \mathcal{D}$ . It satisfies all the required axioms of an external tensor product since  $\otimes_{\mathcal{C}}^{\mathcal{E}}$  is an external tensor product and  $\eta_{\mathcal{E}}$  is a symmetric monoidal functor. The external tensor product  $\otimes_{\mathcal{D}}^{\mathcal{E}}$  satisfies MM1 and  $\eta_{\mathcal{E}}$  preserves cofibrations and acyclic cofibrations, so  $\otimes_{\mathcal{C}}^{\mathcal{D}}$  satisfies also MM1.

We apply this property to  $(\mathcal{E}^I)_{inj}$ ,  $\mathcal{E}$  and  $Ch_{\mathbb{K}}$  to conclude the proof of this proposition.  $\square$

In the injective model structure, cofibrant diagrams are diagrams of cofibrant objects of  $\mathcal{E}$ , that is  $((\mathcal{E}^I)_{inj})^c = ((\mathcal{E}^c)^I)_{inj}$ . We now apply the following homotopy invariance result to  $(\mathcal{E}^I)_{inj}$ :

**Theorem 3.7.** (*Yalin [26]*) *Let  $\mathcal{E}$  be a cofibrantly generated symmetric monoidal model category over  $Ch_{\mathbb{K}}$ . Let  $\varphi : P \xrightarrow{\sim} Q$  be a weak equivalence between two cofibrant props defined in  $Ch_{\mathbb{K}}$ . This morphism  $\varphi$  gives rise to a functor  $\varphi^* : w(\mathcal{E}^c)^Q \rightarrow w(\mathcal{E}^c)^P$  which induces a weak equivalence of simplicial sets  $\mathcal{N}\varphi^* : \mathcal{N}w(\mathcal{E}^c)^Q \rightarrow \mathcal{N}w(\mathcal{E}^c)^P$ .*

If we suppose that  $\mathbb{K}$  is of characteristic zero, then this result can be extended to colored props, i.e. props equipped with a set of colors labeling inputs and outputs of the operations (see [12]).

A weak equivalence of cofibrant dg props  $\varphi : P \xrightarrow{\sim} Q$  induces a weak equivalence of classification spaces

$$\mathcal{N}w((\mathcal{E}^c)^Q)^I = \mathcal{N}w((\mathcal{E}^c)^I)^Q \xrightarrow{\sim} \mathcal{N}w((\mathcal{E}^c)^I)^P = \mathcal{N}w((\mathcal{E}^c)^P)^I,$$

in particular for  $I = [n]$  and for every integer  $n$ , giving rise to the desired Reedy weak equivalence of bisimplicial sets.  $\square$

### Conclusion.

Recall that according to theorem 1.1 of [17], theorem 2.5 can be extended to the colored case when we work over a field of characteristic zero. Let  $P$  be a differential graded colored prop over a field of characteristic zero. The homotopy theory of homotopy  $P$ -algebras does not depend, up to Dwyer-Kan equivalence, on the choice of a cofibrant resolution of  $P$ . This result works also for 1-colored dg props (i.e. the props of definition 2.3) over a field of any characteristic when we consider props with non-empty inputs.

Moreover, this result enables us to give a sense to homotopical realization problems in this setting. Such a realization problem is of the following form : suppose that  $X$  is a chain complex such that  $H_*X$  is endowed with a  $P$ -algebra structure. Then determine the structures of  $P$ -algebra up to homotopy on  $X$  realizing the  $P$ -algebra structure on the homology.

Examples in the 1-colored case include the list of generalized bialgebras described in [18], for instance Hopf algebras, non-commutative Hopf algebras, Lie bialgebras, etc. These bialgebras are determined by a distributive law between operations and cooperations. Other examples of bialgebras which are not of this kind include Frobenius bialgebras and more general topological field theories. Typical examples of colored props are the one encoding diagrams of algebras (see [12]), so this is an invariance result in particular for homotopy diagrams of various kinds of homotopy algebras.

## REFERENCES

- [1] C. Barwick, D. M. Kan, *Relative categories: another model for the homotopy theory of homotopy theories*, Indagationes Mathematicae 23 (2012), 42-68.
- [2] C. Barwick, D. M. Kan, *A characterization of simplicial localization functors and a discussion of DK equivalences*, Indagationes Mathematicae 23 (2012), 69-79.
- [3] C. Berger, I. Moerdijk, *Axiomatic homotopy theory for operads*, Comment. Math. Helv. 78 (2003), 805-831.
- [4] C. Berger, I. Moerdijk, *Resolution of coloured operads and rectification of homotopy algebras*, Categories in algebra, geometry and mathematical physics, Contemp. Math. 431 (2007), 31-58.
- [5] J. Bergner, *A model category structure on the category of simplicial categories*, Trans. Amer. Math. Soc. 359 (2007), 2043-2058.
- [6] J. Bergner, *A survey of  $(\infty, 1)$ -categories*, in J. Baez and J. P. May, Towards Higher Categories, IMA Volumes in Mathematics and Its Applications, Springer, 2010, 69-83.
- [7] J. M. Boardman, R. M. Vogt, *Homotopy invariant algebraic structures on topological spaces*, Lecture Notes in Mathematics 347, Springer-Verlag, 1973.
- [8] W. G. Dwyer, D. M. Kan, *Simplicial localization of categories*, J. Pure and Applied Algebra 17 (1980), 267-284.
- [9] W. G. Dwyer, D. M. Kan, *Calculating simplicial localizations*, J. Pure and Applied Algebra 18 (1980), 17-35.
- [10] W. G. Dwyer, D. M. Kan, *Function complexes in homotopical algebra*, Topology 19 (1980), 427-440.
- [11] B. Enriquez, P. Etingof, *On the invertibility of quantization functors*, Journal of Algebra volume 289 (2005), 321-345.
- [12] Y. Fregier, M. Markl, D. Yau, *The  $L_\infty$ -deformation complex of diagrams of algebras*, New York J. Math. 15 (2009), 353-392.
- [13] B. Fresse, *Props in model categories and homotopy invariance of structures*, Georgian Math. J. 17 (2010), pp. 79-160.
- [14] E. Getzler, J. D. S. Jones, *Operads, homotopy algebra and iterated integrals for double loop spaces*, 1994.
- [15] P. S. Hirschhorn, *Model categories and their localizations*, Mathematical Surveys and Monographs volume 99, AMS (2003).
- [16] Mark Hovey, *Model categories*, Mathematical Surveys and Monographs volume 63, AMS (1999).
- [17] Mark W. Johnson, Donald Yau, *On homotopy invariance for algebras over colored PROPs*, Journal of Homotopy and Related Structures 4 (2009), 275-315.
- [18] J-L. Loday, *Generalized bialgebras and triples of operads*, Astérisque No. 320, SMF (2008).
- [19] J-L. Loday, B. Vallette, *Algebraic Operads*, Grundlehren der mathematischen Wissenschaften, Volume 346, Springer-Verlag (2012).
- [20] J. Lurie, *Higher Topos Theory*, Annals of Mathematics Studies 170, Princeton University Press, 2009.
- [21] S. MacLane, *Categorical algebra*, Bull. Amer. Math. Soc. Volume 71, Number 1 (1965), 40-106.
- [22] M. Markl, *Homotopy algebras are homotopy algebras*, Forum Math. 16 (2004), 129-160.
- [23] C. Rezk, *Spaces of algebra structures and cohomology of operads*, PhD Thesis, Massachusetts Institute of Technology (1996).
- [24] C. Rezk, *A model for the homotopy theory of homotopy theory*, Trans. Amer. Math. Soc., 353 (2001), 973-1007.
- [25] B. Toen, *Vers une axiomatisation de la théorie des catégories supérieures*, K-Theory 34 (2005), no. 3, 233-263.
- [26] S. Yalin, *Classifying spaces and moduli spaces of algebras over a prop*, 2012, arxiv.org/abs/1207.2964 (submitted).

LABORATOIRE PAUL PAINLEVÉ, UNIVERSITÉ DE LILLE 1, CITÉ SCIENTIFIQUE, 59655 VILLENEUVE D'ASCQ CEDEX, FRANCE

E-mail address: Sinan.Yalin@ed.univ-lille1.fr