



HAL
open science

The Moutard transformation and two-dimensional multi-point delta-type potentials

Roman Novikov, Iskander Taimanov

► **To cite this version:**

Roman Novikov, Iskander Taimanov. The Moutard transformation and two-dimensional multi-point delta-type potentials. *Russian Mathematical Surveys*, 2013, 68 (5), pp.957-959. hal-00846512

HAL Id: hal-00846512

<https://hal.science/hal-00846512v1>

Submitted on 19 Jul 2013

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The Moutard transformation and two-dimensional multi-point delta-type potentials

R.G. Novikov ^{*} I.A. Taimanov [†]

Let H be a two-dimensional Schrödinger operator $H = -\Delta + U = -4\bar{\partial}\partial + U$, where $\partial = \frac{1}{2} \left(\frac{\partial}{\partial x} - i \frac{\partial}{\partial y} \right)$, $x, y \in \mathbb{R}$, and let ω be a formal solution to the equation

$$H\omega = 0. \tag{1}$$

The Moutard transformation corresponds to H and ω the operator

$$\tilde{H} = -4\bar{\partial}\partial + \tilde{U} = -4\bar{\partial}\partial + U - 8\bar{\partial}\partial \log \omega \tag{2}$$

such that for every φ meeting the equation $H\varphi = 0$ a function θ satisfying the system

$$(\omega\theta)_z = -i\omega^2 \left(\frac{\varphi}{\omega} \right)_z, \quad (\omega\theta)_{\bar{z}} = i\omega^2 \left(\frac{\varphi}{\omega} \right)_{\bar{z}}, \tag{3}$$

satisfies $\tilde{H}\theta = 0$. The function θ is defined modulo $\frac{1}{\omega}$ due to the integration constant in the right-hand sides of (3).

Recently the Moutard transformation which originates in the surface theory was used for constructing special types of two-dimensional potentials and blowing up solutions of the Novikov–Veselov equation [1, 2].

In difference with [1] which concerns with regular potentials in the present note we deal with multi-point delta-type potentials. We consider also the Faddeev eigenfunctions [3] of the corresponding operators H on the zero energy level. These eigenfunctions are defined by conditions

$$H\psi = 0, \quad \psi(z, \bar{z}, \lambda) = e^{\lambda z} (1 + o(1)) \quad \text{as } z \rightarrow \infty, \quad \lambda \in \mathbb{C} \setminus \{0\}.$$

In addition,

$$\psi = e^{\lambda z} \left(1 + \frac{a(\lambda, \bar{\lambda})}{z} + e^{\bar{\lambda}\bar{z} - \lambda z} \frac{b(\lambda, \bar{\lambda})}{\bar{z}} + o\left(\frac{1}{|z|}\right) \right) \quad \text{as } z \rightarrow \infty,$$

where a, b are the Faddeev generalized "scattering" data on the zero energy level.

^{*}CNRS (UMR 7641), Centre de Mathématiques Appliquées, Ecole Polytechnique, 91128 Palaiseau, France, and Faculty of Control and Applied Mathematics, MIPT, 141700 Dolgoprudny, Russia; e-mail: novikov@cmap.polytechnique.fr

[†]Sobolev Institute of Mathematics, 630090 Novosibirsk, Russia; e-mail: taimanov@math.nsc.ru

Theorem 1 *A formal application of the Moutard transformation to the zero potential $U = 0$ by using a polynomial in z function $\omega = P(z) = \prod_{k=1}^N (z - z_k)$ leads to the multi-point delta-type potential*

$$\tilde{U}(z) = -8\pi \sum_{k=1}^N \delta(z - z_k).$$

For this potential the Faddeev eigenfunctions on the zero energy level take the form

$$\psi = e^{\lambda z} \left(1 + \frac{2}{P} \sum_{k=1}^N \frac{(-1)^k P^{(k)}(z)}{\lambda^k} \right), \quad (4)$$

where $P^{(k)}(z) = \partial^k P(z)$. In addition, for these eigenfunctions $a = -2N/\lambda$ and $b \equiv 0$.

The proof of this theorem is based on solving system (3) with respect to $\psi = \theta$ for $\omega = P(z)$ and $\varphi = ie^{\lambda z}$, and on straightforward computations. However we need to clarify the meaning of Schrödinger operators with such potentials.

Actually in this case we consider the Moutard system (3) with $\omega = P(z)$ as the appropriate regularization of the Schrödinger equation $\tilde{H}\theta = 0$ with the potential \tilde{U} of Theorem 1. In addition, for $N = 1$, the Schrödinger equation $(-\Delta + \tilde{U})\psi = 0$ with \tilde{U} and ψ from Theorem 1 is formally fulfilled under the following conventions:

$$\bar{\partial} \left(e^{\lambda z} \left(\frac{1}{z} \right)^2 \right) = e^{\lambda z} \frac{2}{z} \bar{\partial} \left(\frac{1}{z} \right) = \frac{2\pi e^{\lambda z} \delta(z)}{z}.$$

We remark that the functions ψ of (4) essentially differ from the Faddeev eigenfunctions found in [4, 5] for the Schrödinger operators with multi-point delta-type potentials. The reason is that in [4, 5] the operator with such a potential is replaced by its regularization going back to [6], whereas in the present note we work formally with the original potentials considering the regularization, of the equation $\tilde{H}\theta = 0$, given by the Moutard system (3).

In addition, in view of the property $b \equiv 0$ for ψ of (4) the potentials of theorem 1 may be considered as "reflectionless" in the sense of the Faddeev generalized "scattering" data a, b . In this sense the functions ψ of (4) are similar to the Faddeev eigenfunctions found in [9] for some regular potentials.

In [1, 8] the Moutard transformation is extended to a transformation of solutions of the Novikov–Veselov equation [7]

$$U_t = \partial^3 U + \bar{\partial}^3 U + 3\partial(UV) + 3\bar{\partial}(U\bar{V}) = 0, \quad -4\bar{\partial}V = \partial U. \quad (5)$$

This equation has the Manakov form $H_t = HA + BH$ where A and B are differential operators. If U satisfies (5) and ω meets (1) and the equation

$$(\partial_t + A)\omega = 0, \quad (6)$$

then the extended Moutard transformation of U has the same form (2) and gives a new solution of (5). For the zero potential $U = V = 0$, we have $A = \partial^3 + \bar{\partial}^3$ and $\omega(z, t) = P(z, t) = \prod_{k=1}^N (z - z_k(t))$ satisfies (6) if and only if

$$\frac{\partial P}{\partial t} = \frac{\partial^3 P}{\partial z^3}.$$

The latter equation describes an algebraic dynamics of the zeroes of $P(z, t)$ (such a dynamics for another reason was considered in [1]). A formal application of the extended Moutard transformation leads to the potential

$$\tilde{U}(z, t) = -8\pi \sum_{k=1}^N \delta(z - z_k(t))$$

which apparently may be considered as a formal solution to (5).

Acknowledgement. This work was done during the visit of one of the authors (I.A.T.) to Centre de Mathématique Appliquées of Ecole Polytechnique in July, 2013. The work was partially supported by TFP No 14.A18.21.0866 of Ministry of Education and Sciences of Russian Federation (R.G.N.) and by the program “Fundamental problems of nonlinear dynamics” of the Presidium of RAS and the grant 1431/GF of Ministry of Education and Science of Republic of Kazakhstan (I.A.T.).

References

- [1] Taimanov, I.A., and Tsarev, S.P.: Two-dimensional rational solitons and their blow-up via the Moutard transformation. *Theoret. and Math. Phys.* **157** (2008), 1525-1541.
- [2] Taimanov, I.A., and Tsarev, S.P.: Blowing up solutions of the Novikov-Veselov equation. *Doklady Math.* **77** (2008), 467-468.
- [3] Faddeev, L.D.: Growing solutions of the Schrödinger equation. *Soviet Phys. Dokl.* **10** (1965), 1033–1035.
- [4] Grinevich, P.G., and Novikov, R.G.: Faddeev eigenfunctions for point potentials in two dimensions. *Physics Letters A* **376** (2012), 1102–1106.
- [5] Grinevich, P.G., and Novikov, R.G.: Faddeev eigenfunctions for multi-point potentials. arXiv:1211.0292.
- [6] Berezin, F.A., and Faddeev, L.D.: Remark on Schrödinger equation with singular potential. *Soviet Mathematics* **2** (1961), 372–375.
- [7] Novikov, S.P., and Veselov, A.P.: Finite-zone, two-dimensional potential Schrödinger operators. Explicit formulas and evolution equations. *Soviet Math. Dokl.* **30** (1984), 588–591.

- [8] Hu Heng-Chun, Lou Sen-Yue, and Liu Qing-Ping: Darboux transformation and variable separation approach: the Nizhnik-Novikov-Veselov equation. *Chinese Phys. Lett.* **20** (2003), 1413–1415.
- [9] Taimanov, I.A., and Tsarev, S.P.: Faddeev eigenfunctions for two-dimensional Schrödinger operators via the Moutard transformation. [arXiv:1208.4556](https://arxiv.org/abs/1208.4556).