



HAL
open science

Multi-horizon solar radiation forecasting for Mediterranean locations using time series models

Cyril Voyant, Christophe Paoli, Marc Muselli, Marie Laure Nivet

► **To cite this version:**

Cyril Voyant, Christophe Paoli, Marc Muselli, Marie Laure Nivet. Multi-horizon solar radiation forecasting for Mediterranean locations using time series models. *Renewable and Sustainable Energy Reviews*, 2013. hal-00846823

HAL Id: hal-00846823

<https://hal.science/hal-00846823>

Submitted on 21 Jul 2013

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Multi-horizon solar radiation forecasting for Mediterranean locations using time series models

Cyril Voyant^{1,2*}, Christophe Paoli¹, Marc Muselli¹, Marie-Laure Nivet¹

¹ University of Corsica, CNRS UMR SPE 6134, 20250 Corte, France

² Castelluccio Hospital, Radiotherapy Unit, BP 85, 20177 Ajaccio, France

*Corresponding author, tel/fax +33(0)4 952 936 66/937 97, voyant@univ-corse.fr

Abstract.

Considering the grid manager's point of view, needs in terms of prediction of intermittent energy like the photovoltaic resource can be distinguished according to the considered horizon: following days ($d+1$, $d+2$ and $d+3$), next day by hourly step ($h+24$), next hour ($h+1$) and next few minutes ($m+5$ e.g.). Through this work, we have identified methodologies using time series models for the prediction horizon of global radiation and photovoltaic power. What we present here is a comparison of different predictors developed and tested to propose a hierarchy. For horizons $d+1$ and $h+1$, without advanced ad hoc time series pre-processing (stationarity) we find it is not easy to differentiate between autoregressive moving average (*ARMA*) and multilayer perceptron (*MLP*). However we observed that using exogenous variables improves significantly the results for *MLP*. We have shown that the *MLP* were more adapted for horizons $h+24$ and $m+5$. In summary, our results are complementary and improve the existing prediction techniques with innovative tools: stationarity, numerical weather prediction combination, *MLP* and *ARMA* hybridization, multivariate analysis, time index, etc.

Keywords: Time series, artificial neural networks, stationarity, autoregressive moving average, prediction, global radiation, hybrid model.

1. Introduction

There are lots of alternatives to greenhouse gas emissions generated by fuels combustion [1,2]. It is particularly the case of photovoltaic (*PV*) and wind energy sources, which one of the main advantages is the renewable and inexhaustible aspects and the main disadvantages are related to their intermittencies. This variability is related to winter/summer transition, to day/night transition and to the opacity of atmosphere [3,4]. To overcome these problems, which can be prohibitive, three solutions can be envisaged: split and better distribute the total available power, predict the resource to manage the transition between different energies sources and store the energy excess to redistribute it at the right time [5,6]. This paper deals only with the second solution: the forecasting of the renewable energy sources. The optimization and the management of energy system are really a challenging issue especially when there are insufficient renewable energies to meet the demand. It is essential to anticipate the global radiation decrease (or increase) for an ideal transition. Several methods have been developed by experts around the world and can be divided in two main groups: (i) methods using mathematical formalism of Times Series (*TS*), (ii) numerical weather prediction (*NWP*) model and weather satellite imagery. The technique used depends on considered source, and on their startup delay (from five minutes to 1 hour). Note that for an ideal management it is appreciable to know the eventual fluctuations one or two days ahead. These temporal characteristics define the horizon of the prediction to consider. According to the horizons of these methods are more effective compared to others [8]. Considering the grid manager's point of view, needs in terms of prediction can be distinguished according to the considered horizon: the resource that will be available on the following days ($d+1$, $d+2$ et $d+3$), the next day by hourly step ($h+24$), during the next hour ($h+1$), and in the five next minutes ($m+5$). These horizons allow understanding the various aspects of the prediction: the medium term, the short term and the very short term. The $d+1$ and $d+2$ predictions are important for the manager because they have immediate industrial applications and economic impacts especially in the case of small and relatively isolated electric grids. Indeed, in this case it is essential to organize and anticipate the fossil stocks. Concerning the $h+1$ horizon, it corresponds more or less to the ignition delay of the thermal system. In fact, starting a heat engine takes about 30 minutes; the manager must be able to predict the intermittent energy cuts at least 1 hour in advance. Concerning the $h+24$ prediction, its interest combines the two precedents. The knowledge 24 hours in advance of the renewable energy enables better inventory management concerning fossil fuel, and an anticipation of the critical moments where the grid manager must be vigilant. Finally, the few minutes horizon concerns for example the means of production related to hydroelectric power plants and to gas turbines. Indeed, just a few minutes are necessary to electricity to be available in these cases. We can also note that short term forecasting (now-casting) can be very useful to control indoor climate in buildings with automation system. Thus it seemed interesting to compare different methods based upon the analysis of historical *TS*

of global radiation for several horizons: $d+1$, $h+1$, $h+24$ and $m+5$. In this paper we propose, horizon by horizon, a classification of predictor tested on various Mediterranean towns. Our goal is to provide robust predictors with the most generic approach possible.

In the next, the time series forecasting models proposed in the literature are first reviewed. In section 3 we will detail the methodologies of prediction we have tested, taking care to explain the *TS* formalism dedicated to the global solar radiation modeling and the need to make it stationary (time series pre-processing). Then we will expose the result of comparison between modeling and measure in the daily case, hourly case and five minutes case. Finally we will close the paper with a comparison of the results against those of the literature, emphasizing the link between predictor performance and type of horizon.

2. Review on time series forecasting models

In this section, we present a review of the literature on time series forecasting models for global radiation. Optimal use of renewable energy requires a good characterization and good predictive potential for sized detectors or estimate the potential energy power plants [9,10]. There are a lot of models allowing *TS* predictions. It is possible to list them into four groups [11,12]:

- naive models are essential to verify the relevance of complex models. Include persistence, average or the k -nearest neighbors (k -NN) [13-16];
- conditional probability models are rarely mentioned in the literature regarding global radiation. Include Markov chains and predictions based on Bayesian inference [17-22];
- reference models based on the family of autoregressive moving average, *ARMA* [23,24];
- connectionist models (artificial neural network) and more particularly the Multi-Layer Perceptron (*MLP*) which is the artificial neural network architecture the most often used [25-27].

The following deals with the two last groups: *ARMA* and neural network models. Indeed *ARMA* is the most classical and popular for time series modeling and artificial neural network seems to be the best alternative to conventional approaches. As climate of the earth is dominated by non-linear processes, *ANN* by its non-linear nature is effective to predict cloudy days and so solar radiation. Concerning the prediction of solar radiation, we can cite works of Mellit [26,27] in which it is possible to find a synthesis of the coupling of *MLP* with global radiation. In addition to these works, there are others related to the prediction of weather data such as solar radiation [28-35]. Neural networks have been studied on many sites and researchers have shown the ability of these techniques

to accurately predict the time series of meteorological data [32]. Table 1 presents several representative examples of the use of artificial neural networks (ANN) methods applied to the modeling or prediction of solar radiation and PV energy in the 2000s. For the years prior to 2000, the interested reader may also refer to the article Mellit [26]. For all the articles presented in Table 1 we can see that the errors associated with predictions (monthly, daily, hourly and minute) are between 5% and 10%. However we see that the MLP can be used with exogenous parameters or coupled with other predictors (Markov, Wavelet, etc.). In the Mellit and Kalogirou article review [26], we find that 79% of Artificial Intelligence (AI) methods used in weather prediction data are based on a connectionist approach (ANN). We can also cite the use of fuzzy logic (5%), Adaptive neuro fuzzy inference system (ANFIS) (5%), networks coupling wavelet decomposition and ANN (8%) and mix ANN/Markov chain (3%). In sum, the use of ANN, especially the MLP represents a large majority of research works. This is the most commonly used technique. Other methods are used only sporadically.

authors	topic	location	horizon	error	conclusions
[58] Almonacid(2010)	Estimation of PV energy	Spain (Jaèn)	monthly	$MAPE = 7.3 \%$	<i>MLP</i> better than reference models (bilinear interpolation method and Blaesser's method)
[31] Behrang et al. (2010)	Global radiation modeling with different <i>ANN</i>	Iran (Dezful)	d+1	$MAPE = 5.2 \%$	<i>MLP</i> with exogenous inputs is very efficient (8 models compared)
[56] Benghanem and mellit(2010)	Global radiation modeling with avec <i>RBF, MLP</i> and standard regression	Saudi Arabia (Al-madinah)	d+1	$R^2=0.98$	<i>RBF</i> is the most efficient, moreover the approach is validated on <i>PV</i> system (8 models are compared)
[27] Mellit et Pavan(2010)	Global radiation forecasting at horizon with <i>ANN</i>	Italy (Trieste)	h+24	$R^2>94 \%$	<i>MLP</i> validated on <i>PV</i> wall (no other compared predictors)
[59] Azadeh et al. (2009)	Global radiation modeling with <i>ANN</i>	Iran (6 cities)	monthly	Accuracy = 94 % (error = 6 %)	<i>MLP</i> better than Angström model
[57] Chaabene and Ben Ammar(2008)	Global radiation prediction with hybrid <i>MLP</i> with fuzzy logic, <i>ARMA</i> and Kalman filters	Tunisia (Energy and Thermal Research Centre)	d+1 m+5	$nMBE= -9.11 \%$ $nRMSE< 10 \%$	Dynamic predictions are considered coupling <i>ARMA</i> , Kalman filter and neuro-fuzzy estimators
[60] Jiang (2008)	Diffuse radiation prediction with <i>MLP</i>	China (8 cities)	monthly	Accuracy = 95 %	The methodology is validated on the entire Chinese territory (compared to two empirical models)
[52] Mubiru and Banda (2008)	Global radiation modeling with different <i>MLP</i>	Uganda (4 sites)	d+1	$RMSE = 107 \text{ Wh/m}^2$	<i>MLP</i> better than 5 empirical models
[53] Bosch et al. (2008)	Global radiation modeling	Spain (13 sites)	d+1	$nRMSE = 7.5 \%$	The <i>MLP</i> can be used in the mountainous area. the error is acceptable (no comparison with other methods)
[55] Elminir et al. (2007)	Prediction of diffuse radiation with <i>MLP</i>	Egypt (3 stations)	h+1 d+1	Standard error = 4.2 % Standard error = 9 %	<i>MLP</i> better than 2 linear regressions models
[61] Mellit et al. (2006)	Prediction of global radiation with <i>MLP</i> and wavelets	Algeria (36°43' N; 3°2' E)	d+1	$MAPE< 6 \%$	Method validated for the <i>PV</i> output and various meteorological data. More than 7 models are compared (<i>AR</i> , <i>ARMA</i> , <i>MTM</i> , <i>MLP</i> , <i>RBFN</i> , Wavelet networks, etc.)
[62] Cao and Cao (2005)	Prediction of global radiation with recurrent <i>MLP</i> and wavelets	China (Shanghai)	d+1	$nRMSE = 8 \%$ (with wavelet) and 35 % without wavelet	Wavelet decomposition improves the prediction
[63] Mellit et al. (2005)	Global radiation modeling with <i>MLP</i> and Markov approach	Algeria (4 sites)	d+1	$nRMSE = 8 \%$	<i>MLP</i> better than <i>AR</i> , <i>ARMA</i> and Markov chains
[64] Sozen et al. (2004)	Global radiation modeling with <i>MLP</i>	Turkey (17 stations)	d+1	$MAPE< 7 \%$	Training and test areas are relocated, <i>MLP</i> is robust. The comparison is done with classical regression models
[65] Reddy and Manish (2003)	Global radiation modeling with <i>MLP</i>	India (2 stations)	h+1	$MAPE = 4 \%$	<i>MLP</i> better than 3 classical regression models
[36] Sfetos and Coonick (2000)	Global radiation forecasting with <i>MLP</i>	Corsica (41.55°N, 8.48°E)	h+1	$RMSE = 27.6 \text{ W/m}^2$	Multivariate <i>MLP</i> modeling improves the prediction. 13 Models are tested (<i>ARMA</i> , <i>RNFN</i> , <i>ANFIS</i> , etc.)

Table 1: representative examples of the use of *ANNs* method applied to the modeling or prediction of solar radiation and *PV* energy from 2000s

Also in this literature review [26], the results of different researches considering a lot of places, were compared. The prediction error (MAPE in this case) of monthly global radiation induced by the use of an ANN is estimated between 0.2% and 10.1% depending on the city and the architecture considered (median = 4%). The results presented are so disparate they seem incomparable. However, we must consider that in some locations the cloud occurrences are minimal while others are subject to much less forgiving climates. Concerning the global radiation, Sfetsos [36] has showed that neural networks generated an error of 7% and ARMA methodologies, an error of 8%. Behranget al. [37] have compiled a list of the prediction error with neural networks for global radiation. For identical locations, the errors can double or even triple. The conclusions on the MLP can be generalized to other predictors. According to the literature, the parameters that influence the prediction are manifold, so it is difficult to use the results from other studies. Considering this fact, it may be interesting to test methods or parameters even though they have not necessarily been proven in other studies. Based on the foregoing, all parameters inherent to the MLP or ARMA method must be studied for each tested site.

After literature review and considering the difficulty to make definite conclusion we wanted to study estimators which are little or very rarely studied in the renewable energy field. Thus, we tried a prediction methodology based on Bayesian inferences. There are many works on the coupling with other predictors such as neural networks [38,39] or as discriminant test for variables selection [40]. However, this technique is widely used in econometrics, through very theoretical publications cannot really compare with other prediction methods. We can especially mention Xiang Fei [41], which showed that the Bayesian inferences allow an estimate equal to autoregressive (AR) model with non-stationary variables. The error in the studied series is close to 10% for both models. Concerning Markov chains, they are rarely used in energy, according to the paper of Hoacaoglu [42] there is a prediction error of 6% for daily radiation and for Muselliet al. [43] an error on the PV predicted energy on horizontal surface equal to 10%. Based on these results, we chose to incorporate this type of predictor in our study. The other three studied estimators are persistence, k-NN and average which are easy to implement. Indeed, there is no learning phase, and few constraints are needed to use them (stationarity, pretreatment, assumptions, etc.). Although advanced methods provide better results, we think it is important to keep in mind the balance between model complexity and quality of prediction. For this reason, it is necessary to compare the sophisticated models against "naïve" models [4,15,44,45]. According to the references listed above, the following remarks can be made:

- ANN and ARMA models seem to be the most popular time series predictors;
- it is very difficult to compare or evaluate predictors because many of them looks like to be site and horizon dependant;

- there is no convention dealing with errors estimation tools (e.g. seasonal errors best for certain days), neither than with data test selection.

Considering these limitations we propose for each considered horizon a homogeneous experimental protocol.

3. Materials and methods

The methodology used in this work is based on time series forecasting. A Time Series (*TS*) is intuitively defined as an ordered sequence of past values of the variable that we are trying to predict [24]. Thus, the current value at t of the *TS* x is noted x_t , where t , the time index, is between 1 and n , with n is the total number of observations. We call h the number of values to predict. The prediction of time series from $(n+1)$ to $(n+h)$, knowing the historic from x_1 to x_n , is called the prediction horizon (horizon 1, ..., horizon h). For the horizon 1 (the simplest case), the general formalism of the prediction will be represented by Equation 1 where ϵ represents the error between the prediction and the measurement, f_n the model to estimate and t time index taking the $(n-p)$ following values: $n, n-1, \dots, p+1, p$. Where n is the number of observations and p the number of model parameters (it is assumed that $n \gg p$). [44,45]

$$x_{t+1} = f_n(x_t, x_{t-1}, \dots, x_{t-p+1}) + \epsilon(t+1) \quad \text{Equation 1}$$

Studies in finance and econometrics have yielded many models more or less sophisticated. Some of these models have been applied in the case of the prediction of global solar radiation. To estimate the f_n model, a stationarity hypothesis is often necessary. This result originally shown for *ARMA* methods [23,24] can be also applicable for the study and prediction with neural network [46,47]. We can also note that few authors suggest that periodic nature of a time series can also be captured from the *AI* models like *MLP*, very often with the inclusion of a time indicator [36]. However we have considered that in practice, the input data must be stationary to use an *MLP*. In previous works [44,45], we have developed sophisticated methods to make the global radiation time series stationary. We have demonstrated that the use of the clear sky index (*CSI*) obtained with Solis model [48] is the more reliable in Mediterranean places. As the seasonality is often not completely erased after this operation, we use a method of seasonal adjustments (seasonal variance corrected by periodic coefficients) based on the moving average [24] (CSI^*). The chosen method is essentially interesting for the case of a deterministic nature of the series seasonality (true for the global radiation series) but not for the stochastic seasonality [23]. It is also possible to use a variant of *CSI*, considering only the radiation outside the atmosphere, we obtain in this way the clearness index (k) [49] and k^* with the previous method of seasonal corrections.

Considering the limitations described at the end of the section 2, we decided to establish a homogeneous experimental protocol for each considered horizon. Thereby, for all horizons studied ($d+1$, $h+1$, $h+24$ and $m+5$), we have compared *ARMA* and *MLP* predictors against at least one naive predictor (e.g. persistence). We focused our work on a general methodology for estimating the prediction error:

- test of prediction over a long period, not on "well chosen" days;
- use of *RMSE* to penalize large deviations [50];
- normalization of *RMSE* for comparison on many sites:

$$nRMSE = \sqrt{E[(\hat{x} - x)^2] / \langle x^2 \rangle} \quad \text{Equation 2}$$

- no cumulative predictions except for specific studies which has the effect of average the error and decrease it;
- distribution of errors according to seasons because the energy consumption is not the same throughout the year;
- tests on several locations, in order to avoid phenomena regional climates;
- use of a naive predictor as a reference for prediction to evaluate the proposed methodology (balance between model complexity and quality of prediction);
- use of confidence interval to define margin of error, as e.g. the classical IC95%, in order to provide information on the prediction robustness.

For *ARMA* and *MLP* methods, we have studied the impact of stationary process for the indexes *CSI*, k and relative seasonal adjustments (CSI^* and k^*). Concerning *MLP*, we studied the contribution of exogenous meteorological data (multivariate method) at different time lags and data issued from a numerical weather prediction model (*NWP*). The confidence interval has been calculated after at least six training simulations. We also studied the performance of a hybrid *ARMA/ANN* model from a rule based on the analysis of hourly data series. Finally we evaluated for each method the error estimation for annual and seasonal periods: Winter, Spring, Summer and Autumn. It should be noted that due to the difficulty to obtain data, the protocol could not be followed homogeneously for all data. The following section presents the results and for each horizon in chronological order.

4. Results

Data used in the experiments are related to the French meteorological organization database. As manipulations on horizons proceed, this database was expanded iteratively. Our goal is to provide robust and predictive methodology as generic as possible, avoiding falling into the specifics of a place. The non-homogeneity strict of manipulations is due to this typical construction. In fact it is very difficult to obtain quality data. At the beginning there was not much data available and after first experiments it seemed to be interesting to test our method on a larger sample. The table below lists for each horizon all manipulations performed and the data associated.

Horizon	Manipulations performed			Data associated
	Predictor used	Stationary method	Variable selection	
d+1	Mean, persistence, <i>SARIMA</i> , Bayesian inference, Markov chains, k-NN, <i>ANN</i>	<i>CSI</i> , k, <i>CSI*</i> , k*	<i>PACF</i> , cross correlation	Ajaccio (1971:1989) and Bastia/Ajaccio (1998:2007)
h+1	Mean, persistence, <i>ARIMA</i> , <i>ANN</i>	<i>CSI</i> , k	<i>PACF</i> , cross correlation, linear regression	Ajaccio/ Bastia/ Marseille/ Montpellier/ Nice (1998:2007)
h+24	Persistence, <i>ARMA</i> , <i>ANN</i>	<i>CSI</i> , k	<i>PACF</i> , cross correlation	Ajaccio (1999:2008)
m+5	Persistence, <i>ARMA</i> , <i>ANN</i>	<i>CSI</i> , k	<i>PACF</i> , cross correlation	Ajaccio (2009,2010)

Table 2: list of manipulations performed and data associated with each horizon.

For the most complete horizon (hourly case), the data used to test models are from 5 coastal cities located in the Mediterranean area and near mountains: Montpellier (43°4'N / 3°5'E, 2 m alt), Nice (43°4'N / 7°1'E, 2 m alt), Marseille (43°2'N / 5°2'E, 5 m alt), Bastia (42°3'N / 9°3'E, 10 m alt) and Ajaccio (41°5'N / 8°5'E, 4 m alt). The available data are global radiation, pressure (P , Pa; average and daily gradient¹, measured by numerical barometer during 1 hour), nebulosity (N , Octas), ambient temperature (T , °C; maximum, minimum, average and night², measured done during an half hour), wind speed (Ws , m/s; average at 10 meters, measured during the 10 last minutes of the half hourly step), peak wind speed (PKW , m/s; maximum speed of wind at 10 meters, measured during 30 minutes), wind direction (Wd , deg at 10 meters measured during an half hour), sunshine duration (Su , h, computed with the global radiation series and the power threshold 120 W/m²), relative humidity (RH , % instantaneous measure at the end of the half-hour) and rain precipitations (RP , mm, 5 cumulative measures of 6 minutes during the half-hour). The data are transposed into hourly values by Météo-France organization.

4.1. Daily case

¹ Difference between the mean pressure of day j and day $j-1$

² Measured at 3:00 AM

As the knowledge of the available solar energy for the next days allows fossil energy provision and interconnection energy management, daily horizon is very important. For this horizon and for all studied models, the years 1971-1987 are the basis of learning and the two years from 1988 to 1989 are dedicated to the test of the prediction. With this horizon, the method based on average, Markov chains, k-NN and Bayesian inferences are tested. For all this methodology the results are equivalent, the error ($nRMSE$) is close to 25.5% (from 25.1 for Markov chains to 26.13 for the persistence). Without stationarization and exogenous inputs, the two predictors $ARMA$ and MLP are more efficient than other methods; the errors of prediction are smaller than 22% and relatively close. The MLP is noted as: $(Endo^{N_e}) \times N_h \times I$ where N_e is the number of endogenous nodes and N_h the number of hidden neurons. For this first study, where only endogenous data are considered, these two predictors are equivalent and outperform other approaches. If now we make the TS stationary by using k^* or CSI^* and seasonal adjustments (k^* and CSI^*) we note that the error of prediction decreases. The best results are related to the k^* and CSI^* pretreatments and are shown in the Table 3. With these methodologies the errors are reduced by 1.5 points ($nRMSE = 20.2\%$ for k^* and $nRMSE = 20.3\%$ for CSI^*). Indeed, it is necessary to adapt the models and architectures to the new dynamics of the signal. The optimization leads to use the model $ARMA(2,2)$, while for the MLP configuration remains unchanged.

	Raw data	Statio k^*	Statio CSI^*
$ARMA$	$21.18 \pm 0\%$ $AR(8)$	$20.31 \pm 0\%$ $ARMA(2.2)$	$20.32 \pm 0\%$ $ARMA(2.2)$
PMC	$20.97 \pm 0.15\%$ $Endo^{1-8} \times 3 \times 1$	$20.17 \pm 0.1\%$ $Endo^{1-8} \times 3 \times 1$	$20.25 \pm 0.1\%$ $Endo^{1-8} \times 3 \times 1$

Table 3: prediction errors for $ARMA$ and MLP ($nRMSE \pm IC95\%$). Predictions done for years 1988 and 1989.

For more details on results of other methods (persistence, Bayesian, KNN, etc.), the reader can refer to our previous work [15,44]. Again, the MLP and $ARMA$ methods appear to be equivalent for $d+1$. Indeed, with or without the use of seasonal adjustments, it is very difficult to prioritize them. It seems, in the particular case that we just examined, that MLP based results are also convincing than $ARMA$ based results. Regarding the comparison between the two stationary methodologies (k^* and CSI^*), it is not possible to conclude, averages are not significantly different. However, make stationary the TS improves the prediction error both for $ARMA$ and MLP .

Once finished these first experimentations, we decided to explore the multivariate option. In order to increase the confidence degree of our conclusions we choose to make our test considering two locations: Ajaccio and Bastia (where forecasting is considered to be more difficult). Indeed one of the particularities of the MLP use is based on the possibility to do multivariate regressions. The use of the exogenous data should better model the phenomena. The MLP is noted as: $(Endo^{N_e} E^{Me}) \times N_h \times I$ where N_e and N_h are the numbers of endogenous and exogenous nodes. For Ajaccio for example the better

model of *MLP* with exogenous data is $(Endo^2 Su^1 N^1)_{x3x1}$ while for Bastia it is $(Endo^4 Su^1 RH^1 N^1)_{x3x1}$. As the errors are respectively $21.5\% \pm 0.05\%$ and $25.4 \pm 0.2\%$, we can deduce that the generated error is location-dependent. In addition, we have shown that the use of exogenous variables improved the *MLP* prediction mainly during winter and autumn (gain of 0.7 point). Similar results are obtained with the *PV* energy forecasting [44].

The main conclusions for this d+1 horizon can be resumed as following:

- without the use of exogenous variables, *MLP* are equivalent to *ARMA* ($nRMSE \sim 22\%$);
- for cloudy months (winter and autumn), the use of exogenous variables improves the quality of the prediction (gain of 0.7 point);
- make the *TS* stationary with k^* , or if possible CSI^* is appropriate (gain of 1.5 points);
- persistence is an interesting naive predictor, which gives very good results in spring and summer ($nRMSE = 26.1\%$);
- the prediction methodology is applicable in the global radiation case and *PV* energy case.

This first study on the daily horizon allows us understanding how to use the *MLP* and other predictors studied. We showed that tested predictors like Markov, Bayes and k -NN are relatively equal in terms of prediction. The details of this comparison are given in [44]. These predictors proved to be much less suitable for predicting global radiation as *ARMA* or *MLP*. With this result, we decide in the following to not use the Markov, Bayes and k -NN estimators. For the naive estimator, only the persistence will be used for its ease of use and good results, especially on sunny days ($nRMSE = 19\%$ in May and June).

4.2. Hourly case

For this horizon the CSI^* approach simplifies the *MLP* architecture: one endogenous input and a maximum of 8 hidden neurons for the five *TS* studied. But this does not improve the prediction error, so in the following, the stationary model will not use the periodic coefficients. Performing the same study in the case of *ARMA* predictions, CSI^* and CSI stationarization give similar results. Henceforth, we will therefore use the CSI with these predictors. Note that the clearness index generates less efficient results [45]. The Table 4 presents the comparison of seasonal $nRMSE$ related to estimators for global radiation for the five cities. For predictions with *MLP*, we study the case with only the endogenous variables (*MLP*_{endo}) and the combining of endogenous and exogenous variables (*MLP*_{endo-exo}).

Models	Anual	Winter	Spring	Summer	Autum
Persistence	36.0±2.0	46.4±5.1	36.3±3.4	30.0±3.2	41.5±3.3
<i>ARMA</i>	16.4±0.7	22.2±1.8	15.9±1.2	14.1±0.9	19.4±0.6
<i>MLP</i> endo	17.0±1.2	20.3±2.0	17.0±2.3	15.6±1.2	18.1±1.1
<i>MLP</i> endo-exo	16.8±1.3	20.2±2.1	16.9±2.2	15.5±1.4	17.5±1.4

Table4: performance comparison ($nRMSE$ and confidence interval in %) between different studied models (average on the five cities). Bold characters represent the lowest values

In summer, the interest of methods like *MLP*endo and *MLP*endo-exo is minimal. This is undoubtedly due to the low probability of occurrence of clouds during this period. A linear process like *ARMA* seems best suited. We can probably conclude that the use of *MLP* with endogenous and exogenous variables is interesting when the cloud cover is intense (mainly in autumn and winter). In [45] we have shown that the predictor hybridation (*ARMA* and *MLP*endo-exo) increases the quality of predictions. The method used is based on the following selection rule:

$$\text{if } |\varepsilon^{AR}(t)| \leq |\varepsilon^{PMC}(t)| \text{ then } \hat{X}(t+1) = \hat{X}^{AR}(t+1) \text{ else } \hat{X}(t+1) = \hat{X}^{PMC}(t+1)$$

Equation 3

The Figure 1 shows the average gain (computed on the five cities) of $nRMSE$ obtained by the hybrid method compared to the better *MLP* (grey bars) and the better *ARMA* (dashed bars). The gain is positive when the hybridization is better than traditional methods.




Figure 1: mean gain related to the hybrid model compared to the models *MLP* (grey bars) and *ARMA* (dashed bars)

The maximum gain is observed in winter ($3.8 \pm 0.8\%$ better than the *ARMA* model) and the minimum is in summer, when the hybrid method is as interesting as the *ARMA* method (gain of $0.02 \pm$

0.5%). For all sites, it is clear that the hybrid model approximates correctly the global radiation [45]. In previous study [45] we have shown that exogenous data (meteorological measures) can be replaced by estimation of analytic models like the numerical weather prediction model ALADIN [45]. In this context, the results generated by hybrid *MLP/ARMA*, ALADIN and *CSI** should be different (see the table 5).

	<i>ANNual</i>	<i>Winter</i>	<i>Spring</i>	<i>Summer</i>	<i>Autumn</i>
Ajaccio	14.9(25.1)	19.4(34.7)	15.5(25.2)	11.0(21.4)	17.0(33.9)
Bastia	16.5(27.1)	19.5(35.0)	17.5(27.1)	13.2(22.6)	17.9(34.4)
Montpellier	14.7(26.9)	15.7(32.6)	15.2(25.9)	13.4(24.6)	15.5(33.2)
Marseille	13.4(25.3)	16.6(32.9)	14.8(25.3)	9.3(20.0)	13.8(32.3)
Nice	15.3(26.4)	16.6(32.1)	15.3(24.5)	10.3(21.1)	26.2(37.1)

Table 5: prediction error (*nRMSE* %) for the hybrid model *ARMA*, *MLP*, *ALADIN*, *CSI**, the persistence results are presented between parenthesis.

This hybrid model is very interesting: the 10% threshold has been crossed in Marseille. Although summer is the season where the hybrid methodology is the less interesting, all seasons and cities benefit from this hybridization model. We can note that *MLP* and *ARMA* are very effective alone in summer period. To resume, use of the hybrid method reduces the error by 11% compared to the prediction done by persistence (mean on the five cities).

In summary, the fact to make stationary the global radiation *TS* reduces the error by $0.5 \pm 0.1\%$ for the five locations studied. The use of ALADIN and of hybridization models shows a real potential and a strong interest. This step allows to increase significantly the quality of the prediction (gain close to 3.5 points). In the end, if we compare this approach with a simple prediction such as persistence, there is a reduction of the prediction error of more than 11%.

The methodology of prediction based on *CSI*, ALADIN *MLP* and *ARMA* is certainly complicated to implement, but gives results far superior to those from the other tested techniques. We note that for this horizon, the *CSI* must be used to overcome seasonal variations. In addition, the use of exogenous variables is an added value to the modeling. Forecasts of meteorological variables from ALADIN model offer prediction accuracy. However, the use of meteorological measurements gives also good results, although less efficient. Finally, the combination of all the improvements that were recently proposed amplifies the quality of the prediction.

4.3.24-hours ahead case

This new horizon studied is the prediction for the next day hour by hour [10,51] of the global radiation profile. Unlike hourly, daily or monthly horizons, this horizon is little discussed in the

literature. We may mention the work of Mellit and Pavan [27] which propose to use as input of the prediction tool (*MLP*) the daily mean values of solar radiation and temperature, and the day of the considered month. To satisfy this prediction horizon, we have considered approaches based on the use of *MLP*, following conclusions presented earlier in this paper. As a first step, we focus on the endogenous case, and then we will introduce exogenous parameters. The predictor is a *MLP* like in the previous case, but with multiple outputs (one by hours). Measurements are chronologically positioned in the input vector of *MLP*. We choose to compare the *MLP* results with those obtained by methods of persistence and *ARMA*. The last method we have tested is based on multiple *ARMA* models which each are dedicated to one particular hour. Note that all these methods are compatible with the use of the clearness index (k and k^*) and the clear sky index (*CSI* and CSI^*). Moreover, in $h+1$ and $d+1$ horizons, the seasonal adjustments did not show strong superiority. For these reasons, in the next manipulations only k and *CSI* will be considered. The goal is to find a relatively simple and generalizable methodology taken care of not draw conclusions about data snooping. Results are shown in the table 6.

Type		<i>ANN</i> ual	Winter	Spring	Summer	Autumn
Persistence		35.1	54.8	35.2	28.0	40.4
<i>ARMA</i>	k	29.1	44.6	29.2	24.0	33.2
	<i>CSI</i>	28.6	44.2	28.6	23.1	32.8
<i>MLP</i>	k	27.9	44.2	27.9	22.2	32.7
	<i>CSI</i>	27.8	42.8	28.4	22.0	31.3

Table 6: $nRMSE(\%)$ of predictions realized with the *MLP*. Bold characters represent the best results.

We note that sophisticated approaches as *ARMA* or *MLP* largely outperform naive models especially in winter. Note also that the best predictions are obtained with the use of the clear sky index (*CSI*). Contrary to the previous case ($h+1$ case), the *MLP* is systematically better than *ARMA* model. The interest of a hybrid approach seems for this reason not relevant. However, it is possible to integrate exogenous inputs. After several trials, we found that the more interesting data are the hourly pressure and cloudiness of the last day, and the daily average nebulosity of the two last days. The contribution of these variables is presented in the Table 7 (only the *CSI* methodology is shown because more interesting).

Type		<i>ANN</i> ual	Winter	Spring	Summer	Autumn
Persistence		35.1	54.8	35.2	28.0	40.4
<i>ARMA</i>		28.6	44.2	28.6	23.1	32.8
<i>MLP</i> endo		27.8	42.8	28.4	22.0	31.3
<i>MLP</i> exo		27.3	42.4	27.8	21.7	31.3

Table7: impact of exogenous variables on the prediction quality ($nRMSE$ %). In bold the best results.

For the h+24 horizon the contribution of exogenous variables is less explicit than for previous case studied. These kind of deep horizons (≥ 24 h) modify approach to consider. Thus, this type of prediction is particularly difficult to realize. Search the smoothness of a 24-hours ahead prediction depends on too many parameters to expect to get the same level of results as for horizons $h+1$ or $d+1$. We can conclude that it is valuable to make stationary data ($nRMSE$ gain close to 0.5 point). To do this the use of clear sky index is preferable, even if the clearness index gives results almost similar. The *CSI* allows an $nRMSE$ gain of 0.5 point for *ARMA* and 0.1 point for *MLP* related to the k use. The classical approach involving a single *MLP* with multiple outputs is recommended: $nRMSE$ gain of 0.6 point for k index and 0.4 point for *CSI* index related to a *MLP* committee like described in the *ARMA* case. In the present state of our knowledge, the ratio between performance and complexity induces, to not use exogenous variables (maximal $nRMSE$ gain of 0.6 point in Winter).

4.4. Five minutes case

By its nature this prediction horizon is completely different from what we have studied so far. The originality of this case is the sampling frequency of measurement that is less than the dynamics of cloud occurrence. Thus, in 5 minutes the sky has a high probability of remain identical. Data are available on the PV wall of Vignola laboratory [44]. They cover the period from March 2009 to September 2010. The installation allows identifying three separate areas: 0° , 45° SE and 45° SW tilted at 80° relative to the ground surface.

<i>Orientation / Type</i>		Total	May	June	July	August
<i>SW</i>	<i>MLP</i>	21.4	31.4	20.7	14.2	19.5
	<i>MLP + k</i>	22.5	32.3	20.1	15.4	19.6
	<i>MLP + CSI</i>	22.2	31.9	21.1	16.3	20.0
	<i>Persistence</i>	21.8	32.3	20.9	14.4	19.6
<i>S</i>	<i>MLP</i>	20.2	28.0	22.6	13.5	16.5
	<i>MLP + k</i>	21.7	29.6	23.7	14.8	18.4
	<i>MLP + CSI</i>	21.9	29.7	25.5	17.4	19.5
	<i>Persistence</i>	20.8	28.8	23.2	13.8	17.1
<i>SE</i>	<i>MLP</i>	23.2	31.8	26.5	14.6	20.6
	<i>MLP + k</i>	24.2	32.6	27.6	15.1	21.7
	<i>MLP + CSI</i>	25.6	33.3	28.1	17.8	23.8
	<i>Persistence</i>	24.5	33.3	27.9	14.8	22.0

Table8: Stationary process impact on the error of prediction ($nRMSE$ in %)

The Table 8 shows the impact of the stationary process. Unlike in the daily and hourly case this study does not allow concluding that the use of CSI and k are justified. For this tilt and orientation, the theoretical models are limited. In these configurations the solar shield complicated the phenomena. For this reason, CSI, k, CSI^* and k^* are not used in the following (only raw data).

In fact, in the raw global radiation TS , output of MLP corresponds to an improved persistence. As the prediction seems to be a persistence (delay of 5 min), weights related to the first lag are important and other are close to zero.

Simpler tools, accessible with MLP could improve the prediction results. Indeed, the MLP can alone choose its own stationarity, using as input time indexes, which will enable it to establish a regression on the time of the periodic phenomenon. The two time indices used are related to the hour of the day and day of the year. The transfer function in the hidden layer which gives the best results is the Gaussian function. The use of time index generates an added value to the quality of the prediction. Results are systematically improved by this tool: $nRMSE$ is reduced by 0.7 point for the SW and S orientations and 0.1% in the SE case. The average gain is greater than 0.5 point, ensuring a real advantage in using this stationarization mode. Table 9 shows the results obtained.

	<i>MLP</i>	<i>MLP</i> +time index	persistence
SE	23.2	23.1	24.5
S	20.2	19.5	20.8
SW	21.4	20.7	21.8

Table 9: prediction error ($nRMSE$ in %) related to the MLP and the time index methodology

Note that for this horizon, the use of $ARMA$ is not relevant because the optimization led us to use a simple $AR(1)$ where the regression coefficient of lag 1 is close to 1. This kind of model is in fact persistence. Like MLP is systematically better than persistence, the hybridization of models is not justified. Moreover, the use of exogenous data does not provide benefit for the prediction. Furthermore there are very few measurements with a sampling near 5 minutes. This kind of prediction process is very complicated to construct. In brief, we have seen in this section that methods used to make stationary the TS are not available for this horizon ($nRMSE$ increased by 1 point). It is more appropriate to use the raw series and not the clear sky or clearness index, but the use of time index is interesting to take into account the seasonality. We may also note that the MLP -based methodology improves outcomes ($nRMSE$ improved to more than 1 point) compared to a simpler approach based on persistence.

5. Conclusion

In all bibliographic items related to the estimation of global radiation, we find that the errors associated with predictions (monthly, daily, hourly and minute) differ from sites and from authors. Methodologies of predictions are usually so different that they are difficult to compare. In addition, the estimation errors are heterogeneous: prediction error on certain days or sampled over an extended period, test on the cumulative predictions, use of non-standard error parameters, etc. To overcome all these features we present here a methodology of comparison of different predictors developed and tested to propose a hierarchy. Only the *TS* approach is studied, other weather models using numerical weather prediction models or satellite images are not considered. For horizons $d+1$ and $h+1$, our results are partly consistent with the literature. Indeed, *MLP* are adapted and used to make predictions of global radiation with an acceptable error [52] and are also applicable to mountainous areas [53]. Regarding prioritization of *ARMA* and *MLP*, the results shown here are different from traditional bibliographic results [26,54,55]. In fact, without stationarity we do not think it is easy to differentiate between *ARMA* and *MLP*. Moreover, while *ANN* by its non-linear nature is effective to predict cloudy days, *ARMA* techniques are more dedicated to sunny days without cloud occurrences. However, we agree with Berhanghet al. [37] with the fact that the use of exogenous variables improves the results of *MLP*. As in the literature, we found that the relevant approaches in the case of the prediction of radiation were equally in the case of the prediction of *PV* power [26,56]. Although it is not routinely used in the literature, we believe that persistence can correctly judge the validity of complex technical and we chose as naive predictor. In literature, clear sky model and seasonal adjustments based on periodic coefficients have not often been used with the prediction of global radiation. The views of the results presented here, their investigation looks promising. Finally, for horizons $h+24$ and $m+5$, there are still too few studies using the *MLP*. However, as Mellit and Pavan [27] and Chaabene and Ben Ammar [57] we believe and have shown that the *MLP* were adapted to these situations. In addition, our approach with the use of time index appears to be efficient. In summary, our results are complementary and improve the existing prediction techniques with innovative tools (stationarity, *NWP* combination, *MLP* and *ARMA* hybridization, multivariate analysis, time index, etc.).

Through this work, we have identified some methodologies for the prediction horizon of global radiation. We can conclude that these two types of predictions are relatively equal in the methodology to implement. In Table 10 are listed and summarized *TS* based methods we recommend for different prediction horizons.

Horizons	stationarity	Exogenous data	Required predictors	difficulty	<i>nRMSE</i>
----------	--------------	----------------	---------------------	------------	--------------

d+1	CSI*	Measures: <i>Su.N.RH</i>	<i>MLP</i> (<i>>ARMA>pers</i>)	++	23.4%
h+1	CSI	NWP: <i>N. P. RP</i>	<i>Hybrid MLP+ARMA</i> (<i>>MLP>ARMA>pers</i>)	+++	14.9%
h+24	<i>k</i>	-	<i>MLP multi-outputs</i> (<i>>multiMLP>ARMA>pers</i>)	+	27.3%
m+5	<i>Time index</i>	-	<i>MLP</i> (<i>>ARMA>pers</i>)	+	20.2%

Table10: summary of the results presented in this paper

In view of the previous manipulations, we note that the results can be completely different depending on the time horizon. For this reason, we must pay attention to the methods used and the expected results. What should be sought is a simple method to implement, cost effective and workable in several locations: the selection of data and model parameters must be chosen parsimoniously. To conclude this paper, we believe that the establishment of a benchmark in the areas of renewable energy would allow the community to better share, understand and interpret the results: same data, comparisons of models using the same tools RMSE, *nRMSE*, IC95%, etc. The recent European COST (Cooperation in Science and Technology) initiative called WIRE (Weather Intelligence for Renewable Energies)³ seems to follow this idea and should be encouraged.

Acknowledgement

Thanks to an agreement with Météo-France, which is the French meteorological organization, we had the opportunity to freely access to some of their forecasts and measures.

³ http://www.cost.eu/domains_actions/essem/Actions/ES1002

References

1. Schlink U, Herbarth O, Richter M, Dorling S, Nunnari G, Cawley G, et al. Statistical models to assess the health effects and to forecast ground-level ozone. *Environmental Modelling & Software* 2006;21:547–58.
2. Jenn Jiang H. Policy review of greenhouse gas emission reduction in Taiwan. *Renewable and Sustainable Energy Reviews* 2011;15(2):1392–402.
3. Benghanem M, Mellit A, Alamri SN. ANN-based modelling and estimation of daily global solar radiation data: A case study. *Energy Conversion and Management*. 2009;50(7):1644–55.
4. Voyant C, Muselli M, Paoli C, Nivet M-L. Optimization of an artificial neural network dedicated to the multivariate forecasting of daily global radiation. *Energy* 2011;36(1):348–59.
5. Carl-Jochen W. Hydrogen energy — Abundant, efficient, clean: A debate over the energy-system-of-change. *International Journal of Hydrogen Energy* 2009;34:1–52.
6. Darras C, Muselli M, Poggi P, Voyant C, Hogue JC, Montignac F. PV output power fluctuations smoothing: The MYRTE platform experience. *International Journal of Hydrogen Energy* 2012;37:14015-14025.
7. Do Sacramento EM, Sales AD, de Lima LC, Veziroglu TN. A solar–wind hydrogen energy system for the Ceará state – Brazil. *International Journal of Hydrogen Energy*. 2008;33(20):5304–11.
8. Tamer K, Azah M, Sopian K. A review of solar energy modeling techniques. *Renewable and Sustainable Energy Reviews* 2012;16: 2864-2869.,
9. Cao JC, Cao SH. Study of forecasting solar irradiance using neural networks with preprocessing sample data by wavelet analysis. *Energy* 31;2006:3435-3445.
10. Chaouachi A, Kamel RM, Ichikawa R, Hayashi H, Nagasaka K. Neural Network Ensemble-Based Solar Power Generation Short-Term Forecasting. *World Academy of Science, Engineering and Technology* 54;2009..
11. Brockwell Peter J, Richard A. Davis. *Time series: theory and methods*. 2nd ed. New York: Springer-Verlag, 1991.
12. Faraday J, C Chatfield. *Times Series Forecasting with Neural Networks: A Case Study*. Applied Statistics, 1998.
13. Yakowitz S. Nearest neighbors method for time series analysis, *Journal of Time Series Analysis* 1987;8:235-247.

14. Armstrong JS (Ed.). Principles of forecasting: a handbook for researchers and practitioners. Boston; London, Kluwer Academic, 2001.
15. Paoli, C, Voyant C, Muselli M, Nivet ML. Solar radiation forecasting using ad-hoc time series preprocessing and neural networks. Springer-Verlag, 2009:898:907.
16. Sharif M, Burn D. Simulating climate change scenarios using an improved K-nearest neighbor model, Journal of Hydrology 2006;325:179-196.
17. Celeux G, Nakache J. Analyse discriminante sur variables qualitatives, Polytechnicaproceeding, Paris, France, 1994.
18. Diday E, Lemaire J, Pouget J, Testu F. Éléments d'analyse de données, Dunod, 1982.
19. Muselli M., Poggi P, Notton G, Louche A. First order Markov chain model for generating synthetic “typical days” series of global irradiation in order to design photovoltaic stand alone systems, Energ. Convers & Manage 2011;42:675-687.
20. Logofet D. The mathematics of Markov models: what Markov chains can really predict in forest successions, Ecological Modeling 2000;126:285-298.
21. Sharif M, Burn D. Simulating climate change scenarios using an improved K-nearest neighbor model, Journal of Hydrology 2006;325:179-196.
22. Yakowitz S. Nearest neighbors method for time series analysis, Journal of Time Series Analysis 1987;8:235-247.
23. Hamilton J. Time series analysis, Princeton University Press, 1994.
24. Bourbonnais R , M. Terraza, Analyse des séries temporelles. Application à l 'Economie et à la Gestion, 2 éd., Dunod, Paris, 2008.
25. Ahmed O, Driss Z, Hanane D, Rachid B. Artificial neural network analysis of Moroccan solar potential, Renewable and Sustainable Energy Reviews 2012;16:4876-4889.
26. Mellit AM, Kalogirou S, Hontoria L, Shaari S. Artificial intelligence techniques for sizing photovoltaic systems: A review, Renewable and Sustainable Energy Reviews 2009;13:406-419.
27. Mellit AM, Pavan A. 24-h forecast of solar irradiance using artificial neural network: Application for performance prediction of a grid-connected PV plant at Trieste, Italy, Sol. Energ 2010;8::807-821.
28. Hocaoglu F, Gerek O, Kurban M. Hourly solar radiation forecasting using optimal coefficient 2-D linear filters and feed-forward neural networks. Solar Energy 2008;82:714-726.

29. Fariba B, Dehghan AA, Faghieh AR. Empirical models for estimating global solar radiation: A review and case study. *Renewable and Sustainable Energy Reviews* 2013;21: 798-821.
30. Kalogirou S. Artificial neural networks in renewable energy systems applications: a review. *Renewable and Sustainable Energy Reviews* 2001;5:373-401.
31. Kemmoku Y. Daily insolation forecasting using a multi-stage neural network. *Solar Energy* 1999;66:193-199.
32. Hontoria L, Aguilera J, Zufiria P. Generation of hourly irradiation synthetic series using the neural network multilayer perceptron. *Solar Energy* 2002;72:441-446.
33. Negnevitsky M. An Expert System Application for Clearing Overloads. *International Journal of Power and Energy Systems* 1995;15:9-13.
34. Al-Alawi SM, Al-Hinai HA. An RNA-based approach for predicting global radiation in locations with no direct measurement instrumentation. *Renewable Energy* 1998;14:199-204.
35. Mohandes M, Rehman S, Halawani TO. Estimation of global solar radiation using artificial neural networks. *Renewable Energy* 1998;14:179–184.
36. Sfetsos A, Coonick AH. Univariate and multivariate forecasting of hourly solar radiation with artificial intelligence techniques. *Solar Energy* 2000;68:169-178.
37. Behrang MA, Assareh E, Ghanbarzadeh A, Noghrehabadi AR. The potential of different artificial neural network (ANN) techniques in daily global solar radiation modeling based on meteorological data. *Solar Energy* 2010;84:1468-1480.
38. Lauret P, Fock E, Randrianarivony RN, Manicom-Ramsamy JF. Bayesian neural network approach to short time load forecasting. *Energy Conversion and Management* 2008;49:1156-1166.
39. Bruneau P, Boudet L. « Bayesian Variable Selection in Neural Networks for Short-Term Meteorological Prediction ». In *Neural Information Processing*, édité par Tingwen Huang, Zhigang Zeng, Chuandong Li, et Chi Sing Leung, 289-296. *Lecture Notes in Computer Science* Springer Berlin Heidelberg 2012;7666:289-296.
40. Riviere C, Lauret P, Ramsamy JF, Page Y. A Bayesian neural network approach to estimating the energy equivalent speed. *Accident Analysis & Prevention* 2006;38:248–259.
41. Fei, Xiang, Chung-Cheng Lu, Ke Liu. A bayesian dynamic linear model approach for real-time short-term freeway travel time prediction. *Transportation Research Part C, Emerging Technologies* 2011;19:1306-1318.

42. Naveen S, Siddhartha V. Stochastic techniques used for optimization in solar systems: A review. *Renewable and Sustainable Energy Reviews*2012;16:1399-1411.
43. Muselli, M, P. Poggi, G. Notton, et A. Louche. First order Markov chain model for generating synthetic “typical days” series of global irradiation in order to design photovoltaic stand alone systems. *Energy Conversion and Management* 2001;42:675-687.
44. Paoli C, Voyant C, Muselli M, Nivet ML. Forecasting of preprocessed daily solar radiation time series using neural networks. *Solar Energy* 2010;84:2146-2160.
45. Voyant, Cyril, Marc Muselli, Christophe Paoli, et Marie-Laure Nivet. Numerical weather prediction (NWP) and hybrid *ARMA/ANN* model to predict global radiation. *Energy* 2012;39::341-355.
46. Kim, Tae Yoon, KyongJoo Oh, Chiho Kim, et Jong Doo Do. Artificial neural networks for non-stationary time series. *Neurocomputing*2004;61:439-447.
47. Zhang G. Peter, Min Qi. Neural network forecasting for seasonal and trend time series. *European Journal of Operational Research* 2005;160:501-514.
48. Mueller RW, Dagestad KF, Ineichen P, Schroedter-Homscheidt M, Cros S, Dumortier D, R. Kuhlemann R, et al. Rethinking satellite-based solar irradiance modelling: The SOLIS clear-sky module. *Remote Sensing of Environment* 2004;91:160-174.
49. Badescu V. Modeling solar radiation at the earth’s surface: recent advances. Springer, 2008.
50. Voyant, C. Prédiction de séries temporelles de rayonnement solaire global et de production d’énergie photovoltaïque à partir de réseaux de neurones artificiels. Thesis, Université de Corse (2011).
51. Cococcioni, M, D’Andrea E, Lazzerini B. 24-hour-ahead forecasting of energy production in solar *PV* systems. In 2011 11th International Conference on Intelligent Systems Design and Applications (ISDA) 2011:1276 -1281.
52. Meita R, Abudureyimu A, Nagasaka K. Mapping of solar energy potential in Indonesia using artificial neural network and geographical information system. *Renewable and Sustainable Energy Reviews*2012;16:1437-1449.
53. Bosch J, Lopez G, Batlles F. Daily solar irradiation estimation over a mountainous area using artificial neural networks. *Renewable Energy* 2008;33:1622-1628.
54. Reddy K. Solar resource estimation using artificial neural networks and comparison with other correlation models. *Energy Conversion and Management* 2003;44:2519-2530.

55. Amit KY, Chandel SS. Tilt angle optimization to maximize incident solar radiation: A review. *Renewable and Sustainable Energy Reviews* 2013;23:503-513.
56. Benghanem, Mohamed, Adel Mellit. Radial Basis Function Network-based prediction of global solar radiation data: Application for sizing of a stand-alone photovoltaic system at Al-Madinah, Saudi Arabia. *Energy* 2010;35:3751-3762.
57. Chaabene M, Benammar M. Neuro-fuzzy dynamic model with Kalman filter to forecast irradiance and temperature for solar energy systems. *Renewable Energy* 2008;33:1435-1443.
58. Almonacid F, Rus C, Hontoria L, Muñoz FJ. Characterisation of PV CIS module by artificial neural networks. A comparative study with other methods. *Renewable Energy* 2010;35:973-980.
59. A. Azadeh, A. Maghsoudi, et S. Sohrabkhani, An integrated artificial neural networks approach for predicting global radiation. *Energy Conversion and Management* 2009;50:1497-1505.
60. Jiang Y. Prediction of monthly mean daily diffuse solar radiation using artificial neural networks and comparison with other empirical models. *Energy Policy* 2008;36:3833-3837.
61. Mellit A, Benghanem A, Kalogirou SA. An adaptive wavelet-network model for forecasting daily total solar-radiation. *Applied Energy* 2006;83:705-722.
62. Cao S, Cao J. Forecast of solar irradiance using recurrent neural networks combined with wavelet analysis. *Applied Thermal Engineering* 2005;25:161-172.
63. Mellit A, Benghanem M, Arab AH, Guessoum A. A simplified model for generating sequences of global solar radiation data for isolated sites: Using artificial neural network and a library of Markov transition matrices approach. *Solar Energy* 2005;79:469-482.
64. Sözen A, Arcaklioğlu E, Özalp M. Estimation of solar potential in Turkey by artificial neural networks using meteorological and geographical data. *Energy Conversion and Management* 2004;45:3033-3052.
65. Reddy K, Ranjan M. Solar resource estimation using artificial neural networks and comparison with other correlation models. *Energy Conversion and Management* 2003;44:2519-2530.

List of Tables

Table 1: representative examples of the use of *ANNs* method applied to the modeling or prediction of solar radiation and *PV* energy from

Table 2: list of manipulations performed and data associated with each horizon

Table 3: prediction error for *ARMA* and *MLP* ($nRMSE \pm IC95\%$). Predictions done for the year 1988 and 1889 for the Ajaccio case.

Table 4: performance comparison ($nRMSE$ in %) between different studied models (average on the five cities). Bold characters represent the lowest values

Table 5: prediction error ($nRMSE$ en %) for the hybrid model *ARMA*, *PMC*, *ALADIN*, *CSI**. between parenthesis are presented the persistence results

Table 6: $nRMSE(\%)$ of predictions realized with the *MLP*. Bots characters represent the best results.

Table 7: providing of exogenous variables on the prediction quality. In bold the best $nRMSE$

Table 8: stationary impact on the error of prediction ($nRMSE$ in %)

Table 9: prediction error ($nRMSE$ in %) related to the *MLP* and the time index methodology

Table 10: summary of the better forecasting methodologies and in parentheses the ranking of the *MLP*, *ARMA* and persistence approaches

List of Figure

Figure 1: mean gain related to the hybrid model compared to the models *MLP* (grey bars) and *ARMA* (dashed bars)