



HAL
open science

Higher preprojective algebras and stably Calabi-Yau properties

Claire Amiot, Steffen Oppermann

► **To cite this version:**

Claire Amiot, Steffen Oppermann. Higher preprojective algebras and stably Calabi-Yau properties. 2013. hal-00847027v1

HAL Id: hal-00847027

<https://hal.science/hal-00847027v1>

Submitted on 22 Jul 2013 (v1), last revised 18 Apr 2014 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

HIGHER PREPROJECTIVE ALGEBRAS AND STABLY CALABI-YAU PROPERTIES

CLAIRE AMIOT AND STEFFEN OPPERMANN

ABSTRACT. In this paper, we give sufficient properties for a finite dimensional graded algebra to be a higher preprojective algebra. These properties are of homological nature, they use Gorensteiness and bimodule isomorphisms in the stable category of Cohen-Macaulay modules. We prove that these properties are also necessary for 3-preprojective algebras using [Kel11] and for preprojective algebras of higher representation finite algebras using [Dug12].

CONTENTS

1. Introduction	2
Plan of the paper	3
Notation	3
Acknowledgement	3
2. Preliminaries	3
2.1. Graded algebras and graded modules	3
2.2. Graded Cohen-Macaulay modules	4
2.3. Stably graded Calabi-Yau algebras	6
2.4. Higher preprojective algebras	7
3. Homological characterization of finite dimensional preprojective algebras	7
3.1. Main result and strategy of the proof	7
3.2. The global dimension of Λ	8
3.3. Proof of Theorem 3.1	9
4. The Gorenstein dimension	12
4.1. Gorenstein dimension and cluster tilting subcategories	12
4.2. The selfinjective case	15
5. Bimodule Calabi-Yau properties of preprojective algebras	15
5.1. Classical preprojective algebras	16
5.2. The case $d = 3$	17
5.3. The $(d - 1)$ -representation finite case	20
References	21

Date: July 22, 2013.

Most work on this project was done during visits of the authors to each others universities, funded respectively by Norwegian Research Council project 196600/V30 and the GDR Théorie de Lie algébrique et géométrique. The first author is partially supported by the ANR project ANR-09-BLAN-0039-02.

Key words and phrases. Cohen-Macaulay modules, stable categories, Calabi-Yau categories, preprojective algebras, Calabi-Yau algebras.

2010 *Mathematics Subject Classification.* 16E05, 16E35, 16E65, 16G50, 18E30.

1. INTRODUCTION

Preprojective algebras play an important role in many different parts of mathematics. Such an algebra is associated to a quiver Q without oriented cycles. It has been defined by Gelfand and Ponomarev in the 70's to get a better understanding of the representation theory of the path algebra of the quiver Q . Recently, in the context of higher Auslander-Reiten theory, Iyama generalized the definition of preprojective algebras. If Λ is a finite dimensional algebra of global dimension $d - 1$, its d -preprojective algebra $\Pi_d(\Lambda)$ is defined as the tensor algebra $T_\Lambda \text{Ext}_{\Lambda^e}^{d-1}(\Lambda, \Lambda^e)$ where Λ^e is the enveloping algebra $\Lambda \otimes_k \Lambda^{\text{op}}$. It is naturally a positively graded algebra.

In this paper, we are interested in the properties that characterize finite dimensional preprojective algebras. For $d = 2$, the preprojective algebra $\Pi = \Pi_2(kQ)$ is finite dimensional if and only if Q is a Dynkin quiver, and, in that case, by a classical result, the algebra Π is selfinjective and there is a functorial isomorphism

$$\text{Ext}_{\Pi}^1(X, Y) \cong D \text{Ext}_{\Pi}^1(Y, X)$$

for any $X, Y \in \text{mod } \Pi$. So in other words, the triangulated category $\underline{\text{mod}} \Pi$ is 2-Calabi-Yau. The duality above comes from an isomorphism

$$\text{Hom}_{\Pi^e}(\Pi, \Pi^e) \cong \Omega_{\Pi^e}^3 \Pi(1)$$

in the stable category of graded bimodules $\underline{\text{gr}}\text{-}\Pi^e$ (where $\Pi(1)$ is the graded bimodule Π shifted by 1). This isomorphism can also be written

$$\mathbf{R}\text{Hom}_{\Pi^e}(\Pi, \Pi^e)[3] \cong \Pi(1) \quad \text{in } D^b(\underline{\text{gr}}\text{-}\Pi^e)/\text{gr-perf } \Pi^e$$

using the triangle equivalence $\underline{\text{gr}}\text{-}\Pi^e \cong D^b(\underline{\text{gr}}\text{-}\Pi^e)/\text{gr-perf } \Pi^e$. The main result of this paper is the following.

Theorem 1.1 (Theorem 3.1). *Let $\Pi = \bigoplus_{i \geq 0} \Pi_i$ be a finite dimensional graded algebra satisfying the following properties:*

- (a) $\text{pdim } D\Pi = \text{idim } \Pi \leq d - 2$, that is Π is of Gorenstein dimension $\leq d - 2$;
- (b) $\mathbf{R}\text{Hom}_{\Pi^e}(\Pi, \Pi^e)[d + 1] \cong \Pi(1)$ in $D^b(\underline{\text{gr}}\text{-}\Pi^e)/\text{gr-perf } \Pi^e$;
- (c) $\text{Ext}_{\underline{\text{gr}}\text{-}\Pi^e}^j(\Pi, \Pi^e(i)) = 0$ for all $i < 0$ and $j > 0$.

Then Π is isomorphic as a graded algebra to $\Pi_d(\Lambda)$ for some algebra Λ of global dimension $d - 1$.

Note that the property (b) implies that the stable category of maximal Cohen-Macaulay modules over Π is a d -Calabi-Yau triangulated category.

In a second part of the paper, we show that in certain situations also the converse of the above theorem holds. We prove that properties (a), (b) and (c) are satisfied by the finite dimensional preprojective algebras for $d = 2$ and $d = 3$ (Theorems 5.1 and 5.7). Moreover, using the results of Dugas in [Dug12], we prove that properties (a), (b) and (c) holds for selfinjective d -preprojective algebras for any d . More precisely we prove the following.

Theorem 1.2 (Theorems 4.8 and 5.9). *The map $\Lambda \mapsto \Pi_d(\Lambda)$ gives a one to one correspondence between $(d-1)$ -representation finite algebras Λ and finite dimensional selfinjective graded algebras Π satisfying $\mathbf{R}\text{Hom}_{\Pi^e}(\Pi, \Pi^e)[d + 1] \cong \Pi(1)$ in $D^b(\underline{\text{gr}}\text{-}\Pi^e)/\text{gr-perf } \Pi^e$.*

This result is very similar to Theorem 4.35 of [HIO12] that asserts that the preprojective construction gives a one to one correspondence between $(d-1)$ representation-infinite algebras Λ and homologically smooth algebras Π satisfying $\mathbf{R}\mathrm{Hom}_{\Pi^e}(\Pi, \Pi^e)[d] \cong \Pi(1)$ in $D^b(\mathrm{gr}\text{-}\Pi^e)$.

Plan of the paper. The paper is organized as follows. We start in Section 2 with preliminaries on graded Gorenstein algebras, and Cohen-Macaulay modules, and define the notion of bimodule stably (1)-twisted d -Calabi-Yau algebras. The main result of the paper is proved in Section 3. Section 4 gives an interpretation of the Gorenstein dimension of an d -preprojective algebra $\Pi_d(\Lambda)$ in term of some Hom-vanishing in the category $D^b(\mathrm{mod}\ \Lambda)$. In Section 5, we prove that the converse of Theorem 3.1 is true for $d = 2$, $d = 3$, and for preprojective algebras of $(d-1)$ -representation finite algebras.

Notation. All algebras in this paper are finite dimensional algebras over a field k . For an algebra A , we denote by A^e the tensor algebra $A \otimes A^{\mathrm{op}}$. The dual $\mathrm{Hom}_k(A, k)$ of A is denoted by DA . When nothing else is stated explicitly, tensor products are over the field k .

Acknowledgement. We thank Alex Dugas for helpful hints about his article [Dug12].

2. PRELIMINARIES

2.1. Graded algebras and graded modules. Let $A = \bigoplus_{n \in \mathbb{N}} A_n$ be a positively graded algebra. For a graded A -module $M = \bigoplus_{n \in \mathbb{Z}} M_n$, and for any $p \in \mathbb{Z}$, we denote by $M(p)$ the graded module $\bigoplus_{n \in \mathbb{Z}} M_{n+p}$, that is the degree n part of $M(p)$ is M_{p+n} . We denote by $\mathrm{gr}\text{-}A$ the category of finitely generated graded A -modules. Morphisms in $\mathrm{gr}\text{-}A$ are graded morphisms homogeneous of degree 0. The category $\mathrm{gr}\text{-}A$ is an abelian Krull-Schmidt category.

By an abuse of notation, for $M \in \mathrm{gr}\text{-}A$ we will denote by $M \in \mathrm{mod}\ A$ its image in $\mathrm{mod}\ A$ under the forgetful functor $\mathrm{gr}\text{-}A \rightarrow \mathrm{mod}\ A$. Note that the A^{op} -module $\mathrm{Hom}_A(M, A)$ has a natural structure of graded A^{op} -module given by $\bigoplus_{p \in \mathbb{Z}} \mathrm{Hom}_{\mathrm{gr}\text{-}A}(M, A(p))$.

We write

$$\begin{aligned} \mathrm{gr}\text{-}\mathrm{proj}_{\leq 0} A &= \mathrm{add}\{A(i) \mid i \geq 0\}, \text{ and} \\ \mathrm{gr}\text{-}\mathrm{proj}_{> 0} A &= \mathrm{add}\{A(i) \mid i < 0\} \end{aligned}$$

for the subcategories of $\mathrm{gr}\text{-}\mathrm{proj}\ A$ of projectives generated in positive, respectively in non-positive, degrees.

For an additive category \mathcal{A} , we denote by $\mathbf{K}^{b,-}(\mathcal{A})$ (resp. $\mathbf{K}^b(\mathcal{A})$) the homotopy category of right bounded (resp. bounded) complexes of objects in \mathcal{A} .

Proposition 2.1. *Let A be a positively graded algebra, such that A_0 has finite global dimension. Then $D^b(\mathrm{gr}\text{-}A) = \mathbf{K}^{b,-}(\mathrm{gr}\text{-}\mathrm{proj}\ A)$ has a semiorthogonal decomposition*

$$\mathbf{K}^{b,-}(\mathrm{gr}\text{-}\mathrm{proj}\ A) = \left\langle \mathbf{K}^b(\mathrm{gr}\text{-}\mathrm{proj}_{\leq 0} A), \mathbf{K}^{b,-}(\mathrm{gr}\text{-}\mathrm{proj}_{> 0} A) \right\rangle.$$

That is, we have $\mathrm{Hom}_{\mathbf{K}^{\mathrm{b},-}(\mathrm{gr}\text{-}\mathrm{proj} A)}(\mathbf{K}^{\mathrm{b}}(\mathrm{gr}\text{-}\mathrm{proj}_{\leq 0} A), \mathbf{K}^{\mathrm{b},-}(\mathrm{gr}\text{-}\mathrm{proj}_{> 0} A)) = 0$, and for $X \in \mathbf{K}^{\mathrm{b},-}(\mathrm{gr}\text{-}\mathrm{proj} A)$ there is a triangle

$$\underbrace{X_{\leq 0}}_{\in \mathbf{K}^{\mathrm{b}}(\mathrm{gr}\text{-}\mathrm{proj}_{\leq 0} A)} \longrightarrow X \longrightarrow \underbrace{X_{> 0}}_{\in \mathbf{K}^{\mathrm{b},-}(\mathrm{gr}\text{-}\mathrm{proj}_{> 0} A)} \longrightarrow$$

Remark 2.2. It follows that the triangle in Proposition 2.1 is functorial in X . We will denote by $(-)>_0$ and $(-)\leq_0$ the functors in the triangle.

This semi-orthogonal decomposition has already been used in [Orl05].

Proof of Proposition 2.1. Since A is positively graded the space $\mathrm{Hom}_{\mathrm{gr}\text{-}A}(A(i), A(j))$ vanishes whenever $j < i$. It follows that

$$\mathrm{Hom}_{\mathrm{gr}\text{-}\mathrm{proj} A}(\mathrm{gr}\text{-}\mathrm{proj}_{\leq 0} A, \mathrm{gr}\text{-}\mathrm{proj}_{> 0} A) = 0,$$

and thus we have the Hom-vanishing of the proposition.

To obtain the triangle, we observe that any graded projective is the direct sum of a graded projective generated in non-positive degree and a graded projective generated in positive degree. Choosing such decompositions for all terms of a right bounded complex X of graded projective A -modules gives rise to a short exact sequence $X_{\leq 0} \twoheadrightarrow X \twoheadrightarrow X_{> 0}$ and thus to a triangle in the homotopy category, where $X_{\leq 0} \in \mathbf{K}^{\mathrm{b},-}(\mathrm{gr}\text{-}\mathrm{proj}_{\leq 0} A)$ and $X_{> 0} \in \mathbf{K}^{\mathrm{b},-}(\mathrm{gr}\text{-}\mathrm{proj}_{> 0} A)$. Finally observe that, since A_0 has finite global dimension, we may assume that only finitely many terms of the complex are not generated in positive degrees, that is $X_{\leq 0}$ is also left bounded. \square

2.2. Graded Cohen-Macaulay modules.

Definition 2.3. An algebra A is said to be *Gorenstein* if its injective dimension (denoted $\mathrm{id} A$) and the projective dimension of its dual (denoted $\mathrm{pd} DA$) are both finite. For such an algebra, the *Gorenstein dimension* of A is the integer $\mathrm{id} A = \mathrm{pd} A$. We define the category of (*maximal*) *Cohen-Macaulay* modules by:

$$\mathrm{CM}(A) := \{X \in \mathrm{mod} A \mid \mathrm{Ext}_A^i(X, A) = 0 \text{ for } i > 0\}$$

If moreover A is a graded algebra we denote by

$$\mathrm{gr}\text{-}\mathrm{CM}(A) := \{X \in \mathrm{gr}\text{-}A \mid \mathrm{Ext}_A^i(X, A) = 0 \text{ for } i > 0\}$$

the category of graded (*maximal*) Cohen-Macaulay A -modules.

The next result is the graded version of a famous triangle equivalence [Buc87, Theorem 4.4.1] (see also [KV87, Ric89]).

Theorem 2.4. *Let A be a graded Gorenstein algebra, then $\mathrm{gr}\text{-}\mathrm{CM}(A)$ is a Frobenius category and there is a triangle equivalence*

$$\mathrm{D}^{\mathrm{b}}(\mathrm{gr}\text{-}A) / \mathrm{gr}\text{-}\mathrm{perf} A \xrightarrow{\sim} \underline{\mathrm{gr}\text{-}\mathrm{CM}(A)}.$$

Because of this equivalence, we will use the notation $\underline{\mathrm{gr}\text{-}\mathrm{CM}(A)}$ for the category $\mathrm{D}^{\mathrm{b}}(\mathrm{gr}\text{-}A) / \mathrm{gr}\text{-}\mathrm{perf}(A)$. That is we may write $M \cong N$ in $\underline{\mathrm{gr}\text{-}\mathrm{CM}(A)}$ even if M and N are not Cohen-Macaulay modules.

The next two lemmas are classical results on maximal Cohen-Macaulay modules that will be used in this paper.

Lemma 2.5. *Let A be a graded Gorenstein algebra of dimension g . For $X \in \text{gr-}A$, its g -th syzygy $\Omega^g(X)$ is Cohen-Macaulay.*

Lemma 2.6. *Let A be a graded Gorenstein ring, M a graded A -module. Then there is a short exact sequence*

$$K \twoheadrightarrow {}_{\text{CM}}M \twoheadrightarrow M$$

with $\text{pd } K < \infty$ and ${}_{\text{CM}}M \in \text{gr-CM } A$.

In this situation ${}_{\text{CM}}M$ is called a Cohen Macaulay replacement of M .

Proof. This fact is well-known, but we include a sketch of the proof for the convenience of the reader.

Sufficiently high syzygies of M are Cohen-Macaulay by Lemma 2.5, so the claim trivially holds for them. Assume we already found the upper sequence in the diagram below.

$$\begin{array}{ccccc}
 K_i & \twoheadrightarrow & {}_{\text{CM}}(\Omega^i M) & \twoheadrightarrow & \Omega^i M \\
 & & \downarrow \text{approx.} & & \downarrow \\
 & & \bar{P}_{i-1} & \dashrightarrow & P_{i-1} \\
 & & \downarrow & & \downarrow \\
 K_{i-1} & \dashrightarrow & {}_{\text{CM}}(\Omega^{i-1} M) & \dashrightarrow & \Omega^{i-1} M
 \end{array}$$

The right vertical sequence is the defining sequence of $\Omega^i M$, and the middle one is obtained by taking a left projective approximation of ${}_{\text{CM}}(\Omega^i M)$ and its cokernel. We obtain an induced map as indicated by the upper dashed arrow, which can be assumed to be split epi (by enlarging \bar{P}_{i-1} if necessary). Then we obtain the desired sequence in the lower row of the diagram. Indeed, K_{i-1} has finite projective dimension since both the kernel of $\bar{P}_{i-1} \twoheadrightarrow P_{i-1}$ and K_i do. \square

Remark 2.7. The proof above also shows that, provided M is generated in degree 0, we may choose ${}_{\text{CM}}M$ to only have projective summands which are generated in degree 0.

Lemma 2.8. *In the setup of Lemma 2.6 the following are equivalent:*

- (1) $\forall j > 0 \forall i < 0: \text{Ext}_{\text{gr-}A}^j(M, A(i)) = 0$, and
- (2) K can be chosen such that it has a projective resolution in $\text{gr-proj}_{\leq 0} A$.

Proof. Since ${}_{\text{CM}}M$ is Cohen-Macaulay, applying the functor $\text{Hom}_{\text{gr-}A}(-, A(i))$ to the short exact sequence, we obtain isomorphisms

$$\text{Ext}_{\text{gr-}A}^j(M, A(i)) \cong \text{Ext}_{\text{gr-}A}^{j-1}(K, A(i)) \quad \text{for all } j > 1$$

and an exact sequence

$$\begin{array}{ccccc}
 \text{Hom}_{\text{gr-}A}(M, A(i)) & \twoheadrightarrow & \text{Hom}_{\text{gr-}A}({}_{\text{CM}}M, A(i)) & \twoheadrightarrow & \\
 \longrightarrow & & \text{Hom}_{\text{gr-}A}(K, A(i)) & \twoheadrightarrow & \text{Ext}_{\text{gr-}A}^1(M, A(i)).
 \end{array}$$

(1) \implies (2): Pick a minimal projective resolution of K

$$0 \longrightarrow P_d \longrightarrow \dots \longrightarrow P_0 \longrightarrow K.$$

Let j be the leftmost position such that P_j is not contained in $\text{gr-proj}_{\leq 0} A$ (assuming such a term exists). It follows that $\text{Ext}_{\text{gr-}A}^j(K, A(i)) \neq 0$ for some $i < 0$. By (1) and the discussion above it follows that $j = 0$. Thus K has a non-zero direct summand in $\text{gr-proj}_{> 0} A$. Since $\text{Ext}_{\text{gr-}A}^1(M, \text{gr-proj}_{> 0} A) = 0$ such a summand can be split off of the short exact sequence.

(2) \implies (1): Since K has a projective resolution in $\text{gr-proj}_{\leq 0} A$, we have $\text{Ext}_{\text{gr-}A}^j(K, A(i)) = 0$ for all $i < 0$ and all j . The claim then immediately follows from the discussion above. \square

2.3. Stably graded Calabi-Yau algebras.

Definition 2.9. A Gorenstein graded algebra A is called *stably* (p) -*twisted* d -*Calabi-Yau* if there is a functorial isomorphism

$$D\underline{\text{Ext}}_{\text{gr-}A}^i(X, Y) \cong \underline{\text{Ext}}_{\text{gr-}A}^{d-i}(Y(p), X)$$

for any $X, Y \in \text{gr-CM} A$ and for any $i \in \mathbb{Z}$.

Definition 2.10. A graded algebra A is called *bimodule stably* (p) -*twisted* d -*Calabi-Yau* if there is an isomorphism

$$\mathbf{R}\text{Hom}_{A^e}(A, A^e)[d+1] \cong A(p) \text{ in } \underline{\text{gr-CM}}(A^e).$$

The integer p is called the Gorenstein parameter.

Remark 2.11. Recall that a bimodule $(d+1)$ -Calabi-Yau algebra is a homologically smooth algebra satisfying

$$\mathbf{R}\text{Hom}_{A^e}(A, A^e)[d+1] \cong A \text{ in } \text{D}^b(\text{gr-}A^e).$$

So the choice of d for the Calabi-Yau dimension in our definition could seem strange. But the reason of our choice is motivated by the following result.

Theorem 2.12. *Let A be a finite dimensional graded Gorenstein algebra which is bimodule stably (p) -twisted d -Calabi-Yau, then A is stably (p) -twisted d -Calabi-Yau.*

Proof. This can be shown using an Auslander-Reiten formula in $\text{gr-CM} A$ as it is done in [Yos90, 3.10] and [IY08, Theorem 8.3] for local isolated singularities. Here we give a different argument using the description of the category $\underline{\text{gr-CM}} A$ as the localisation $\text{D}^b(\text{gr-}A)/\text{gr-perf} A$.

From [Kel08, Lemma 4.1] we know that for any $X, Y \in \text{D}^b(\text{gr-}A)$ there is a functorial isomorphism

$$D \text{Hom}_{\text{D}^b(\text{gr-}A)}(X, Y) \cong \text{Hom}_{\text{D}^b(\text{gr-}A)}(\mathbf{R}\text{Hom}_{A^e}(A, A^e) \overset{\mathbf{L}}{\otimes}_A Y, X).$$

Moreover since the algebra A is Gorenstein, the functor $\mathbf{R}\text{Hom}_{A^e}(A, A^e) \overset{\mathbf{L}}{\otimes}_A -$ sends any perfect complex to a perfect complex. Therefore we can apply [Ami08, Proposition 4.3.1] to deduce that the category $\text{D}^b(\text{gr-}A)/\text{gr-perf} A$ has a Serre functor whose inverse is given by $\mathbf{R}\text{Hom}_{A^e}(A, A^e)[1] \overset{\mathbf{L}}{\otimes}_A -$.

Now for any objects $X, Y \in \text{gr-CM } A$ and for any $i \in \mathbb{Z}$ we have functorial isomorphisms

$$\begin{aligned}
 D\underline{\text{Ext}}_{\text{gr-}A}^i(X, Y) &\cong D\underline{\text{Hom}}_{\text{gr-}A}(X, Y[i]) \\
 &\cong \underline{\text{Hom}}_{\text{gr-}A}(\mathbf{R}\text{Hom}_{A^e}(A, A^e)[i+1] \overset{\mathbf{L}}{\otimes}_A Y, X) \\
 &\cong \underline{\text{Hom}}_{\text{gr-}A}(A[i-d](p) \overset{\mathbf{L}}{\otimes}_A Y, X) \\
 &\cong \underline{\text{Hom}}_{\text{gr-}A}(Y(p), X[d-i]) \\
 &\cong \underline{\text{Ext}}_{\text{gr-}A}^{d-i}(Y(p), X).
 \end{aligned}$$

□

Note that if A is bimodule stably (p) -twisted d -Calabi-Yau, then it is bimodule stably d -Calabi-Yau as an ungraded algebra. Therefore, by the same argument the stable category $\underline{\text{CM}} A$ is d -Calabi-Yau.

2.4. Higher preprojective algebras. Let Λ be a finite dimensional algebra of global dimension $d-1$. We denote by $\mathbb{S}_{d-1} = - \overset{\mathbf{L}}{\otimes}_{\Lambda} D\Lambda[-d+1]$ the composition of the Serre functor with the $(d-1)$ desuspension of the bounded derived category $\text{D}^b(\text{mod } \Lambda)$. It is an autoequivalence of $\text{D}^b(\text{mod } \Lambda)$. We denote by τ_{d-1} the composition

$$\text{mod } \Lambda \hookrightarrow \text{D}^b(\text{mod } \Lambda) \xrightarrow{\mathbb{S}_{d-1}} \text{D}^b(\text{mod } \Lambda) \xrightarrow{H^0} \text{mod } \Lambda$$

The algebra Λ is called τ_{d-1} -finite if the functor τ_{d-1} is nilpotent.

Definition 2.13. [IO13] The d -preprojective algebra of Λ is defined to be the tensor algebra

$$\Pi_d(\Lambda) := T_{\Lambda} \text{Ext}_{\Lambda}^{d-1}(D\Lambda, \Lambda).$$

It is immediate to see that we have $\Pi_d \Lambda \cong \bigoplus_{p \geq 0} \tau_{d-1}^{-p} \Lambda$ as Λ -bimodules. Hence the algebra Λ is τ_{d-1} -finite if and only if $\Pi_d(\Lambda)$ is finite dimensional.

3. HOMOLOGICAL CHARACTERIZATION OF FINITE DIMENSIONAL PREPROJECTIVE ALGEBRAS

In this section we prove the main result of the paper, that gives a sufficient condition for a finite dimensional graded algebra Π to be the d -preprojective algebra of its degree zero subalgebra.

3.1. Main result and strategy of the proof.

Theorem 3.1. *Let Π be a finite dimensional positively graded algebra, and an integer $d \geq 2$, such that*

- (1) Π is Gorenstein of dimension at most $d-2$, that is $\text{id } \Pi = \text{pd } D\Pi = g$, for some $g \leq d-2$;
- (2) Π is bimodule stably (1) -twisted d -Calabi-Yau;
- (3) $\text{Ext}_{\text{gr-}\Pi^e}^j(\Pi, \Pi^e(i)) = 0$ for all $i < 0$ and $j > 0$.

Then $\Lambda = \Pi_0$ is a τ_{d-1} -finite algebra of global dimension $\leq d-1$, and Π is the d -preprojective algebra of Λ .

The main ingredient of the proof is the triangle

$$\Pi_{\leq 0} \longrightarrow \Pi \longrightarrow \Pi_{> 0} \longrightarrow \quad \text{in } D^b(\text{gr-}\Pi^e)$$

given by Proposition 2.1, that is a decomposition of the projective bimodule resolution of Π according to the degrees the projective bimodules are generated in. More precisely the strategy of the proof consists of the computation of the cohomology groups of the triangle

$$\Pi_{\leq 0} \otimes_{\Pi}^{\mathbf{L}} \Lambda \longrightarrow \Pi \otimes_{\Pi}^{\mathbf{L}} \Lambda \longrightarrow \Pi_{> 0} \otimes_{\Pi}^{\mathbf{L}} \Lambda \longrightarrow \quad \text{in } D^b(\text{gr-}\Pi \otimes \Lambda^{\text{op}}).$$

Let us start by reformulating some of the conditions. By Lemmas 2.6 and 2.8 and Remark 2.7 there is a short exact sequence of Π^e -modules

$$K \twoheadrightarrow M \twoheadrightarrow \Pi,$$

such that K has a finite projective resolution, with terms generated in non-positive degree, and M is a Cohen-Macaulay Π^e -module, all of whose projective summands are generated in degree zero.

We get the following immediate reformulation of Condition (2) in terms of M :

Observation 3.2. We may reformulate the condition that Π is stably (1)-twisted d -Calabi-Yau (Condition (2)) by adding the following isomorphisms (in $\text{gr-CM}(\Pi^e)$) in front of and after the defining one:

$$\mathbf{R}\text{Hom}_{\Pi^e}(M, \Pi^e)[d+1] \cong \mathbf{R}\text{Hom}_{\Pi^e}(\Pi, \Pi^e)[d+1] \stackrel{\text{def}}{\cong} \Pi(1) \cong \Omega_{\Pi^e}^{d+1} \Pi[d+1](1).$$

Note that since M is Cohen-Macaulay we may drop the \mathbf{R} in the left-most term. Moreover by Condition (1) we have that also $\Omega_{\Pi^e}^{d+1} \Pi$ is Cohen-Macaulay. Thus, under Condition (1), Condition (2) is equivalent to

$$\text{Hom}_{\Pi^e}(M, \Pi^e)(-1) \cong \Omega_{\Pi^e}^{d+1} \Pi$$

up to projective summands generated in degree 1.

3.2. The global dimension of Λ .

Proposition 3.3. *In the setup of Theorem 3.1 we have $\text{gl.dim } \Lambda \leq d - 1$.*

Proof. It suffices to show that $\text{pd}_{\Lambda^e} \Lambda \leq d - 1$. We have the following isomorphisms of Λ^e -modules:

$$\begin{aligned} \Omega_{\Lambda^e}^d \Lambda &\cong (\Omega_{\Pi^e}^d \Pi)_0 \\ &\cong (\Omega_{\Pi^e}^{-1} \Omega_{\Pi^e}^{d+1} \Pi)_0 && \text{since } \Omega_{\Pi^e}^d \Pi \text{ is CM (Assumption (1))} \\ &\cong (\Omega_{\Pi^e}^{-1} \text{Hom}_{\Pi^e}(M, \Pi^e)(-1))_0 && \text{by Observation 3.2} \\ &\cong (\text{Hom}_{\Pi^e}(\Omega_{\Pi^e} M, \Pi^e)(-1))_0 \\ &= \text{Hom}_{\text{gr-}\Pi^e}(\Omega_{\Pi^e} M, \Pi^e(-1)) \end{aligned}$$

From the short exact sequence $K \twoheadrightarrow M \twoheadrightarrow \Pi$, it is easy to construct a short exact sequence $K' \twoheadrightarrow \Omega_{\Pi^e} M \twoheadrightarrow \Omega_{\Pi^e} \Pi$ with K' having a projective resolution in $\text{gr-proj}_{\leq 0} \Pi^e$. Thus $\text{Hom}_{\text{gr-}\Pi^e}(K', \Pi^e(-1))$ vanishes and we have

$$\text{Hom}_{\text{gr-}\Pi^e}(\Omega_{\Pi^e} M, \Pi^e(-1)) \cong \text{Hom}_{\text{gr-}\Pi^e}(\Omega_{\Pi^e} \Pi, \Pi^e(-1)).$$

Therefore we have

$$\Omega_{\Lambda^e}^d(\Lambda) \cong \text{Ext}_{\text{gr-}\Pi^e}^1(\Pi, \Pi^e(-1)) = 0$$

by assumption (3). \square

3.3. Proof of Theorem 3.1. We start with a technical lemma that will be useful.

Lemma 3.4. *In the setup of Theorem 3.1 we have an isomorphism*

$$H^i(\Pi_{>0} \otimes_{\Pi}^{\mathbf{L}} \Lambda) \cong H^{i+d}(\Pi(-1) \otimes_{\Lambda}^{\mathbf{L}} \mathbf{R}\text{Hom}_{\Lambda}(D\Lambda, \Lambda)) \quad \text{in } \text{gr-}(\Pi \otimes \Lambda^{\text{op}})$$

for all $i \geq -d + g + 1$, where g is the Gorenstein dimension of Π .

In particular

$$H^{-1}(\Pi_{>0} \otimes_{\Pi}^{\mathbf{L}} \Lambda) \cong \Pi(-1) \otimes_{\Lambda} \text{Ext}_{\Lambda}^{d-1}(D\Lambda, \Lambda)$$

and

$$H^i(\Pi_{>0} \otimes_{\Pi}^{\mathbf{L}} \Lambda) = 0 \quad \forall i \geq 0.$$

Proof. Since K has a projective resolution with terms in $\text{gr-proj}_{\leq 0} \Pi^e$ we have

$$\Pi_{>0} \cong M_{>0} \quad \text{in } D^b(\text{gr-}\Pi^e).$$

From Observation 3.2 we know that the graded Cohen-Macaulay Π^e -modules M and $\text{Hom}_{\Pi^e}(\Omega_{\Pi^e}^{d+1} \Pi, \Pi^e)(-1)$ are isomorphic up to projective summands generated in degree 0. Thus

$$\Pi_{>0} \cong (\text{Hom}_{\Pi^e}(\Omega_{\Pi^e}^{d+1} \Pi, \Pi^e)(-1))_{>0}.$$

Next we denote by P the complex formed by the first $d + 1$ terms of a graded projective resolution of Π as Π^e -module. Thus we have a triangle

$$\Omega_{\Pi^e}^{d+1} \Pi[d] \longrightarrow P \longrightarrow \Pi \longrightarrow \quad \text{in } D^b(\text{gr-}\Pi^e).$$

Applying the functor $(\mathbf{R}\text{Hom}_{\Pi^e}(-[-d], \Pi^e)(-1))_{>0}$ we obtain

$$(\mathbf{R}\text{Hom}_{\Pi^e}(\Pi[-d], \Pi^e)(-1))_{>0} \longrightarrow (\text{Hom}_{\Pi^e}(P[-d], \Pi^e)(-1))_{>0} \longrightarrow \Pi_{>0} \longrightarrow .$$

Observe that since $\Omega_{\Pi^e}^g \Pi$ is Cohen-Macaulay, the complex $\mathbf{R}\text{Hom}_{\Pi^e}(\Omega_{\Pi^e}^g \Pi, \Pi^e)$ is concentrated in homological degree 0, hence the complex $\mathbf{R}\text{Hom}_{\Pi^e}(\Pi, \Pi^e)$ is concentrated in homological degrees $0, \dots, g$. Therefore the complex

$$\mathbf{R}\text{Hom}_{\Pi^e}(\Pi[-d], \Pi^e)(-1) = \mathbf{R}\text{Hom}_{\Pi^e}(\Pi, \Pi^e)[d](-1)$$

is concentrated in homological degrees $-d, \dots, -d+g$. Thus also $\mathbf{R}\text{Hom}_{\Pi^e}(\Pi[-d], \Pi^e)(-1) \otimes_{\Pi}^{\mathbf{L}} \Lambda$ is concentrated in homological degrees $\leq -d + g$.

It follows that for $i \geq -d + g + 1$ we have

$$(3.1) \quad \begin{aligned} H^i(\Pi_{>0} \otimes_{\Pi}^{\mathbf{L}} \Lambda) &\cong H^i(\text{Hom}_{\Pi^e}(P[-d], \Pi^e)(-1))_{>0} \otimes_{\Pi}^{\mathbf{L}} \Lambda \\ &\cong H^{i+d}((\text{Hom}_{\Pi^e}(P, \Pi^e)(-1))_{>0} \otimes_{\Pi}^{\mathbf{L}} \Lambda). \end{aligned}$$

Moreover observe that

$$(\text{Hom}_{\Pi^e}(P, \Pi^e)(-1))_{>0} \cong \text{Hom}_{\Pi^e}(P, \Pi^e)_{>-1}(-1) \cong \text{Hom}_{\Pi^e}(P_{\leq 0}, \Pi^e)(-1).$$

Since Π is a positively graded algebra, $P_{\leq 0}$ is generated in degree 0. Hence $P_{\leq 0} = \Pi \otimes Q \otimes \Pi$, where Q is the complex formed by the first $(d + 1)$ terms

of a projective resolution of $\Pi_0 = \Lambda$ as Λ^e -module. Since $\text{gl.dim } \Lambda \leq d - 1$, Q is a projective resolution of Λ as Λ^e -module, this leads to

$$(3.2) \quad (\text{Hom}_{\Pi^e}(P, \Pi^e)(-1))_{>0} \cong \Pi \otimes_{\Lambda} \text{Hom}_{\Lambda^e}(Q, \Lambda^e) \otimes_{\Lambda} \Pi(-1).$$

Combining (3.1) and (3.2) we obtain

$$\begin{aligned} \mathbf{H}^i(\Pi_{>0} \otimes_{\Pi}^{\mathbf{L}} \Lambda) &\cong \mathbf{H}^{i+d}((\text{Hom}_{\Pi^e}(P, \Pi^e)(-1))_{>0} \otimes_{\Pi}^{\mathbf{L}} \Lambda) \\ &\cong \mathbf{H}^{i+d}(\Pi \otimes_{\Lambda} \text{Hom}_{\Lambda^e}(Q, \Lambda^e) \otimes_{\Lambda} \Pi(-1) \otimes_{\Pi} \Lambda) \\ &\cong \mathbf{H}^{i+d}(\Pi(-1) \otimes_{\Lambda}^{\mathbf{L}} \mathbf{R}\text{Hom}_{\Lambda^e}(\Lambda, \Lambda^e)) \\ &\cong \mathbf{H}^{i+d}(\Pi(-1) \otimes_{\Lambda}^{\mathbf{L}} \mathbf{R}\text{Hom}_{\Lambda}(D\Lambda, \Lambda)) \end{aligned}$$

for $i \geq -d + g + 1$.

Now since $g \leq d - 2$, we have $g - d + 1 \leq -1$. Moreover the global dimension of Λ is at most $d - 1$, so the complex $\mathbf{R}\text{Hom}_{\Lambda}(D\Lambda, \Lambda)$ is concentrated in degrees $\leq d - 1$. Hence we have

$$\begin{aligned} \mathbf{H}^{i+d}(\Pi(-1) \otimes_{\Lambda}^{\mathbf{L}} \mathbf{R}\text{Hom}_{\Lambda}(D\Lambda, \Lambda)) &\cong \Pi(-1) \otimes_{\Lambda} \text{Ext}^{d-1}(D\Lambda, \Lambda) && \text{for } i = -1 \\ &= 0 && \text{for } i \geq 0. \end{aligned}$$

This finishes the proof. \square

We now obtain the following short exact sequence, which is an essential ingredient to our proof.

Proposition 3.5. *In the setup of Theorem 3.1 we have a short exact sequence of graded $\Pi^{\text{op}} \otimes \Lambda$ -modules*

$$\Pi(-1) \otimes_{\Lambda} \text{Ext}_{\Lambda}^{d-1}(D\Lambda, \Lambda) \twoheadrightarrow \Pi \twoheadrightarrow \Lambda,$$

where $\Pi \twoheadrightarrow \Lambda$ is the natural projection.

Proof. Consider Π as object in $\text{D}^b(\text{gr-}\Pi^e)$, and the triangle

$$\Pi_{\leq 0} \longrightarrow \Pi \longrightarrow \Pi_{>0} \longrightarrow$$

given by Proposition 2.1. Applying the functor $-\otimes_{\Pi}^{\mathbf{L}} \Lambda$ we obtain a triangle

$$\Pi_{\leq 0} \otimes_{\Pi}^{\mathbf{L}} \Lambda \longrightarrow \Pi \otimes_{\Pi}^{\mathbf{L}} \Lambda \longrightarrow \Pi_{>0} \otimes_{\Pi}^{\mathbf{L}} \Lambda \longrightarrow \quad \text{in } \text{D}^b(\text{gr-}\Pi \otimes \Lambda^{\text{op}}).$$

Recall that from Observation 3.2, $\Omega_{\Pi^e}^{d+1}\Pi$ and $\text{Hom}_{\Pi^e}(M, \Pi^e)(-1)$ are isomorphic up to projective summands generated in degree 1. Hence we have the following isomorphisms in $\text{D}^b(\text{gr-}\Pi^e)$:

$$\begin{aligned} (\Omega_{\Pi^e}^{d+1}\Pi)_{\leq 0} &\cong (\text{Hom}_{\Pi^e}(M, \Pi^e)(-1))_{\leq 0} \\ &\cong (\text{Hom}_{\Pi^e}(\Pi, \Pi^e)(-1))_{\leq 0} && \text{since the projective resolution} \\ & && \text{of } K \text{ is generated in non-positive} \\ & && \text{degrees} \\ &= 0 && \text{since } \Pi \text{ is a positively graded algebra.} \end{aligned}$$

So we get $\Pi_{\leq 0} \cong P_{\leq 0} = \Pi \otimes_{\Lambda} Q \otimes \Pi$ where Q is a projective resolution of Λ as Λ^e -module. Therefore we obtain $\Pi_{\leq 0} \otimes_{\Pi}^{\mathbf{L}} \Lambda \cong \Pi$.

Since we clearly have $\Pi \otimes_{\Pi}^{\mathbf{L}} \Lambda = \Lambda$ we obtain an exact sequence

$$\mathbf{H}^{-1}(\Pi_{>0} \otimes_{\Pi}^{\mathbf{L}} \Lambda) \twoheadrightarrow \Pi \twoheadrightarrow \Lambda \twoheadrightarrow \mathbf{H}^0(\Pi_{>0} \otimes_{\Pi}^{\mathbf{L}} \Lambda).$$

Now the claim follows from the “in particular” part of Lemma 3.4 above. \square

By Proposition 3.5 we have an isomorphism of graded $\Pi \otimes \Lambda$ -modules

$$\Pi(-1) \otimes_{\Lambda} \text{Ext}_{\Lambda}^{d-1}(D\Lambda, \Lambda) \cong \Pi_{>0}.$$

The next Proposition shows that this is enough to identify Π as the tensor algebra $T_{\Lambda} \text{Ext}_{\Lambda}^{d-1}(D\Lambda, \Lambda)$ and thus finishes the proof of Theorem 3.1.

Proposition 3.6. *Let Π be a positively graded ring, $\Lambda = \Pi_0$, and X a $\Lambda \otimes \Lambda^{\text{op}}$ -module. Assume there is an isomorphism*

$$\Pi(-1) \otimes_{\Lambda} X \cong \Pi_{>0}$$

of graded $\Pi \otimes \Lambda^{\text{op}}$ -modules.

Then, as graded rings,

$$\Pi \cong T_{\Lambda} X.$$

Proof. Let $h: \Pi(-1) \otimes_{\Lambda} X \xrightarrow{\cong} \Pi_{>0}$ as in the assumption.

We define an isomorphism of graded $\Lambda \otimes \Lambda^{\text{op}}$ -modules

$$\varphi: T_{\Lambda} X \longrightarrow \Pi$$

iteratedly by $\varphi_0 = \text{id}_{\Lambda}$ and by letting φ_n be the composition

$$(T_{\Lambda} X)_n = X^{\otimes_{\Lambda} n} \xrightarrow{\varphi_{n-1} \otimes \text{id}_X} \Pi_{n-1} \otimes_{\Lambda} X \cong \Pi_n,$$

where the last isomorphism is the degree n -part of h .

It only remains to check that φ respects the ring-multiplication. It suffices to check that

$$\varphi_{n+m}(f \otimes g) = \varphi_n(f) \varphi_m(g)$$

for any $f \in X^{\otimes_{\Lambda} n}$ and $g \in X^{\otimes_{\Lambda} m}$. We show this by induction on m . For $m = 0$ this is just the Λ -linearity of φ_n . For $m = 1$ we have

$$\varphi_{n+1}(f \otimes g) \stackrel{\text{def}}{=} h(\varphi_n(f) \otimes g) = \varphi_n(f) h(1 \otimes g) \stackrel{\text{def}}{=} \varphi_n(f) \varphi_1(g).$$

For $m > 1$ we may assume that $g = x \otimes g'$ for some $x \in X$. (An arbitrary element is a sum of elementary tensors, but since all maps involved are linear it suffices to consider a single elementary tensor.) Now

$$\begin{aligned} \varphi_{n+m}(f \otimes g) &= \varphi_{n+m}(f \otimes x \otimes g') \\ &= \varphi_{n+1}(f \otimes x) \varphi_{m-1}(g') && \text{by inductive assumption} \\ &= \varphi_n(f) \varphi_1(x) \varphi_{m-1}(g') && \text{by the case } m = 1 \\ &= \varphi_n(f) \varphi_1(x \otimes g') && \text{by inductive assumption} \\ &= \varphi_n(f) \varphi_m(g). \end{aligned} \quad \square$$

Observation 3.7. Since Π is a finite-dimensional algebra, it follows that Λ is τ_{d-1} -finite, since this is equivalent to $T_{\Lambda} \text{Ext}_{\Lambda}^{d-1}(D\Lambda, \Lambda)$ being finite dimensional.

4. THE GORENSTEIN DIMENSION

4.1. Gorenstein dimension and cluster tilting subcategories. In this section, we express the Gorenstein dimension of the algebra Π of Theorem 3.1 in terms of certain vanishing of extensions in the derived category $D^b(\text{mod } \Lambda)$. We start with the following result/definition:

Theorem 4.1 ([Ami08, 5.4.2][Iya11, 1.22]). *Let Λ be a τ_{d-1} -finite algebra. Then the category*

$$\mathcal{U} = \text{add}\{\mathbb{S}_{d-1}^i \Lambda \mid i \in \mathbb{Z}\} \subseteq D^b(\text{mod } \Lambda)$$

is a $(d-1)$ -cluster tilting subcategory, that is

$$\begin{aligned} \mathcal{U} &= \{X \in D^b(\text{mod } \Lambda) \mid \text{Ext}_{\Lambda}^i(\mathcal{U}, X) = 0 \ \forall i = 1, \dots, d-2\} \\ &= \{X \in D^b(\text{mod } \Lambda) \mid \text{Ext}_{\Lambda}^i(X, \mathcal{U}) = 0 \ \forall i = 1, \dots, d-2\}. \end{aligned}$$

Note that if Π is an algebra as in Theorem 3.1, its degree zero subalgebra Λ is always τ_{d-1} -finite, so the result above applies. The aim here is to express the Gorenstein dimension g of Π using the subcategory \mathcal{U} . More precisely the main result of this section is the following.

Theorem 4.2. *In the setup of Theorem 3.1 the Gorenstein-dimension g of Π is given by*

$$g = d - 1 + \max\{i < 0 \mid \text{Hom}_{D^b(\text{mod } \Lambda)}(\mathcal{U}, \mathcal{U}[i]) \neq 0\}.$$

The proof of this theorem consists of two main steps: First, in Lemma 4.3 and Proposition 4.4, we calculate g in terms of the non-vanishing of homologies of the complex $\Pi \otimes_{\Lambda}^{\mathbf{L}} \mathbf{R}\text{Hom}_{\Lambda}(D\Lambda, \Lambda)$. Second we show that this description coincides with the right hand side term given in the theorem.

Lemma 4.3. *In the setup of Theorem 3.1 we have*

$$H^i(\Pi \otimes_{\Lambda}^{\mathbf{L}} \mathbf{R}\text{Hom}_{\Lambda}(D\Lambda, \Lambda)) = 0 \quad \forall i \in \{g+1, \dots, d-2\}.$$

Proof. By Lemma 3.4 we have

$$H^i(\Pi \otimes_{\Lambda}^{\mathbf{L}} \mathbf{R}\text{Hom}_{\Lambda}(D\Lambda, \Lambda)) = H^{i-d}(\Pi_{>0} \otimes_{\Pi}^{\mathbf{L}} \Lambda) \quad \forall i \geq g+1.$$

Looking at the proof of Proposition 3.5 we see that

$$H^i(\Pi_{>0} \otimes_{\Pi}^{\mathbf{L}} \Lambda) = 0 \quad \forall i \neq -1.$$

The claim follows from these two statements. \square

We now prove a converse of Lemma 4.3.

Proposition 4.4. *In the setup of Theorem 3.1 we have*

$$g = \max\{i \leq d-2 \mid H^i(\Pi \otimes_{\Lambda}^{\mathbf{L}} \mathbf{R}\text{Hom}_{\Lambda}(D\Lambda, \Lambda)) \neq 0\}.$$

Proof. We have the inequality “ \geq ” by Lemma 4.3. It remains to show that $H^g(\Pi \otimes_{\Lambda}^{\mathbf{L}} \mathbf{R}\text{Hom}_{\Lambda}(D\Lambda, \Lambda)) \neq 0$. To do so, we analyse what happens in the proof of Lemma 3.4 for $i = -d + g$. As in the proof there, we obtain the triangle

$$(\mathbf{R}\text{Hom}_{\Pi^e}(\Pi[-d], \Pi^e)(-1))_{>0} \otimes_{\Pi}^{\mathbf{L}} \Lambda \longrightarrow (\text{Hom}_{\Pi^e}(P[-d], \Pi^e)(-1))_{>0} \otimes_{\Pi}^{\mathbf{L}} \Lambda \longrightarrow \Pi_{>0} \otimes_{\Pi}^{\mathbf{L}} \Lambda \longrightarrow$$

where P is the complex formed by the first $(d+1)$ terms of a projective resolution of Π as a graded Π -bimodule. Since $H^i(\Pi_{>0} \otimes_{\Pi}^{\mathbf{L}} \Lambda) = 0$ for $i \neq -1$ it follows that

$$H^i((\mathbf{R}\mathrm{Hom}_{\Pi^e}(\Pi[-d], \Pi^e)(-1))_{>0} \otimes_{\Pi}^{\mathbf{L}} \Lambda) \cong H^i((\mathrm{Hom}_{\Pi^e}(P[-d], \Pi^e)(-1))_{>0} \otimes_{\Pi}^{\mathbf{L}} \Lambda)$$

whenever $i < -1$. In particular

$$\begin{aligned} H^g((\mathbf{R}\mathrm{Hom}_{\Pi^e}(\Pi, \Pi^e)(-1))_{>0} \otimes_{\Pi}^{\mathbf{L}} \Lambda) &= H^{g-d}((\mathbf{R}\mathrm{Hom}_{\Pi^e}(\Pi[-d], \Pi^e)(-1))_{>0} \otimes_{\Pi}^{\mathbf{L}} \Lambda) \\ &\cong H^{g-d}((\mathrm{Hom}_{\Pi^e}(P[-d], \Pi^e)(-1))_{>0} \otimes_{\Pi}^{\mathbf{L}} \Lambda) \\ &\quad (\text{since } g-d \leq -2) \\ &= H^g((\mathrm{Hom}_{\Pi^e}(P, \Pi^e)(-1))_{>0} \otimes_{\Pi}^{\mathbf{L}} \Lambda). \end{aligned}$$

As in the proof of Lemma 3.4 we observe that

$$\begin{aligned} (\mathrm{Hom}_{\Pi^e}(P, \Pi^e)(-1))_{>0} \otimes_{\Pi}^{\mathbf{L}} \Lambda &\cong (\mathrm{Hom}_{\Pi^e}(P_{\leq 0}, \Pi^e)(-1)) \otimes_{\Pi}^{\mathbf{L}} \Lambda \\ &\cong \Pi(-1) \otimes_{\Lambda}^{\mathbf{L}} \mathrm{Hom}_{\Lambda^e}(Q, \Lambda^e) \\ &\cong \Pi(-1) \otimes_{\Lambda}^{\mathbf{L}} \mathbf{R}\mathrm{Hom}_{\Lambda}(D\Lambda, \Lambda) \end{aligned}$$

where Q is a projective resolution of Λ as a Λ -bimodule. Hence it suffices to show that

$$H^g((\mathbf{R}\mathrm{Hom}_{\Pi^e}(\Pi, \Pi^e)(-1))_{>0} \otimes_{\Pi}^{\mathbf{L}} \Lambda) \neq 0.$$

Since the complex $\mathbf{R}\mathrm{Hom}_{\Pi^e}(\Pi, \Pi^e)$ is concentrated in homological degrees at most g we have

$$H^g((\mathbf{R}\mathrm{Hom}_{\Pi^e}(\Pi, \Pi^e)(-1))_{>0} \otimes_{\Pi}^{\mathbf{L}} \Lambda) = H^g((\mathbf{R}\mathrm{Hom}_{\Pi^e}(\Pi, \Pi^e)(-1))_{>0}) \otimes_{\Pi} \Lambda.$$

Tensoring with Λ cannot kill a finitely generated Π -module, so it suffices to show that

$$H^g((\mathbf{R}\mathrm{Hom}_{\Pi^e}(\Pi, \Pi^e)(-1))_{>0}) \neq 0.$$

By assumption (3) the homology of the complex of graded Π^e -modules $\mathbf{R}\mathrm{Hom}_{\Pi^e}(\Pi, \Pi^e)(-1)$ is concentrated in positive degrees, and hence we have

$$(\mathbf{R}\mathrm{Hom}_{\Pi^e}(\Pi, \Pi^e)(-1))_{>0} = \mathbf{R}\mathrm{Hom}_{\Pi^e}(\Pi, \Pi^e)(-1).$$

Thus it suffices that

$$\underbrace{H^g(\mathbf{R}\mathrm{Hom}_{\Pi^e}(\Pi, \Pi^e)(-1))}_{=\mathrm{Ext}_{\Pi^e}^g(\Pi, \Pi^e)(-1)} \neq 0,$$

which holds by definition of g . \square

Therefore to prove Theorem 4.2 it is sufficient to prove

$$\begin{aligned} &\max\{i \leq d-2 \mid H^i(\Pi \otimes_{\Lambda}^{\mathbf{L}} \mathbf{R}\mathrm{Hom}_{\Lambda}(D\Lambda, \Lambda)) \neq 0\} \\ &= d-1 + \max\{i < 0 \mid \mathrm{Hom}_{\mathrm{D}^b(\mathrm{mod} \Lambda)}(\mathcal{U}, \mathcal{U}[i]) \neq 0\}. \end{aligned}$$

For the proof, we prepare the following two lemmas.

Lemma 4.5. *Let Λ be an algebra of global dimension at most $d-1$. Assume for some $j \geq 0$ and some $p \leq -1$ we have*

$$H^i(\mathbb{S}_{d-1}^{-j}(\Lambda)) = 0 \quad \forall i \in \{p, \dots, -1\}.$$

Then

$$H^i(\mathbb{S}_{d-1}^{-(j+1)}(\Lambda)) \cong H^i(\mathbb{S}_{d-1}^{-1}(\tau_{d-1}^{-j}\Lambda)) \quad \forall i \geq p.$$

Proof. Since Λ has global dimension $\leq d - 1$ an easy induction shows that the functor \mathbb{S}_{d-1}^{-j} preserves the left aisle $D^b(\text{mod } \Lambda)^{\leq 0}$ of the canonical t -structure of $D^b(\text{mod } \Lambda)$. Therefore $\mathbb{S}_{d-1}^{-j}\Lambda$ is in negative degrees and we can consider the triangle

$$\text{trunc}^{<0}(\mathbb{S}_{d-1}^{-j}(\Lambda)) \longrightarrow \mathbb{S}_{d-1}^{-j}(\Lambda) \longrightarrow \underbrace{\mathrm{H}^0(\mathbb{S}_{d-1}^{-j}(\Lambda))}_{=\tau_{d-1}^{-j}\Lambda} \longrightarrow,$$

where $\text{trunc}^{<0}X$ is the usual truncation of the complex X . Applying \mathbb{S}_{d-1}^{-1} to it we obtain the triangle

$$\mathbb{S}_{d-1}^{-1}(\text{trunc}^{<0}(\mathbb{S}_{d-1}^{-j}(\Lambda))) \longrightarrow \mathbb{S}_{d-1}^{-(j+1)}(\Lambda) \longrightarrow \mathbb{S}_{d-1}^{-1}(\tau_{d-1}^{-j}\Lambda) \longrightarrow.$$

By assumption we have that $\text{trunc}^{<0}(\mathbb{S}_{d-1}^{-j}(\Lambda))$ is concentrated in degrees $\leq p - 1$. Hence $\mathbb{S}_{d-1}^{-1}(\text{trunc}^{<0}(\mathbb{S}_{d-1}^{-j}(\Lambda)))$ is also concentrated in degrees $\leq p - 1$. Thus the triangle gives the desired isomorphism of homologies. \square

Lemma 4.6. *Let Λ be an algebra of global dimension at most $d - 1$. The following are equivalent:*

- (1) $\forall i \in \{p, \dots, -1\} \forall j \geq 0: \mathrm{H}^i(\mathbb{S}_{d-1}^{-j}(\Lambda)) = 0;$
- (2) $\forall i \in \{p, \dots, -1\} \forall j \geq 0: \mathrm{H}^i(\mathbb{S}_{d-1}^{-1}(\tau_{d-1}^{-j}\Lambda)) = 0.$

Proof. (1) \implies (2) follows immediately from Lemma 4.5.

(2) \implies (1) follows from Lemma 4.5 by induction on j . \square

Now we are ready to prove Theorem 4.2.

Proof of Theorem 4.2. We first note that $\mathbf{R}\mathrm{Hom}_{\Lambda}(D\Lambda, \Lambda) \otimes_{\Lambda}^{\mathbf{L}} -$ is the inverse Serre functor on $D^b(\text{mod } \Lambda)$, and that Π is isomorphic to $\bigoplus_{j \geq 0} \tau_{d-1}^{-j}\Lambda$ as a Λ -module. Hence we get the following equivalences for $\ell \leq d - 2$:

$$\begin{aligned} \mathrm{H}^i(\mathbf{R}\mathrm{Hom}_{\Lambda}(D\Lambda, \Lambda) \otimes_{\Lambda}^{\mathbf{L}} \Pi) &= 0 \quad \forall i \in \{\ell, \dots, d - 2\} \\ \iff \mathrm{H}^i(\mathbb{S}^{-1}(\Pi)) &= 0 \quad \forall i \in \{\ell, \dots, d - 2\} \\ \iff \mathrm{H}^i(\mathbb{S}^{-1}(\tau_{d-1}^{-j}\Lambda)) &= 0 \quad \forall i \in \{\ell, \dots, d - 2\} \forall j \geq 0 \\ \iff \mathrm{H}^i(\mathbb{S}_{d-1}^{-1}(\tau_{d-1}^{-j}\Lambda)) &= 0 \quad \forall i \in \{-d + 1 + \ell, \dots, -1\} \forall j \geq 0 \\ \iff \mathrm{H}^i(\mathbb{S}_{d-1}^{-(j+1)}(\Lambda)) &= 0 \quad \forall i \in \{-d + 1 + \ell, \dots, -1\} \forall j \geq 0, \end{aligned}$$

where the last equivalence is Lemma 4.6 for $p = -d + 1 + \ell$.

We may drop the restriction to non-negative j , since $\mathbb{S}_{d-1}^{-j}(\Lambda)$ is concentrated in positive degrees for negative j . So we have

$$\begin{aligned} \mathrm{H}^i(\mathbf{R}\mathrm{Hom}_{\Lambda}(D\Lambda, \Lambda) \otimes_{\Lambda}^{\mathbf{L}} \Pi) &= 0 \quad \forall i \in \{\ell, \dots, d - 2\} \\ \iff \mathrm{H}^i(\mathbb{S}_{d-1}^{-j}(\Lambda)) &= 0 \quad \forall i \in \{-d + 1 + \ell, \dots, -1\} \forall j \\ \iff \mathrm{Hom}_{D^b(\text{mod } \Lambda)}(\Lambda, \mathcal{U}[i]) &= 0 \quad \forall i \in \{-d + 1 + \ell, \dots, -1\}. \end{aligned}$$

Therefore we get

$$\max\{i \leq d - 2 \mid \mathrm{H}^i(\Pi \otimes_{\Lambda}^{\mathbf{L}} \mathbf{R}\mathrm{Hom}_{\Lambda}(D\Lambda, \Lambda)) \neq 0\} = d - 1 + \max\{i < 0 \mid \mathrm{Hom}_{D^b(\text{mod } \Lambda)}(\mathcal{U}, \mathcal{U}[i]) \neq 0\}$$

which finishes the proof of Theorem 4.2. \square

4.2. The selfinjective case. The situation of Theorem 3.1 is especially nice when the Gorenstein dimension g of the algebra Π is 0, that is when Π is selfinjective. In that case we prove that the algebra Π is the preprojective algebra of a $(d - 1)$ -representation finite algebra.

Definition 4.7 ([IO11, Def 2.2], [IO13, Thm 3.1]). A τ_{d-1} -algebra Λ is said to be $(d - 1)$ -representation finite if the subcategory $\mathcal{U} \subset \mathrm{D}^b(\mathrm{mod} \Lambda)$ is stable under the Serre functor, that is $\mathbb{S}\mathcal{U} = \mathcal{U}$.

Theorem 4.8. *Let Π be a finite dimensional selfinjective positively graded algebra which is bimodule stably (1) -twisted d -Calabi-Yau. Then $\Lambda = \Pi_0$ is a $(d - 1)$ -representation finite algebra and there is an isomorphism of graded algebra $\Pi \cong \Pi_d(\Lambda)$.*

In order to prove this result, we introduce a technical definition.

Definition 4.9. [IO13] Let Λ be a τ_{d-1} -finite algebra. We say that Λ has the vanishing of small negative extensions property (**vosnex** for short) if

$$\mathrm{Hom}_{\mathrm{D}^b(\mathrm{mod} \Lambda)}(\mathcal{U}, \mathcal{U}[i]) = 0 \quad \forall i \in \{-(d - 3), \dots, -1\}.$$

From Theorem 4.2, we immediately deduce the following result, giving an equivalent but more transparent characterization of what it means for an algebra to satisfy the vosnex property.

Corollary 4.10. *In the setup of Theorem 3.1 the following are equivalent:*

- (a) *the graded algebra Π has Gorenstein dimension ≤ 1*
- (b) *the algebra $\Lambda = \Pi_0$ has the vosnex property.*

Using this corollary together with Theorem 3.1 and some results in [IO13], we achieve the proof of Theorem 4.8.

Proof of Theorem 4.8. First note that an algebra is selfinjective if and only if it is Gorenstein of dimension 0. Moreover if Π is selfinjective, then it clearly satisfies hypothesis (3) of Theorem 3.1. Hence Π is the d -preprojective algebra of its degree zero part Λ . Moreover, by [IO13, Corollary 3.7], the vosnex property implies that Λ is a $(d - 1)$ -representation finite algebra. \square

5. BIMODULE CALABI-YAU PROPERTIES OF PREPROJECTIVE ALGEBRAS

Throughout this section k is assumed to be an algebraically closed field.

By a classical result due to Ringel [Rin98], if Q is an acyclic quiver, the 2-preprojective algebra Π of the hereditary algebra kQ is the usual preprojective algebra of Q . If Q is Dynkin, then it is well-known that Π is selfinjective, finite-dimensional and that $\underline{\mathrm{mod}} \Pi$ is 2-Calabi-Yau.

On the other hand, if Λ is a τ_2 -finite algebra of global dimension 2, its preprojective algebra $\Pi = \Pi_3(\Lambda)$ is the endomorphism algebra of a cluster-tilting object in a 2-Calabi-Yau category [Ami09, Theorem 4.10]. Hence by [KR07, Theorem 3.3], the algebra Π is Gorenstein and the stable category $\underline{\mathrm{CM}} \Pi$ is 3-Calabi-Yau.

More generally, if Λ is an τ_{d-1} -finite algebra of global dimension $d - 1$, its preprojective algebra $\Pi = \Pi_d(\Lambda)$ is the endomorphism algebra of a $(d - 1)$ -cluster-tilting object in a $(d - 1)$ -Calabi-Yau category by [Guo11, Theorem 4.9]. If moreover Λ is $(d - 1)$ -representation finite, the stable

category $\underline{\text{mod}}\Pi$ is d -Calabi-Yau by [IO13, Corollary 4.6] (see also [Dug12, Proposition 3.3]).

In this section, we prove that these Calabi-Yau properties can be deduced from bimodule properties of the preprojective algebra. More precisely, we prove that in the above cases, the preprojective algebra satisfies the properties (1), (2) and (3) of Theorem 3.1.

5.1. Classical preprojective algebras. Let Q be an acyclic quiver. Then the 2-preprojective algebra $\Pi_2(kQ) = \text{T}_{kQ} \text{Ext}_{kQ}^1(DkQ, kQ)$ is given by the double quiver \bar{Q} , obtained from Q by adding for any $a: i \rightarrow j$ an arrow $\bar{a}: j \rightarrow i$, with the preprojective relations: $\sum_{a \in Q_1} a\bar{a} - \bar{a}a$. The functor τ_1 is isomorphic to the Auslander-Reiten translation of $\text{D}^b(\text{mod } kQ)$. Thus kQ is τ_1 -finite if and only if the quiver Q is of Dynkin type.

Using this description, we prove the converse of Theorem 3.1 for the case $d = 2$.

Theorem 5.1. *Let Λ be a basic τ_1 -finite algebra of global dimension ≤ 1 . Then the 2-preprojective algebra $\Pi := \Pi_2(\Lambda)$ satisfies the following properties:*

- (1) Π is selfinjective (=Gorenstein of dimension 0);
- (2) Π is bimodule stably (1)-twisted 2-Calabi-Yau.

In particular, $\underline{\text{mod}}\Pi$ is a 2-Calabi-Yau category.

Note that the selfinjectivity of Π immediately implies that $\text{Ext}_{\text{gr-}\Pi^e}^j(\Pi, \Pi^e(i))$ vanishes for all i and all $j > 0$, so condition (3) of Theorem 3.1 is automatically satisfied.

Proof. (1) is well known.

The beginning of the minimal projective resolution of Π as a graded Π -bimodule is of the following form:

$$P_\bullet := \bigoplus_{i \in Q_0} \Pi e_i \otimes e_i \Pi(-1) \xrightarrow{d_2} \bigoplus_{a \in Q_1} (\Pi e_{\bar{t}(a)} \otimes e_{\bar{s}(a)} \Pi(-1) \oplus \Pi e_{t(a)} \otimes e_{s(a)} \Pi) \xrightarrow{d_1} \bigoplus_{i \in Q_0} \Pi e_i \otimes e_i \Pi,$$

where the maps d_1 and d_2 are given on components by

$$d_1^{a \rightarrow i} : \Pi e_{t(a)} \otimes e_{s(a)} \Pi \longrightarrow \Pi e_i \otimes e_i \Pi$$

$$e_{t(a)} \otimes e_{s(a)} \longmapsto a e_i \otimes e_i - e_i \otimes e_i a$$

and

$$d_2^{i \rightarrow a} : \Pi e_i \otimes e_i \Pi \longrightarrow \Pi e_{\bar{t}(a)} \otimes e_{\bar{s}(a)} \Pi \oplus \Pi e_{t(a)} \otimes e_{s(a)} \Pi$$

$$e_i \otimes e_i \longmapsto \sum_{a, s(a)=i} e_i \otimes e_{\bar{s}(a)} a + \bar{a} e_{t(a)} \otimes e_i$$

(cf [Scho, ES98] for a non graded version).

Then one easily checks that $\text{Hom}_{\Pi^e}(d_2, \Pi^e) \cong d_1(1)$, that is $\text{Hom}_{\Pi^e}(P_\bullet, \Pi^e)[2] \cong P_\bullet(1)$ in $\text{K}^b(\text{gr-proj } \Pi^e)$. Hence taking H^{-2} , one obtains $\text{Hom}_{\Pi^e}(\Pi, \Pi^e) \cong \Omega_{\Pi^e}^3(\Pi)(1)$ in $\text{gr-}\Pi^e$, which implies (2) by Observation 3.2 (note that Π is Cohen-Macaulay by (1)). \square

5.2. **The case $d = 3$.** For the case $d = 3$, we use a result due to Keller which shows that any 3-preprojective algebra is given by a quiver with potential. So we start by recalling some definitions due to Derksen, Weyman and Zelevinsky [DWZ08].

Definition 5.2. Let Q be a quiver, and W a potential, that is a (possibly infinite) linear combination of cycles in Q . Then the associated *Jacobian algebra* is

$$J = \text{Jac}(Q, W) = \widehat{kQ}/(\partial_a W \mid \varphi \in Q_1),$$

where \widehat{kQ} is the completion of path algebra kQ , and ∂_a is the unique linear map such that $\partial_a p = \sum_{p=ua_v} v u$ for a path p .

Observation 5.3. Let (Q, W) be a quiver with potential. We have

$$\begin{aligned} kQ_0 &= J/\text{Rad } J, \text{ and} \\ kQ_1 &= \text{the } kQ_0 \otimes kQ_0^{\text{op}}\text{-module generated by the arrows of } Q. \end{aligned}$$

The complex

$$(5.1) \quad \bigoplus_{a \in Q_1} J e_{s(a)} \otimes e_{t(a)} J \xrightarrow{d_2} \bigoplus_{a \in Q_1} J e_{t(a)} \otimes e_{s(a)} J \xrightarrow{d_1} \bigoplus_{i \in Q_0} J e_i \otimes e_i J$$

is the beginning of a projective resolution of J as J^e -module.

Here the maps are given on components by

$$\begin{aligned} d_1^{a \rightarrow i} : J e_{t(a)} \otimes e_{s(a)} J &\longrightarrow J e_i \otimes e_i J \\ e_{t(a)} \otimes e_{s(a)} &\longmapsto a e_i \otimes e_i - e_i \otimes e_i a \end{aligned}$$

and $d_2^{a \rightarrow b} = \partial_{a,b} W$, where for a cyclic path p we define

$$\begin{aligned} \partial_{a,b}(p) : J e_{s(a)} \otimes e_{t(a)} J &\longrightarrow J e_{t(b)} \otimes e_{s(b)} J \\ e_{s(a)} \otimes e_{t(a)} &\longmapsto \sum_{p=u_1 a u_2 b u_3} e_{s(a)} u_2 e_{t(b)} \otimes e_{s(b)} u_3 u_1 e_{t(a)} \\ &\quad + \sum_{p=u_1 b u_2 a u_3} e_{s(a)} u_3 u_1 e_{t(b)} \otimes e_{s(b)} u_2 e_{t(a)} \end{aligned}$$

Observation 5.4. Let Π be a finite dimensional algebra, and let $\{e_i \mid i \in \{1, \dots, n\}\}$ be a complete set of idempotents. Then

$$\begin{aligned} \Pi e_i \otimes e_j \Pi &\longrightarrow \text{Hom}_{\Pi^e}(\Pi e_j \otimes e_i \Pi, \Pi^e) \\ a_1 e_i \otimes e_j a_2 &\longmapsto \left[b_1 e_j \otimes e_i b_2 \longmapsto b_1 e_j a_2 \otimes a_1 e_i b_2 \right] \end{aligned}$$

is an isomorphism for any $i, j \in \{1, \dots, n\}$.

Lemma 5.5. Let (Q, W) be a quiver with potential. Let a and b be two arrows of Q .

Then, with the identification of Observation 5.4,

$$\text{Hom}_{J^e}(\partial_{a,b} W, J^e) = \partial_{b,a} W.$$

Proof. It is enough to check this for a cyclic path p . More precisely we have to check that

$$i \circ \partial_{b,a}(p) = \text{Hom}_{\mathbf{J}^e}(\partial_{a,b}(p), \mathbf{J}^e) \circ i.$$

where i is the isomorphism of Observation 5.4. This can be verified by a straight-forward calculation. \square

There is a link between 3-preprojective algebras and Jacobian algebras given by the following result.

Theorem 5.6. [Kel11, Theorem6.12 a)] *Let Λ be a basic finite dimensional algebra of global dimension ≤ 2 . Let Q be the quiver of Λ , and let R be a minimal set of relations, such that $\Lambda \cong kQ/(R)$ and such that R is the disjoint union of sets representing a basis of the Ext_{Λ}^2 -space between any two simple Λ -modules.*

Then there is an isomorphism $T_{\Lambda} \text{Ext}_{\Lambda}^2(D\Lambda, \Lambda) \cong \text{Jac}(\bar{Q}, W)$, where \bar{Q} is obtained by adding to Q an arrow $a_r : t(r) \rightarrow s(r)$ for each $r \in R$, and $W = \sum_{r \in R} a_r r$. The grading on $T_{\Lambda} \text{Ext}_{\Lambda}^2(D\Lambda, \Lambda)$ is given by the arrows of Q having degree 0, and the arrows corresponding to the relations having degree 1.

Using Keller's description of 2-preprojective algebras as Jacobian algebras, we can prove the converse of Theorem 3.1 for the case $d = 3$.

Theorem 5.7. *Let Λ be a τ_2 -finite algebra of global dimension ≤ 2 . Let $\Pi = T_{\Lambda} \text{Ext}_{\Lambda}^2(D\Lambda, \Lambda)$ be the associated 3-preprojective algebra. Then*

- (1) Π is Gorenstein of dimension ≤ 1 ;
- (2) Π is bimodule stably (1)-twisted 3-Calabi-Yau;
- (3) $\text{Ext}_{\Pi^e}^j(\Pi, \Pi^e(i)) = 0$ for all $i < 0$ and all $j > 0$.

In particular, the category $\underline{\text{CM}}\Pi$ is 3-Calabi-Yau.

We start by the following lemma which gives an equivalent condition for (2) in terms of the second syzygy of Π .

Lemma 5.8. *A finite dimensional graded algebra of Gorenstein dimension ≤ 1 is bimodule stably (1)-twisted 3-Calabi-Yau if and only if there is an isomorphism $\text{Hom}_{\Pi^e}(\Omega_{\Pi^e}^2 \Pi, \Pi^e) \cong \Omega_{\Pi^e}^2 \Pi(1)$ in $\text{gr-CM } \Pi^e$.*

Proof. First note that since the Gorenstein dimension of Π is ≤ 1 , $\Omega_{\Pi^e} \Pi$ is Cohen-Macaulay by Lemma 2.5. Therefore $\Omega_{\Pi^e}^2(\Pi)$ is also Cohen-Macaulay and does not have any projective direct summands. Then we observe that

$$\begin{aligned} & \mathbf{R}\text{Hom}_{\Pi^e}(\Pi, \Pi^e)[4] \cong \Pi(1) \text{ in } \underline{\text{gr-CM}}(\Pi^e) \\ \iff & \mathbf{R}\text{Hom}_{\Pi^e}(\Pi[-2], \Pi^e) \cong \Pi[-2](1) \text{ in } \underline{\text{gr-CM}}(\Pi^e) \\ \iff & \text{Hom}_{\Pi^e}(\Omega_{\Pi^e}^2 \Pi, \Pi^e) \cong \Omega_{\Pi^e}^2 \Pi(1) \text{ in } \underline{\text{gr-CM}}(\Pi^e) \\ \iff & \text{Hom}_{\Pi^e}(\Omega_{\Pi^e}^2 \Pi, \Pi^e) \cong \Omega_{\Pi^e}^2 \Pi(1) \text{ in } \text{gr-CM}(\Pi^e). \end{aligned}$$

The last equivalence holds since both $\text{Hom}_{\Pi^e}(\Omega_{\Pi^e}^2 \Pi, \Pi^e)$ and $\Omega_{\Pi^e}^2 \Pi(1)$ are Cohen-Macaulay without projective summands. \square

Proof of Theorem 5.7. (1) holds by [KR07, Proposition 2.1].

By Theorem 5.6, there exists a quiver with potential (\bar{Q}, W) with an isomorphism $\Pi \cong \text{Jac}(\bar{Q}, W)$. We consider the graded version of the exact sequence in (5.1), and its terms by

$$\begin{aligned} P_2 &= \bigoplus_{a \in \bar{Q}_1} \Pi e_{s(a)} \otimes e_{t(a)} \Pi(-1 + \deg a), \\ P_1 &= \bigoplus_{a \in \bar{Q}_1} \Pi e_{t(a)} \otimes e_{s(a)} \Pi(-\deg a), \text{ and} \\ P_0 &= \bigoplus_{i \in \bar{Q}_0} \Pi e_i \otimes e_i \Pi(0). \end{aligned}$$

By Lemma 5.5, keeping track of the external grading, we have

$$(5.2) \quad \text{Hom}_{\Pi^e}(d_2, \Pi^e) = d_2(1).$$

We denote $(-)^{\vee} = \text{Hom}_{\Pi^e}(-, \Pi^e)$. Since the Gorenstein dimension of Π is ≤ 1 , the cokernel of d_2 is Cohen-Macaulay, and we have an isomorphism

$$(\text{Coker } d_2)^{\vee} \cong \text{Ker}(d_2^{\vee}).$$

Using (5.2) we get the following isomorphisms

$$\begin{aligned} (\Omega_{\Pi^e}^2 \Pi)^{\vee} &\cong (\text{Im } d_2)^{\vee} \cong \text{Coker}((\text{Coker } d_2)^{\vee}) \xrightarrow{\quad} P_1^{\vee} \\ &\cong \text{Coker}(\text{Ker}(d_2^{\vee})) \xrightarrow{\quad} P_1^{\vee} \\ &\cong \text{Coker}(\text{Ker}(d_2(1))) \xrightarrow{\quad} P_2(1) \\ &\cong \text{Im } d_2(1) \cong \Omega_{\Pi^e}^2 \Pi(1) \end{aligned}$$

Hence we get (2) by Lemma 5.8.

By (1), $\text{Ext}_{\Pi^e}^j(\Pi, \Pi^e)$ vanishes for $j \geq 2$.

Denote by N the maximal summand of $\Omega_{\Pi^e} \Pi$ without projective summands. The module N is Cohen-Macaulay and we have $\Omega_{\Pi^e} \Pi \cong N \oplus P$. The projective module P is a summand of P_1 and the induced map $P_2 \rightarrow P$ vanishes. Since the arrows of degree 1 correspond to minimal relations of Λ , and hence also to certain minimal relations of $T_{\Lambda} \text{Ext}_{\Lambda}^2(D\Lambda, \Lambda)$, P is generated in degree 0, that is $P \in \text{add } \Pi^e(0)$.

Now since Π is bimodule stably (1)-twisted 3-Calabi-Yau, we have isomorphisms

$$\text{Hom}_{\Pi^e}(N, \Pi^e) \cong \Omega_{\Pi^e}^2 N(1) \cong \Omega_{\Pi^e}^3 \Pi(1) \text{ in } \text{gr-CM } \Pi^e$$

The right isomorphism holds since $\Omega_{\Pi^e}^3 \Pi$ does not have any projective summands.

Now we have

$$\begin{aligned} \text{Ext}_{\text{gr-}\Pi^e}^1(\Pi, \Pi^e(i)) &\cong \text{Hom}_{\text{gr-}\Pi^e}(\Omega_{\Pi^e} \Pi, \Pi^e(i)) \\ &\cong (\text{Hom}_{\Pi^e}(\Omega_{\Pi^e} \Pi, \Pi^e))_i \\ &\cong (\text{Hom}_{\Pi^e}(N, \Pi^e))_i \oplus (\text{Hom}_{\Pi^e}(P, \Pi^e))_i \\ &\cong (\Omega_{\Pi^e}^3 \Pi)_{i+1} \oplus (\text{Hom}_{\Pi^e}(P, \Pi^e))_i \end{aligned}$$

Now if $i < 0$, $(\mathrm{Hom}_{\Pi^e}(P, \Pi^e))_i$ clearly vanishes since P , hence $\mathrm{Hom}_{\Pi^e}(P, \Pi^e)$ are generated in degree 0. Moreover since $\mathrm{gl.dim} \Lambda \leq 2$ the minimal projective resolution of Π is generated in strictly positive degrees from position 3 on, that is $(\Omega_{\Pi^e}^3 \Pi)_{\leq 0}$ vanishes and thus claim (3) here holds. \square

5.3. The $(d - 1)$ -representation finite case.

Theorem 5.9. *Let Λ be a $(d - 1)$ -representation finite algebra. Then its d -preprojective algebra Π is finite dimensional selfinjective and stably bimodule (1)-twisted d -Calabi-Yau. In particular, $\underline{\mathrm{mod}} \Pi$ is d -Calabi-Yau.*

Proof. Finite dimensionality and selfinjectivity are proved in [IO13, Corollary 3.4].

Let Λ be a $(d - 1)$ -representation-finite algebra. Then the category \mathcal{U} is a $(d - 1)$ -cluster tilting subcategory of $\mathrm{D}^b(\mathrm{mod} \Lambda)$, which satisfies $\mathcal{U} = \mathcal{U}[d - 1]$. Moreover the category $\mathrm{D}^b(\mathrm{mod} \Lambda)$ is an algebraic triangulated category, that is it is equivalent to the stable category of some Frobenius category. Hence we can apply Theorem 3.2 of [Dug12] and we obtain an isomorphism

$$\Omega_{\mathcal{U} \otimes \mathcal{U}^{\mathrm{op}}}^{d+1}(\mathrm{Hom}_{\mathcal{U}}(-, -)) \cong \mathrm{Hom}_{\mathcal{U}}(-, -[-d + 1])$$

in $\mathrm{mod}(\mathcal{U} \otimes \mathcal{U}^{\mathrm{op}})$ up to projective summands.

So we have the following isomorphisms in $\underline{\mathrm{gr}}\text{-}\Pi^e$:

$$\begin{aligned} \Omega_{\Pi^e}^{d+1}(\Pi) &\cong \bigoplus_{i \geq 0} \mathrm{Hom}_{\mathrm{D}^b(\mathrm{mod} \Lambda)}(\Lambda, \mathbb{S}_{d-1}^{-i} \Lambda[-d + 1]) \\ &\cong \bigoplus_{i \geq 0} \mathrm{Hom}_{\mathrm{D}^b(\mathrm{mod} \Lambda)}(\mathbb{S} \Lambda, \mathbb{S}_{d-1}^{-i+1} \Lambda) \end{aligned}$$

On the other hand

$$\begin{aligned} \mathrm{Hom}_{\Pi^e}(\Pi, \Pi^e) &\cong \mathrm{Hom}_{\Pi}(D\Pi, \Pi) \\ &\cong \bigoplus_{i \in \mathbb{Z}} \mathrm{Hom}_{\mathrm{D}^b(\mathrm{mod} \Lambda)}(D\Lambda, \mathbb{S}_{d-1}^{-i} \Lambda). \end{aligned}$$

Combining these two isomorphisms, and observing that $\mathbb{S}\Lambda = D\Lambda$, we obtain $\Omega_{\Pi^e}^{d+1}(\Pi) \cong \mathrm{Hom}_{\Pi^e}(\Pi, \Pi^e)(-1)$ in $\underline{\mathrm{gr}}\text{-}\Pi^e$. Since Π is selfinjective, Π is Cohen-Macaulay, so Π is bimodule stably (1)-twisted d -Calabi-Yau by Observation 3.2. \square

Remark 5.10. Let Λ be an algebra which is τ_{d-1} -finite and satisfying the vosnex property. In [IO13], the authors prove that the algebra $\Pi = \Pi_d(\Lambda)$ is Gorenstein of dimension ≤ 1 and that the stable category $\underline{\mathrm{CM}} \Pi$ is d -Calabi-Yau (Theorem 1.2(1)). It follows from the proof (see [IO13, Theorem 5.11]) that moreover the category of graded Cohen-Macaulay modules $\underline{\mathrm{gr}}\text{-}\underline{\mathrm{CM}} \Pi$ is (1)-twisted d -Calabi-Yau.

Unfortunately, it is not clear from the proof that this Calabi-Yau property comes from a bimodule property. So we cannot use the results in [IO13] to prove a result similar to Theorem 5.9 in the case where the algebra Π has Gorenstein dimension 1.

REFERENCES

- [Ami08] C. Amiot, *Sur les petites catégories triangulées*, Ph.D. thesis (2008), <http://www-fourier.ujf-grenoble.fr/~amiot/these.pdf>.
- [Ami09] C. Amiot, *Cluster categories for algebras of global dimension 2 and quivers with potential*, *Ann. Inst. Fourier (Grenoble)*, **59** (2009), 2525–2590. MR 2640929.
- [Buc87] R. O. Buchweitz, *Maximal Cohen-Macaulay modules and Tate cohomology over Gorenstein rings*, unpublished manuscript, 155 pages, 1987.
- [DWZ08] H. Derksen, J. Weyman, and A. Zelevinsky, *Quivers with potentials and their representations. I. Mutations*, *Selecta Math. (N.S.)* **14** (2008), no. 1, 59–119.
- [Dug12] A. Dugas, *Periodicity of d -cluster-tilted algebras.*, *J. Algebra* **368** (2012), 40–52.
- [ES98] K. Erdmann and N. Snashall, *On Hochschild cohomology of preprojective algebras I*, *Journal of Algebra* **205** (2), (1998), 391–412.
- [Guo11] L. Guo, *Cluster tilting objects in generalized higher cluster categories*, *J. Pure Appl. Algebra* **215** (2011), no 9, 2055–2071.
- [Gin06] V. Ginzburg, *Calabi-Yau algebras*, arXiv:math/0612139.
- [Hap88] D. Happel, *Triangulated categories in the representation theory of finite-dimensional algebras*, Cambridge University Press, Cambridge 1988.
- [HIO12] M. Herschend, O. Iyama and S. Oppermann, *n -representation infinite algebras*, arXiv:1205.1272, to appear in *Adv. Math.*
- [Iya11] O. Iyama, *Cluster-tilting for higher Auslander algebras*, *Adv. Math.* **226** (2011), no. 1, 1–61.
- [IY08] O. Iyama, Y. Yoshino, *Mutation in triangulated categories and rigid Cohen-Macaulay modules*, *Invent. Math.* **172** (2008), no. 1, 117–168.
- [IO11] O. Iyama, S. Oppermann, *n -representation finite algebras and n -APR tilting*, *Trans. Amer. Math. Soc.* **363** (2011), no. 12, 6575–6614.
- [IO13] O. Iyama, S. Oppermann, *Stable categories of higher preprojective algebras*, *Adv. Math.* **244** (2013), 23–68.
- [Kel08] B. Keller, *Calabi-Yau triangulated categories*, *Trends in representation theory of algebras and related topics*, 467–489, EMS Ser. Congr. Rep., Eur. Math. Soc., Zurich, 2008.
- [Kel11] B. Keller, *Deformed Calabi-Yau completions*, with appendix by Michel Van den Bergh, *J. Reine Angew. Math.* **654** (2011), 125–180.
- [KV87] B. Keller and Dieter Vossieck, *Sous les catégories dérivées (French) [Beneath the derived categories]* *C. R. Acad. Sci., Paris Sér. I Math.* **305** (1987), no. 6, 225–228.
- [KR07] B. Keller, I. Reiten, *Cluster-tilted algebras are Gorenstein and stably Calabi-Yau*, *Adv. Math.* **211** (2007), no. 1, 123–151.
- [Orl05] D. Orlov, *Derived categories of coherent sheaves and triangulated categories of singularities*, *Algebra, arithmetic, and geometry: in honor of Yu. I. Manin. Vol. II*, 503–531, *Progr. Math.* **270**, Birkhauser Boston, Inc., Boston, MA, 2009.
- [Ric89] J. Rickard, *Derived categories and stable equivalence*, *J. Pure Appl. Algebra*, **61** (1989), no. 3, 303317.
- [Rin98] Ringel, C. M. *The preprojective algebra of a quiver*. (English summary) *Algebras and modules, II (Geiranger, 1996)*, 467–480, *CMS Conf. Proc.* **24**, Amer. Math. Soc., Providence, RI, 1998.
- [Scho] Schofield, A. *Wild algebras with periodic Auslander-Reiten translate*, unpublished manuscript.
- [Yos90] Y. Yoshino, *Cohen-Macaulay modules over Cohen-Macaulay rings*, *London Mathematical Society Lecture Note Series*, **146**. Cambridge University Press, Cambridge, 1990.

INSTITUT FOURIER, 100 RUE DES MATHÉMATIQUES, 38402 SAINT MARTIN D’HÈRES, FRANCE

E-mail address: Claire.Amiot@ujf-grenoble.fr

INSTITUTT FOR MATEMATISKE FAG, NTNU, 7491 TRONDHEIM, NORWAY

E-mail address: steffen.oppermann@math.ntnu.no