



HAL
open science

Energy, decay rate, and effective masses for a moving polaron in a Fermi sea: Explicit results in the weakly attractive limit

Christian Trefzger, Yvan Castin

► **To cite this version:**

Christian Trefzger, Yvan Castin. Energy, decay rate, and effective masses for a moving polaron in a Fermi sea: Explicit results in the weakly attractive limit. 2013. hal-00847562v2

HAL Id: hal-00847562

<https://hal.science/hal-00847562v2>

Preprint submitted on 31 Oct 2013 (v2), last revised 20 Dec 2013 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Energy, decay rate, and effective masses for a moving polaron in a Fermi sea: Explicit results in the weakly attractive limit

CHRISTIAN TREFZGER AND YVAN CASTIN

Laboratoire Kastler Brossel, École Normale Supérieure and CNRS, UPMC, 24 rue Lhomond, 75231 Paris, France

PACS 03.75.Ss – Degenerate Fermi gases

Abstract – We study the properties of an impurity of mass M moving through a spatially homogeneous three-dimensional fully polarized Fermi gas of particles of mass m . In the weakly attractive limit, where the effective coupling constant $g \rightarrow 0^-$ and perturbation theory can be used, both for a broad and a narrow Feshbach resonance, we obtain an explicit analytical expression for the complex energy $\Delta E(\mathbf{K})$ of the moving impurity up to order two included in g . This also gives access to its longitudinal and transverse effective masses $m_{\parallel}^*(\mathbf{K})$, $m_{\perp}^*(\mathbf{K})$, as functions of the impurity wave vector \mathbf{K} . Depending on the modulus of \mathbf{K} and on the impurity-to-fermion mass ratio M/m we identify four regions separated by singularities in derivatives with respect to \mathbf{K} of the second-order term of $\Delta E(\mathbf{K})$, and we discuss the physical origin of these regions. Remarkably, the second-order term of $m_{\parallel}^*(\mathbf{K})$ presents points of non-differentiability, replaced by a logarithmic divergence for $M = m$, when \mathbf{K} is on the Fermi surface of the fermions. We also discuss the third-order contribution and relevance for cold atom experiments.

Introduction. – Recent cold atom experiments have reached an unprecedented accuracy in measuring the equation of state of an interacting Fermi gas [1–4]. This has allowed to confirm that in strongly spin-polarized configurations the minority atoms dressed by the Fermi sea of the majority atoms form a normal gas of quasiparticles called Fermi polarons [5–7].

While the problem of a single impurity at rest in a Fermi sea has been thoroughly studied [5–13], previous works on a moving impurity have focused only on its decay rate [10, 14]. However, the real part of the polaronic complex energy is also of theoretical interest and is experimentally accessible [15]. Whereas most theories for the impurity at rest successfully used a variational ansatz [6], this ansatz is not reliable anymore for a moving polaron with momentum $\hbar\mathbf{K}$ since at low K it wrongly predicts a zero decay rate in contrast to the K^4 -law found in [10, 14].

In this work, we focus on the weakly attractive regime $k_F a \rightarrow 0^-$, where a is the s -wave scattering length of a minority atom and a fermion and k_F is the Fermi wave number of the majority atoms. Using a systematic expansion in powers of $k_F a$, we go beyond the Fermi liquid description of [14]: We determine not only the decay rate, but also the real part of the polaronic complex energy, and being not restricted to low momenta, we have access to momentum-dependent effective masses of the polaron.

Within our microscopic approach we have a complete description of the system: We determine regions of parameters separated by singularities in derivatives of the second-order term of the polaronic complex energy, as shown in Fig.1. We discuss the physical origin of the singularities and the relevance for current cold atom experiments.

The model. – At zero temperature, we consider in three dimensions an ideal Fermi gas of particles of same spin state and mass m perturbed by the presence of a moving impurity of mass M and momentum $\hbar\mathbf{K}$. While we assume no interactions among the fermions (contrary to [16]), the impurity interacts with each fermion through a s -wave interaction of negligible range. The system is enclosed in a quantization volume \mathcal{V} with periodic boundary conditions, and is described by the Hamiltonian $\hat{H} = \hat{H}_0 + \hat{V}$:

$$\hat{H}_0 = \sum_{\mathbf{k}} \left(\varepsilon_{\mathbf{k}} \hat{u}_{\mathbf{k}}^{\dagger} \hat{u}_{\mathbf{k}} + E_{\mathbf{k}} \hat{d}_{\mathbf{k}}^{\dagger} \hat{d}_{\mathbf{k}} \right), \quad (1)$$

$$\hat{V} = \frac{g_0}{\mathcal{V}} \sum_{\substack{\mathbf{k}_1, \mathbf{k}_2 \\ \mathbf{k}_3, \mathbf{k}_4}} \delta_{\mathbf{k}_1 + \mathbf{k}_2, \mathbf{k}_3 + \mathbf{k}_4}^{\text{mod}} \hat{d}_{\mathbf{k}_4}^{\dagger} \hat{u}_{\mathbf{k}_3}^{\dagger} \hat{u}_{\mathbf{k}_2} \hat{d}_{\mathbf{k}_1}, \quad (2)$$

where $\hat{u}_{\mathbf{k}}$ ($\hat{d}_{\mathbf{k}}$) annihilates a fermion (impurity) with wave vector \mathbf{k} and kinetic energy $\varepsilon_{\mathbf{k}} = \frac{\hbar^2 k^2}{2m}$ ($E_{\mathbf{k}} = \frac{\hbar^2 k^2}{2M}$). To avoid ultraviolet divergences we use a standard lat-




Fig. 1: Regions of the moving polaron in the $M/m - K/k_F$ plane, where the second-order correction to the complex energy is a smooth function of K/k_F . On the dashed line, $K = k_F$, and on the solid line, $K = k_F M/m$, a logarithmic divergence (discontinuity) appears in the n th order derivative with respect to K of the real part (imaginary part) of the second-order term of the polaronic complex energy (9): $n = 3$ for the dashed line, $n = 4$ for the solid line, $n = 2$ at the point $(M = m, K = k_F)$.

tice model where the positions of the particles are discretized on a cubic lattice of elementary step $b \ll k_F^{-1}$ and binary interactions between the impurity and a fermion take place at the same lattice site with a bare coupling constant g_0 [17]. In momentum space this takes the form Eq. (2). The model provides automatically a cutoff in the Fourier space since wave vectors are restricted to the first Brillouin zone. In Eq. (2), the modified Kronecker delta is to be understood modulo a vector of the reciprocal lattice and ensures conservation of the quasi-momentum. The bare coupling constant g_0 is then related to the scattering length a through the formula

$$\frac{1}{g_0} = \frac{1}{g} - \int_{\text{FBZ}} \frac{d^3 k}{(2\pi)^3} \frac{2\mu}{\hbar^2 k^2} \quad \text{and} \quad g = \frac{2\pi\hbar^2 a}{\mu}, \quad (3)$$

where the integral is taken in the first Brillouin zone, $\text{FBZ} = [-\pi/b, \pi/b]^3$, and $\mu = mM/(m+M)$ is the reduced mass. In practice, at the end of the perturbative calculations to come, we will take the thermodynamic limit, where $\mathcal{V} \rightarrow +\infty$, and also the continuous space limit $b \rightarrow 0^+$, so that the first Brillouin zone will converge to the whole Fourier space.

The goal is to calculate the complex energy of an impurity of wave vector \mathbf{K} moving in the zero temperature Fermi gas of N particles. The ground state of the N fermions is the usual Fermi sea [FS] of energy $E_{\text{FS}}(N)$. In the absence of interactions the ground state of the system is given by the free impurity immersed in the Fermi sea

$$|\psi^{(0)}(\mathbf{K})\rangle = \hat{d}_{\mathbf{K}}^\dagger |\text{FS}\rangle, \quad (4)$$

with an energy, measured with respect to the energy of

the Fermi sea, given by

$$\Delta E^{(0)}(\mathbf{K}) = \langle \psi^{(0)}(\mathbf{K}) | \hat{H}_0 | \psi^{(0)}(\mathbf{K}) \rangle - E_{\text{FS}}(N) = E_{\mathbf{K}}. \quad (5)$$

We now consider the weakly interacting case $a \rightarrow 0^-$, where a quasiparticle polaronic ground state is known to exist [5–7, 11]. For a fixed lattice spacing b , in the limit where $|a| \ll b$ we can Taylor expand the bare coupling constant (3):

$$g_0 \underset{g \rightarrow 0^-}{=} g + g^2 \int_{\text{FBZ}} \frac{d^3 k}{(2\pi)^3} \frac{2\mu}{\hbar^2 k^2} + O(g^3), \quad (6)$$

and in this weakly attractive limit one can treat \hat{V} in the Hamiltonian with usual perturbation theory. Up to second order in \hat{V} , we find in the thermodynamic limit that the energy of the moving polaron is complex:

$$\Delta E(\mathbf{K}) \underset{a \rightarrow 0^-}{=} E_{\mathbf{K}} + \rho g_0 - \int'_{\text{FBZ}^2} \frac{d^3 k d^3 q}{(2\pi)^6} \frac{g_0^2}{F_{\mathbf{k},\mathbf{q}}(\mathbf{K})} + O(g_0^3), \quad (7)$$

where the prime on the integral means that it is restricted to $0 < q < k_F$ and to $k > k_F$, and we have introduced the mean density of the Fermi sea $\rho = N/\mathcal{V}$ related to the Fermi wave number k_F by the usual relation $k_F = (6\pi^2 \rho)^{1/3}$, $E_F = \frac{\hbar^2 k_F^2}{2m}$, and we have defined

$$F_{\mathbf{k},\mathbf{q}}(\mathbf{K}) = E_{\mathbf{K}+\mathbf{q}-\mathbf{k}} + \varepsilon_{\mathbf{k}} - \varepsilon_{\mathbf{q}} - E_{\mathbf{K}} - i\eta, \quad (8)$$

with $\eta \rightarrow 0^+$. Substituting Eq. (6) into Eq. (7) and keeping only terms up to order g^2 , as announced we get a finite expression for a vanishing lattice spacing

$$\Delta E(\mathbf{K}) \underset{g \rightarrow 0^-}{=} E_{\mathbf{K}} + \rho g + \frac{(\rho g)^2}{E_F} f(K/k_F) + O(g^3). \quad (9)$$

The dimensionless complex function $f(K/k_F)$ is defined as follows:

$$f\left(\frac{K}{k_F}\right) = \frac{E_F}{\rho^2} \int' \frac{d^3 q}{(2\pi)^3} \int \frac{d^3 k}{(2\pi)^3} \left[\frac{2\mu}{\hbar^2 k^2} - \frac{Y(k - k_F)}{F_{\mathbf{k},\mathbf{q}}(\mathbf{K})} \right], \quad (10)$$

where $Y(x)$ is the Heaviside function, and the integral over \mathbf{k} is now taken over the whole Fourier space.

Results. – The integral in Eq. (10) can be calculated exactly, as it will be detailed elsewhere [18]. Using the Dirac relation $\lim_{\eta \rightarrow 0^+} (x - i\eta)^{-1} = P \frac{1}{x} + i\pi\delta(x)$, where P is the principal value and $\delta(x)$ the Dirac distribution, we can separate the integral into its real and imaginary parts.

The imaginary part of Eq. (9) reveals that due to scattering of the impurity with the fermions, a moving polaron radiates particle-hole pairs [14] and thus decays out of its initial momentum channel, at a rate Γ_0 given by

$$\frac{\hbar\Gamma_0}{2} \underset{g \rightarrow 0^-}{=} -\frac{(\rho g)^2}{E_F} \Im f(K/k_F) + O(g^3). \quad (11)$$

This polaron decay rate coincides with the Fermi golden




Fig. 2: (Color online) Function $-\Im f(\kappa)$ entering the imaginary part of Eq. (9) (row 1), and its second derivative entering the imaginary part of the inverse effective mass Eq. (20) (row 2), for various mass ratios. Dashed lines correspond to the asymptote of Eq. (15). Red vertical lines correspond to the boundaries at $\kappa = 1$ and at $\kappa = M/m$.

rule result, as best seen from Eq. (7). It is an isotropic function of the impurity wave vector \mathbf{K} . Depending on the impurity-to-fermion mass ratio $r = M/m$ and on the modulus of \mathbf{K} , $\Im f(K/k_F)$ assumes different piecewise smooth (indefinitely derivable) functional forms on the four different regions represented in Fig. 1. (i) The low- K region, $0 < K/k_F < \min(1, r)$, where

$$-\Im f(\kappa) \underset{\text{region 1}}{=} \frac{3\pi}{20r} \kappa^4. \quad (12)$$

As expected, $\Im f(0) = 0$. (ii) The $r < 1$ (light impurity) crossover region, $\min(1, r) < K/k_F < \max(1, r)$, where

$$-\Im f(\kappa) \underset{\text{region 2}}{=} \frac{3\pi r}{20(r^2 - 1)^2} \times [(r^2 - 2)\kappa^4 + 10\kappa^2 - 20r\kappa + 15r^2 - 4r^3/\kappa]. \quad (13)$$

(iii) The $r > 1$ (heavy impurity) crossover region, $\min(1, r) < K/k_F < \max(1, r)$, where

$$-\Im f(\kappa) \underset{\text{region 3}}{=} \frac{3\pi r}{20(r^2 - 1)^2} \times \left[\frac{\kappa^4}{r^2} - 10\kappa^2 + 10(r^2 + 1)\kappa - 15r^2 + \frac{6r^2 - 2}{\kappa} \right]. \quad (14)$$

There is no crossover region for a unit impurity-to-fermion mass ratio, $r = 1$. (iv) Finally the high- K region, $\max(1, r) < K/k_F$, where

$$-\Im f(\kappa) \underset{\text{region 4}}{=} \frac{3\pi r}{10(r+1)^2} \frac{5\kappa^2 - 2r - 1}{\kappa}. \quad (15)$$

Eqs. (12,15) generalize to $r \neq 1$ the results of [10]. Eq. (12) reproduces the Fermi liquid theory result [14] obtained at




Fig. 3: (Color online) Function $\Re f(\kappa)$ entering the real part of Eq. (9) (row 1), and its second derivative entering the real part of the inverse effective mass Eq. (20) (row 2), for various mass ratios. Dashed lines: Asymptotic behavior (18). Red vertical lines: Boundaries at $\kappa = 1$ and at $\kappa = M/m$. Insets: Enlargements showing the non-differentiability of the second-order term of $M/m_{\parallel}^*(\mathbf{K})$ at $K = k_F$.

vanishing κ , for a vanishing density of impurities. In Fig. 2 we plot the function $-\Im f(\kappa)$ for different values of the mass ratio between the impurity and the fermions.

For a large impurity wave vector, $K/k_F \gg \max(1, r)$, the asymptotic value of $-\Im f(\kappa)$ is given by the asymptotic expansion of Eq. (15), and is linear in K . This can be quantitatively recovered with a simple kinetic argument: For $K \gg (r+1)k_F$, due to conservation of energy and momentum, the fermionic wave number k after scattering with the impurity is $\gg k_F$ except in a small cone in the forward direction, so that the Pauli blocking effect may be neglected. Also the impurity-to-fermion relative velocity is $v \simeq \hbar K/M$, which leads to a collision rate $\Gamma_0 \simeq \rho \sigma v$, where ρ is the fermionic density and $\sigma = 4\pi a^2$ is impurity-to-fermion scattering cross section.

Let us now discuss the real part of the polaronic complex energy (9) for which much less was known. We find that $\Re f(K/k_F)$ is an isotropic function of \mathbf{K} with the following analytic expression

$$\begin{aligned} \Re f(\kappa) = & \frac{3r}{r+1} - \frac{3r}{20(r^2-1)^2} \left\{ \frac{(r^2-1)}{2} (\kappa^2 + 20r - 9) \right. \\ & + \frac{(\kappa-1)^3}{\kappa} [\kappa(\kappa+3)(r^2-2) + 6r^2 - 2] \ln \left| \frac{\kappa-1}{\kappa} \right| \\ & \left. + \frac{(\kappa-r)^4}{\kappa} \left(\frac{\kappa+4r}{r^2} \right) \ln \left| \frac{\kappa-r}{\kappa} \right| + (\kappa \rightarrow -\kappa) \right\}. \quad (16) \end{aligned}$$

This function is plotted in Fig. 3 for different values of the impurity-to-fermion mass ratio r . The locus of points where the moduli in the argument of the \ln functions vanish forms the boundaries of the same physical regions dis-

cussed for $\Im f(K/k_F)$, and sketched in Fig. 1. We have numerically observed that Eq. (16) is a monotonically decreasing function of κ . It has a vanishing derivative at the origin, $\Re f'(0) = 0$, where it reaches its maximum value

$$\Re f(0) = \frac{9r(1-r^2+2r^2 \ln r)}{4(r^2-1)^2}. \quad (17)$$

An expression equivalent to (17) was obtained in [8, 9] in the context of a Λ -particle in a Fermi sea of nucleons with $\nu = 4$ spin-isospin states, hence an extra factor ν in those references. Eq. (16) vanishes at large K/k_F :

$$\Re f(\kappa) \underset{\kappa \gg 1+r}{=} \frac{r}{4\kappa^2} + O(1/\kappa)^4. \quad (18)$$

This indicates that, for large K/k_F and up to second-order in g , the dispersive effect of the Fermi sea on the energy of the impurity reduces to the mean-field term ρg . Remarkably, Eq. (16) has a finite limit for a unit mass ratio

$$\lim_{r \rightarrow 1} \Re f(\kappa) = \frac{3}{2} - \frac{3}{40} \left[\frac{11}{2} + \kappa^2 + (2\kappa^3 + 4\kappa^2 + 6\kappa + 3) \times \frac{(\kappa-1)^2}{\kappa} \ln |\kappa-1| - 2\kappa^4 \ln |\kappa| + (\kappa \rightarrow -\kappa) \right], \quad (19)$$

and as well as for $\Im f(K/k_F)$ the crossover region is absent in this special case $M = m$. In the weakly attractive limit, $g \rightarrow 0^-$, the second-order effect of the Fermi sea on the energy of the impurity moving through it can then be summarized as follows for an arbitrary mass ratio: On the real (imaginary) part of the resulting complex energy the effect of the medium is maximal (minimal) when the impurity is at rest, $\mathbf{K} = \mathbf{0}$, while the effect of the medium is minimal (maximal) when the impurity is moving fast.

A related observable of interest is the effective mass of the polaron. Since the impurity is moving with momentum $\hbar \mathbf{K}$, one can define the effective mass $m_{\parallel}^*(\mathbf{K})$ along the direction of \mathbf{K} , and the effective mass $m_{\perp}^*(\mathbf{K})$ along the direction perpendicular to \mathbf{K} [19]. $m_{\parallel}^*(\mathbf{K})$ is related to the second derivative of the complex energy Eq. (9) with respect to the impurity wave number $K = \kappa k_F$ as follows:

$$\frac{M}{m_{\parallel}^*(\mathbf{K})} \underset{g \rightarrow 0^-}{=} 1 + \frac{r}{2} \left(\frac{\rho g}{E_F} \right)^2 f''(\kappa) + O(g^3), \quad (20)$$

and $m_{\perp}^*(\mathbf{K})$ is related to the first derivative:

$$\frac{M}{m_{\perp}^*(\mathbf{K})} \underset{g \rightarrow 0^-}{=} 1 + \frac{r}{2} \left(\frac{\rho g}{E_F} \right)^2 \frac{f'(\kappa)}{\kappa} + O(g^3). \quad (21)$$

When $K \rightarrow 0$, Eqs. (20, 21) have the same limit, which coincides with the result of [10], and one recovers the usual rotational symmetry of the effective mass tensor. At non-zero K , Eq. (21) is a differentiable function of K , whereas Eq. (20) presents interesting singularities and is non-differentiable in $\kappa = 1$ as shown in Figs. 2, 3. Furthermore, for $r = 1$, $\Re f''(\kappa)$ presents a logarithmic

divergence $\propto \ln |\kappa - 1|$ in $\kappa = 1$, as can be seen in Eq. (19). This suggests that for $r = 1$, $M/m_{\parallel}^*(\mathbf{K})$ cannot be Taylor expanded in powers of g at the point $\kappa = 1$, as e.g. the function $g^2 \ln |g|$, and a non-perturbative approach must be used [18]. Instead, $\Im f''(\kappa)$ has a jump $J = \Im f''(1^+) - \Im f''(1^-) = 9\pi/4$ in $\kappa = 1$ for $r = 1$.

Physical interpretation. – We now interpret the boundaries between regions in Fig. 1. Intuitively, singularities in the polaron complex energy may originate from the fact that (i) the energy denominator in Eq. (7) can vanish, and (ii) the domains of variation of \mathbf{k} and \mathbf{q} in Eq. (7) have sharp boundaries $k = q = k_F$. One thus investigates the form of the Fourier space domain where the energy denominator vanishes for \mathbf{k} and \mathbf{q} both at the Fermi surface. For $K < k_F$, it is found that this domain supports all values of $\theta \in [0, \pi/2]$, where θ is the angle between \mathbf{K} and $\mathbf{k} - \mathbf{q}$. For $K > k_F$, on the contrary, the allowed values of θ range from $\arccos(k_F/K)$ to $\pi/2$. This designates $K = k_F$ as a first peculiar line in the plane $M/m - K/k_F$. Note that $\pi - 2\theta$ is the scattering angle of an incoming impurity of wave vector \mathbf{K} on the Fermi sea when both the resulting scattered fermion and hole are at the Fermi surface. To obtain the second peculiar line in Fig. 1, we introduce the additional idea that the energy denominator as well as its first order derivatives with respect to \mathbf{k} and to \mathbf{q} vanish at $k = q = k_F$, which indeed leads to $K = rk_F$.

Narrow Feshbach resonance. – In a recent experiment [15] impurities of ^{40}K were immersed in a Fermi sea of ^6Li , where the s -wave interaction between the impurity and each fermion has a non-negligible range due to the presence of a narrow Feshbach resonance. Such a situation differs from the one we have considered so far: A new length appears in the system, namely the Feshbach length R_* , which at the experimental density [15] was of the order of $R_* \simeq k_F^{-1}$ and thus not negligible. This physical situation can be described by a two-channel model (see e.g. [21] and references therein). For simplicity we keep in the two-channel model the same lattice model as for the single-channel Hamiltonian (1,2). The Hamiltonian of the system then becomes $\hat{\mathcal{H}} = \hat{\mathcal{H}}_0 + \hat{W} + E_{\text{mol}} \hat{N}_{\text{mol}}$, with

$$\hat{\mathcal{H}}_0 = \hat{H}_0 + \sum_{\mathbf{k}} \frac{\varepsilon_{\mathbf{k}}}{1+r} \hat{b}_{\mathbf{k}}^\dagger \hat{b}_{\mathbf{k}}, \quad (22)$$

$$\hat{W} = \frac{\Lambda}{\sqrt{V}} \sum_{\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3} \delta_{\mathbf{k}_1 + \mathbf{k}_2, \mathbf{k}_3}^{\text{mod}} \left(\hat{b}_{\mathbf{k}_3}^\dagger \hat{u}_{\mathbf{k}_2} \hat{d}_{\mathbf{k}_1} + \text{H.c.} \right) \quad (23)$$

and $\hat{N}_{\text{mol}} = \sum_{\mathbf{k}} \hat{b}_{\mathbf{k}}^\dagger \hat{b}_{\mathbf{k}}$ is the number of closed-channel molecules operator. Here $\hat{b}_{\mathbf{k}}$ annihilates a closed-channel molecule with wave vector \mathbf{k} , kinetic energy $\hbar^2 k^2 / [2(M+m)] = \varepsilon_{\mathbf{k}} / (1+r)$ and internal energy E_{mol} . The relation between E_{mol} , the interchannel coupling Λ , the Feshbach length R_* and the s -wave scattering length a is

$$E_{\text{mol}} = -\frac{\Lambda^2}{g_0} \quad \text{and} \quad R_* = \frac{\pi \hbar^4}{\Lambda^2 \mu^2}, \quad (24)$$

where g_0 is still given by Eq. (3). For a fixed lattice spacing b and Feshbach length R_* , one sees that in the weakly attractive limit $a \rightarrow 0^-$ the energy of the closed-channel molecule diverges, $E_{\text{mol}} \rightarrow +\infty$. Therefore, in such a limit we introduce the hermitian projector \hat{P} onto the low-energy subspace in which closed-channel molecules are absent, and the projector $\hat{Q} = \hat{1} - \hat{P}$. Within this low-energy subspace one can derive an exact effective Hamiltonian depending on a complex energy z [20]:

$$\hat{\mathcal{H}}_{\text{eff}}(z) = \hat{P}\hat{\mathcal{H}}\hat{P} + \hat{P}\hat{W}\hat{Q} \frac{\hat{Q}}{z\hat{Q} - \hat{Q}\hat{\mathcal{H}}\hat{Q}} \hat{Q}\hat{W}\hat{P}. \quad (25)$$

The complex energy $E_{\text{FS}}(N) + \Delta E$ of the impurity must then be (in the thermodynamic limit and under appropriate analytic continuation) an eigenvalue of $\hat{\mathcal{H}}_{\text{eff}}(E_{\text{FS}}(N) + \Delta E)$, which constitutes an implicit equation. Here we solve it perturbatively up to order g_0^2 by making an asymptotic expansion of (25) in powers of E_{mol} to obtain in the subspace of momentum $\hbar\mathbf{K}$:

$$\begin{aligned} \hat{\mathcal{H}}_{\text{eff}}(E_{\text{FS}}(N) + \Delta E) &= \hat{P}\hat{H}\hat{P} \\ &- \frac{1}{E_{\text{mol}}^2} \hat{P}\hat{W}\hat{Q}[E_{\mathbf{K}} + E_{\text{FS}}(N) - \hat{\mathcal{H}}_0]\hat{Q}\hat{W}\hat{P} + O(g_0^3), \end{aligned} \quad (26)$$

where we used $-E_{\text{mol}}^{-1}\hat{P}\hat{W}\hat{Q}\hat{W}\hat{P} = \hat{P}\hat{V}\hat{P}$ and the fact that $\Delta E = E_{\mathbf{K}} + O(g_0)$. Remarkably, one recovers as the first term the single-channel model Hamiltonian (1,2) modified by the second term of (26), which is an effective-range correction proportional to R_* , that vanishes, as it should, when $R_* \rightarrow 0$. It remains to apply the usual second-order perturbation theory in g_0 to the effective Hamiltonian (26). This amounts to adding to the already calculated single channel result the contribution of the second term of (26) simply treated to first order in perturbation theory, which is real since this term is hermitian. After thermodynamic and zero-lattice spacing limits we obtain:

$$\Delta E(\mathbf{K}) \underset{g \rightarrow 0^-}{=} E_{\mathbf{K}} + \rho g + \frac{(\rho g)^2}{E_{\text{F}}} \tilde{f}(K/k_{\text{F}}) + O(g^3), \quad (27)$$

where

$$\tilde{f}(\kappa) = f(\kappa) - \frac{3\pi}{2} \frac{r}{(1+r)^3} \left(\kappa^2 + \frac{3r^2}{5} \right) k_{\text{F}} R_*. \quad (28)$$

The correction to $f(\kappa)$ is a smooth function of κ so that the regions of Fig. 1 remain unchanged for \tilde{f} .

At this point a comparison with the hard-sphere result [10] at $K = 0$ is possible with some care. First, one formally identifies the effective range $r_e = -2R_*$ of the two-channel model with the one $2a/3$ of the hard-sphere model, which turns the second term of (28) into an energy correction of order a^3 . Second, this energy correction must appear in the theory of [10] as a single T -matrix vertex, that is in the contribution ϵ_1 , see Eq. (3.5) of [10]. Third, one must include the fact that the hard-sphere interaction

also scatters in the p -wave channel, with a scattering volume $\mathcal{V}_s = a^3/3$ for the convention of [22], which leads to an energy shift given by first order perturbation theory applied to the pseudo-potential of [23]. Then we find consistency between our result and the one of [10] at $K = 0$.

Experimental feasibility. – To determine the experimental relevance of the second-order correction to the polaronic complex energy predicted in this work we refer to the experimental conditions [15], where for $k_{\text{F}}|a| < 2$ the real part (imaginary part) of the polaronic complex energy can be measured with a precision of $5 \times 10^{-3}E_{\text{F}}$ ($10^{-4}E_{\text{F}}$). Then our theory for a moving polaron may be tested experimentally for a finite scattering length when two conditions on $k_{\text{F}}|a|$ hold simultaneously: (i) the second-order correction in Eq. (27) must be sufficiently large to be observed, (ii) the third-order correction, which is neglected in Eq. (27), must be sufficiently small. To test the second condition one has to calculate the third-order correction $\Delta E^{(3)}$ to Eq. (27). Extending the perturbative method used to derive Eq. (26) up to order g_0^3 we obtain

$$\begin{aligned} \Delta E^{(3)}(\mathbf{K}) &= g^3 \int' \frac{d^3q}{(2\pi)^3} \left\{ \frac{1}{\Lambda^2} \left(E_{\mathbf{K}} + \varepsilon_{\mathbf{q}} - \frac{\varepsilon_{\mathbf{K}+\mathbf{q}}}{1+r} \right) \right. \\ &- \int \frac{d^3k}{(2\pi)^3} \left[\frac{2\mu}{\hbar^2 k^2} - \frac{Y(k - k_{\text{F}})}{F_{\mathbf{k},\mathbf{q}}(\mathbf{K})} \right] \left. \right\}^2 - g^3 \frac{\rho^2}{\Lambda^2} \\ &- g^3 \int \frac{d^3k}{(2\pi)^3} \left[\int' \frac{d^3q}{(2\pi)^3} \frac{Y(k - k_{\text{F}})}{F_{\mathbf{k},\mathbf{q}}(\mathbf{K})} \right]^2, \end{aligned} \quad (29)$$

with $F_{\mathbf{k},\mathbf{q}}$ defined in Eq. (8). The first (third) contribution originate from the subspace with one hole and two particles (one particle and two holes), and the second contribution originates from a conspiracy between the mean-field and the effective-range terms within the subspace with one hole. At $\mathbf{K} = \mathbf{0}$ and $R_* = O(a)$, Eq. (29) reproduces [10].

We have performed explicit calculations for $k_{\text{F}}a = -0.46$, which is intermediate between weakly and strongly interacting regime: In Fig. 4b we show that, for the real part of the complex energy, the condition of experimental observability is indeed satisfied over the useful range $0 \leq K \leq 2k_{\text{F}}$, with a reasonably small correction due to $\Re\Delta E^{(3)}(\mathbf{K})$; we further test the real part of our perturbative result up to $n = 3$ (solid line) against the real part resulting from the single particle-hole ansatz (12) of [21] (circles), showing reasonable agreement. Instead for the imaginary part in Fig. 4a, while $\Im\Delta E^{(2)}(\mathbf{K})$ is measurable within the experimental precision [15] for $K > 0.5k_{\text{F}}$, the correction due to $\Im\Delta E^{(3)}(\mathbf{K})$ is not negligible, and the full perturbative result up to $n = 3$ differs qualitatively from the ansatz (12) of [21]. The reason for this difference has to be sought in the ansatz, which wrongly predicts a zero decay rate of a moving impurity with wave number $K < K_c$, due to the fact that the effect of the interaction is incorrectly described in the particle-hole continuum, which gives a wrong (zero) value to the lower border of that continuum [21]. To find K_c , one simply note that




Fig. 4: For a ^{40}K impurity interacting with a Fermi sea of ^6Li ($r = 6.6439$) on a narrow Feshbach resonance with $k_{\text{F}}R_* = 1$ as in [15], and $k_{\text{F}}a = -0.46$. Imaginary part (a), and real part (b) of the complex energy of the impurity counted with respect to the energy of the Fermi sea, as a function of its reduced wave number $\kappa = K/k_{\text{F}}$. In (b) the mean-field prediction $E_{\mathbf{K}} + \rho g$ was subtracted for clarity. Dashed (solid) lines: Perturbative calculation up to order two (three) in g . Circles: The single particle-hole ansatz. On the left of the vertical dotted line of (a) the ansatz incorrectly predicts a zero imaginary part. The inset in (a) is a magnification with the vertical axis in log-scale.

within the ansatz approach a real solution to the moving polaron is always possible if $\Delta E(\mathbf{K}) < 0$ [21], which at the mean-field level leads to

$$K_c \underset{a \rightarrow 0^-}{\sim} k_{\text{F}} \sqrt{-(r+1) \frac{2k_{\text{F}}a}{3\pi}}, \quad (30)$$

already a good approximation for Fig. 4.

Validity conditions. – For our perturbative treatment to apply, the impurity-to-fermion scattering must be in the Born regime, with a scattering amplitude $f_{k_{\text{rel}}} = -a/[1 + a(ik_{\text{rel}} + k_{\text{rel}}^2 R_*)] \simeq -a$ at the relative wave number $k_{\text{rel}} = \mu|\mathbf{K}/M - \mathbf{q}/m| \leq (K + rk_{\text{F}})/(1+r)$, so that

$$k_{\text{rel}}|a| \ll 1 \quad \text{and} \quad k_{\text{rel}}^2|a|R_* \ll 1. \quad (31)$$

However, this is not sufficient when $\tilde{K} \equiv K/(1+r) \gg k_{\text{F}}$: In this regime one can use the T -matrix formalism [10] to first order, neglecting the initial momentum of the fermions, to obtain $\Delta E(\mathbf{K}) - E_{\mathbf{K}} \simeq -\rho(2\pi\hbar^2/\mu)f_{\tilde{K}}$. Requiring that, in the real part of this result, the terms of order three in a are small as compared to the a^2 ones gives

$$a^2 \tilde{K}^2 |1 - \tilde{K}^2 R_*^2| \ll \left| \frac{k_{\text{F}}a}{6\pi(1+r)} \frac{k_{\text{F}}^2}{\tilde{K}^2} - a\tilde{K}^2 R_* \right|, \quad (32)$$

the first term in the right-hand side coming from Eq. (18).

Conclusion. – In the weakly attractive limit, both for broad and narrow Feshbach resonances, we have calculated analytically, up to second order in the interaction strength g , the energy, the decay rate, and the effective masses of an impurity moving with momentum $\hbar\mathbf{K}$ through a fully polarized Fermi sea, extending known results for $K = 0$ to finite momenta. At that order, our results show the existence of four regions separated by singularities of system's

observables, as e.g. the effective mass. We have characterized the degree and the physical origin of these singularities, which lays in the impurity-to-fermion scattering at the Fermi surface. For realistic values of the parameters we have shown that our second order correction is experimentally observable with the radio-frequency spectroscopic technique. We have also given an integral expression for the third-order complex energy, which allows us to estimate the neglected terms in the second-order theory.

We thank M. Zaccanti and M. Cetina for useful discussions. We acknowledge financial support from the ERC Project FERLODIM N.228177. C. T. acknowledges support from a Marie Curie Intra European grant INTERPOL N.298449.

REFERENCES

- [1] N. Navon *et al.*, *Science* **328**, 729 (2010).
- [2] S. Nascimbene *et al.*, *Nature (London)* **463**, 1057 (2010).
- [3] Mark Ku, A.T. Sommer, L.W. Cheuk, M.W. Zwierlein, *Science* **335**, 563 (2012).
- [4] K. Van Houcke *et al.*, *Nature Phys.* **8**, 366 (2012).
- [5] C. Lobo, A. Recati, S. Giorgini, S. Stringari, *Phys. Rev. Lett.* **97**, 200403 (2006).
- [6] F. Chevy, *Phys. Rev. A* **74**, 063628 (2006); R. Combescot, A. Recati, C. Lobo, F. Chevy, *Phys. Rev. Lett.* **98**, 180402 (2007).
- [7] N. Prokof'ev, B. Svistunov, *Phys. Rev. B* **77**, 020408(R) (2008); *ibid.*, *Phys. Rev. B* **77**, 125101 (2008).
- [8] J.D. Walecka, *Nuovo Cimento* **16**, 342 (1960).
- [9] R.F. Bishop, *Nucl. Phys. B* **17**, 573 (1970).
- [10] R.F. Bishop, *Ann. Phys.* **78**, 391 (1973).
- [11] R. Combescot, S. Giraud, *Phys. Rev. Lett.* **101**, 050404 (2008).
- [12] P. Massignan, G.M. Bruun, *Eur. Phys. J. D* **65**, 83 (2011).
- [13] J. Vlietinck, J. Ryckebusch, K. Van Houcke, *Phys. Rev. B* **87**, 115133 (2013).
- [14] G.M. Bruun *et al.*, *Phys. Rev. Lett.* **100**, 240406 (2008).
- [15] C. Kohstall *et al.*, *Nature* **485**, 615 (2012).
- [16] Yusuke Nishida, *Phys. Rev. A* **85**, 053643 (2012).
- [17] C. Mora, Y. Castin, *Phys. Rev. A* **67**, 053615 (2003); L. Pricoupenko, Y. Castin, *J. Phys. A* **40**, 12863 (2007); E. Burovski, N. Prokof'ev, B. Svistunov, M. Troyer, *New J. Phys.* **8**, 153 (2006); O. Juillet, *New J. Phys.* **9**, 163 (2007).
- [18] C. Trefzger, Y. Castin, in preparation.
- [19] N.W. Ashcroft, N.D. Mermin, in *Solid State Physics*, (HRW International Editions, 1976).
- [20] C. Cohen-Tannoudji, J. Dupont-Roc, G. Grynberg, in *Processus d'interaction entre photons et atomes* (InterEditions and Editions du CNRS, Paris, 1988).
- [21] C. Trefzger, Y. Castin, *Phys. Rev. A* **85**, 053612 (2012).
- [22] M. Jona-Lasinio, L. Pricoupenko, Y. Castin, *Phys. Rev. A*, **77**, 043611 (2008).
- [23] N. Regnault, Th. Jolicœur, *Phys. Rev. B*, **70**, 241307(R) (2004).