



HAL
open science

Existence of Gibbsian point processes with geometry-dependent interactions

David Dereudre, Rémy Drouilhet, Hans Otto Georgii

► **To cite this version:**

David Dereudre, Rémy Drouilhet, Hans Otto Georgii. Existence of Gibbsian point processes with geometry-dependent interactions. *Probability Theory and Related Fields*, 2012, 153 (3-4), pp.643-670. 10.1007/s00440-011-0356-5 . hal-00850422

HAL Id: hal-00850422

<https://hal.science/hal-00850422>

Submitted on 5 Jul 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Existence of Gibbsian point processes with geometry-dependent interactions

David Dereudre*, Remy Drouilhet[†] and Hans-Otto Georgii[‡]

November 1, 2018

Abstract

We establish the existence of stationary Gibbsian point processes for interactions that act on hyperedges between the points. For example, such interactions can depend on Delaunay edges or triangles, cliques of Voronoi cells or clusters of k -nearest neighbors. The classical case of pair interactions is also included. The basic tools are an entropy bound and stationarity.

Keywords. Gibbs measure, hypergraph, Delaunay mosaic, Voronoi tessellation, entropy.

MSC.Primary 60K35; Secondary: 60D05, 60G55, 82B21.

1 Introduction

Recent developments in statistical physics, stochastic geometry and spatial statistics involve Gibbs point processes with interactions depending on the local geometry of configurations in \mathbb{R}^d . A prominent class of such interactions is based on the nearest-neighbor graph coming from the Delaunay triangulation. In biology, such systems are used to model interacting cells in tissues or foams [11], [20]. In spatial statistics and stochastic geometry, structured point patterns, point processes lying along fibers or regular Delaunay tessellations have been studied via geometric Gibbs modifications; see [1], [6] and [21]. A probabilistic motivation comes from stationary renewal processes: Since these can be characterized as Gibbs processes for interactions between nearest-neighbor pairs of points [16, Section 6], Gibbs processes on \mathbb{R}^d with Delaunay tile interaction can be viewed as a multi-dimensional counterpart [8].

In this paper we consider general geometry-dependent interactions that are defined on a hypergraph structure \mathcal{E} . For every point configuration ω in \mathbb{R}^d , $\mathcal{E}(\omega)$ denotes a set of hyperedges on ω , and the formal Hamiltonian of ω is given by

$$H(\omega) = \sum_{\eta \in \mathcal{E}(\omega)} \varphi(\eta, \omega)$$

¹LAMAV, Université de Valenciennes et du Hainaut-Cambrésis, Le Mont Houy 59313 Valenciennes Cedex 09, France. E-mail: david.dereudre@univ-valenciennes.fr

²LJK, Université de Grenoble, B.S.H.M., 1251 Av. Centrale, BP 47 38040 Grenoble Cedex 9, France. E-mail: Remy.Drouilhet@upmf-grenoble.fr

³Mathematisches Institut der Universität München, Theresienstraße 39, 80333 München, Germany. E-mail: georgii@math.lmu.de

with a potential $\varphi(\eta, \omega)$ which only depends on η and the points of ω in some neighborhood of η . This locality of φ will be called the finite horizon property, see (2.1) below. This general setting includes all the above-mentioned cases and also the classical many-body interactions of finite range that are familiar from statistical mechanics [22].

There is a principal difference between the geometric interactions considered here and the classical many-body interactions. Namely, suppose a particle configuration ω is augmented by a new particle at x . In the case of a many-body interaction, this is only influenced by an additional interaction term between x and the particles of ω , and the interaction between the particles of ω is not affected by x . In other words, the classical many-body interactions are additive. In our setting, the new particle at x typically alters the hyperedges around x completely: some hyperedges are created and some others are destroyed. This means that both $\mathcal{E}(\omega \cup \{x\}) \setminus \mathcal{E}(\omega)$ and $\mathcal{E}(\omega) \setminus \mathcal{E}(\omega \cup \{x\})$ are non-empty, so that $H(\omega \cup \{x\})$ and $H(\omega)$ each contain terms that are not present in the other. This phenomenon blurs the usual distinction between attractive and repulsive interactions. Moreover, if the potential $\varphi(\eta, \omega)$ is allowed to take the value ∞ (hard-exclusion case), we arrive at the so-called non-hereditary situation that a configuration ω is excluded although $\omega \cup \{x\}$ is possible. This last case makes it difficult to use an infinitesimal characterization of Gibbs measures in terms of their Campbell measures and Papangelou intensities. Nevertheless, such an infinitesimal approach was first used to prove the existence of Gibbs measures for Delaunay interactions by requiring geometric constraints on the interaction [1, 2, 4, 6]. A quite different global approach, first used in [14] and based on stationarity and thermodynamic quantities such as pressure and free energy density, recently allowed to prove the existence of Gibbsian Delaunay tessellations for general bounded interactions without any geometric restrictions [8], and a similar approach could also be applied to quermass-interaction processes [7].

In this paper we address the existence problem for the general formalism of hypergraph interactions introduced here. Our approach is global as in [7, 8] and leads to a significant improvement of the existing results. In particular, we establish the existence of Gibbsian Delaunay tessellations for non-bounded and hard-exclusion potentials. In the classical context of stable many-body interaction of finite range, our results permit to relax the superstability assumption. Basic ingredients of the proof are an entropy bound to exploit the compactness of the level sets of the entropy density, and a somewhat delicate control of the range of the interaction, which takes advantage of stationarity.

The general setting of Gibbs measures for hypergraph interactions is introduced in Section 2. Section 3 contains our assumptions and the existence theorems. Section 4 offers a series of examples that includes many-body interactions of finite range as well as interactions on Delaunay tiles and Voronoi cells and between k -nearest neighbors. The proofs of the main results follow in Section 5, and an appendix is devoted to measurability questions.

2 Preliminaries

2.1 Point configurations and hyperedge interactions

Consider the Euclidean space \mathbb{R}^d of arbitrary dimension $d \geq 1$. Subregions of \mathbb{R}^d will typically be denoted by Λ or Δ and will always be assumed to be Borel with positive

Lebesgue measure $|\Lambda|$ resp. $|\Delta|$. We write $\Delta \Subset \mathbb{R}^d$ if Δ is bounded. A *configuration* is a subset ω of \mathbb{R}^d which is locally finite, in that $\omega \cap \Delta$ has finite cardinality $N_\Delta(\omega) = \#(\omega \cap \Delta)$ for all $\Delta \Subset \mathbb{R}^d$. The space Ω of all configurations is equipped with the σ -algebra \mathcal{F} that is generated by the counting variables N_Δ with $\Delta \Subset \mathbb{R}^d$. It will often be convenient to write ω_Δ in place of $\omega \cap \Delta$. As usual, we take as reference measure on (Ω, \mathcal{F}) the Poisson point process Π^z of an arbitrary intensity $z > 0$. Recall that Π^z is the unique probability measure on (Ω, \mathcal{F}) such that the following holds for all $\Delta \Subset \mathbb{R}^d$: (i) N_Δ is Poisson distributed with parameter $z|\Delta|$, and (ii) conditional on $N_\Delta = n$, the n points in Δ are independent with uniform distribution on Δ , for each integer $n \geq 1$.

Next, let $\Omega_f \subset \Omega$ denote the set of all finite configurations ω (which means that $\#(\omega) < \infty$), and $\mathcal{F}_f = \mathcal{F}|_{\Omega_f}$ the trace σ -algebra of \mathcal{F} on Ω_f . The product space $\Omega_f \times \Omega$ carries the product σ -algebra $\mathcal{F}_f \otimes \mathcal{F}$. For each $\Lambda \subset \mathbb{R}^d$ we write $\Omega_\Lambda = \{\omega \in \Omega : \omega \subset \Lambda\}$ for the set of all configurations in Λ , $\text{pr}_\Lambda : \omega \rightarrow \omega_\Lambda = \omega \cap \Lambda$ for the projection from Ω to Ω_Λ , $\mathcal{F}'_\Lambda = \mathcal{F}|_{\Omega_\Lambda}$ for the trace σ -algebra of \mathcal{F} on Ω_Λ , and $\mathcal{F}_\Lambda = \text{pr}_\Lambda^{-1} \mathcal{F}'_\Lambda \subset \mathcal{F}$ for the σ -algebra of all events that happen in Λ only. The reference measure on $(\Omega_\Lambda, \mathcal{F}'_\Lambda)$ is $\Pi_\Lambda^z := \Pi^z \circ \text{pr}_\Lambda^{-1}$. Finally, let $\Theta = (\vartheta_x)_{x \in \mathbb{R}^d}$ be the shift group, where $\vartheta_x : \Omega \rightarrow \Omega$ is the translation by the vector $-x \in \mathbb{R}^d$. By definition, $N_\Delta(\vartheta_x \omega) = N_{\Delta+x}(\omega)$ for all $\Delta \Subset \mathbb{R}^d$.

The interaction of points to be considered in this paper will depend on the geometry of their location. This geometry will be described in terms of a hypergraph, and the interaction potential will be defined on the hyperedges.

Definition.

- A *hypergraph structure* is a measurable subset \mathcal{E} of $\Omega_f \times \Omega$ such that $\eta \subset \omega$ for all $(\eta, \omega) \in \mathcal{E}$. If $(\eta, \omega) \in \mathcal{E}$, we say that η is a *hyperedge* of ω , and we write $\eta \in \mathcal{E}(\omega)$.
- A *hyperedge potential* is a measurable function φ from a hypergraph structure \mathcal{E} to $\mathbb{R} \cup \{\infty\}$.
- A hyperedge potential φ (or, more explicitly, the pair (\mathcal{E}, φ)) is called *shift-invariant* if

$$(\vartheta_x \eta, \vartheta_x \omega) \in \mathcal{E} \text{ and } \varphi(\vartheta_x \eta, \vartheta_x \omega) = \varphi(\eta, \omega) \text{ for all } (\eta, \omega) \in \mathcal{E} \text{ and } x \in \mathbb{R}^d.$$

- Let us say that φ (or the pair (\mathcal{E}, φ)) satisfies the *finite horizon property* if for each $(\eta, \omega) \in \mathcal{E}$ there exists some $\Delta \Subset \mathbb{R}^d$ such that

$$(2.1) \quad (\eta, \tilde{\omega}) \in \mathcal{E} \text{ and } \varphi(\eta, \tilde{\omega}) = \varphi(\eta, \omega) \text{ when } \tilde{\omega} = \omega \text{ on } \Delta.$$

We will assume throughout this paper that the hyperedge potential φ under consideration is shift-invariant and exhibits the finite horizon property. Moreover, for notational convenience we set $\varphi = 0$ on \mathcal{E}^c . Since \mathcal{E} is measurable, this does not affect the measurability of φ .

The domain \mathcal{E} of φ can be considered as a rule that turns each configuration ω into a hypergraph $(\omega, \mathcal{E}(\omega))$. Both \mathcal{E} and φ are not affected by translations. Moreover, the presence of a hyperedge $\eta \in \mathcal{E}(\omega)$ and the value of $\varphi(\eta, \omega)$ can be determined by looking at ω in a (sufficiently large but) bounded neighborhood Δ of η , called the *horizon of η in ω* , which in general depends on both η and ω . Note that in general there is no minimal such

horizon. To obtain a standard choice of Δ one can take the closed ball $B_{\eta,\omega} = \bar{B}(g_\eta, r_{\eta,\omega})$ with center at the gravicenter g_η of η and radius $r_{\eta,\omega}$ chosen smallest possible. Finally, we note that the concept of hypergraph structure is similar to that of a cluster property as introduced in [25]. Here are two examples the reader might keep in mind. Further examples will follow in Section 4.

Example 2.1. Many-body interactions of bounded range. Let $r > 0$ and

$$\text{LC}_r = \{(\eta, \omega) : \eta \subset \omega, \text{diam}(\eta) \leq r, \omega \in \Omega\}$$

be the locally complete graph. Thus, for each $\omega \in \Omega$, $\text{LC}_r(\omega)$ consists of all hyperedges between points of distance at most r . If we assume that $\varphi(\eta, \omega)$ only depends on η , we are in the classical situation of many-body interactions of range r . The finite horizon property for $(\eta, \omega) \in \text{LC}_r$ then holds for arbitrary $\Delta \supset \eta$. If LC_r is restricted to hyperedges of cardinality two, we arrive at the familiar pair interactions of statistical mechanics.

Example 2.2. Delaunay potentials. The set Del of Delaunay hyperedges consists of all pairs (η, ω) with $\eta \subset \omega$ for which there exists an open ball $B(\eta, \omega)$ with $\partial B(\eta, \omega) \cap \omega = \eta$ that contains no points of ω . For $k = 1, \dots, d+1$ we write $\text{Del}_k = \{(\eta, \omega) \in \text{Del} : \#\eta = k\}$ for the set of all Delaunay simplices with k vertices. Clearly, Del and Del_k are hypergraph structures. It is possible that the convex hull of a set $\eta \in \text{Del}(\omega)$ is not a simplex, namely when η consists of four or more points on a sphere with no point inside. However, this is an exceptional case, which occurs only with probability zero for our Poisson reference measure Π^z . Note that $B(\eta, \omega)$ is only uniquely determined when $\#\eta = d+1$ and η is affinely independent.

The simplest class of Delaunay hyperedge potentials consists of pair interactions of the form $\varphi(\eta, \omega) = \phi(|x - y|)$ for $\eta = \{x, y\} \in \text{Del}_2(\omega)$. Such a φ satisfies the finite horizon property (2.1) with $\Delta = \bar{B}(\eta, \omega)$ for any ball $B(\eta, \omega)$ as above.

An example of a potential $\varphi(\eta, \omega)$ on Del_2 which does not only depend on η but also on ω is $\varphi(\eta, \omega) = \phi(\text{Vor}_\omega(x), \text{Vor}_\omega(y))$. Here we write $\text{Vor}_\omega(x)$ for the Voronoi cell associated to a point $x \in \omega$, viz. the set

$$(2.2) \quad \text{Vor}_\omega(x) := \{y \in \mathbb{R}^d : |x - y| \leq |\tilde{x} - y| \quad \forall \tilde{x} \in \omega\}$$

of all points of \mathbb{R}^d which are closer to x than to all other points of ω . It is well-known that the Voronoi cells form a tessellation [24]. Also, any two points of a configuration are connected by a Delaunay edge if and only if their Voronoi cells have a non-trivial intersection. That is,

$$(2.3) \quad \{x, y\} \in \text{Del}_2(\omega) \iff \#(\text{Vor}_\omega(x) \cap \text{Vor}_\omega(y)) > 1.$$

This reveals that the Delaunay graph is a nearest-neighbor graph. The potential above satisfies the finite horizon property (2.1) with Δ equal to the closure of the set $\bigcup_{\xi \in \text{Del}(\omega), \xi \cap \eta \neq \emptyset} B(\xi, \omega)$, provided the cells $\text{Vor}_\omega(x)$ and $\text{Vor}_\omega(y)$ are bounded. The proviso is necessary because unbounded Voronoi cells, which can occur at the ‘‘boundary’’ of ω , are not protected by the points in Δ . So we must exclude from $\text{Del}_2(\omega)$ all edges $\{x, y\}$ for which $\text{Vor}_\omega(x) \cup \text{Vor}_\omega(y)$ is unbounded.

2.2 Gibbs measures for hyperedge potentials

Our objective here is to introduce the concept of a Gibbsian point process for an activity $z > 0$ and a hyperedge potential φ defined on a hypergraph structure \mathcal{E} . First we will introduce the Hamiltonian for a bounded region $\Lambda \Subset \mathbb{R}^d$ with configurational boundary condition $\omega \in \Omega$. This requires to consider the set of hyperedges η in a configuration ω for which either η itself or $\varphi(\eta, \omega)$ depends on the points of ω in Λ . Specifically, we set

$$(2.4) \quad \mathcal{E}_\Lambda(\omega) = \{\eta \in \mathcal{E}(\omega) : \varphi(\eta, \zeta \cup \omega_{\Lambda^c}) \neq \varphi(\eta, \omega) \text{ for some } \zeta \in \Omega_\Lambda\}.$$

Recall the convention that $\varphi = 0$ on \mathcal{E}^c . So, if $\eta \in \mathcal{E}(\omega) \setminus \mathcal{E}(\zeta \cup \omega_{\Lambda^c})$ for some $\zeta \in \Omega_\Lambda$ then either $\eta \notin \mathcal{E}_\Lambda(\omega)$, or η is irrelevant for ω , in that $\varphi(\eta, \omega) = 0$. The *Hamiltonian in Λ with boundary condition ω* is then given by the formula

$$(2.5) \quad H_{\Lambda, \omega}(\zeta) := \sum_{\eta \in \mathcal{E}_\Lambda(\zeta \cup \omega_{\Lambda^c})} \varphi(\eta, \zeta \cup \omega_{\Lambda^c}) \quad \text{for } \zeta \in \Omega_\Lambda,$$

provided this sum is well-defined. As usual, we also consider the associated partition function

$$Z_{\Lambda, \omega}^z := \int e^{-H_{\Lambda, \omega}(\zeta)} \Pi_\Lambda^z(d\zeta).$$

To ensure that these quantities are well-defined, we impose the following condition on the boundary condition ω . Let $\varphi^- = (-\varphi) \vee 0$ be the negative part of φ .

Definition. A configuration $\omega \in \Omega$ is called *admissible for a region $\Lambda \Subset \mathbb{R}^d$ and an activity $z > 0$* if

$$H_{\Lambda, \omega}^-(\zeta) := \sum_{\eta \in \mathcal{E}_\Lambda(\zeta \cup \omega_{\Lambda^c})} \varphi^-(\eta, \zeta \cup \omega_{\Lambda^c}) < \infty \quad \text{for } \Pi_\Lambda^z\text{-almost all } \zeta \in \Omega_\Lambda$$

(so that $H_{\Lambda, \omega}$ is almost surely well-defined), and $0 < Z_{\Lambda, \omega}^z < \infty$. We write $\Omega_*^{\Lambda, z}$ for the set of all these ω .

We note that, in contrast to the standard setting of statistical mechanics, the partition function is not automatically positive because, in the present setting, $H_{\Lambda, \omega}(\emptyset)$ is not necessarily finite. For $\omega \in \Omega_*^{\Lambda, z}$, we can define the Gibbs distribution for $(\mathcal{E}, \varphi, z)$ in a region $\Lambda \Subset \mathbb{R}^d$ with boundary condition ω as usual by

$$(2.6) \quad G_{\Lambda, \omega}^z(F) = \int_{\Omega_\Lambda} \mathbf{1}_F(\zeta \cup \omega_{\Lambda^c}) e^{-H_{\Lambda, \omega}(\zeta)} \Pi_\Lambda^z(d\zeta) / Z_{\Lambda, \omega}^z,$$

where $F \in \mathcal{F}$ is arbitrary.

Definition. Let \mathcal{E} be a hypergraph structure, φ a hyperedge potential, and $z > 0$ an activity. A probability measure P on (Ω, \mathcal{F}) is called a *Gibbs measure for \mathcal{E} , φ and z* if $P(\Omega_*^{\Lambda, z}) = 1$ and

$$(2.7) \quad \int f dP = \int_{\Omega_*^{\Lambda, z}} \frac{1}{Z_{\Lambda, \omega}^z} \int_{\Omega_\Lambda} f(\zeta \cup \omega_{\Lambda^c}) e^{-H_{\Lambda, \omega}(\zeta)} \Pi_\Lambda^z(d\zeta) P(d\omega)$$

for every $\Lambda \Subset \mathbb{R}^d$ and every measurable $f : \Omega \rightarrow [0, \infty[$.

The equations (2.7) are known as the DLR equations (after Dobrushin, Lanford and Ruelle). They express that $G_{\Lambda,\omega}^z(F)$ is a version of the conditional probability $P(F | \mathcal{F}_{\Lambda^c})(\omega)$. We will be particularly interested in Gibbs measures that are stationary, that is, invariant under the shift group $\Theta = (\vartheta_x)_{x \in \mathbb{R}^d}$. We write \mathcal{P}_Θ for the set of all Θ -invariant probability measures P on (Ω, \mathcal{F}) with finite intensity $i(P) = \int N_{[0,1]^d} dP$, and $\mathcal{G}_\Theta(\varphi, z)$ for the set of all Gibbs measures for φ and z that belong to \mathcal{P}_Θ . We conclude this section with a discussion of measurability questions.

Remark 2.1. Measurability. The quantities introduced above are not measurable with respect to the underlying σ -algebras defined so far, but only with respect to their universal completion. Specifically, for each σ -algebra \mathcal{A} let \mathcal{A}^* be the associated σ -algebra of all universally measurable sets, i.e., of the sets which belong to the P -completion of \mathcal{A} for all probability measures P on \mathcal{A} ; see [5, pp. 36 & 280]. It is then the case that, for each $\Lambda \Subset \mathbb{R}^d$, the Hamiltonian $(\zeta, \omega) \mapsto H_{\Lambda,\omega}(\zeta)$ is measurable with respect to $(\mathcal{F}'_\Lambda \otimes \mathcal{F}_{\Lambda^c})^*$. Likewise, the partition function $\omega \rightarrow Z_{\Lambda,\omega}^z$ is measurable with respect to $\mathcal{F}_{\Lambda^c}^*$, and $\Omega_*^{\Lambda,z} \in \mathcal{F}_{\Lambda^c}^*$. Moreover, $(\omega, F) \rightarrow G_{\Lambda,\omega}^z(F)$ is a probability kernel from $(\Omega_*^{\Lambda,z}, \mathcal{F}_{\Lambda^c}^* |_{\Omega_*^{\Lambda,z}})$ to (Ω, \mathcal{F}) . All this will be proved in the appendix. *We will therefore identify all probability measures in this paper with their respective complete extension.* This convention underlies already the preceding definition of a Gibbs measure.

3 Hypotheses and results

Having defined the concept of Gibbs measure for a hyperedge potential we now turn to our main theme, the existence of such Gibbs measures. Let us state the conditions we need. In the subsequent section we will provide a series of examples for which these conditions are met.

We begin with an assumption which says that hyperedges with a large horizon require the existence of a large ball with only a few points. This will imply that the Hamiltonian $H_{\Lambda,\omega}$ depends only on the points of ω in a bounded region $\partial\Lambda(\omega)$, and can be viewed as a sharpening of the finite horizon property (2.1).

- (R)** *The range condition.* There exist constants $\ell_R, n_R \in \mathbb{N}$ and $\delta_R < \infty$ such that for all $(\eta, \omega) \in \mathcal{E}$ one can find a horizon Δ as in (2.1) satisfying the following:
- For every $x, y \in \Delta$, there exist ℓ open balls B_1, \dots, B_ℓ (with $\ell \leq \ell_R$) such that
 - the set $\cup_{i=1}^\ell \bar{B}_i$ is connected and contains x and y , and
 - for each i , either $\text{diam } B_i \leq \delta_R$ or $N_{B_i}(\omega) \leq n_R$.

Note that **(R)** is trivially satisfied when all horizon sets can be chosen to have uniformly bounded diameters. For instance, this holds in Example 2.1. The use of the range condition **(R)** will be revealed by Proposition 3.1 below, which states that the following finite range property holds almost surely for nondegenerate $P \in \mathcal{P}_\Theta$.

Definition. Let $\Lambda \Subset \mathbb{R}^d$ be given. We say a configuration $\omega \in \Omega$ *confines the range of φ from Λ* if there exists a set $\partial\Lambda(\omega) \Subset \mathbb{R}^d$ such that $\varphi(\eta, \zeta \cup \tilde{\omega}_{\Lambda^c}) = \varphi(\eta, \zeta \cup \omega_{\Lambda^c})$ whenever $\tilde{\omega} = \omega$ on $\partial\Lambda(\omega)$, $\zeta \in \Omega_\Lambda$ and $\eta \in \mathcal{E}_\Lambda(\zeta \cup \omega_{\Lambda^c})$. In this case we write $\omega \in \Omega_{\text{ct}}^\Lambda$. $\partial\Lambda(\omega)$ is called the ω -boundary of Λ , and we use the abbreviation $\partial_\Lambda \omega = \omega_{\partial\Lambda(\omega)}$.

Given any $\omega \in \Omega_{\text{cr}}^\Lambda$, we assume in the following that $\partial\Lambda(\omega) = \Lambda^r \setminus \Lambda$, where Λ^r is the closed r -neighborhood of Λ and $r = r_{\Lambda,\omega}$ is chosen as small as possible. Moreover, for $\omega \in \Omega_{\text{cr}}^\Lambda$ we have

$$(3.1) \quad H_{\Lambda,\omega}(\zeta) = \sum_{\eta \in \mathcal{E}_\Lambda(\zeta \cup \partial_\Lambda \omega)} \varphi(\eta, \zeta \cup \partial_\Lambda \omega),$$

and this sum extends over a finite set. This means that the first assumption in the definition of admissibility for Λ is satisfied. Here is the proposition announced above. It will follow from Proposition 5.4 below.

Proposition 3.1. *Under (R), for each $\Lambda \Subset \mathbb{R}^d$ there exists a set $\hat{\Omega}_{\text{cr}}^\Lambda \in \mathcal{F}_{\Lambda^c}$ such that $\hat{\Omega}_{\text{cr}}^\Lambda \subset \Omega_{\text{cr}}^\Lambda$ and $P(\hat{\Omega}_{\text{cr}}^\Lambda) = 1$ for all $P \in \mathcal{P}_\Theta$ with $P(\{\emptyset\}) = 0$.*

Our next assumption is stability, the standard assumption that ensures the finiteness of all partition functions. In our setting, a somewhat modified definition turns out to be suitable.

(S) *Stability.* The hyperedge potential φ is called *stable* if there exists a constant $c_S \geq 0$ such that

$$(3.2) \quad H_{\Lambda,\omega}(\zeta) \geq -c_S \#(\zeta \cup \partial_\Lambda \omega)$$

for all $\Lambda \Subset \mathbb{R}^d$, $\zeta \in \Omega_\Lambda$ and $\omega \in \Omega_{\text{cr}}^\Lambda$.

In Remark 3.4 below we will show that this definition is a natural extension of the familiar concept of stability in statistical mechanics. Complementary to the lower bound provided by stability, we will also need a further condition that provides at least a partial upper bound for the Hamiltonians. This is because in the extreme case when φ is constantly equal to ∞ we have $Z_{\Lambda,\cdot}^z \equiv 0$, so that the definition of Gibbs measures is meaningless.

Let $M \in \mathbb{R}^{d \times d}$ be an invertible $d \times d$ matrix and consider for each $k \in \mathbb{Z}^d$ the cell

$$(3.3) \quad C(k) := \{Mx \in \mathbb{R}^d : x - k \in [-1/2, 1/2]^d\}.$$

These cells together constitute a periodic partition of \mathbb{R}^d into parallelepipeds. For example, the columns M_1, \dots, M_d of M might form an orthogonal basis of \mathbb{R}^d or, for $d = 2$, define the sides of an equilateral triangle. For brevity we write $C = C(0)$. Let Γ be a measurable subset of $\Omega_C \setminus \{\emptyset\}$ and

$$(3.4) \quad \bar{\Gamma} = \left\{ \omega \in \Omega : \vartheta_{Mk}(\omega_{C(k)}) \in \Gamma \text{ for all } k \in \mathbb{Z}^d \right\}$$

the set of all configurations whose restriction to an arbitrary cell $C(k)$, when shifted back to C , belongs to Γ . We call each $\omega \in \bar{\Gamma}$ pseudo-periodic. The required control of the Hamiltonian from above will then be achieved by the following assumption on the joint behavior of \mathcal{E} , φ and z . Recall that $r_{\Lambda,\omega}$ was defined before (3.1).

(U) *Upper regularity.* M and Γ can be chosen so that the following holds.

(U1) *Uniform confinement:* $\bar{\Gamma} \subset \Omega_{c_\Gamma}^\Lambda$ for all $\Lambda \in \mathbb{R}^d$, and

$$r_\Gamma := \sup_{\Lambda \in \mathbb{R}^d} \sup_{\omega \in \bar{\Gamma}} r_{\Lambda, \omega} < \infty.$$

(U2) *Uniform summability:* $c_\Gamma^+ := \sup_{\omega \in \bar{\Gamma}} \sum_{\eta \in \mathcal{E}(\omega): \eta \cap C \neq \emptyset} \frac{\varphi^+(\eta, \omega)}{\#(\hat{\eta})} < \infty$,

where $\hat{\eta} := \{k \in \mathbb{Z}^d : \eta \cap C(k) \neq \emptyset\}$ and φ^+ is the positive part of φ .

(U3) *Strong non-rigidity:* $e^{z|C|} \Pi_C^z(\Gamma) > e^{c_\Gamma}$, where c_Γ is defined as in (U2) with φ in place of φ^+ .

Hypothesis (U1) states that the configurations in $\bar{\Gamma}$ confine the range of φ in a uniform way. So, for $\omega \in \bar{\Gamma}$, the ω -boundary $\partial\Lambda(\omega)$ of a set $\Lambda \in \mathbb{R}^d$ is contained in the r_Γ -boundary $\partial^\Gamma\Lambda := \Lambda^{r_\Gamma} \setminus \Lambda$, and the cardinality of $\partial_\Lambda\omega$ is not larger than that of $\partial_\Lambda^\Gamma\omega := \omega_{\partial^\Gamma\Lambda}$. On the other hand, condition (U2) provides a uniform upper bound for the local Hamiltonians H_Λ on $\bar{\Gamma}$. This bound is of order $c_\Gamma|\Lambda|$, cf. (5.8) below. Finally, hypothesis (U3) holds for all z in an interval of the form $]z_0, \infty[$, provided that $\Pi_C^z(\Gamma) > 0$ for some (and thus all) $z > 0$. Indeed, since $\emptyset \notin \Gamma$ it follows that

$$(3.5) \quad e^{z|C|} \Pi_C^z(\Gamma) = \sum_{k=1}^{\infty} \frac{z^k}{k!} \int_C \cdots \int_C \mathbb{1}_\Gamma(\{x_1, \dots, x_k\}) dx_1 \dots dx_k$$

is then a strictly increasing function of z . We emphasize that condition (U) imposes conflicting demands on Γ . While (U1) and (U2) suggest to choose the set Γ as small as possible, (U3) requires that Γ is not too small. The point will be to choose a set Γ that satisfies all requirements simultaneously. Here is our main existence theorem.

Theorem 3.2. *For every hypergraph structure \mathcal{E} , hyperedge potential φ and activity $z > 0$ satisfying (S), (R) and (U) there exists at least one Gibbs measure $P \in \mathcal{G}_\Theta(\varphi, z)$.*

In some cases it is difficult to satisfy hypothesis (U3) when z is small. This occurs, for example, when a hard-exclusion hyperedge interaction enforces a minimal number of points per cell; see Proposition 4.2. But a slight variation of proof allows to establish the existence of a Gibbs measure for every $z > 0$ also in this case. Assumptions (U1) and (U3) are replaced by conditions ($\hat{U}1$) and ($\hat{U}3$) as follows.

(\hat{U}) *Alternative upper regularity.* M and Γ can be chosen so that the following holds.

($\hat{U}1$) *Lower density bound:* There exist constants $a, b > 0$ such that $\#(\zeta) \geq a|\Lambda| - b$ whenever $\zeta \in \Omega_f$ is such that $H_{\Lambda, \omega}(\zeta) < \infty$ for some $\zeta \subset \Lambda \in \mathbb{R}^d$ and some $\omega \in \bar{\Gamma}$.

($\hat{U}2$) = (U2) *Uniform summability.*

($\hat{U}3$) *Weak non-rigidity:* $\Pi_C^z(\Gamma) > 0$.

Here is the modified existence theorem.

Theorem 3.3. *A Gibbs measure $P \in \mathcal{G}_\Theta(\varphi, z)$ exists also under the hypotheses **(S)**, **(R)** and $(\hat{\mathbf{U}})$.*

It is often natural to choose Γ as the set of all configurations that consist of a single point in some Borel set $A \subset C$. So we define

$$(3.6) \quad \Gamma^A = \{\zeta \in \Omega_C : \zeta = \{x\} \text{ for some } x \in A\}.$$

For $\Gamma = \Gamma^A$, the assumptions **(U)** and $(\hat{\mathbf{U}})$ are respectively called **(U^A)** and $(\hat{\mathbf{U}}^A)$. In particular, **(U2^A)** and **(U3^A)** take the simpler form

$$(U2^A) \quad c_A^+ := \sup_{\omega \in \bar{\Gamma}^A} \sum_{\eta \in \mathcal{E}(\omega) : \eta \cap C \neq \emptyset} \frac{\varphi^+(\eta, \omega)}{\#(\eta)} < \infty \quad \text{and}$$

$$(U3^A) \quad z|A| > e^{c_A}.$$

We then have the following corollary of Theorems 3.2 and 3.3.

Corollary 3.4. *A Gibbs measure $P \in \mathcal{G}_\Theta(\varphi, z)$ exists under the hypotheses **(S)**, **(R)** and either **(U^A)** or $(\hat{\mathbf{U}}^A)$.*

We conclude this section with a series of comments on our assumptions and on the extension to marked particles.

Remark 3.1. Bounded horizons. The conditions **(R)** and **(U^A)** hold as soon as $\varphi(\{0\}, \{0\})$ is finite and (\mathcal{E}, φ) has bounded horizons, in that there exists some $r_\varphi < \infty$ such that $r_{\eta, \omega} \leq r_\varphi$ for all $(\eta, \omega) \in \mathcal{E}$. (The notation $r_{\eta, \omega}$ was introduced after (2.1).) Indeed, condition **(R)** holds trivially with $\delta_R = 2r_\varphi$. As for **(U^A)**, let $M = aE$, where $a > 2r_\varphi$ and E is the identity matrix. Let $A = B(0, b)$ be a centered ball of radius $b < a/2 - r_\varphi$. For $\Gamma = \Gamma^A$, condition **(U1)** holds with $r_\Gamma = r_\varphi$. Moreover, by the choice of a and b , each $\eta \in \mathcal{E}(\omega)$ with $\omega \in \bar{\Gamma}^A$ must be a singleton $\{x\}$, so that $\varphi(\eta, \omega) = \varphi(\{x\}, \{x\})$. In view of the shift-invariance of φ , this means that **(U2^A)** holds with $c_A^+ = \varphi^+(\{0\}, \{0\}) < \infty$. Finally, **(U3^A)** holds if a and b are in fact chosen so large that also $\pi z b^2 > e^{\varphi(\{0\}, \{0\})}$.

Remark 3.2. Scale-invariant potentials. Suppose \mathcal{E} and φ are scale-invariant in the sense that $(r\eta, r\omega) \in \mathcal{E}$ and $\varphi(r\eta, r\omega) = \varphi(\eta, \omega)$ for all $(\eta, \omega) \in \mathcal{E}$ and $r > 0$. Here, $r\eta = \{rx : x \in \eta\}$ and $r\omega = \{rx : x \in \omega\}$. Then Theorem 3.2 is still valid when assumption **(U3)** is replaced by $(\hat{\mathbf{U}}3)$. Indeed, the scale invariance of φ implies that the image of a Gibbs measure for φ and z under the rescaling $\omega \rightarrow r\omega$ is a Gibbs measure for φ and zr^{-d} . So, it is sufficient to have the existence of a Gibbs measure for large z , and this follows from the remark around (3.5).

Remark 3.3. Stability via sublinearity of the hypergraph. We say that a hypergraph \mathcal{E} is sublinear if there exists a constant $C < \infty$ such that $\#\mathcal{E}(\omega) \leq C \#(\omega)$ for every finite configuration ω . In this case, the stability is ensured by requiring that the hyperedge potential φ is bounded below, in that

$$(3.7) \quad \varphi(\eta, \omega) \geq -c_\varphi$$

for some $c_\varphi < \infty$. If the sublinearity of the hypergraph structure fails, the stability can simply be achieved by requiring that the potential φ is nonnegative (i.e., $c_\varphi = 0$). For example, for $d = 2$ it follows from Euler's formula that the cardinalities of the Delaunay edges and triangles are sublinear [8], so that the stability follows directly from (3.7).

Remark 3.4. Stability: comparison with the classical case. Consider the hypergraph structure $\mathcal{E} = \text{LC}_r$ of Example 2.1 describing many-body interactions of range r . In contrast to Example 2.2 where the Delaunay tiles depend on the presence of further particles, it is then meaningful to define the energy of a finite configuration $\zeta \in \Omega_f$ by $H(\zeta) = \sum_{\eta \in \mathcal{E}(\zeta)} \varphi(\eta)$. The classical stability condition asserts that $H(\zeta) \geq -c_S \#(\zeta)$ for all $\zeta \in \Omega_f$; see [22], for example. This follows from **(S)** by choosing $\Lambda \supset \zeta$ and $\omega = \emptyset$. Conversely, the Hamiltonian (2.5) is equivalent to the Hamiltonian $\tilde{H}_{\Lambda, \omega}(\zeta) := H(\zeta \cup \partial_{\Lambda} \omega)$ in the sense that the associated Gibbs distributions coincide (at least when $\varphi < \infty$), and the classical stability assumption for H gives **(S)** for $\tilde{H}_{\Lambda, \omega}$. This shows that hypothesis **(S)** is essentially equivalent to the classical concept of stability.

Remark 3.5. Sub-hypergraph potentials. Consider a shift-invariant sub-hypergraph structure $\mathcal{E}' \subset \mathcal{E}$ of \mathcal{E} and a hyperedge potential φ on \mathcal{E} , and let φ' be its restriction to \mathcal{E}' . In general, φ' does not satisfy the finite horizon property, but let us assume it does. Which of the assumptions **(R)**, **(S)**, **(U)** and **(\hat{U})** on φ are inherited by φ' ? It is clear that assumptions **(R)**, (U1), (U2) and (\hat{U} 3) are hereditary. Assumption (U3) remains also valid for φ' , but for a different range of values of z because the constant c_{Γ} is different in general. Assumption **(S)** is lost, but a positive exception is the case of Remark 3.3 when stability follows from the sublinearity of \mathcal{E} and the lower boundedness of φ ; these properties are obviously inherited by φ' . Assumption (\hat{U} 1) is lost in general.

Remark 3.6. Upper regularity in Delaunay models. For potentials acting on the Delaunay graph, the matrix \mathbf{M} and the set Γ in hypotheses **(\mathbf{U}^A)** and **($\hat{\mathbf{U}}^A$)** will be chosen as follows. Let \mathbf{M} be such that $|\mathbf{M}_i| = a > 0$ for $i = 1, \dots, d$ and $\angle(\mathbf{M}_i, \mathbf{M}_j) = \pi/3$ for $i \neq j$, and let $\Gamma = \Gamma^A$ with $A = B(0, b)$. If $b \leq \varrho_0 a$ for some sufficiently small constant $\varrho_0 > 0$, the Delaunay neighborhood of a point x in a configuration $\omega \in \bar{\Gamma}$ contains a minimal number of points denoted by γ_d . For $d = 2$ one can take $\varrho_0 = \sqrt{3}/6$ and has $\gamma_2 = 6$, and the Delaunay neighborhood of the unique point x_k in $\omega \cap C(k)$ consists of the unique points x_l in $\omega \cap C(l)$ with $l - k \in \{(-1, 0), (-1, 1), (0, 1), (1, 0), (1, -1), (0, -1)\}$. For $d > 2$, γ_d is less easy to determine but it is clearly not larger than $3^d - 1$, the value corresponding to the case $\mathbf{M} = a\mathbf{E}$.

Remark 3.7. Extension to the marked case. The preceding results can be easily extended to the case of particles with internal degrees of freedom, or marks. Let Σ be an arbitrary Polish space with Borel σ -algebra \mathcal{S} and reference probability measure μ . Σ serves as the space of marks. That is, each marked point is represented by a position $x \in \mathbb{R}^d$ and a mark $\sigma \in \Sigma$, and each configuration ω is a countable subset of $\mathbb{R}^d \times \Sigma$ having a locally finite projection onto \mathbb{R}^d . The role of the reference measure on the configuration space Ω is taken over by the Poisson point process Π^z with intensity measure $z\lambda \otimes \mu$, where λ is Lebesgue measure on \mathbb{R}^d . The translations ϑ_x act only on the positions of the particles and leave their marks untouched. We do not discuss the further formal details here, which are standard and can be found in [19] or [15], for example. What we want to emphasize here is that all definitions and results above carry over to this setting without any change, provided it is understood that all regions Λ or Δ in \mathbb{R}^d always refer to the positional part of a configuration. For example: the notation ω_{Δ} now stands for $\omega \cap (\Delta \times \Sigma)$; a set $\Delta \Subset \mathbb{R}^d$ is the horizon of $(\eta, \omega) \in \mathcal{E}$ if $(\eta, \tilde{\omega}) \in \mathcal{E}$ and $\varphi(\eta, \tilde{\omega}) = \varphi(\eta, \omega)$ whenever $\tilde{\omega} = \omega$ on $\Delta \times \Sigma$; and the condition in Remark 3.1 should now read $\sup_{\sigma \in S} \varphi(\{(0, \sigma), \{(0, \sigma)\})\}) < \infty$ for some Borel set $S \subset \Sigma$ with $\mu(S) > 0$.

4 Examples

In this section we present a series of examples to illustrate our general existence results. These examples satisfy the assumptions of Theorem 3.2 or 3.3 and many of them have been introduced in practical or theoretical papers without justification or with justification in some partial cases.

To sort the examples of this section we will distinguish whether or not the potential $\varphi(\eta, \omega)$ depends explicitly on ω . If not, we speak of a *pure hyperedge potential*. Otherwise, the finite horizon property implies that $\varphi(\eta, \omega)$ actually depends only on some points of $\omega \setminus \eta$ close to η , which is expressed by speaking of a *neighborhood-dependent hyperedge potential*. But note that this distinction is merely a matter of how the interaction is represented. In the pure case, the extended potential $(\eta, \omega) \rightarrow \mathbb{1}_{\mathcal{E}}(\eta, \omega) \varphi(\eta)$ on $\Omega_f \times \Omega$ clearly does depend on ω , while in the other case one can often include the neighboring points of a hyperedge into an enlarged hyperedge to obtain a pure hyperedge potential.

Most of the following examples are based on the Delaunay graph. For simplicity, we will then often confine ourselves to the case $d = 2$, in which the stability is ensured by Remark 3.3 as soon as φ is bounded from below. But the reader should note that analogous results hold also in higher dimensions when φ is nonnegative, so that **(S)** is trivial.

4.1 Pure hyperedge interactions

In this subsection we consider examples of hyperedge potentials φ which only depend on the first parameter, so that $\varphi(\eta, \omega) = \varphi(\eta, \eta) =: \varphi(\eta)$.

4.1.1 Many-body interactions of finite range

Let $r > 0$ and $\mathcal{E} = \text{LC}_r$ be the locally complete graph of Example 2.1, and suppose that $\varphi(\eta, \omega) = \varphi(\eta)$. Remark 3.1 then shows that a Gibbs measure exists as soon as the potential φ is stable and $\varphi(\{0\}) < \infty$. By Remark 3.4, the first condition is equivalent to the classical stability assumption, and the second is necessary for defining Gibbs measures. So, as was observed first in [14, Remark 4.2], the techniques used here allow to weaken the superstability assumption of Ruelle's classical existence result [23], provided the interaction has finite range. But our techniques neither allow to treat the case of infinite range nor to rederive Ruelle's probability estimates.

An example of a stable but not superstable many-body interaction is the so-called quermass interaction. In space dimension $d = 2$, the associated Hamiltonian for a configuration ω is a linear combination of area, perimeter and Euler-Poincaré characteristic of the union of all discs of fixed radius that are centered at the points of ω . By the additivity of the Minkowski functionals, this Hamiltonian can be expressed in terms of a many-body interaction of finite range. Its stability is proved in [18], but the superstability fails. The existence of Gibbs processes for this interaction has been proved in [7] by the same methods used here.

4.1.2 Delaunay edge interactions

Here we consider two classes of potentials φ on the hypergraph structure Del_2 in \mathbb{R}^2 which are bounded below and depend only on the length of the Delaunay edges, in that

$\varphi(\{x, y\}) := \phi(|x - y|)$ when $\{x, y\} \in \text{Del}_2$. Such potentials have been studied in [1, 2, 3, 4]. We improve the existence results of these papers.

4.1.2.1 Polynomially increasing edge interactions. Suppose that

$$(4.1) \quad \phi(\ell) \leq \kappa_0 + \kappa_1 \ell^\alpha \quad \text{for some constants } \kappa_0 \geq 0, \kappa_1 \geq 0 \text{ and } \alpha > 0 .$$

Recall the definition of ϱ_0 in Remark 3.6.

Proposition 4.1. *Let $d = 2$ and φ be a pure edge potential on Del_2 which is bounded below and satisfies (4.1). Then there exists at least one Gibbs measure for φ and every activity*

$$z > (1 + 2\varrho_0)e^{3\kappa_0}(3\alpha e^2 \kappa_1 / 2)^{1/\alpha} / (\pi \varrho_0^2).$$

Proof. We apply Corollary 3.4 with the assumption (\mathbf{U}^A) . The stability condition (\mathbf{S}) holds because of Remark 3.3. We know further from Example 2.2 that every edge $\eta \in \text{Del}_2(\omega)$ has the finite horizon $\bar{B}(\eta, \omega)$. This shows that assumption (\mathbf{R}) is satisfied. Concerning assumption (\mathbf{U}^A) , let \mathbf{M} and Γ^A be defined as in Remark 3.6 with $b = \varrho_0 a$ and a to be chosen later. The assumptions $(\mathbf{U}1^A)$ and $(\mathbf{U}2^A)$ are then trivial. We also find that $c_A \leq 3(\kappa_0 + \kappa_1 a^\alpha (1 + 2\varrho_0)^\alpha)$. So, $(\mathbf{U}3^A)$ holds as soon as $z > K_0 e^{K_1 a^\alpha} / a^2$, where $K_0 = e^{3\kappa_0} / (\pi \varrho_0^2)$ and $K_1 = 3\kappa_1 (1 + 2\varrho_0)^\alpha$. Optimizing over a we find the value $a = (2/\alpha K_1)^{1/\alpha}$, which gives the sufficient condition $z > K_0 (\alpha K_1 e^2 / 2)^{1/\alpha}$. \square \square

The bound for z in Proposition 4.1 is certainly not the best possible. However, in the bounded case $\kappa_1 = 0$ it reduces to the trivial condition $z > 0$. As for condition (4.1), at first sight it might seem surprising that $\phi(\ell)$ does not necessarily converge to zero when ℓ tends to infinity, but is even allowed to converge to ∞ . However, $\phi(\ell)$ should not be compared with a pairwise interaction that must decay over large distances, but with a nearest-neighbor interaction between oscillators that form an elastic lattice. A potential like the harmonic interaction $\phi(\ell) = \ell^2$ is therefore quite natural.

4.1.2.2 Long-edge exclusion. Suppose there are constants $0 \leq \ell_0 < \ell_1 \leq \ell_2$ such that

$$(4.2) \quad \sup_{\ell_0 \leq \ell \leq \ell_1} \phi(\ell) < \infty \quad \text{and} \quad \phi(\ell) = \infty \quad \text{if } \ell > \ell_2.$$

Proposition 4.2. *Let $d = 2$ and φ be a pure edge potential on Del_2 which is bounded below and satisfies (4.2). Then there exists at least one Gibbs measure for φ and every $z > 0$.*

Proof. We apply Corollary 3.4 with the assumption $(\hat{\mathbf{U}}^A)$. Hypotheses (\mathbf{S}) and (\mathbf{R}) follow as in Proposition 4.1. Condition $(\hat{\mathbf{U}}1^A)$ is satisfied since the long-edge exclusion condition ensures a minimal density of points. The values a and b in the definition of \mathbf{M} and Γ are now chosen such that each $\{x, y\} \in \text{Del}_2(\omega)$ (with $\omega \in \bar{\Gamma}^A$) satisfies $\ell_0 < |x - y| < \ell_1$. Then $(\hat{\mathbf{U}}2^A)$ follows, and $(\hat{\mathbf{U}}3^A)$ is obvious. \square \square

Since $\phi(\ell)$ may be equal to ∞ for ℓ less than some $r_0 < \ell_0$, the present example includes the classical case of a hard-core interaction which forces the points to keep distance at least $r_0 > 0$ from each other.

Remark 4.1. Let us consider whether or not Propositions 4.1 and 4.2 remain valid when the Delaunay graph is replaced by a subgraph, as was discussed in Remark 3.5. One possible subgraph is the Gabriel graph. By definition, the Gabriel edge set $\text{Gab}_2(\omega)$ consists of all edges $\{x, y\} \in \text{Del}_2(\omega)$ for which the open disc with center $(x + y)/2$ and radius $|x - y|/2$ contains no point of ω . Since Gab_2 is shift-invariant and local, the restriction of φ to Gab_2 remains a shift-invariant hyperedge potential satisfying the finite horizon property (2.1). By Remark 3.3, **(S)** is inherited when we pass from Del_2 to Gab_2 . In view of our choice of M and Γ , the constant c_Γ remains unchanged, so that **(U3)** is also inherited without modification of the valid range for z . This means that Proposition 4.1 holds also for the Gabriel graph. Unfortunately, condition $(\hat{U}1^A)$ is lost, so that Proposition 4.2 does not carry over to the Gabriel graph. Another example of a Delaunay subgraph is the minimum spanning tree graph. This is tailor-made to be non-local, so that our results cannot be applied.

4.1.3 Delaunay tile interactions

In this subsection we deal with potentials on the hypergraph structure Del_3 of all Delaunay triangles in \mathbb{R}^2 , and we still assume that φ is bounded below and depends only on the hyperedge and not on the remaining configuration. Such models have been considered recently in [6] and [8]. We improve the existence results given there.

4.1.3.1 Polynomially increasing triangle interactions. The first example is specified by assuming that the interaction of a Delaunay triangle $\eta \in \text{Del}_3$ is controlled by its size as in Proposition 4.1:

$$(4.3) \quad \varphi(\eta) \leq \kappa_0 + \kappa_1 \delta(\eta)^\alpha \text{ for some constants } \kappa_0 \geq 0, \kappa_1 \geq 0 \text{ and } \alpha > 0,$$

where $\delta(\eta)$ is the diameter of the circumcircle of η . We then have a similar result.

Proposition 4.3. *Let $d = 2$ and φ be a pure triangle potential on Del_3 which is bounded below and satisfies (4.3). Then there exists at least one Gibbs measure for φ and every*

$$z > ((2/\sqrt{3}) + 2\rho_0)e^{2\kappa_0}(\alpha e^2 \kappa_1)^{1/\alpha} / (\pi \rho_0^2).$$

If φ is bounded above, so that $\kappa_1 = 0$, we recover the existence result of [8].

Proof. The proof is identical to that of Proposition 4.1. One simply has to note that $\delta(\eta) \leq ((2/\sqrt{3}) + 2\rho_0)a$ when $\eta \in \text{Del}_3(\omega)$ for some $\omega \in \bar{\Gamma}$. \square \square

4.1.3.2 Shape-dependent triangle interactions. The shape of a Delaunay triangle can be captured by its angles. So let $\beta(\eta)$ and $\gamma(\eta)$ respectively denote the smallest and the largest interior angle of a triangle η . We assume that φ has the form

$$(4.4) \quad \varphi(\eta) := \phi(\beta(\eta), \gamma(\eta)).$$

Proposition 4.4. *Let $d = 2$ and φ be a pure triangle potential on Del_3 bounded below and satisfying (4.4). Suppose there exists some $\delta > 0$ such that*

$$\sup_{\gamma \geq \beta > (\pi/3) - \delta} \phi(\beta, \gamma) < \infty.$$

Then there exists at least one Gibbs measure for φ and every $z > 0$.

Proof. We can again apply Corollary 3.4 with the assumption (\mathbf{U}^A) . The hypotheses (\mathbf{S}) and (\mathbf{R}) hold for the same reasons as in Proposition 4.1. The matrix \mathbf{M} and the set Γ^A can be defined as before, except that b is now chosen so small that $\beta(\eta) > (\pi/3) - \delta$ for all $\eta \in \text{Del}_3(\omega)$ with $\omega \in \bar{\Gamma}^A$. Conditions $(\mathbf{U}1^A)$ and $(\mathbf{U}2^A)$ are then obvious. By Remark 3.2, it only remains to check $(\hat{\mathbf{U}}3^A)$ because φ is scale invariant. But this is trivial. \square \square

The conditions of Proposition 4.4 include the *hard-equilaterality model*, in which

$$(4.5) \quad \phi(\beta, \gamma) = \begin{cases} \infty & \text{if } \beta \leq (\pi/3) - \delta, \\ 0 & \text{otherwise} \end{cases}$$

for some $0 < \delta < \pi/3$. The associated Gibbs measure produces a random Delaunay triangulation for which each triangle is almost equilateral in the sense that every angle exceeds $\pi/3 - \delta$. This model is considered in [10] (Model 1, Section 2.3) as the crystallized triangulation model.

4.2 Neighborhood-dependent hyperedge interactions

We now turn to examples of potentials φ for which $\varphi(\eta, \omega)$ does not only depend on η but also on the points of ω in some neighborhood of η . In two of these examples, the underlying hypergraph structure is based on the singleton graph

$$(4.6) \quad \text{SG} = \{(\{x\}, \omega) : x \in \omega \in \Omega\}$$

for which all hyperedges are singletons. Since this might seem trivial, we add immediately that more complex hyperedges will be hidden in the horizons of the potentials.

4.2.1 Forced-clustering k -nearest neighbor interactions

In the k -nearest neighbors model on the space \mathbb{R}^d of any dimension d , each point $x \in \omega$ interacts with the k points of ω that are closest to x . This model was introduced in [1, 3]. We focus here on a non-hereditary variant, in which the k -nearest neighbors of x are forced to keep within distance $\delta > 0$ from each other. This model was mentioned in [9] without precise justification of the proof, which is given here.

Let $k \geq 1$ be some fixed positive integer. For each integer $1 \leq i \leq k$, each configuration $\omega \in \Omega$ with $\#(\omega) \geq k+1$, and each $x \in \omega$ we let $x^{i:\omega}$ be the i -th nearest neighbor of x in ω . More precisely, $x^{i:\omega}$ is defined as the i -th element of ω in the total order $<_x$ on \mathbb{R}^d , in which $y_1 <_x y_2$ if and only if

$$|y_1 - x| < |y_2 - x| \text{ or } (|y_1 - x| = |y_2 - x| \text{ and } y_1 <_d y_2).$$

Here, $<_d$ stands for the lexicographic order on \mathbb{R}^d . We also set $x^{0:\omega} = x$.

Now let $\text{SG}_k = \{(\{x\}, \omega) \in \text{SG} : \#(\omega) \geq k+1\}$. We consider hyperedge potentials φ on SG_k of the form

$$(4.7) \quad \varphi(\{x\}, \omega) = \begin{cases} \phi(x^{0:\omega}, \dots, x^{k:\omega}) & \text{if } \text{diam}\{x^{0:\omega}, \dots, x^{k:\omega}\} < \delta, \\ \infty & \text{otherwise,} \end{cases}$$

where $\phi : (\mathbb{R}^d)^{k+1} \rightarrow \mathbb{R}$ is any *bounded* measurable function and $\delta > 0$ a fixed constant. It is clear that this interaction is non-hereditary in the sense that the removal of a particle from

an allowed configuration can lead to a forbidden configuration. Since no kind of symmetry is required of ϕ , the present formalism includes both the directed and undirected k -nearest neighbor graphs.

Proposition 4.5. *In the forced-clustering k -nearest neighbor model described above, there exists at least one Gibbs measure for φ and every $z > 0$.*

Proof. Let us apply Theorem 3.2. The hypergraph structure \mathbf{SG}_k is clearly sublinear and φ is bounded below. The stability assumption **(S)** thus follows from Remark 3.3. Next, for given $(\{x\}, \omega) \in \mathbf{SG}_k$, the closed ball $\bar{B}(x, |x^{k:\omega} - x|)$ with center x and radius $|x^{k:\omega} - x|$ can serve as a horizon of $(\{x\}, \omega)$. Since the corresponding open ball contains at most k points of ω , assumption **(R)** follows immediately. Concerning **(U)**, let $b > 0$ be a number to be specified later and $\mathbf{M} = a\mathbf{E}$ as in Remark 3.1 for some $a > 2(b+\delta)$. We set

$$\Gamma = \{\omega = \{x_0, x_1, \dots, x_k\} : x_0 \in B(0, b), x_i \in B(x_0, \delta/2) \ \forall i = 1, \dots, k\}.$$

For each $\omega \in \bar{\Gamma}$ and $x \in \omega$ it then follows that $|x - x^{k:\omega}| < \delta$. So (U1) holds with $r_\Gamma = \delta$, and (U2) follows with $c_\Gamma^\dagger \leq (k+1)\|\phi\|_\infty$. Finally, we have

$$\begin{aligned} e^{z|C|} \Pi_C^z(\Gamma) &= \frac{z^{k+1}}{(k+1)!} \int_C \dots \int_C \mathbb{1}_\Gamma(\{x_0, \dots, x_k\}) dx_0 \dots dx_k \\ &\geq \frac{z^{k+1}}{(k+1)!} \nu_d^{k+1} b^d (\delta/2)^{kd}, \end{aligned}$$

where ν_d is the volume of the unit ball in \mathbb{R}^d . This shows that (U3) holds as soon as b is chosen large enough. □ □

4.2.2 Voronoi cell interactions

Here we consider potentials that depend on the structure of the Voronoi cells. This type of model was introduced first by Ord; see the discussion in [21]. The spatial dimension d is arbitrary here. Specifically, let

$$\mathbf{SG}_b = \{(\{x\}, \omega) \in \mathbf{SG} : \text{Vor}_\omega(x) \text{ is bounded}\}$$

be the hypergraph structure of singletons with bounded Voronoi cells and φ be of the form

$$\varphi(\{x\}, \omega) = \phi(\text{Vor}_\omega(x)) \quad \text{for } (\{x\}, \omega) \in \mathbf{SG}_b.$$

For instance, ϕ might depend on the number of faces, the volume, or the surface area of the Voronoi cells. The necessity of passing from \mathbf{SG} to \mathbf{SG}_b was discussed in Example 2.2. Each $(\{x\}, \omega) \in \mathbf{SG}_b$ has as finite horizon the bounded Voronoi flower

$$(4.8) \quad \Delta = \bigcup_{\xi \in \text{Del}(\omega): \xi \ni x} \bar{B}(\xi, \omega).$$

The following proposition includes a result of [1].

Proposition 4.6. *In the Voronoi cell interaction model with bounded ϕ , there exists at least one Gibbs measure for φ and every $z > 0$.*

Proof. Let us apply Corollary 3.4 again. The stability comes from the sublinearity of $\text{SG}_b(\omega)$ and the lower boundedness of φ , cf. Remark 3.3. The proof of (\mathbf{U}^A) is essentially the same as in Proposition 4.1 with $\kappa_1 = 0$. Concerning the range condition (\mathbf{R}) , let $\{x\} \in \text{SG}_b(\omega)$ and y_1, y_2 any two points of the set Δ defined in (4.8). Then there exist $\xi_1, \xi_2 \in \text{Del}(\omega)$ such that $y_1 \in \bar{B}(\xi_1, \omega)$, $y_2 \in \bar{B}(\xi_2, \omega)$ and $x \in \bar{B}(\xi_1, \omega) \cap \bar{B}(\xi_2, \omega)$. The latter means that the union of these two balls is connected. By definition, $B(\xi_1, \omega)$ and $B(\xi_2, \omega)$ contain no point of ω . So, condition (\mathbf{R}) holds with $\ell_R = 2$, $n_R = 0$ and arbitrary δ_R . \square \square

In the above, the boundedness of ϕ was only assumed for simplicity. In analogy to Propositions 4.1 and 4.3, one can consider potentials φ that are polynomially increasing in the diameter of the cell's flower (4.8). The existence result then holds only for sufficiently large activity z . It is also straightforward to consider hard-exclusion models as in Propositions 4.2 and 4.4. For example, let $d = 2$ and

$$(4.9) \quad \phi(\text{Vor}_\omega(x)) = \begin{cases} 0 & \text{if } \text{Vor}_\omega(x) \text{ has six edges (and vertices),} \\ \infty & \text{otherwise.} \end{cases}$$

It is then clear that Theorem 3.2 applies, and by scale invariance a Gibbsian point process exists for all $z > 0$. The typical configurations of such a Gibbs process preserve the topology of the triangular lattice but are less regular than those of the hard-equilaterality model (4.5) for small δ . The model (4.9) may therefore be called the *randomly distorted triangular lattice*.

4.2.3 Adjacent Voronoi cell interactions

Here we reconsider the potential presented in Example 2.2, which describes an interaction between two adjacent Voronoi cells. That is, let $d = 2$ and the set of hyperedges of a configuration $\omega \in \Omega$ be given by

$$\text{Del}_{2,b}(\omega) := \{\{x, y\} \in \text{Del}_2(\omega) : \text{Vor}_\omega(x) \text{ and } \text{Vor}_\omega(y) \text{ are bounded}\}.$$

Suppose the potential φ has the form

$$\varphi(\eta, \omega) = \phi(\text{Vor}_\omega(x), \text{Vor}_\omega(y)) \quad \text{for } \eta = \{x, y\} \in \text{Del}_{2,b}(\omega).$$

For instance, $\varphi(\eta, \omega)$ can either depend on the length of the common edge or on the area ratio of the cells $\text{Vor}_\omega(x)$ and $\text{Vor}_\omega(y)$. As noticed before, the Voronoi “doubleflower”

$$\Delta = \bigcup_{\xi \in \text{Del}(\omega), \xi \cap \eta \neq \emptyset} \bar{B}(\xi, \omega).$$

then serves as finite horizon of $(\eta, \omega) \in \text{Del}_{2,b}$. In this setting we have the following result which can be proved in the same way as Proposition 4.6.

Proposition 4.7. *In the adjacent Voronoi cell interaction model in two dimensions with bounded ϕ , there exists at least one Gibbs measure for φ and every $z > 0$.*

As in the case of Proposition 4.6, this example can easily be extended to the case when ϕ is polynomially increasing in the diameter of the associated doubleflower, or when ϕ exhibits a hard exclusion that permits the configurations in $\bar{\Gamma}^A$ for a suitable choice of A . A particular adjacent Voronoi cell interaction with hard exclusion was proposed in [10], Model 3, Section 2.3. In this model, a hard exclusion forces the cells to be neither too small nor too large, and a smooth contribution induces a competition between the areas of adjacent cells.

4.2.4 Conclusion

The preceding series of examples presents only a selection of possible models and could easily be extended. For instance, having dealt with interactions acting on single Voronoi cells or pairs of adjacent Voronoi cells, we could proceed to triples of Voronoi cells with a common point, which are indexed by Delaunay triangles, or even larger clusters of Voronoi cells. On the other hand, the preceding interactions can be combined (i.e., added up) to obtain models with a richer interaction structure. A further universe of models opens up if one passes to marked configurations as in Remark 3.7. Future will show which kind of interaction will prove suitable for modeling geometric structures in the plane or in space that occur in the sciences. In any case, it should be evident that the conditions of our theorems are flexible enough to guarantee the existence of Gibbsian point processes for a large variety of geometric interactions.

5 Proofs of the theorems

Let \mathcal{E} , φ and z be fixed throughout this section. Before we enter into the proofs of the theorems, it is necessary to verify a basic ingredient of the theory of Gibbs measures, namely the consistency of the finite-volume Gibbs distributions.

Lemma 5.1. *Let $\Lambda \subset \Delta \Subset \mathbb{R}^d$ and $\omega \in \Omega_*^{\Delta, z}$. Then*

$$G_{\Delta, \omega}^z(\Omega_*^{\Lambda, z}) = 1 \quad \text{and} \quad \int f dG_{\Delta, \omega}^z = \int \left(\int f dG_{\Lambda, \tilde{\omega}}^z \right) G_{\Delta, \omega}^z(d\tilde{\omega})$$

for all measurable functions $f : \Omega \rightarrow [0, \infty[$.

Proof. Let Λ , Δ and ω be fixed. Since $G_{\Delta, \omega}^z$ does not depend on ω_Δ , we can assume for notational convenience that $\omega \subset \Delta^c$. Consider any two configurations $\zeta \in \Omega_\Lambda$ and $\xi \in \Omega_{\Delta \setminus \Lambda}$. It follows straight from the definition that $\mathcal{E}_\Lambda(\zeta \cup \xi \cup \omega) \subset \mathcal{E}_\Delta(\zeta \cup \xi \cup \omega)$. Hence, $H_{\Delta, \omega}(\zeta \cup \xi)$ is the sum of $H_{\Lambda, \xi \cup \omega}(\zeta)$ and a term in $\mathbb{R} \cup \{\infty\}$ which does not depend on ζ . So, $H_{\Lambda, \xi \cup \omega}^-(\zeta) < \infty$ when $H_{\Delta, \omega}^-(\zeta \cup \xi) < \infty$, and $H_{\Lambda, \xi \cup \omega}(\zeta) < \infty$ when $H_{\Delta, \omega}(\zeta \cup \xi) < \infty$. On the other hand, it follows that

$$H_{\Delta, \omega}(\zeta \cup \xi) + H_{\Lambda, \xi \cup \omega}(\zeta') = H_{\Delta, \omega}(\zeta' \cup \xi) + H_{\Lambda, \xi \cup \omega}(\zeta)$$

for all $\zeta' \in \Omega_\Lambda$. Taking the negative exponential and integrating over ζ' we thus find that

$$(5.1) \quad e^{-H_{\Delta, \omega}(\zeta \cup \xi)} Z_{\Lambda, \xi \cup \omega}^z = e^{-H_{\Lambda, \xi \cup \omega}(\zeta)} Z_{\Lambda, \Delta, \omega}^z(\xi),$$

where $Z_{\Lambda, \Delta, \omega}^z(\xi) = \int e^{-H_{\Delta, \omega}(\zeta' \cup \xi)} \Pi_{\Lambda}^z(d\zeta')$ is the partial partition function for which the configuration ξ is held fixed. Since $G_{\Delta, \omega}^z$ is concentrated on the set $\{H_{\Delta, \omega}^- < \infty, H_{\Delta, \omega} < \infty\}$ which is contained in $\{H_{\Lambda, \cdot \cup \omega}^- < \infty, H_{\Lambda, \cdot \cup \omega} < \infty\}$, we can conclude that

$$\begin{aligned} G_{\Delta, \omega}^z(Z_{\Lambda, \cdot}^z = 0) &= G_{\Delta, \omega}^z \circ \text{pr}_{\Delta \setminus \Lambda}^{-1}(Z_{\Lambda, \Delta, \omega}^z(\cdot) = 0) \\ &= (Z_{\Delta, \omega}^z)^{-1} \int Z_{\Lambda, \Delta, \omega}^z(\cdot) \mathbb{1}_{\{Z_{\Lambda, \Delta, \omega}^z(\cdot) = 0\}} d\Pi_{\Delta \setminus \Lambda}^z = 0. \end{aligned}$$

Likewise, $G_{\Delta, \omega}^z(Z_{\Lambda, \cdot}^z = \infty) = 0$ because $\int Z_{\Lambda, \Delta, \omega}^z(\cdot) d\Pi_{\Delta \setminus \Lambda}^z = Z_{\Lambda, \omega}^z < \infty$. Combining these results we obtain the first assertion of the lemma. As a consequence, we can divide Eq. (5.1) by $Z_{\Delta, \omega}^z Z_{\Lambda, \xi \cup \omega}^z$ to obtain the consistency equation $G_{\Delta, \omega}^z = \int G_{\Lambda, \tilde{\omega}}^z G_{\Delta, \omega}^z(d\tilde{\omega})$. \square

We now turn to the proof of Theorem 3.2. So we assume that the hypotheses **(S)**, **(R)** and **(U)** are satisfied. As usual, we construct a Gibbs measure as a limit of Gibbs distributions in suitable boxes. We choose M and Γ as in hypothesis **(U)** and consider for each $n \geq 1$ the paralleloptope

$$\Lambda_n = \bigcup_{k \in L_n} C(k),$$

where $L_n = \{-n, \dots, n\}^d$. Let $\bar{\omega} \in \bar{\Gamma}$ be a fixed pseudo-periodic configuration with $\sup_{k \in \mathbb{Z}^d} N_{C(k)}(\bar{\omega}) < \infty$; the last condition can be satisfied by letting $\bar{\omega}$ be periodic. Assumption **(U1)** implies that $\bar{\omega} \in \Omega_{\text{cr}}^{\Lambda_n}$. Combined with (3.1), **(S)** and **(U2&3)**, this shows that $\bar{\omega}$ is admissible for Λ_n and z ; cf. (5.8) below. So we can define the Gibbs distribution

$$G_n = G_{\Lambda_n, \bar{\omega}}^z \circ \text{pr}_{\Lambda_n}^{-1}$$

in Λ_n with boundary condition $\bar{\omega}$ and activity z , projected to Λ_n . Since we aim at constructing a shift-invariant Gibbs measure, we will introduce a spatial averaging of G_n , and it is convenient to work directly on the set of all shift-invariant probability measures on Ω .

So, let P_n be the probability measure on (Ω, \mathcal{F}) relative to which the configurations in the disjoint blocks $\Lambda_n + (2n+1)Mk, k \in \mathbb{Z}^d$, are independent with identical distribution G_n . We consider the averaged measure

$$(5.2) \quad \hat{P}_n = \frac{1}{v_n} \int_{\Lambda_n} P_n \circ \vartheta_x^{-1} dx,$$

where $v_n = |\Lambda_n|$ is the volume of Λ_n . By the periodicity of P_n , \hat{P}_n is shift-invariant. Moreover, the intensity $i(\hat{P}_n) = \int N_{\Lambda_n} dG_n / v_n$ of \hat{P}_n is finite because

$$\int N_{\Lambda_n} e^{-H_{\Lambda_n, \bar{\omega}}} d\Pi_{\Lambda_n}^z \leq e^{c_S \#(\partial \Lambda_n \bar{\omega})} \int N_{\Lambda_n} e^{c_S N_{\Lambda_n}} d\Pi_{\Lambda_n}^z < \infty$$

by (3.1) and **(S)**. So, $\hat{P}_n \in \mathcal{P}_{\Theta}$.

We will show that the sequence (\hat{P}_n) has an accumulation point in a suitable topology. As in [14], we will take the required compactness from the compactness of the level sets of the specific entropy. Let us recall the necessary concepts.

A measurable function $f : \Omega \rightarrow \mathbb{R}$ is called local and tame if

$$f(\omega) = f(\omega_{\Lambda}) \quad \text{and} \quad |f(\omega)| \leq a N_{\Lambda}(\omega) + b$$

for all $\omega \in \Omega$, some $\Lambda \in \mathbb{R}^d$ and suitable constants $a, b \geq 0$. Let \mathcal{L} be the set of all local and tame functions. The *topology of local convergence*, or \mathcal{L} -topology, on \mathcal{P}_Θ is then defined as the weak* topology induced by \mathcal{L} , i.e., as the smallest topology for which the mappings $P \mapsto \int f dP$ (with $f \in \mathcal{L}$) are continuous. Note that the intensity $P \mapsto i(P)$ is continuous in the \mathcal{L} -topology.

Next, for any $P \in \mathcal{P}_\Theta$ let $P_{\Lambda_n} = P \circ \text{pr}_{\Lambda_n}^{-1}$ be the projection of P to Ω_{Λ_n} and

$$I(P_{\Lambda_n} | \Pi_{\Lambda_n}^z) = \begin{cases} \int f \ln f d\Pi_{\Lambda_n}^z & \text{if } P_{\Lambda_n} \ll \Pi_{\Lambda_n}^z \text{ with density } f, \\ \infty & \text{otherwise} \end{cases}$$

the relative entropy of P_{Λ_n} with respect to $\Pi_{\Lambda_n}^z$; here, “ \ll ” stands for absolute continuity. The *specific entropy of P* (relative to Π^z) is then defined by

$$(5.3) \quad I^z(P) = \lim_{n \rightarrow \infty} v_n^{-1} I(P_{\Lambda_n} | \Pi_{\Lambda_n}^z);$$

see [12] and [15] for the existence of the limit and further properties of I^z . Our key tool is the following result of [13, Lemma 3.4], which is based on [15, Prop. 2.6].

Proposition 5.2. *For all $c_1, c_2 \geq 0$ and $z > 0$, the set*

$$\{P \in \mathcal{P}_\Theta : I^z(P) - c_1 i(P) \leq c_2\}$$

is relatively sequentially compact in the \mathcal{L} -topology.

In view of this fact, the following entropy bound implies that the sequence (\hat{P}_n) has a convergent subsequence.

Proposition 5.3. *In the limit $n \rightarrow \infty$ we have*

$$(5.4) \quad I^z(\hat{P}_n) - c_S i(\hat{P}_n) \leq |C|^{-1} (c_\Gamma - \ln \Pi_C^z(\Gamma)) + o(1).$$

Proof. First of all, the definition of \hat{P}_n readily implies that

$$(5.5) \quad I^z(\hat{P}_n) = v_n^{-1} I(G_n | \Pi_{\Lambda_n}^z);$$

see the proof of [12, Proposition (16.34)]. Likewise, $i(\hat{P}_n) = v_n^{-1} \int N_{\Lambda_n} dG_n$. By the definition of G_n , we know further that

$$(5.6) \quad I(G_n | \Pi_{\Lambda_n}^z) = - \int H_{\Lambda_n, \bar{\omega}} dG_n - \ln Z_{\Lambda_n, \bar{\omega}}^z.$$

So we need to estimate the two terms on the right-hand side. As for the first term, hypotheses **(S)** and **(U1)** give

$$(5.7) \quad \int H_{\Lambda_n, \bar{\omega}} dG_n \geq -c_S \int N_{\Lambda_n} dG_n - c_S \#(\partial_{\Lambda_n}^\Gamma \bar{\omega}),$$

and the assumption on $\bar{\omega}$ implies that $\#(\partial_{\Lambda_n}^\Gamma \bar{\omega}) = o(v_n)$.

It remains to estimate the partition function $Z_{\Lambda_n, \bar{\omega}}^z$. By **(U1)** we can find a number $m \geq 1$ such that $\partial \Lambda_n^\Gamma \subset \Lambda_{n+m}$ for all $n \geq 1$. Fix any n and let $\zeta \in \Omega_{\Lambda_n}$ be such that $\bar{\zeta} := \zeta \cup \bar{\omega}_{\Lambda_n^c} \in \bar{\Gamma}$. We claim that

$$(5.8) \quad H_{\Lambda_n, \bar{\omega}}(\zeta) \leq c_\Gamma \#L_n + o(v_n),$$

where the error term is uniform in ζ . Indeed, since $\eta \in \mathcal{E}_{\Lambda_n}(\bar{\zeta})$ when $\eta \in \mathcal{E}(\bar{\zeta})$ and $k \in \hat{\eta} \cap L_n$, we can write

$$H_{\Lambda_n, \bar{\omega}}(\zeta) = \sum_{k \in L_n} \sum_{\eta \in \mathcal{E}(\bar{\zeta}): \hat{\eta} \ni k} \frac{\varphi(\eta)}{\#\hat{\eta}} + \sum_{k \in L_{n+m} \setminus L_n} \sum_{\eta \in \mathcal{E}_{\Lambda_n}(\bar{\zeta}): \hat{\eta} \ni k} \frac{\varphi(\eta)}{\#\hat{\eta}}.$$

In view of (U2) and translation invariance, the first term on the right is not larger than $c_\Gamma \#L_n$. Likewise, the second term is dominated by $c_\Gamma^+ \#(L_{n+m} \setminus L_n)$. This proves (5.8) and leads us to the estimate

$$(5.9) \quad Z_{\Lambda_n, \bar{\omega}}^z \geq \int \mathbf{1}_{\bar{\Gamma}(\bar{\zeta})} e^{-H_{\Lambda_n, \bar{\omega}}(\zeta)} \Pi_{\Lambda_n}^z(d\zeta) \geq e^{-c_\Gamma \#L_n - o(v_n)} \Pi_C^z(\Gamma)^{\#L_n}.$$

Combining this with (5.5) to (5.7) we end up with (5.4). \square

The two propositions above imply that the sequence (\hat{P}_n) admits a subsequence that converges to some $\hat{P} \in \mathcal{P}_\Theta$ in the \mathcal{L} -topology. The limit \hat{P} is non-degenerate, in that $\hat{P} \neq \delta_\emptyset$. Indeed, in view of the lower semicontinuity of I^z (implied by Proposition 5.2) and the continuity of the intensity i we obtain from (5.4) that

$$(5.10) \quad I^z(\hat{P}) - c_S i(\hat{P}) \leq |C|^{-1} (c_\Gamma - \ln \Pi_C^z(\Gamma)).$$

But hypothesis (U3) ensures that the quantity on the right-hand side is strictly less than $z = I^z(\delta_\emptyset) - c_S i(\delta_\emptyset)$.

It is natural to expect that \hat{P} is the Gibbs measure we are looking for. Unfortunately, however, we are unable to show that \hat{P} is concentrated on the admissible configurations. However, since \hat{P} is non-degenerate, we can consider the conditioned measure $P = \hat{P}(\cdot | \{\emptyset\}^c) \in \mathcal{P}_\Theta$ with $P(\{\emptyset\}) = 0$ and apply Proposition 3.1. Let us give a more precise statement of this proposition.

Let ℓ_R, n_R, δ_R be the constants introduced in condition **(R)**. Also, let δ_- and δ_+ be the diameters of the largest open ball in C and of the smallest closed ball containing C , respectively. Fix an integer $m \geq 6\ell_R\delta_+/\delta_-$. For each $n \geq 1$, we decompose the parallelotope $\hat{\Lambda}_n := \Lambda_{n+(2n+1)m}$ into the $(2m+1)^d$ translates $\Lambda_n^k := \Lambda_n + (2n+1)\mathbf{M}k$ of Λ_n , where $k \in L_m$. For any $\Lambda \in \mathbb{R}^d$ let $n_\Lambda \geq 1$ be the smallest number with $\Lambda_{n_\Lambda} \supset \Lambda$ and $n_\Lambda \geq \delta_R/6\delta_+$. For all $n \geq n_\Lambda$ we consider the events

$$(5.11) \quad \hat{\Omega}_{\text{cr}}^{\Lambda, n} = \left\{ \min_{0 \neq k \in L_m} N_{\Lambda_n^k} > n_R \right\} \in \mathcal{F}_{\hat{\Lambda}_n \setminus \Lambda}$$

as well as $\hat{\Omega}_{\text{cr}}^\Lambda = \bigcup_{n \geq n_\Lambda} \hat{\Omega}_{\text{cr}}^{\Lambda, n} \in \mathcal{F}_{\Lambda^c}$. We then have the following result.

Proposition 5.4. *Given any $\Lambda \in \mathbb{R}^d$, we have $\hat{\Omega}_{\text{cr}}^\Lambda \subset \Omega_{\text{cr}}^\Lambda$ and $\partial\Lambda(\omega) \subset \hat{\Lambda}_n$ when $\omega \in \hat{\Omega}_{\text{cr}}^{\Lambda, n}$ for some $n \geq n_\Lambda$. Moreover, $P(\hat{\Omega}_{\text{cr}}^\Lambda) = 1$ for all $P \in \mathcal{P}_\Theta$ with $P(\{\emptyset\}) = 0$.*

Proof. Let Λ and $n \geq n_\Lambda$ be fixed and consider any $\omega \in \Omega$ for which the range of φ from Λ is not confined within $\hat{\Lambda}_n$. Then there exists a configuration $\tilde{\omega} \in \Omega$ with $\tilde{\omega} = \omega$ on $\hat{\Lambda}_n \setminus \Lambda$, a configuration $\zeta \in \Omega_\Lambda$ and a hyperedge $\eta \in \mathcal{E}_\Lambda(\zeta \cup \omega_{\Lambda^c})$ such that $\varphi(\eta, \zeta \cup \omega_{\Lambda^c}) \neq \varphi(\eta, \zeta \cup \tilde{\omega}_{\Lambda^c})$. So, every horizon Δ of $(\eta, \zeta \cup \omega_{\Lambda^c})$ as in (2.1) hits $\hat{\Lambda}_n$. By (2.4), Δ hits Λ too. Now let Δ be chosen as in **(R)**. We pick some $x \in \Lambda \cap \Delta$ and $y \in \Delta \setminus \hat{\Lambda}_n$ and, as in **(R)**, a chain of $\ell \leq \ell_R$ balls that hit each other successively and run from x to y .

There is a first ball $B_{\ell'}$ hitting $\hat{\Lambda}_n^c$. Shrinking $B_{\ell'}$ if necessary, we find a connected chain $B_1, \dots, B_{\ell'}$ of at most ℓ_R balls that are all contained in $\hat{\Lambda}_n$, connect x to $\partial\hat{\Lambda}_n$, and have either diameter at most δ_R or contain at most n_R particles of $\zeta \cup \omega_{\Lambda^c}$. Since the distance between x and $\partial\hat{\Lambda}_n$ is at least $m(2n+1)\delta_-$, at least one ball B_i has a diameter exceeding $3(2n+1)\delta_+$. As $n \geq \delta_R/6\delta_+$, this bound is larger than both δ_R and $3 \text{diam } \Lambda_n$. So there exist at least two indices $k, k' \in L_m$ such that Λ_n^k and $\Lambda_n^{k'}$ are included in B_i and thus hold at most n_R points of $\zeta \cup \omega_{\Lambda^c}$. At least one of these parallelotopes is different from Λ_n , say Λ_n^k . This proves that $\omega \notin \hat{\Omega}_{\text{cr}}^{\Lambda, n}$ and completes the proof of the first statement.

To prove the second claim let $P \in \mathcal{P}_{\Theta}$ be such that $P(\{\emptyset\}) = 0$. We have

$$1 - P(\hat{\Omega}_{\text{cr}}^{\Lambda}) = P\left(\bigcap_{n \geq n_{\Lambda}} (\hat{\Omega}_{\text{cr}}^{\Lambda, n})^c\right) \leq \inf_{n \geq n_{\Lambda}} \sum_{0 \neq k \in L_m} P(N_{\Lambda_n^k} \leq n_R).$$

By translation invariance, the last expression is equal to

$$(\#L_m - 1) \inf_{n \geq n_{\Lambda}} P(N_{\Lambda_n} \leq n_R).$$

But this term vanishes because

$$P(N_{\Lambda_n} \leq n_R) \rightarrow P(N_{\mathbb{R}^d} \leq n_R) = P(\{\emptyset\}) = 0$$

as $n \rightarrow \infty$. The next to last identity comes from the well-known fact [19, 6.1.3] that $P(0 < N_{\mathbb{R}^d} < \infty) = 0$ when P is translation invariant. The proof is therefore complete. \square

The final step in the proof of Theorem 3.2 is as follows.

Proposition 5.5. *The conditional probability $P = \hat{P}(\cdot | \{\emptyset\}^c) \in \mathcal{P}_{\Theta}$ is a Gibbs measure for \mathcal{E} , φ and z .*

Proof. Since $\hat{P} \in \mathcal{P}_{\Theta}$ with $\hat{P}(\{\emptyset\}) < 1$, P is well-defined and belongs to \mathcal{P}_{Θ} . To show that P is a Gibbs measure we fix some $\Lambda \Subset \mathbb{R}^d$ and consider the sets $\hat{\Omega}_{\text{cr}}^{\Lambda, p}$ defined in (5.11) for $p \geq n_{\Lambda}$. We also set $\hat{\Omega}_{\text{cr}}^{\Lambda, \leq p} = \bigcup_{n=n_{\Lambda}}^p \hat{\Omega}_{\text{cr}}^{\Lambda, n}$. It is sufficient to show that

$$(5.12) \quad \int_{\hat{\Omega}_{\text{cr}}^{\Lambda, \leq p}} f d\hat{P} = \int_{\hat{\Omega}_{\text{cr}}^{\Lambda, \leq p} \cap \Omega_{*}^{\Lambda, z}} f_{\Lambda} d\hat{P}$$

whenever $f : \Omega \rightarrow [0, 1]$ is a local function and $p \geq n_{\Lambda}$ is so large that f is $\mathcal{F}_{\hat{\Lambda}_p}$ -measurable. Here, f_{Λ} is defined by

$$f_{\Lambda}(\omega) := \int f dG_{\Lambda, \omega}.$$

Indeed, letting $p \rightarrow \infty$ and setting $f = 1$ we then find that $\hat{P}(\hat{\Omega}_{\text{cr}}^{\Lambda} \cap \Omega_{*}^{\Lambda, z}) = \hat{P}(\hat{\Omega}_{\text{cr}}^{\Lambda})$ and thus $P(\Omega_{*}^{\Lambda, z}) = 1$ by Proposition 5.4. For arbitrary f we obtain further that $P = \int G_{\Lambda, \omega}^z P(d\omega)$. Since Λ is arbitrary, this means that P is a Gibbs measure.

To prove (5.12) let f and $p \geq n_{\Lambda}$ be fixed. It will be convenient to replace the sequence (\hat{P}_n) introduced in (5.2) by an alternative sequence of measures with the same limit \hat{P} . Suppose n is so large that $\hat{\Lambda}_p \subset \Lambda_n$ and let

$$\Lambda_n^{\circ} = \{x \in \mathbb{R}^d : \hat{\Lambda}_p + x \subset \Lambda_n\}$$

be the “ $\hat{\Lambda}_p$ -interior” of Λ_n , which coincides with the closure of $\Lambda_{n-p-(2p+1)m}$. We define the (subprobability) measure

$$\bar{G}_n := \frac{1}{v_n} \int_{\Lambda_n^\circ} G_{\Lambda_n, \bar{\omega}}^z \circ \vartheta_x^{-1} dx = \frac{1}{v_n} \int_{\Lambda_n^\circ} G_{\Lambda_n - x, \vartheta_x \bar{\omega}}^z dx;$$

the equality comes from the shift-invariance of φ and the symmetry of Λ_n° . The argument in [15, Lemma 5.7] then shows that $\int f d\hat{P}_n - \int f d\bar{G}_n \rightarrow 0$ for all $f \in \mathcal{L}$. This means that \hat{P} can also be viewed as an accumulation point of the sequence (\bar{G}_n) . Now let $x \in \Lambda_n^\circ$, so that $\hat{\Lambda}_p \subset \Lambda_n - x$. Using the consistency lemma 5.1 and the fact that $\hat{\Omega}_{\text{cr}}^{\Lambda, \leq p} \in \mathcal{F}_{\hat{\Lambda}_p \setminus \Lambda} \subset \mathcal{F}_{(\Lambda_n - x) \setminus \Lambda}$ we find

$$\int_{\hat{\Omega}_{\text{cr}}^{\Lambda, \leq p}} f dG_{\Lambda_n - x, \vartheta_x \bar{\omega}}^z = \int_{\Omega_*^{\Lambda, z} \cap \hat{\Omega}_{\text{cr}}^{\Lambda, \leq p}} \left(\int f dG_{\Lambda, \omega}^z \right) G_{\Lambda_n - x, \vartheta_x \bar{\omega}}^z(d\omega),$$

and averaging over x yields

$$(5.13) \quad \int_{\hat{\Omega}_{\text{cr}}^{\Lambda, \leq p}} f d\bar{G}_n = \int_{\Omega_*^{\Lambda, z} \cap \hat{\Omega}_{\text{cr}}^{\Lambda, \leq p}} f_\Lambda d\bar{G}_n.$$

The integrand of the integral on the left is measurable with respect to $\mathcal{F}_{\hat{\Lambda}_p \setminus \Lambda}$ and thus belongs to \mathcal{L} . By (A.4) in the appendix, the integrand on the right of (5.13) is measurable with respect to the universal completion $\mathcal{F}_{\hat{\Lambda}_p \setminus \Lambda}^*$ and thus can be squeezed between two functions in \mathcal{L} which coincide \hat{P} -almost surely; cf. [5], Proposition 2.2.3. So, (5.13) gives (5.12) in the limit when n runs through a subsequence for which \bar{G}_n tends to \hat{P} in the \mathcal{L} -topology. \square

The proof of Theorem 3.3 requires only two minor observations. First, we note that $(\hat{U}1)$ and $(\hat{U}2)$ together with (\mathbf{R}) imply $(U1)$. Indeed, let $\omega \in \bar{\Gamma}$. Condition $(\hat{U}2)$ then shows that $H_{\Lambda_m, \omega}(\omega_{\Lambda_m}) < \infty$ for all $m \geq 1$. If m is so large that $av_m - b > n_R$ for the constant n_R in (\mathbf{R}) , the lower density bound $(\hat{U}1)$ thus gives that each translate Λ_m^k , $k \in \mathbb{Z}^d$, contains more than n_R points of ω . Hence, every ball with at most n_R points has a diameter no larger than $2 \text{diam}(\Lambda_m)$. Invoking the range condition (\mathbf{R}) , we can therefore conclude that $(U1)$ holds with $r_\Gamma \leq \ell_R \max(\delta_R, 2 \text{diam}(\Lambda_m))$.

Next we note that the non-rigidity condition in its strong form $(U3)$ was only used below (5.10) when we showed that the accumulation point \hat{P} is non-degenerate; for all other purposes, the weak form $(\hat{U}3)$ was sufficient. However, the non-degeneracy of \hat{P} is trivial under $(\hat{U}1)$ because $i(\hat{P}_n) \geq a - b/v_n$ and thus $i(\hat{P}) \geq a > 0$ by the continuity of i . This completes the proof of Theorem 3.3.

Appendix: Measurability

Here we collect and prove the measurability properties we have used and add a further comment on measurability. Let $\Lambda \Subset \mathbb{R}^d$ be fixed.

Claim A.1. $\mathcal{E}_\Lambda := \{(\eta, \omega) : \eta \in \mathcal{E}_\Lambda(\omega)\} \in (\mathcal{F}_f \otimes \mathcal{F})^*$.

Indeed, consider the measurable functions $f_\Lambda(\eta, \zeta, \omega) = (\eta, \zeta \cup \omega_{\Lambda^c})$ and $g(\eta, \zeta, \omega) = (\eta, \omega)$ from $\Omega_f \times \Omega_\Lambda \times \Omega$ to $\Omega_f \times \Omega$. Since \mathcal{E} and φ are measurable by assumption, the event $\bar{\mathcal{E}}_\Lambda := \{\varphi \circ g \neq \varphi \circ f_\Lambda\}$ then belongs to $\mathcal{F}_f \otimes \mathcal{F}_\Lambda \otimes \mathcal{F}$, and \mathcal{E}_Λ is equal to the projection image $g(\bar{\mathcal{E}}_\Lambda)$. Since \mathcal{F}_Λ is known [17, 19] to be the Borel σ -algebra for a Polish topology on Ω_Λ , one can apply Prop. 8.4.4 of [5] to conclude that \mathcal{E}_Λ is universally measurable, as claimed.

Claim A.2. *The functions $(\zeta, \omega) \rightarrow H_{\Lambda, \omega}(\zeta)$ and $(\zeta, \omega) \rightarrow H_{\Lambda, \omega}^-(\zeta)$ are measurable with respect to $(\mathcal{F}'_\Lambda \otimes \mathcal{F}_{\Lambda^c})^*$.*

To see this, we observe first that the mapping f_Λ in Claim A.1 is measurable from $\mathcal{F}_f \otimes \mathcal{F}'_\Lambda \otimes \mathcal{F}_{\Lambda^c}$ to $\mathcal{F}_f \otimes \mathcal{F}$, and therefore also from $(\mathcal{F}_f \otimes \mathcal{F}'_\Lambda \otimes \mathcal{F}_{\Lambda^c})^*$ to $(\mathcal{F}_f \otimes \mathcal{F})^*$; see [5], Lemma 8.4.6. In view of Claim A.1, this means that the indicator function $\mathbb{1}_{\mathcal{E}_\Lambda}(\eta, \zeta \cup \omega_{\Lambda^c})$ is measurable with respect to $(\mathcal{F}_f \otimes \mathcal{F}'_\Lambda \otimes \mathcal{F}_{\Lambda^c})^*$. Given any probability measure on $\Omega_f \times \Omega_\Lambda \times \Omega_{\Lambda^c}$, we can therefore squeeze this indicator function between two $\mathcal{F}_f \otimes \mathcal{F}'_\Lambda \otimes \mathcal{F}_{\Lambda^c}$ -measurable functions which coincide almost surely; cf. Proposition 2.2.3 of [5]. Writing

$$H_{\Lambda, \omega}(\zeta) = \sum_{\eta \subset \omega} \mathbb{1}_{\mathcal{E}_\Lambda}(\eta, \zeta \cup \omega_{\Lambda^c}) \varphi(\eta, \zeta \cup \omega_{\Lambda^c}),$$

applying Theorem 5.1.2 of [19] repeatedly when the indicator function is replaced by one of the squeezing functions and using Proposition 2.2.3 of [5] in the converse direction we get the result.

Claim A.3. *The partition function $\omega \rightarrow Z_{\Lambda, \omega}^z$ is measurable with respect to $\mathcal{F}_{\Lambda^c}^*$, $\Omega_*^{\Lambda, z} \in \mathcal{F}_{\Lambda^c}^*$, and $G_{\Lambda, \omega}^z(F)$ is a probability kernel from $(\Omega_*^{\Lambda, z}, \mathcal{F}_{\Lambda^c}^* |_{\Omega_*^{\Lambda, z}})$ to (Ω, \mathcal{F}) .*

Let P be an arbitrary probability measure on \mathcal{F}_{Λ^c} . As in Claim A.2, we can squeeze the function $e^{-H_{\Lambda, \omega}(\zeta)}$ between two $\mathcal{F}'_\Lambda \otimes \mathcal{F}_{\Lambda^c}$ -measurable functions which coincide $\Pi_\Lambda^z \otimes P$ -almost surely. Integrating these functions over ζ with respect to Π_Λ^z we obtain two functions of ω , which squeeze $Z_{\Lambda, \omega}^z$, are \mathcal{F}_{Λ^c} -measurable by the measurability part of Fubini's theorem, and coincide P -almost surely. As P was arbitrary, the first result follows. In the same way one finds that the function $\omega \rightarrow \Pi_\Lambda^z(H_{\Lambda, \omega}^- < \infty)$ is $\mathcal{F}_{\Lambda^c}^*$ -measurable. Hence

$$\Omega_*^{\Lambda, z} = \{\omega \in \Omega : \Pi_\Lambda^z(H_{\Lambda, \omega}^- < \infty) = 1, 0 < Z_{\Lambda, \omega}^z < \infty\} \in \mathcal{F}_{\Lambda^c}^*.$$

One also finds that the integral in (2.6) depends $\mathcal{F}_{\Lambda^c}^*$ -measurably on ω , which proves the last statement.

Claim A.4. *Let $p \geq n_\Lambda$ be fixed and suppose condition **(R)** holds. Claims A.2 and A.3 remain valid with $\mathcal{F}_{\hat{\Lambda}_p \setminus \Lambda}$ in place of \mathcal{F}_{Λ^c} as soon as all quantities are restricted to the set $\hat{\Omega}_{\text{cr}}^{\Lambda, p}$ defined in (5.11). In particular, $\Omega_*^{\Lambda, z} \cap \hat{\Omega}_{\text{cr}}^{\Lambda, p} \in \mathcal{F}_{\hat{\Lambda}_p \setminus \Lambda}^*$, and $G_{\Lambda, \omega}^z(F)$ is a probability kernel from $(\Omega_*^{\Lambda, z} \cap \hat{\Omega}_{\text{cr}}^{\Lambda, p}, \mathcal{F}_{\hat{\Lambda}_p \setminus \Lambda}^* |_{\Omega_*^{\Lambda, z} \cap \hat{\Omega}_{\text{cr}}^{\Lambda, p}})$ to (Ω, \mathcal{F}) .*

Indeed, by Proposition 5.4 and (3.1) we have

$$H_{\Lambda, \omega}(\zeta) = \sum_{\eta \in \mathcal{E}_\Lambda(\zeta \cup \omega_{\hat{\Lambda}_p \setminus \Lambda})} \varphi(\eta, \zeta \cup \omega_{\hat{\Lambda}_p \setminus \Lambda}) \quad \text{when } \omega \in \hat{\Omega}_{\text{cr}}^{\Lambda, p}.$$

The counterpart of Claim A.2 is therefore obvious, and the analog of Claim A.3 follows as before.

Claim A.5. *The use of universal measurability could be avoided by modifying the definition of \mathcal{E}_Λ .*

Namely, $\mathcal{E}_\Lambda(\omega)$ could be defined as the set of all $\eta \in \mathcal{E}(\omega)$ for which either $\eta \cap \Lambda \neq \emptyset$ or

$$\Pi_\Lambda^z \otimes \Pi_\Lambda^z \left((\zeta_1, \zeta_2) \in \Omega_\Lambda^2 : \varphi(\eta, \zeta_1 \cup \omega_{\Lambda^c}) \neq \varphi(\eta, \zeta_2 \cup \omega_{\Lambda^c}) \right) > 0.$$

Then $\mathcal{E}_\Lambda \in \mathcal{F}_f \otimes \mathcal{F}$ by the measurability part of Fubini's theorem, and Claims A.2, A.3 and A.4 would follow without the stars referring to universal extensions. This modified definition, however, is less intuitive and destroys the simple monotonicity of $\mathcal{E}_\Lambda(\omega)$ in Λ which was used in the proof of Lemma 5.1. One can still show that the required monotonicity holds for $\Pi_{\Delta \setminus \Lambda}^z$ -almost all ξ , but this is more involved. It is also necessary to redefine $\Omega_{\text{cr}}^\Lambda$ to obtain Proposition 5.4. We therefore decided to make use of universal measurability.

References

- [1] Bertin, E., Billiot, J.-M., Drouilhet, R.: Existence of “nearest-neighbour” spatial Gibbs models, *Adv. in Appl. Probab.* **31**, 895–909 (1999)
- [2] Bertin, E., Billiot, J.-M., Drouilhet, R.: Existence of Delaunay pairwise Gibbs point process with superstable component, *J. Statist. Phys.* **95**, 719–744 (1999)
- [3] Bertin, E., Billiot, J.-M., Drouilhet, R.: k -nearest-neighbour Gibbs point processes, *Markov Proc. Relat. Fields* **5**, 219–234 (1999)
- [4] Bertin, E., Billiot, J.-M., Drouilhet, R.: R -local Delaunay inhibition model, *J. Stat. Phys.* **132**, 649–667 (2008)
- [5] Cohn, D. L.: *Measure Theory*, Birkhäuser, Boston (1980)
- [6] Dereudre, D.: Gibbs Delaunay tessellations with geometric hard core conditions, *J. Statist. Phys.* **131**, 127–151 (2008)
- [7] Dereudre, D.: The existence of quermass-interaction processes for nonlocally stable interaction and nonbounded convex grains, *Adv. Appl. Prob.* **41**, 664–681 (2009)
- [8] Dereudre, D., Georgii, H.-O.: Variational characterisation of Gibbs measures with Delaunay triangle interaction, *Electron. J. Probab.* **14**, 2438–2462 (2009)
- [9] Dereudre, D., Lavancier, F.: Campbell equilibrium equation and pseudo-likelihood estimation for non-hereditary Gibbs point processes, *Bernoulli* **15-4**, 1368–1396 (2009)
- [10] Dereudre, D., Lavancier, F.: Practical simulation and estimation for Gibbs Delaunay-Voronoi tessellations with geometric hardcore interaction, hal-00402029 (2009)
- [11] Farhadifar, R., Röper, J.-C., Aigouy, B., Eaton, S., Jülicher, F.: The influence of cell mechanics, cell-cell interactions, and proliferation on epithelial packing, *Current Biology* **17-24**, 2095–2104 (2007)

- [12] Georgii, H.-O.: Gibbs Measures and Phase Transitions, de Gruyter, Berlin (1988)
- [13] Georgii, H.-O.: Large deviations and the equivalence of ensembles for Gibbsian particle systems with superstable interaction, *Probab. Theory Relat. Fields* **99**, 171–195 (1994)
- [14] Georgii, H.-O., Häggström, O.: Phase transition in continuum Potts models, *Commun. Math. Phys.* **181**, 507–528 (1996)
- [15] Georgii, H.-O., Zessin, H.: Large deviations and the maximum entropy principle for marked point random fields, *Probab. Theory Relat. Fields* **96**, 177–204 (1993)
- [16] Holley, R., Stroock, D. W.: Nearest neighbour birth and death processes on the real line, *Acta Math.* **140**, 103–154 (1978)
- [17] Kallenberg, O.: Random measures, 3rd edition, Akademie Verlag, Berlin (1983)
- [18] Kendall, W. S., Van Lieshout, M. N. M., Baddeley, A. J.: Quermass-interaction processes: conditions for stability, *Adv. Appl. Prob.* **31**, 315–342 (1999)
- [19] Matthes, K., Kerstan, J., Mecke, J.: *Infinitely Divisible Point Processes*, Wiley, Chichester (1978)
- [20] Lautensack, C., Sych, T.: 3D image analysis of open foams using random tessellations, *Image Anal Stereol* **25**, 87–93 (2006).
- [21] Ripley, B. D.: Modelling spatial patterns (with discussion), *J. Royal Statist. Soc.* **B 39**, 172–212 (1977)
- [22] Ruelle, D.: *Statistical Mechanics. Rigorous results*, Benjamin, New York (1969)
- [23] Ruelle, D.: Superstable interactions in classical statistical mechanics. *Commun. Math. Phys.* **18**, 127–159 (1970).
- [24] Schneider, R., Weil, W.: *Stochastic and integral geometry*, Springer, Berlin (2008)
- [25] Zessin, H.: The Gibbs cluster process, preprint (2005).