



**HAL**  
open science

# Characterization of Enhancers Active in the Mouse Embryonic Cerebral Cortex Suggests Sox/Pou cis-Regulatory Logics and Heterogeneity of Cortical Progenitors

Amandine Bery, Ben Martynoga, François Guillemot, Jean-Stéphane Joly, Sylvie Rétaux

## ► To cite this version:

Amandine Bery, Ben Martynoga, François Guillemot, Jean-Stéphane Joly, Sylvie Rétaux. Characterization of Enhancers Active in the Mouse Embryonic Cerebral Cortex Suggests Sox/Pou cis-Regulatory Logics and Heterogeneity of Cortical Progenitors. *Cerebral Cortex*, 2014, 24 (11), pp.2822-2834. 10.1093/cercor/bht126 . hal-00850761

**HAL Id: hal-00850761**

**<https://hal.science/hal-00850761>**

Submitted on 22 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Characterization of Enhancers Active in the Mouse Embryonic Cerebral Cortex Suggests Sox/Pou *cis*-Regulatory Logics and Heterogeneity of Cortical Progenitors

Amandine Bery<sup>1</sup>, Ben Martynoga<sup>3</sup>, François Guillemot<sup>3</sup>, Jean-Stéphane Joly<sup>2</sup> and Sylvie Rétaux<sup>1</sup>

<sup>1</sup>Equipe Development and Evolution of the Forebrain, <sup>2</sup>Equipe Morphogenesis of the Chordate Nervous System, UPR3294 N&D, Institut de Neurobiologie Alfred Fessard, CNRS, Gif-sur-Yvette, France and <sup>3</sup>MRC-NIMR, London, UK

Address correspondence to S. Rétaux, UPR 3294-NeD, Institut de Neurobiologie Alfred Fessard, CNRS, Avenue de la Terrasse, 91198 Gif-sur-Yvette cedex, France. Email: retaux@inaf.cnrs-gif.fr

**We aimed to identify *cis*-regulatory elements that control gene expression in progenitors of the cerebral cortex. A list of 975 putative enhancers were retrieved from a ChIP-Seq experiment performed in NS5 mouse stem cells with antibodies to Sox2, Brn2/Pou3f2, or Brn1/Pou3f3. Through a selection pipeline including gene ontology and expression pattern, we reduced the number of candidate enhancer sequences to 20. Ex vivo electroporation of green fluorescent pProtein (GFP) reporter constructs in the telencephalon of mouse embryos showed that 35% of the 20 selected candidate sequences displayed enhancer activity in the developing cortex at E13.5. In silico transcription factor binding site (TFBS) searches and mutagenesis experiments showed that enhancer activity is related to the presence of Sox/Pou TFBS pairs in the sequence. Comparative genomic analyses showed that enhancer activity is not related to the evolutionary conservation of the sequence. Finally, the combination of in utero electroporation of GFP reporter constructs with immunostaining for Tbr2 (basal progenitor marker) and phospho-histoneH3 (mitotic activity marker) demonstrated that each enhancer is specifically active in precise subpopulations of progenitors in the cortical germinal zone, highlighting the heterogeneity of these progenitors in terms of *cis*-regulation.**

**Keywords:** *cis*-regulation, ChIP-Seq, electroporation, Pou, Sox

## Introduction

The mammalian cortex is a finely organized structure that is thought to have facilitated the emergence of higher cognitive functions during evolution. The development of such a structure depends on tight regulation of the properties and the behavior of cortical progenitors during embryogenesis, through the activation of gene regulatory networks and the combinatorial activity of multiple transcription factors (TFs). Such factors involved in the expansion and differentiation of neural progenitors have been identified. But how they act in space and time is poorly understood because the distant *cis*-regulatory sequences that they bind to regulate gene expression remain mostly unknown. In this study, we set out to identify *cis*-regulatory elements that control gene expression, hence cellular behavior, of cortical progenitors.

During corticogenesis, neuroepithelial cells initially undergo exclusively proliferative, symmetric divisions, expanding the progenitor pool. They then switch to neurogenic divisions that give rise to the earliest born cortical neurons and to other progenitor populations distributing in layered subdivisions of the neuroepithelium (Boulder committee 1970; Bystron et al. 2008). Classically, 2 types of progenitors are recognized: (1) The apical progenitors (APs) including radial glial cells and short neuronal progenitors residing in the ventricular zone (VZ), and (2) the basal progenitors (BPs) that originate as daughters of APs and are located in the subventricular

zone (SVZ; Malatesta et al. 2000; Miyata et al. 2001; Noctor et al. 2001, 2004; Haubensak et al. 2004; Gal et al. 2006; Stancik et al. 2010; reviewed in Mérot et al. 2009). APs and BPs are distinguished on the basis of molecular markers such as *Pax6* or *Tbr2*, respectively (Hartfuss et al. 2001; Tarabykin et al. 2001; Heins et al. 2002; Nieto et al. 2004; Zimmer et al. 2004; Englund et al. 2005; Gal et al. 2006). APs and BPs also differ in their mitotic positions and patterns (reviewed in Sidman and Rakic 1973; Noctor et al. 2008; Taverna and Huttnner 2010). Of note, based on the varied behaviors and molecular properties exhibited by cells in the germinal zone, some authors have proposed that cortical progenitors are more heterogeneous than initially thought, even in lissencephalic or nonprimate cortices (Gal et al. 2006; Kowalczyk et al. 2009; Stancik et al. 2010; Shitamukai et al. 2011; Wang et al. 2011; Kelava et al. 2012; Reillo and Borrell 2012).

Transcription factors expressed in cortical progenitors critically influence their behavior. Among them, *Sox2* is involved in the acquisition and maintenance of neuronal progenitor identity (Bylund et al. 2003; Graham et al. 2003; Ellis et al. 2004). Together with Class III POU (Brn) TFs, *Sox2* binds the Nestin neural enhancer and regulates Nestin expression (Tanaka et al. 2004; Jin et al. 2009) in a cell cycle-dependent manner, thereby orchestrating gene regulation that correlates with the 3-dimensional morphological changes in neural progenitors in vivo (Sunabori et al. 2008). Another example is given by *Pax6*, which expression level controls the balance between neural stem cell self-renewal and neurogenesis (Sansom et al. 2009). Thus, these TFs are major effectors steering the dynamic progression of regulatory states during the generation of cell types.

Only few enhancers, active in neural stem cells, have been identified. The *Sox2* enhancer, SRR2, is active specifically in neural stem/progenitor cells and recruits Sox and POU TFs in telencephalic progenitors (Miyagi et al. 2006). More recently, Visel et al. (2013) have released an atlas of 329 enhancers active in the developing mouse telencephalon, some being active in the cortical germinal zone. From this dataset, they deduced *cis*-regulatory logics that confers regional identity in the telencephalic neuroepithelium in its tangential dimension (e.g., pallium vs. subpallium). Here, and in a similar manner, we sought to describe *cis*-regulatory logics underlying the behavior and possible heterogeneity of a radial compartment of the developing cortex: The progenitors of the germinal zone.

To address this question, we used ChIP-seq followed by functional assay in the embryonic mouse cortex to identify enhancers on which 3 specific cortical progenitor-expressed TFs bind. We report the identification and functional characterization of 7 sequences that behave as active enhancers in distinct categories of cortical progenitors in vivo.

## Materials and Methods

### ChIP-Seq

NS5 cells were kindly provided by Austin Smith and cultured under standard conditions (Conti et al. 2005). Cells were dual fixed with disuccinimidyl glutarate (2 nM) and 1% formaldehyde and then lysed and sonicated as previously described (Castro et al. 2011). Chromatin immunoprecipitation was performed according to the standard Upstate/Millipore protocol using specific antibodies to Sox2 (Santa Cruz sc-17320), Brn2/Pou3f2 (Santa Cruz sc-6029), or Brn1/Pou3f3 (kindly provided by Dr Dies Meijer). Immunoprecipitated material was sequenced according to Illumina's standard protocols. Reads were mapped with Bowtie to the mm9 genome assembly and significant peaks called with MACs v1.3 (Zhang et al. 2008). We defined 2892 regions significantly bound by Sox2 compared with input chromatin and 7176 regions bound by Brn1 and/or Brn2 (these were pooled due to predicted functional redundancy of these factors). Intersection of these lists of ChIP-seq peak coordinates revealed 925 genomic regions bound by Sox2 and Brn1 and/or Brn2. Of note, both Sox2 and Brn TFs are expressed homogeneously in NS5 cells, in a manner that is analogous to their expression throughout the cortical VZ in vivo (immunohistochemistry, data not shown). These data will be described more thoroughly in a manuscript currently in preparation in the Guillemot lab.

### In Silico Analyses of Candidate Enhancer Sequences

#### Evolutionary Conservation

The conservation of the potential candidate enhancers was studied through a multiple alignment program for comparative genomics, MLAGAN (Brudno et al. 2003), and the results were visualized via the VISTA server (Mayor et al. 2000). The different vertebrate species used in the alignment were mouse (*Mus musculus*, baseline), human (*Homo sapiens*), dog (*Canis familiaris*), platypus (*Ornithorhynchus anatinus*), gibbon (*Nomascus leucogenys*), chicken (*Gallus gallus*), xenopus (*Xenopus laevis*), medaka (*Oryzias latipes*), stickleback (*Gasterosteus aculeatus*), and fugu (*Takifugu rubripes*).

#### TFBS Search

p300 binding sites were retrieved from Visel et al. (2009) and compared with chromosomal positions of candidate enhancer sequences. Sox and Pou TF binding motifs were searched on the sequences as follows: [A/G]ACAA[T/A] and C[A/T]TTG[T/A][G/T/A] for Sox (Harley et al. 1994; Tomioka et al. 2002); AT[A/T]GCAT for POU (Schöler et al. 1989; Petryniak et al. 1990; Tomioka et al. 2002; Castro et al. 2006; Cook and Sturm 2008; Ryan and Rosenfeld 2011). Each sequence was scanned with both the forward and the reverse sequences of each motif, the distribution was visualized using the Regulatory Sequence Analysis Tool (van Helden 2003), and frequencies of occurrence were calculated.

#### PWM Calculation

The position weight matrix (PWM) for Sox and Pou transcription factor binding sites (TFBSs) present in the 7 active enhancers was calculated from the alignment of the different consensus motifs found in the enhancers (<http://genome.crg.es/courses/cshl2007/T2/MakeProfile.html>). Then, both PWM were visualized and identified through the JASPAR database (Wasserman and Sandelin 2004).

### Cloning of Target Enhancer Regions and Vector Constructs

Candidate enhancer sequences were amplified as attB3/attB4 PCR fragment using proper primers, AccuprimeTaq (Invitrogen) and mouse genomic DNA as template. Each amplified sequence was inserted in a plasmid to get a library of putative enhancers. This library was generated by Gateway cloning using a 3–4 donor vector and a customized destination vector (p336bglbGFP34HR3R4), where green fluorescent pProtein (GFP) reporter expression is under control of a  $\beta$ -globin minimal promoter. DNA preps for electroporation were endotoxin-free (Qiagen).

For mutagenesis experiments, the TFBSs of Pou (del1) or Sox (del2), or both (del3), were deleted by using the GENEART site-directed mutagenesis system and specific primers (sequences available in Supplementary Table 1).

### Electroporations

Time-pregnant Swiss mice were used, with the midpoint of the dark period as embryonic day 0 (E0). Animals were treated according to the regulations and laws of the European Union (86/609/EEC) for care and handling of animals in research, using procedure ref10/version1 approved by the Ethics Committee CEEA-Paris1. SR's authorization for animal experimentation is 91-329.

### Ex Vivo Electroporation and Cortical Slice Culture

E13.5 embryos were obtained by cesarian opening after cervical dislocation of the pregnant mother. One to 1.5  $\mu$ L of plasmid DNA (1  $\mu$ g/ $\mu$ L) mixed with Fast green was injected into the lateral ventricle, using a beveled and calibrated glass micropipette (pulled on Harvard apparatus) and a microinjector (Femtojet, Eppendorf). For electroporation, 2  $\times$  35 ms pulses of 35 mV with a 1-ms interval were delivered with two 5-mm electrode paddles positioned on either side of the head (BTX, ECM830 Harvard apparatus). Throughout procedure, the embryos were bathed in an ice-cold 1 $\times$  Krebs solution (for 10 $\times$ : 1.26 mM NaCl, 25 mM KCl, 12 mM NaH<sub>2</sub>PO<sub>4</sub>, 12 mM MgCl<sub>2</sub>, 25 mM CaCl<sub>2</sub>) plus 1 g glucose and 1.05 g NaHCO<sub>3</sub>.

Immediately after ex vivo electroporation, the brains of embryos were dissected out in ice-cold 1 $\times$  Krebs solution, embedded in 2% low-melting agarose, and sectioned at 250  $\mu$ m using a vibratome (VT1000S Leica Microsystems). Brain slices were transferred onto a slice culture insert (Millicell) in a plastic tissue culture dish with culture medium containing: 47.5 mL Neurobasal (Invitrogen/Gibco), 1 mL B27 (Invitrogen/Gibco), 0.5 mL 50% glucose, and 0.5 mL 100 $\times$  glutamine. Cultures were maintained in a humidified incubator at 37°C with 5%CO<sub>2</sub> supply. Pictures were taken with an Olympus binocular microscope.

### In Utero Electroporation

In utero electroporation was performed as described by Shimogori and Ogawa (2008) including few modifications. A time-pregnant Swiss mouse was anesthetized with a solution of Ketamine (Merial) and Xylazine (Bayer). The uterine horns containing E13.5 embryos were exposed. One to 1.5  $\mu$ L of plasmid DNA (1  $\mu$ g/ $\mu$ L) mixed with Fast green was manually microinjected through the uterine wall into the lateral ventricle, using a beveled and calibrated glass micropipette. Thirty-five-millisecond pulses of 35 mV with a 1-ms interval were delivered across the uterus with two 5-mm electrodes paddles positioned on either side of the head (BTX, ECM830 Harvard apparatus). Throughout the surgical procedure, the animal was placed on a heater block at 37°C and the uterus was bathed with warm phosphate buffer saline (PBS) (pH 7.4). After the procedure, the uterus was placed back in the abdominal cavity and the wound was surgically sutured. Twenty hours after electroporation the embryonic brains (E14.5) were dissected and fixed in 4% paraformaldehyde in PBS (pH7.4) for subsequent analyses.

### Immunohistochemistry

Fixed E14.5 brains were embedded in 2% agarose (made in MilliQ H<sub>2</sub>O) and sectioned at 80  $\mu$ m using a vibratome (VT1000S Leica Microsystems). The sections were washed several times in PBS/0.5%/Triton X-100/0.5% bovine serum albumine. Incubation with primary and secondary antibodies was performed at 4°C overnight. The primary antibodies utilized were as follows: Rabbit antiphospho-histone H3 (pH3; 1:500, Upstate Biotechnology), rabbit anti-Tbr2 (1:500, Abcam), and mouse anti-GFP monoclonal antibody (1:500, Roche). Before incubation with rabbit anti-Tbr2, the sections were blocked in PBS/2% Tween/2% Triton X-100 with 10% of blocking reagent (Roche) and 10% of normal goat serum (Sigma-Aldrich). Secondary antibodies were goat antirabbit IgG (H + L) Alexa 594 and goat antimouse IgG1 Alexa 488 (Molecular Probes). Finally, sections were washed and mounted in

ProLong Gold antifade reagent with 4',6-diamidino-2-phenylindole (Invitrogen). Photographs were taken with an ApoTome (Zeiss).

### Probes and In Situ Hybridization

cDNA probes for the 7 active enhancers were cloned by RT-PCR from E13.5 brain total RNA. The resulting DNA fragments were inserted in pCRII-TOPO (Invitrogen). After linearization, digoxigenin-labeled antisense RNA probes were generated by T3 or T7 RNA polymerase, and in situ hybridization (ISH) was performed as previously described (Abellan et al. 2010).

### Statistics

The statistical analyses were performed on Statview4.57 (Abacus Concept, Berkeley, CA, USA). The nonparametric Mann-Whitney test was used to compare the 2 populations of enhancers (GFP<sup>+</sup>/active and GFP<sup>-</sup>/inactive in cortical progenitors).

## Results

### Selection of Potential Candidate Enhancers Active in Cortical Progenitors From ChIP-Seq Data

A ChIP-seq experiment on mouse NS5 cells with antibodies specific for Sox2, Brn1, or Brn2 TFs provided a list of 925 potential candidate enhancer sequences, which bind these TFs in this neural stem cell line (Fig. 1). To shorten this list and select sequences to be tested for enhancer activity in the cortical germinal zone, 2 filters were applied. We reasoned that the closest coding gene to the candidate enhancer (found using the NCBI database) was most likely to be the one regulated by that enhancer. We therefore selected sequences that closest genes were:

1. Annotated with a gene ontology (GO) that was related to nervous system development and/or proliferation control. Selected GO terms for flanking genes were: Nervous system, forebrain, neurons, synapses, and axon guidance ( $n = 36$ ); cell proliferation, differentiation, and apoptosis ( $n = 35$ ); cell differentiation and nervous system development ( $n = 30$ ); cell proliferation and nervous system development ( $n = 9$ ); cell cycle ( $n = 8$ ); apoptosis of neurons only ( $n = 2$ ); and cell cycle and nervous system development ( $n = 1$ ). This first “GO” filter shortened down the list to 120 candidate enhancer sequences.
2. Expressed in the developing cortex, according to the following databases: GenePaint, Genesat, and Allen Brain Atlas. This filter further reduced the number of potential candidate enhancer sequences to 20, which are listed in Table 1.

### Features and Evolutionary Conservation of the 20 Candidate Enhancer Sequences


Although the selected 20 ChIP-seq-generated sequences were relatively evenly distributed around the closest neighboring genes, a slight bias toward an intronic localization was observed. Eight were intronic, 6 in the 5' region, and 6 in the 3' region (Table 1). Consistent with potential enhancers, selected sequences were located within 1–200 kb of the nearest gene. For those that are not intronic, their distance to the first exon of closest gene ranges from 2189 to 160 948 bp, with an average of 52 810 bp. The sequences sizes ranged from 232 to 1074 bp, with an average of 504 bp.

The evolutionary conservation of the 20 mouse candidate enhancer sequences was tested on genomic alignments from 10 vertebrate species, including mammals, birds, amphibians, and fishes, with the mouse genomic sequence as baseline (Fig. 2 and ninth column of Table 1). Only 1 of the 20 sequences, enh18(Pou2f1), showed conservation among all species including fish (Fig. 2A). The other sequences were either conserved between mammals only (such as enh20(Sox2), Fig. 2B) or else conserved between tetrapods only (such as enh06(Fubp3); Fig. 2C). Other, specific, conservation patterns were also found, as well as no conservation at all (ex: enh07(Fzd9) or enh12(Sox11); Table 1).

Most of these 20 putative enhancer sequences are novel and not characterized. Indeed, a search in the online database “VISTA Enhancer Browser” (Visel et al. 2007) retrieved none of the sequences listed in Table 1. Although this database contains enhancers for about half of the flanking genes presented in Table 1 (*Btg1*, *Cdb4*, *Gli3*, *Nr2f1*, *Sox11*, *Tle3*, *Sall3*, *Pou2f1*, and *Sox2*), none of them corresponds to sequences found in our study. Only 2 of our 20 potential enhancer sequences have already been reported as confirmed active enhancers: enh19(Nes) corresponds to the *nestin* enhancer called Nes30 (Tanaka et al. 2004); and enh20(Sox2) is the *Sox2* enhancer called SRR2 (Miyagi et al. 2006), and both are described as active in cortical progenitors (Table 1).

### Seven of 20 Candidate Enhancers Are Active in the Developing Cortex

To test the activity of the 18 novel putative enhancer sequences in cortical progenitors in vivo, these sequences were amplified from genomic DNA to generate a library of enhancers and were subcloned into an enhancer-reporter vector containing a minimal  $\beta$ -globin promoter (Fig. 1).



**Figure 1.** Pipeline for the selection, analysis, and testing of enhancers. The strategy used to narrow down the list of 925 putative enhancer sequences to a “testable” number of 20 is summarized, as well as the types of functional analyses performed to test and characterize these 20 sequences.

**Table 1**  
List of Candidate Enhancer Sequences Studied in this Paper

Enhancer	N°ENSEMBL of the closest flanking gene	Name of the flanking gene	Gene ID of closest flanking gene	Enhancer size (bp)	Chromosomal position	Position (5'/3'/intrinsic)	Distance to first exon of closest gene	Evolutionary conservation in vertebrates (mouse-based)	Activity in electroporated cortical slices
enh01	ENSMUSG00000030519	Apba2	11784	429	chr7:71779339-71779768	Intronic	132747	Human only	–
enh02	ENSMUSG00000036478	Btg1	12226	317	chr10:95771677-95771993	5'	99679	Mammals	–
enh03	ENSMUSG00000031871	Cdh5	12562	331	chr8:106488526-106488857	5'	136742	Mammals	–
enh04	ENSMUSG00000027490	E2f1	13555	682	chr2:154402926-154403608	3'	7338	Mammals	–
enh05	ENSMUSG00000041773	Enc1	13803	442	chr13:98022508-98022950	Intronic	11448	Tetrapods	–
enh06	ENSMUSG00000026843	Fubp3	320267	342	chr2:31465402-31465744	Intronic	29875	Tetrapods	–
enh07	ENSMUSG00000049551	Fzd9	14371	362	chr5:135747232-135747594	3'	20132	Not conserved	–
enh08	ENSMUSG00000021318	Gli3	14634	603	chr13:15716504-15717107	Intronic	160948	Human only	–
enh09	ENSMUSG00000029570	Lfn3	16848	551	chr5:141087050-141087600	Intronic	4000	Human only	–
enh10	ENSMUSG000000069171	Nr2f1	13865	355	chr13:78566962-78567317	3'	67217	Mammals	–
enh11	ENSMUSG00000022978	2610039C10Rik	66578	530	chr16:90709954-90710484	5'	17132	–	–
enh12	ENSMUSG00000063632	Sox11	20666	672	chr12:28260349-28261021	3'	55647	Not conserved	–
enh13	ENSMUSG00000032280	Tle3	21887	929	chr9:61377823-61378752	3'	157158	Mammals	–
enh14	ENSMUSG00000003050	Cdh4	12561	361	chr2:179272065-179272426	Intronic	75846	Human only	+
enh15	ENSMUSG00000024565	Sall3	20689	667	chr18:81448646-81449314	3'	26000	Chick only	+
enh16	ENSMUSG00000053129	Gsh1	14842	282	chr5:147967097-147967379	Intronic	32893	Fugu only	+
enh17	ENSMUSG00000068748	Ptprz1	19283	263	chr6:22872987-22873250	5'	47485	Human and chick	+
enh18	ENSMUSG00000026565	Pou2f1	18986	1045	chr1:167864977-167866022	5'	67217	All	+
enh19	ENSMUSG00000004891	Nes	18008	232	chr3:87778258-87778490	Intronic	3243	Mammals	(+) Josephson et al. (1998)
enh20	ENSMUSG00000074637	Sox2	20674	227	chr3:34552864-34553090	3'	3937	Mammals	(+) Zappone et al. (2000); Miyagi et al. (2006)

First, to validate the enhancer-reporter vector, 2 control *ex vivo* electroporation experiments were performed in the E13.5 mouse embryonic cortex. (1) Coelectroporation of pCMV-mCherry and the reporter vector without enhancer showed mCherry expression in the embryonic cortex, validating the electroporation and cortical slice culture procedures; but did not show GFP expression, validating that the minimal  $\beta$ -globin promoter had no basal activity without enhancer (Fig. 3A,B). (2) Coelectroporation of pCMV-mCherry and the reporter vector containing a known *Rnd2* enhancer (Heng et al. 2008) showed both mCherry and GFP expression, thus validating the use of the chosen vector for testing enhancer function in the cortex (data not shown).

When tested in the *ex vivo* coelectroporation assay with pCMV-mCherry, 5 of the 18 putative enhancers cloned in the reporter vector (27%) were active in the embryonic cortex between E13.5 and E14.5 (Fig. 3A,C, see also last column in Table 1, and Supplementary Figure 1 for inactive sequences electroporation tests). The 5 sequences that act as active enhancers in the mouse cortex are: enh14(Cdh4), enh15(Sall3), enh16(Gsh1), enh17(Ptprz1), and enh18(Pou2f1). Two of them are intronic, 2 are located in the 5' of the closest gene and 1 in the 3' of the closest gene. We also noticed that enh18 (Pou2f1), the only enhancer exhibiting pan-vertebrate conservation in the candidate list, is active and drives strong GFP expression (Table 1).

### TFBS Composition and Mutagenesis in Enhancers with Activity in the Developing Cortex

We then checked whether particular combinations of TFBSs correlate with active enhancers in the cortex.


First, the consensus TFBSs corresponding to Sox and Pou factors used in the ChIP-seq experiment were searched for in the 20 candidate sequences of Table 1 (see Materials and Methods), and their distributions were compared between active ( $n = 7$ , Fig. 4A) and inactive ( $n = 13$ , Fig. 4B) sequences.

This analysis highlighted several differences between the 2 categories of enhancers. First, Pou-type TFBSs (but not Sox) were more represented in active enhancer sequences (normalized to sequence length; Fig. 4C). Secondly, the average distance between TFBSs in Pou/Sox pairs were twice shorter in active enhancers ( $16 \pm 6$  bp) than in inactive sequences ( $40 \pm 8$  bp;  $P = 0.0446$ ). Thirdly, we also checked for the presence of p300 transcriptional coactivator binding sites: 71% of active enhancers sequences (i.e., 5 of 7 sequences) had a p300 binding site (source: Visel et al. 2009; forebrain dataset), while only 30% of inactive sequences showed such p300 peak (Fig. 4D).

In addition, the occurrences of Pou and Sox consensus TFBS in active enhancers were compared with a random segment of the genome. Sox TFBSs were 2.1 $\times$  more frequent and Pou TFBSs were 14 $\times$  more frequent in our 7 active enhancers than in mouse chromosome 16, showing a significant enrichment in these 2 TFBSs in the studied sequences, particularly for Pou sites (and see above, Fig. 4C). Finally, the PWMs for Sox and Pou TFBSs present in the 7 active enhancers were generated from the alignment of all the different consensus motifs scanned in these 7 enhancers (Fig. 4E). This shows that, in our system, the Sox-bound motifs rather correspond to a 7-bp sequence.

To test the hypothesis that a combination of Sox and Pou TFBSs is indeed important for the enhancer activity, we next performed enhancer mutagenesis by the deletion of TFBS motifs on 2 selected enhancers. We chose enh14 (Cdh4) because it harbors only 2 TFBS of interest, 1 Sox and 1 Pou; and we chose enh18 (Pou2f1) because it is on the contrary the most TFBS-rich among active sequences, containing 7 Sox and 5 Pou binding sites.

When tested in the *ex vivo* electroporation assay, enh14 (Cdh4) Pou-deleted (del1) or Sox-deleted (del2) were both totally inactive in the embryonic cortex (Fig. 5A), suggesting that the combination and interaction of the 2 binding sites are required to confer enhancer activity.



**Figure 2.** Evolutionary conservation of candidate enhancers. Genomic multialignments (Vista plots) are shown for enh18(Pou2f1) (A), enh20(Sox2) (B), and enh06(Fubp3) (C), with the mouse (*M. musculus*) sequence as baseline. As indicated in the left, alignments show 10 vertebrate species including mammals (*H. sapiens*, *C. familiaris*, *N. leucogenys*, and *O. anatinus -a monotreme*), birds (*G. gallus*), amphibians (*Xenopus*


We also used enh18 (Pou2f1) to challenge our above observation that the distance between Sox and Pou TFBSs may matter to confer enhancer activity. We therefore generated constructs with single or double deletions of the 2 closest Sox and Pou binding sites in enh18 (Fig. 5B, schema). Both one-Pou-deleted (del1) and one-Sox-deleted (del2) versions of enh18 were still active in the embryonic cortex. However, enh18del1 showed much weaker activity than the intact enh18 sequence [compare 75% of GFP-positive slices after intact enh18 electroporation with 31% with enh18(del1); Fig. 5B]. This suggests that adjacent TFBS may partly compensate for the deleted Pou site. Finally, the deletion of both TFBSs (del3) further reduced enh18 activity: We observed 27% of GFP-positive slices after enh18(del3) electroporation; and in all cases, the GFP-positive zone was very reduced when compared with the electroporated area visualized by mCherry (Fig. 5B). Thus, in enh18 (Pou2f1), the deletion of the 2 closest Sox and Pou binding sites strongly reduces the activity of the enhancer.

### Enhancer Activity in Cortical Progenitors

We next used in utero electroporation to further characterize the activity of selected enhancers in cortical progenitors at the cellular level and with a good anatomical resolution. We selected the 3 elements with strongest activities ex vivo, enh14 (Cdh4), enh15(Sall3), and enh18(Pou2f1) to perform reporter assays at E13.5. One day after in utero electroporation, embryos were removed and systematic double-labeling for GFP and either pH3 (a marker for mitotic cells) or *Tbr2* (a specific marker for cortical BPs) was performed in order to characterize the proliferative activity and the type of progenitor cells in which the enhancers were active, respectively (Fig. 6). pCMVCAGGS-GFP electroporation was taken as a control, as the strong and ubiquitous promoter drives GFP expression in all electroporated cells and their progeny, without cell type selectivity.

First, the position of GFP+ cells (i.e. the electroporated cells themselves and/or their progeny) in the cortex was studied (Figs 6 and 7A,B). Twenty-four hours after electroporation of control pCMVCAGGS-GFP (gray bars in Fig. 7), some of the GFP+ cells still resided in the VZ (14.3%, most probably corresponding to APs), while a large majority had migrated to the SVZ (62.2%) and some had already reached the cortical plate (CP, 22.6%). The distribution of GFP+ cells was markedly different from the pCMVCAGGS-GFP control for the 3 tested enhancers, highlighting the specific activity of each enhancer in given cell populations. Enh14(Cdh4) and enh15(Sall3) activities were almost in mirror image of enh18(Pou2f1) activity. Indeed, enh18(Pou2f1) was mostly active in the VZ-SVZ, while enh15(Sall3) and enh14(Cdh4) were active in the SVZ and CP, but not (or weakly) in the VZ. These significantly different activity patterns suggest that enh14(Cdh4) and enh15(Sall3) are activated specifically in cells only when they are engaged in the differentiation pathway, whereas cells in which enh18(Pou2f1) is active are likely to be germinal zone

*tropicalis*), and fishes (medaka: *O. latipes*; fugu: *T. rubripes*; stickleback: *G. aculeatus*). Percent identity (shown between 50% and 100%) is indicated in the right. Conservation in coding exonic regions is colored in blue, and that in noncoding intronic regions is colored in red. The last line of each panel corresponds to the enhancer sequence itself and is used to calibrate the multialignments (hence shows 100% conservation with the mouse baseline sequence).



**Figure 3.** Ex vivo electroporation of putative enhancer sequences. (A) Schema of an embryonic telencephalic hemisphere. The dark square and green cortical area is the region targeted for the ex vivo electroporation test. (B and C) Fluorescence photographs of E14.5 live slices of embryonic cortex cultured on membranes, 24 h after coelectroporation of GFP enhancer-reporter constructs (green) and pCMV-mCherry (red, control for electroporation). (B) Coelectroporation of p336bglbGFP34HR3R4 without enhancer (left panel, green) and pCMV-mCherry (middle panel, red). The right panel shows the merged images of green and red fluorescence. (C) Coelectroporation of p336bglbGFP34HR3R4 with the indicated enhancer and pCMV-mCherry. For each enhancer, the left image shows a low-power magnification of the electroporated telencephalic hemisphere (GFP, green) and the second (GFP, green), third (m-Cherry, red), and fourth (merged) images show higher-power magnifications on the electroporated area. Scale bar: 250  $\mu$ m.

progenitors.

We next examined the proliferative status of cells in which these 3 enhancers are active, though pH3 immunofluorescence staining (Figs 6A and 7C,D). In control pCMVCAGGS-GFP electroporated embryos, 5% of GFP+ cells underwent mitosis in the cortical germinal zones (pH3 total, Fig. 7C). In contrast, enh15(Sall3)-positive cells were never pH3-positive ( $n = 204$  electroporated cells from 4 embryos and 4 experiments), suggesting that this enhancer is never activated in mitotically

active cells. The percentages of pH3+ cells were not statistically different from the pCMVCAGGS-GFP control value for enh14 (Cdh4) and enh18(Pou2f1) electroporations. For these 2 latter enhancers, among the GFP+/pH3+ cells, we further distinguished between those lining the ventricular surface and those located above or far from the ventricular surface, up to the SVZ (see schema in Fig. 7A). In pCMVCAGGS-GFP electroporated cells, the majority of the mitoses occurred along the ventricle in a very apical position, strongly suggesting that they occur in APs, while a few GFP+ cells also underwent mitoses higher in the cortical depth. The same pattern was observed for cells with enh14(Cdh4) activity. In contrast, enh18(Pou2f1) was only active in mitotic cells that divided along the ventricular surface (Fig. 7D).

In addition, we sought to characterize the activity of the 3 enhancers in SVZ progenitors. More specifically, we assessed the BP identity of GFP+ cells through *Tbr2* immunofluorescence staining (Figs 6B and 7E). In control pCMVCAGGS-GFP electroporated embryos, about half of the SVZ GFP+ cells were *Tbr2* immunopositive. In contrast, after electroporation of the 3 enhancer-reporter constructs, only approximately 15% of the GFP+ cells located in the SVZ were *Tbr2*-positive, therefore suggesting that the population of *Tbr2*+ cells is heterogeneous.


#### mRNA Expression of Enhancer's Nearest Flanking Genes

To further refine the study of enh14(Cdh4), enh15(Sall3), and enh18(Pou2f1) patterns of activity, we asked whether these 3 enhancers reproduce entirely, partially, or not at all, the mRNA expression patterns of their flanking genes (of note, there is no direct evidence that these enhancers actually regulate the expression of the nearby genes). To this end, we performed ISH experiments at E11.5, E12.5, and E13.5 (Fig. 8A).

Along these 3 stages, *Cdh4* mRNA was expressed dynamically, first in the upper part of the VZ at E11.5, then in the intermediate zone (IZ) and at lower levels in the VZ/SVZ at E12.5, and finally in only the upper part of the IZ at E13.5 (Fig. 8A1). *Sall3* and *Pou2f1* mRNAs showed more prominent expression in the germinal zones of the cortex at the 3 studied stages. They were both strongly expressed in the VZ/SVZ, with *Pou2f1* (but not *Sall3*) also showing expression in the IZ at E12.5–E13.5 (Fig. 8A2,A3). In summary, the mRNAs for the flanking genes showed globally wider expression than did the activity of their respective putative tested enhancers.

#### Discussion

We have selected 20 sequences with potential enhancer activity in cortical progenitor cells from a ChIP-seq dataset, using a selection pipeline taking into account function and expression of the closest flanking genes. Seven of these 20 sequences (35%) are active enhancers, and this was correlated to a specific signature in terms of TFBS composition, but not evolutionary conservation. Further characterization of cell types in which 3 of these enhancers are active highlights a possible regulatory basis for the heterogeneity of cortical progenitors.




**Figure 4.** In silico analysis of the 20 candidate enhancer sequences. (*A* and *B*) Distribution of Sox and Pou TFBSs along the active (*A*) and inactive (*B*) sequences. Regulatory sequence analysis tools visualization shows Sox (red) and Pou (blue) TFBS as vertical lines along the sequences. Note the various lengths of the sequences (scale bar on top). (*C*) Histograms showing the mean frequency of nucleotides involved in Sox (red) and Pou (blue) TFBSs for active (act) enhancers (bright color) versus inactive (inact) sequences (dim color). The 2 asterisks for Pou TFBSs indicate  $P < 0.01$  (Mann–Whitney). (*D*) Histogram showing the percentage of sequences with p300 TFBS in active (black) and inactive sequences (gray). (*E*) Sox and Pou binding sites in cortical progenitors. First line: consensus TFBS from the literature. Second line: PWMs elicited from this study. Third line: comparison with general Uniprobe PWM.

### Predictive Value of the Selection Pipeline on the ChIP-seq Dataset

The ChIP-seq experiment that served as a basis of this study was performed on NS5 cells, a mouse cell line derived from embryonic stem cells (Conti et al. 2005). To select sequences with potential enhancer activity in cortical progenitors, we applied 2 filters, not on the sequences themselves, but on the closest flanking gene. The first filter was a functional annotation (GO term) including neuron/nervous system or proliferation, and the second one was an expression pattern in the

developing cortex, thereby conferring (1) neuronal progenitor and (2) regional brain specificity to the selected sequences to be tested. Of these 20 resulting sequences, about a third (7 of 20, 35%) behave as enhancers in cortical cells, thus showing a very significant enrichment in “cortical enhancers.” In comparison, using a similar approach with a selection pipeline on 3100 noncoding sequences based on human–fugu conservation and the presence of putative forebrain motifs, Pennacchio et al. (2006) showed that 17% of the tested sequences (4 of 23) are active in the mouse forebrain. Further, the sequences that






**Figure 5.** Mutagenesis by the deletion of Sox and/or Pou TFBSs on enh14(Cdh4) and enh18(Pou2f1). (A) enh14(Cdh4) mutagenesis. (B) enh18(Pou2f1) mutagenesis. On each panel, the top left photograph shows merged GFP and mCherry fluorescence images after electroporation of the intact enhancer. The top right panel shows the Pou (blue) and Sox (red) organization of TFBS on the sequence, as well as the TFBS-deleted constructs that were generated by mutagenesis. The bottom panels show GFP, mCherry, and merged fluorescence photographs after electroporation of mutagenized constructs. Left column, low magnification showing the entire slice/hemisphere. Other columns, higher magnification on the electroporated zone. The *N* given at the bottom left corner of each low magnification picture gives the number of slices in which GFP could be detected in the electroporated area, among all the correctly electroporated, RFP+, slices. Scale bar: 250  $\mu$ m.

were then characterized with some details through in utero electroporation indeed showed enhancer activity in cortical progenitors. This suggest that our selection pipeline confers a high probability for a selected sequence to actually be an active enhancer in a chosen cell type of a specific brain region and may be valuably applied in other cases. In addition, it also points that such a strategy could be advantageously used to find specific drivers for subsequent functional and transgenesis experiments. In the case of the cortex, very few developmental enhancers specifically active in particular cell populations have been characterized. These include the E1 enhancer element of the *Ngn2* gene that activity is confined to a subpopulation of progenitors predominantly in the region of the ventral and lateral pallium (Scardigli et al. 2001; Berger et al. 2004), or an intronic nestin enhancer active in the dorsal telencephalic germinal zone (Walker et al. 2010). More recently, a series of enhancers that can be used as tissue-specific reagents in different dorso-ventral domains of the telencephalon have been reported (Visel et al. 2013). Here, we provide 3 novel enhancers/drivers, with specific activities in certain types of cortical progenitors, and that may be used to study the biology of these progenitors through time-lapse imaging and gene function analyses, or to analyze their progeny through lineage studies. In a recent review on mammalian neural stem cells, Basak and Taylor (2009) indeed stressed the need for

“clean lineage tracing experiments,” hence the need for specific enhancers/promoters active in neural progenitor cells.

#### **TFBS Signature and Regulatory Logics for Enhancers Active in the Cortical Germinal Zone**

When compared with the 13 inactive sequences, the 7 active enhancer sequences showed a higher frequency of Pou TFBS; their TFBS organized in Sox/Pou pairs were close (16 bp) on the sequence; and 71% (5 of 7) presented a p300 ChIP-seq peak. p300 is a transcriptional coactivator that is specifically recruited at enhancers, and was therefore expected as a mark for active enhancers. Accordingly, only 4 of 13 (30%) of inactive sequences had a p300 binding site. However, this criterion does not appear to be strict, and we would actually have “missed” 2 active sequences (i.e., enh14(cdh4) and enh15 (Sall3)) if the criterion had been included in the selection pipeline; Visel et al. 2009). In the same line, the data from the DNase hypersensitivity sites (DHS) experiments (Sabo et al. 2004, 2006; John et al. 2011) of the ENCODE project (ENCODE Project Consortium et al. 2012) are relevant, but not a strict criterion, to be used to identify TFBSs. Indeed, DHS is observed for 6 of 7 (85%) of our active enhancers, and enh17(Ptprz1) would have been discarded from our selection if we had followed this criterion to select the sequences. In addition, DHS is present in 54% of our inactive sequences. These elements of




**Figure 6.** Characterization of cortical progenitors in which enh14(Cdh4), enh15(Sall3), and enh18(Pou2f1) are active after in utero electroporation. (A and B) Photographs of fixed cortical slices after anti-GFP (green), and anti-pH3 (red) (A) or anti-Tbr2 (red) (B) double immunofluorescence staining, 1 day after in utero electroporation in the telencephalon of E13.5 mouse embryos. For each panel, the electroporated construct is indicated on the left. In A and B panels, the first column shows a general view of the electroporated cortical area, and the second column shows a high magnification on the zone in white square, with arrowheads indicating colocalization of GFP and the marker. Scale bar: 50  $\mu$ m.

discussion strengthen the validity and accuracy of the criteria (GO term and expression pattern) we used in our selection pipeline.

Sox2, known as an essential player in establishing and maintaining neuronal progenitors in vertebrates (Mizuseki et al. 1998; Bylund et al. 2003; Graham et al. 2003; Uchikawa et al. 2003), is known to pair off with specific partners to regulate gene transcription (Kamachi et al. 2000; Kondoh and Kamachi 2010). In particular, Sox/Pou TF dimerization and complex recruitment on enhancers are classically reported in the literature, including on the enh19(Nestin) and enh20 (Sox2) sequences reported herein. Tanaka et al. (2004) have shown that Sox and Pou proteins work synergistically to activate the intronic Nes30 enhancer (present study enh19 (Nestin)) in the VZ and SVZ of the mouse embryonic spinal cord. More recently, Walker et al. (2010) showed that the rat nestin enhancer (96% identical to Nes30) is also active in the VZ and SVZ of the developing cortex. Concerning the Sox2 enhancer, gel-shift analyses have identified a binding site for a Sox/Pou complex in SRR2 (present study enh20(Sox2); Tomioka et al. 2002). This SRR2 enhancer sequence

functionally plays an important role in maintaining stem cells identity (Zappone et al. 2000) and drives transcription in neural stem/progenitor cells (Miyagi et al. 2006). Other examples of Sox/Pou complex formation exist, some in the context of stem cell biology: There is a precise correlation between the ability of Pou proteins to form a complex with Sox2 on the enhancer of undifferentiated transcription factor 1 (UTF1), which belongs to the core transcriptional network characterizing pluripotency and the ability to maintain the stem cell state in embryonic stem cells (Nishimoto et al. 1999). The same type of Pou/Sox complex binds to enhancers of Fgf4 (Reményi et al. 2003). Pou/Sox dimerization is also necessary for melanocyte development; and in this case, an interaction with p300 was also reported (Smit et al. 2000); or for *Drosophila* embryogenesis, in which case Sox, Pou but also bHLH, factors interact on a single enhancer of the slit gene (Ma et al. 2000). Our data are in line with these bibliographic elements. Further, we propose that a special regulatory logic may exist specifically in cortical progenitors, which includes a strong control by Pou TFs, as well as a crucial role for Pou/Sox TFs complexes. This logic is fully supported by our data:



**Figure 7.** Quantification of pH3-positive and *Tbr2*-positive cortical progenitors in which *enh14*(*Cdh4*), *enh15*(*Sall3*), and *enh18*(*Pou2f1*) are active. (A) Schema of the different layers/zones of the E14.5 developing cortex used for quantification. GFP-positive cells (green dots) were counted in the CP/IZ, in the subventricular zone (SVZ) defined by dense *Tbr2* immunostaining (gray), and in the ventricular zone (VZ). In the latter, the ventricular surface (corresponding to the thickness of 2 cell diameters) was distinguished from the rest of the VZ for pH3 analysis. (B) Distribution of GFP-positive cells in various cortical layers/germinal zones 1 day after in utero electroporation of GFP reporter-enhancer constructs. The color code for enhancers is indicated: CMV control: gray; *enh14*(*Cdh4*): purple; *enh15*(*Sall3*): turquoise, and *enh18*(*Pou2f1*): dark blue. (C and D) Quantification of GFP-positive cells with regard to pH3 immunoreactivity. The histogram in C concerns all layers, whereas the histogram in D distinguishes cells at the ventricular surface from others (see A). (E) Quantification of GFP-positive cells with regard to *Tbr2* immunoreactivity. In all panels: \* $P < 0.05$ ; \*\* $P < 0.01$ ; \*\*\* $P < 0.001$ ; \*\*\*\* $P < 0.0001$  (Mann–Whitney).

1. *enh15*(*Sall3*) is the only active enhancer without Pou TFBS on its sequence and is also never active in the VZ where self-renewing, stem cell-like radial glia progenitors reside,
2. *enh14*(*Cdh4*) and *enh18*(*Pou2f1*) do present Pou/Sox TFBS pair(s) on their sequence, and they are both active in proliferating VZ progenitors,
3. single or double Sox/Pou site deletions on *enh14* and *enh18* abolishes or strongly reduces enhancer activity, indicating that an interaction between the 2 TFBSs is required for enhancer activity, and
4. in *enh18*(*Pou2f1*) that contains multiple Sox and Pou binding sites, the single deletion of one Pou (but not one Sox) binding site decreases enhancer activity, pointing the importance of Pou TFBS for transcriptional control.

We have also found a tendency for the Sox/Pou TFBS pairs in active enhancers to be spaced by about 16 bp. This is to be compared with a distance of 3 or 0 bp on *Fgf4* or *UTF1* enhancers, respectively (Reményi et al. 2003), or of 54 or 28 bp on the *Drosophila* slit enhancer (Ma et al. 2000), and is therefore in the range of described distances allowing functional interaction of the 2 types of TFs on DNA. We have tested this distance issue through mutagenesis on *enh18*(*pou2f1*), choosing the 2 closest Pou and Sox sites on the sequence (28 bp between the selected pair). The severe reduction in enhancer activity observed after the double deletion of the closest


Pou/Sox pair on *enh18* suggests that the distance is probably a criterion to take into account when considering the functional/physical interactions of these 2 particular TFs on DNA.

### Enhancer Activity: cis-Regulatory Logics Underlying Cortical Progenitor Heterogeneity

We have characterized with some details the activity of *enh14*(*Cdh4*), *enh15*(*Sall3*), and *enh18*(*Pou2f1*). Strikingly, these 3 enhancers are active in clearly different types of progenitors, highlighting heterogeneity of progenitor cells in terms of gene cis-regulation (summarized in Fig. 8B). While *enh18*(*Pou2f1*) is active in proliferating APs and nonproliferating SVZ progenitors, *enh15*(*Sall3*) is on the contrary never active in APs—although the mRNAs of their closest flanking gene are expressed throughout the VZ/SVZ in a very similar pattern. Many recent neurodevelopment biology studies as well as evolutionary comparative analyses on cerebral cortex development converge on the idea that the AP/BP classification of cortical progenitors, although very useful and generally pertinent, does not reflect the actual heterogeneity of these cells. For example, the observation of a unique proliferative compartment in the monkey embryonic cortex, represented by a very large outer SVZ (OSVZ), first suggested a potential mechanism for primate cortical expansion (Smart et al. 2002). These OSVZ progenitors are heterogeneous and include both AP/radial glia-like stem cells (*Pax6*<sup>+</sup>, self-renewing, stem cell-like) and BP (*Tbr2*<sup>+</sup>, transit amplifying) cell types (Fietz et al. 2010; Hansen et al. 2010). Such OSVZ-like progenitors are in fact found in all mammals, both gyrencephalic (Reillo et al. 2011) and lissencephalic including the mouse (Kelava et al. 2011; Shitamukai et al. 2011; Wang et al. 2011). Ongoing studies on ferrets (nonprimate, gyrencephalic) have begun to uncover subtle differences when compared with primates (Reillo and Borrell 2012). Among these differences, some are quantitative (radial glia abundance), some are qualitative (genetic programs controlling cell cycle kinetics and dynamics of self-renewal), and some are time-dependent. Here, we bring some evidences that the heterogeneity and diversity in mouse cortical progenitor types may lie in a finely tuned differential transcriptional control, with cortical progenitor subtypes having specific regulatory signatures and activities. With this regard, it is also striking that the evolutionary conservation of active mouse enhancer sequences is relatively poor with other mammals. *Enh14*(*cdh4*), which is active in both AP- and BP-like progenitors, is conserved between mouse and human, but not dog. *Enh15*(*Sall3*), which is active only in the SVZ, is conserved only between mouse and chick (but note that birds do not have a laminated pallium). These features suggest that fine regulation of gene expression and progenitor behavior differ between closely related mammalian species, and such differences may underlie subtle evolutionary variations in cortical architecture across mammals. Only *enh18*(*Pou2f1*), active in proliferating VZ progenitors, is conserved across all vertebrates. This suggests that this regulatory input may be shared by self-renewing, stem cell-like radial glial cells of the (dorsal) telencephalon in all vertebrate species. Such a hypothesis will have to be tested in birds and fishes in a near future.

### Enhancer Activity versus Expression Pattern of the Flanking Gene

The activity of the 3 enhancers that we have described with some details does not recapitulate totally the expression



**Figure 8.** mRNA expression versus enhancer activity. (A) ISH (purple signal) for *Cdh4* (A1), *Sall3* (A2), and *Pou2f1* transcripts at the indicated stages (E11.5–E13.5) in the embryonic mouse cortex. For each image, layers are indicated. CP: cortical plate; MZ: marginal zone; IZ: intermediate zone; PP: preplate; SP: subplate; SVZ: subventricular zone; VZ: ventricular zone. (B) Corresponding putative enhancer activity, as summarized from results in Figures 5 and 6. Green dots represent cells in which the enhancer is active. Representative numbers and proportions of cells are drawn in each layer for each enhancer and also includes indications on the proliferative status of the cells (black mitotic figure = pH3 immunoreactivity) and the category of progenitor (red dot = *Tbr2* immunoreactivity; see inset).

pattern of the closest flanking gene (Fig. 8). We cannot rule out that other genes (i.e., not the closest flanking gene but a more distal one) may be regulated by these enhancers. Yet, the comparison between enhancer activity and flanking gene expression pattern is quite satisfactory: The activity of *enh14* (*Cdh4*) in cells in the differentiation/migration to the CP process is correlated with low levels of *Cdh4* mRNA in the cortical germinal zones; the activity of *enh15*(*Sall3*), excluded from the VZ where *Sall3* mRNA is expressed, suggests that other enhancer(s) are responsible for *Sall3* expression in APs. Conversely, *enh18*(*Pou2f1*) activity accounts for *Pou2f1* mRNA expression in the germinal zones, but not in the lower part of the IZ/CP. In all cases, it appears that the regulation of mRNA expression is modular, with different enhancers involved at different steps of corticogenesis, underscoring the crucial importance of timing and dynamics of gene regulation in this developmental process.

## Conclusion

Through the window of transcriptional *cis*-regulation, we have found that cortical progenitor types in the mouse embryonic cortex are heterogeneous. They nevertheless share a common *cis*-regulatory logic, involving Pou TFBS together with Sox TFBS, to control their gene expression. Importantly, our approach also provides novel enhancer sequences that can be used as drivers for future analyses of corticogenesis and cortical progenitor biology.

## Supplementary Material

Supplementary material can be found at: <http://www.cercor.oxfordjournals.org/>.

## Funding

This work was supported by the FP7 European project CISSTEM.

## Notes

The authors thank to Yan Jaszczyszyn and Aurélie Heuzé for their help with Gateway cloning and for modifying the p336bglobGFP34H vector, a gift of François Spitz. Many thanks to Yohann Mérot who established *ex vivo* and *in utero* electroporations in the group and gave valuable advices on these methods. *Conflict of Interest:* None declared.

## References

- Abellan A, Menuet A, Dehay C, Medina L, Rétaux S. 2010. Differential expression of LIM-homeodomain factors in Cajal-Retzius cells of primates, rodents, and birds. *Cerebr Cortex*. 20(8):1788–1798.
- Basak O, Taylor V. 2009. Stem cells of the adult mammalian brain and their niche. *Cell Mol Life Sci*. 66(6):1057–1072.
- Berger J, Eckert S, Scardigli R, Guillemot F, Gruss P, Stoykova A. 2004. E1-Ngn2/Cre is a new line for regional activation of Cre recombinase in the developing CNS. *Genesis*. 40(4):195–199.
- Boulder committee. 1970. Embryonic vertebrate central nervous system: revised terminology. *Anat Rec*. 166(2):257–261.

- Brudno M, Do C, Cooper G, Kim MF, Davydov E, Green ED, Sidow A, Batzoglou S. 2003. LAGAN and Multi-LAGAN: efficient tools for large-scale multiple alignment of genomic DNA. *Genome Res.* 13(4):721–731.
- Bylund M, Andersson E, Novitsch BG, Muhr J. 2003. Vertebrate neurogenesis is counteracted by Sox1-3 activity. *Nat Neurosci.* 6:1162–1168.
- Bystron I, Blakemore C, Rakic P. 2008. Development of the human cerebral cortex: Boulder Committee revisited. *Nat Rev Neurosci.* 9(2):110–122.
- Castro DS, Martynoga B, Parras C, Ramesh V, Pacary E, Johnston C, Drechsel D, Lebel-Potter M, Galinanes Garcia L, Hunt C et al. 2011. A novel function of the proneural factor Ascl1 in progenitor proliferation identified by genome-wide characterization of its targets. *Genes Develop.* 25:930–945.
- Castro DS, Skowronska-Krawczyk D, Armant O, Donaldson IJ, Parras C, Hunt C, Critchley JA, Nguyen L, Gossler A, Göttgens B et al. 2006. Proneural bHLH and Brn proteins coregulate a neurogenic program through cooperative binding to a conserved DNA motif. *Development Cell.* 11:831–844.
- Conti L, Pollard SM, Gorba T, Reitano E, Toselli M, Biella G, Sun Y, Sanzone S, Ying GL, Cattaneo E et al. 2005. Niche-independent symmetrical self-renewal of a mammalian tissue stem cell. *PLoS Biol.* 3(9):e282.
- Cook AL, Sturm RA. 2008. POU domain transcription factors: BRN2 as a regulatory of melanocytic growth and tumorigenesis. *Pigment Cell Melanoma Res.* 21(6):611–626.
- Ellis P, Fagan BM, Magness ST, Hutton S, Taranova O, Hayashi S, McMahon A, Rao M, Pevny L. 2004. SOX2, a persistent marker for multipotential neural stem cells derived from embryonic stem cells, the embryo or the adult. *Development Neurosci.* 26(2–4):148–165.
- ENCODE Project Consortium, Dunham I, Kundaje A, Aldred SF, Collins PJ, Davis CA, Doyle F, Epstein CB, Fietze S, Harrow J, Kaul R et al. 2012. An integrated encyclopedia of DNA elements in the human genome. *Nature.* 489(7414):57–74.
- Englund C, Fink A, Lau C, Pham D, Daza RA, Bulfone A, Kowalczyk T, Hevner RF. 2005. Pax6, Tbr2, and Tbr1 are expressed sequentially by radial glia, intermediate progenitors and postmitotic neurons in developing neocortex. *J Neurosci.* 25(1):247–251.
- Fietz SA, Kelava I, Vogt J, Wilsch-Bräuninger M, Stenzel D, Fish JL, Corbeil D, Riehn A, Distler W, Nitsch R et al. 2010. OSVZ progenitors of human and ferret neocortex are epithelial-like and expand by integrin signaling. *Nat Neurosci.* 13(6):690–699.
- Gal JS, Morozov YM, Ayoub AE, Chatterjee M, Rakic P, Haydar TF. 2006. Molecular and morphological heterogeneity of neural precursors in the mouse neocortical proliferative zones. *J Neurosci.* 26:1045–1056.
- Graham V, Khudyakov J, Ellis P, Pevny L. 2003. SOX2 functions to maintain neural progenitor identity. *Neuron.* 39:749–765.
- Hansen DV, Lui JH, Parker PR, Kriegstein AR. 2010. Neurogenic radial glia in the outer subventricular zone of human neocortex. *Nature.* 464(7288):554–561.
- Harley VR, Lovell-Badge R, Goodfellow PN. 1994. Definition of a consensus DNA binding site for SRY. *Nucleic Acids Res.* 22(8):1500–1501.
- Hartfuss E, Galli R, Heins N, Gotz M. 2001. Characterization of CNS precursor subtypes and radial glia. *Development Biol.* 229:15–30.
- Haubensak W, Attardo A, Denk W, Huttner WB. 2004. Neurons arise in the basal neuroepithelium of the early mammalian telencephalon: a major site of neurogenesis. *Proc Natl Acad Sci USA.* 101:3196–3201.
- Heins N, Malatesta P, Ceconi F, Nakafuku M, Tucker KL, Hack MA, Chapput P, Barde YA, Götz M. 2002. Glial cells generate neurons: the role of the transcription factor Pax6. *Nat Neurosci.* 5(4):308–315.
- Heng JI, Nguyen L, Castro DS, Zimmer C, Wildner H, Armant O, Skowronska-Krawczyk D, Bedogni F, Matter JM, Hevner R et al. 2008. Neurogenin2 controls cortical neuron migration through regulation of Rnd2. *Nature.* 455(7209):114–118.
- Jin Z, Liu L, Bian W, Chen Y, Xu G, Cheng L, Jing N. 2009. Different transcription factors regulate nestin gene expression during P19 cell neural differentiation and central nervous system development. *J Biol Chem.* 284(12):8160–8173.
- John S, Sabo PJ, Thurman RE, Sung MH, Biddie SC, Johnson TA, Hager GL, Stamatoyannopoulos JA. 2011. Chromatin accessibility pre-determines glucocorticoid receptor binding patterns. *Nat Genet.* 43(3):264–268.
- Josephson R, Müller T, Pickel J, Okabe S, Reynolds K, Turner PA, Zimmer A, McKay RD. 1998. POU transcription factors control expression of CNS stem cell-specific genes. *Development.* 125(16):3087–3100.
- Kamachi Y, Uchikawa M, Kondoh H. 2000. Pairing Sox off: with partners in the regulation of embryonic development. *Trends Genet.* 16(4):182–187.
- Kelava I, Reillo I, Murayama AY, Kalinka AT, Stenzel D, Tomancak P, Matsuzaki F, Lebrand C, Sasaki E, Schwamborn JC et al. 2011. Abundant occurrence of basal radial glia in the subventricular zone of embryonic neocortex of a lissencephalic primate, the common marmoset *Callithrix jacchus*. *Cereb Cortex.* 22(2):469–481.
- Kelava I, Reillo I, Murayama AY, Kalinka AT, Stenzel D, Tomancak P, Matsuzaki F, Lebrand C, Sasaki E, Schwamborn JC et al. 2012. Abundant occurrence of basal radial glia in the subventricular zone of embryonic neocortex of a lissencephalic primate, the common marmoset *Callithrix jacchus*. *Cereb Cortex.* 22(2):469–481.
- Kondoh H, Kamachi Y. 2010. SOX-partner code for cell specification: Regulation target selection and underlying molecular mechanisms. *Int J Biochem Cell Biol.* 42(3):391–399.
- Kowalczyk T, Pontious A, Englund C, Daza RA, Bedogni F, Hodge R, Attardo A, Bell C, Huttner WB, Hevner RF. 2009. Intermediate neuronal progenitors (basal progenitors) produce pyramidal-projection neurons for all layers of cerebral cortex. *Cerebr Cortex.* 19(10):2439–2450.
- Ma Y, Certel K, Gao Y, Niemitz E, Mosher J, Mukherjee A, Mutsuddi M, Huseinovic N, Crews ST, Johnson WA et al. 2000. Functional interactions between *Drosophila* bHLH/PAS, Sox, and POU transcription factors regulate CNS midline expression of the slit gene. *J Neurosci.* 20(12):4596–4605.
- Malatesta P, Hartfuss E, Götz M. 2000. Isolation of radial glial cells by fluorescent-activated cell sorting reveals a neuronal lineage. *Development.* 127:5253–5263.
- Mayor C, Brudno M, Schwartz JR, Poliakov A, Rubin EM, Frazer KA, Pachter LS, Dubchak I. 2000. VISTA: visualizing global DNA sequence alignments of arbitrary length. *Bioinformatics.* 16:1046.
- Mérot Y, Rétaux S, Heng JI. 2009. Molecular mechanisms of projection neuron production and maturation in the developing cerebral cortex. *Semin Cell Develop Biol.* 20(6):726–734.
- Miyagi S, Nishimoto M, Saito M, Sawamoto K, Okano H, Muramatsu M, Oguro H, Iwana A, Okuda A. 2006. The Sox2 regulatory region 2 functions as a neural stem cell-specific enhancer in the telencephalon. *J Biol Chem.* 281(19):13374–13381.
- Miyata T, Kawaguchi A, Okano H, Ogawa M. 2001. Asymmetric inheritance of radial glial fibers by cortical neurons. *Neuron.* 31:727–741.
- Mizuseki K, Kishi M, Matsui M, Nakanishi S, Sasai Y. 1998. Xenopus Zic-related-1 and Sox-2, two factors induced by chordin, have distinct activities in the initiation of neural induction. *Development.* 122:2449–2464.
- Nieto M, Monuki ES, Tang H, Imitola J, Haubst N, Khoury SJ, Cunningham J, Gotz M, Walsh CA. 2004. Expression of Cux-1 and Cux-2 in the subventricular zone and upper layers II–IV of the cerebral cortex. *J Comp Neurol.* 479:168–180.
- Nishimoto M, Fukushima A, Okuda A, Muramatsu M. 1999. The gene for the embryonic stem cell coactivator UTF1 carries a regulatory element which selectively interacts with a complex composed of Oct-3/4 and Sox-2. *Mol Cell Biol.* 19(8):5453–5465.
- Noctor SC, Flint AC, Weissman TA, Dammerman RS, Kriegstein AR. 2001. Neurons derived from radial glial cells establish radial units in neocortex. *Nature.* 409:714–720.
- Noctor SC, Martinez-Cerdeno V, Ivic L, Kriegstein AR. 2004. Cortical neurons arise in symmetric and asymmetric division zones and migrate through specific phases. *Neurosci.* 7:136–144.
- Noctor SC, Martinez-Cerdeño V, Kriegstein AR. 2008. Distinct behaviors of neural stem and progenitor cells underlie cortical neurogenesis. *J Comp Neurol.* 508(1):28–44.
- Pennacchio LA, Ahituv N, Moses AM, Prabhakar S, Nobrega MA, Shoukry M, Minovitsky S, Dubchak I, Holt A, Lewis KD et al. 2006. In vivo enhancer analysis of human conserved non-coding sequences. *Nature.* 444:499–502.

- Petryniak B, Staudt LM, Postema CE, McCormack WT, Thompson CB. 1990. Characterization of chicken octamer-binding proteins demonstrates that POU domain-containing homeobox transcription factors have been highly conserved during vertebrate evolution. *Proc Natl Acad Sci.* 87:1099–1103.
- Reillo I, Borrell V. 2012. Germinal zones in the developing cerebral cortex of ferret: ontogeny, cell cycle kinetics, and diversity of progenitors. *Cereb Cortex.* 22(9):2039–2054.
- Reillo I, de Juan Romero C, García-Cabezas MÁ, Borrell V. 2011. A role for intermediate radial glia in the tangential expansion of the mammalian cerebral cortex. *Cereb Cortex.* 21(7):1674–1694.
- Reményi A, Lins K, Nissen LJ, Reinbold R, Schöler HR, Wilmanns M. 2003. Crystal structure of a POU/HMG/DNA ternary complex suggests differential assembly of Oct4 and Sox2 on two enhancers. *Genes Develop.* 17(16):2048–2059.
- Ryan AK, Rosenfeld MG. 2011. POU domain family values: flexibility, partnerships, and developmental codes. *Genes Develop.* 11:1207–1225.
- Sabo PJ, Hawrylycz M, Wallace JC, Humbert R, Yu M, Shafer A, Kawamoto J, Hall R, Mack J, Dorschner MO et al. 2004. Discovery of functional noncoding elements by digital analysis of chromatin structure. *Proc Natl Acad Sci.* 101(48):16837–16842.
- Sabo PJ, Kuehn MS, Thurman R, Johnson BE, Johnson EM, Cao H, Yu M, Rosenzweig E, Goldy J, Haydock A et al. 2006. Genome-scale mapping of DNase I sensitivity in vivo using tiling DNA microarrays. *Nat Methods.* 3(7):511–518.
- Sansom SN, Griffiths DS, Faedo A, Kleinjan DJ, Ruan Y, Smith J, van Heyningen V, Rubenstein JL, Livesey FJ. 2009. The level of the transcription factor Pax6 is essential for controlling the balance between neural stem cell self-renewal and neurogenesis. *PLoS Genet.* 5(6):e1000511.
- Scardigli R, Schuurmans C, Gradwohl G, Guillemot F. 2001. Crossregulation between Neurogenin2 and pathways specifying neuronal identity in the spinal cord. *Neuron.* 31(2):203–217.
- Schöler HR, Hatzopoulos AK, Balling R, Suzuki N, Gruss P. 1989. A family of octamer-specific proteins during mouse embryogenesis: evidence for germline-specific expression of an Oct factor. *EMBO J.* 8(9):2543–2550.
- Shimogori T, Ogawa M. 2008. Gene application with in utero electroporation in mouse embryonic brain. *Develop Growth Differ.* 50(6):499–506.
- Shitamukai A, Konno D, Matsuzaki F. 2011. Oblique radial glial divisions in the developing mouse neocortex induce self-renewing progenitors outside the germinal zone that resemble primate outer subventricular zone progenitors. *J Neurosci.* 31(10):3683–3695.
- Sidman RL, Rakic P. 1973. Neuronal migration, with special reference to developing human brain: a review. *Brain Res.* 62(1):1–35.
- Smart IH, Dehay C, Giroud P, Berland M, Kennedy H. 2002. Unique morphological features of the proliferative zones and postmitotic compartments of the neural epithelium giving rise to striate and extrastriate cortex in the monkey. *Cerebr Cortex.* 12(1):37–53.
- Smit DJ, Smith AG, Parsons PG, Muscat GE, Sturm RA. 2000. Domains of Brn-2 that mediate homodimerization and interaction with general and melanocytic transcription factors. *Eur J Biochem.* 267(21):6413–6422.
- Stancik EK, Navarro-Quiroga I, Sellke R, Haydar TF. 2010. Heterogeneity in ventricular zone neural precursors contributes to neuronal fate diversity in the postnatal neocortex. *J Neurosci.* 30:7028–7036.
- Sunabori T, Tokunaga A, Nagai T, Sawamoto K, Okabe M, Miyawaki A, Matsuzaki Y, Miyata T, Okano H. 2008. Cell-cycle-specific nestin expression coordinates with morphological changes in embryonic cortical neural progenitors. *J Cell Sci.* 121(8):1204–1212.
- Tanaka S, Kamachi Y, Tanouchi A, Hamada H, Jing N, Kondoh H. 2004. Interplay of SOX and POU factors in regulation of the nestin in neural primordial cells. *Mol Cell Biol.* 24(20):8834–8846.
- Tarabykin V, Stoykova A, Usman N, Gruss P. 2001. Cortical upper layer neurons derive from the subventricular zone as indicated by Svet1 gene expression. *Development.* 128:1983–1993.
- Taverna R, Huttner WB. 2010. Neural progenitor nuclei IN motin. *Neuron.* 67(6):906–914.
- Tomioka M, Nishimoto M, Miyagi S, Katayanagi T, Fukui N, Niwa H, Muramatsu M, Okuda A. 2002. Identification of Sox-2 regulatory region which is under control of Oct-3/4-Sox-2 complex. *Nucleic Acids Res.* 30(14):3202–3212.
- Uchikawa M, Ishida Y, Takemoto T, Kamachi Y, Kondoh H. 2003. Functional analysis of chicken Sox2 enhancers highlights an array of diverse regulatory elements that are conserved in mammals. *Development.* 130(4):509–519.
- Van Helden J. 2003. Regulatory sequence analysis tools. *Nucleic Acids Res.* 31:3593–3596.
- Visel A, Blow MJ, Li Z, Zhang T, Akiyama JA, Holt A, Plajzer-Frick I, Shoukry M, Wright C, Chen F et al. 2009. ChIP-seq accurately predicts tissue-specific activity of enhancers. *Nature.* 457(7231):854–858.
- Visel A, Minovitsky S, Dubchak I, Pennacchio LA. 2007. VISTA enhancer browser—a database of tissue-specific human enhancers. *Nucleic Acids Res.* 35:D88–D92.
- Visel A, Taher L, Girgis H, May D, Golonzhka O, Hoch RV, McKinsey GL, Pattabiraman K, Silberberg SN, Blow MJ et al. 2013. A high-resolution enhancer atlas of the developing telencephalon. *Cell.* 152(4):895–908.
- Walker AS, Goings GE, Kim Y, Miller RJ, Chenn A, Szele FG. 2010. Nestin reporter transgene labels multiple central nervous system precursor cells. *Neural Plast.* 2010:894374.
- Wang W, Tsai JW, LaMonica B, Kriegstein AR. 2011. A new subtype of progenitor cell in the mouse embryonic neocortex. *Nat Neurosci.* 14(5):555–561.
- Wasserman WW, Sandelin A. 2004. Applied bioinformatics for the identification of regulatory elements. *Nat Rev Genet.* 5(4):276–287.
- Zappone MS, Galli R, Catena R, Meani N, De Biasi S, Mattei E, Tiveron C, Vescovi AL, Lovell-Badge R, Ottolenghi S et al. 2000. Sox2 regulatory sequences direct expression of a  $\beta$ -geo transgene to telencephalic neural stem cells and precursors of the mouse embryo, revealing regionalization of gene expression in CNS stem cells. *Development.* 127:2367–2382.
- Zhang Y, Liu T, Meyer CA, Eeckhoute J, Johnson DS, Bernstein BE, Nussbaum C, Myers RM, Brown M, Li W et al. 2008. Model-based analysis of ChIP-Seq (MACS). *Genome Biol.* 9(9):R137.
- Zimmer C, Tiveron MC, Bodmer R, Cremer H. 2004. Dynamics of *Cux2* expression suggests that an early pool of SVZ precursors is fated to become upper cortical layer neurons. *Cereb Cortex.* 14:1408–1420.