



HAL
open science

Secondary metabolites of *Bagassa guianensis* Aubl. wood: A study of the chemotaxonomy of the Moraceae family

Mariana Royer, Gaëtan Herbette, Véronique Eparvier, Jacques Beauchêne,
Bernard Thibaut, Didier Stien

► **To cite this version:**

Mariana Royer, Gaëtan Herbette, Véronique Eparvier, Jacques Beauchêne, Bernard Thibaut, et al..
Secondary metabolites of *Bagassa guianensis* Aubl. wood: A study of the chemotaxonomy of the
Moraceae family. *Phytochemistry*, 2010, 71, pp.1708-1713. 10.1016/j.phytochem.2010.06.020 . hal-
00857200

HAL Id: hal-00857200

<https://hal.science/hal-00857200>

Submitted on 3 Sep 2013

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22

Secondary metabolites of *Bagassa guianensis* Aubl. wood, a study of the chemotaxonomy of the Moraceae family.

Mariana Royer¹; Gaëtan Herbette²; Véronique Eparvier¹; Jacques Beauchêne³; Bernard Thibaut¹; Didier Stien^{1,*}

Affiliations

¹ CNRS, UMR Ecofog, Université des Antilles et de la Guyane, Cayenne, France

² Spectropole, FR 1739 – Université d’Aix-Marseille, Faculté de Saint-Jérôme, service 511, Avenue Escadrille Normandie Niémen, 13397 Marseille cedex 20, France

³ Cirad, UMR Ecofog, BP 709, F-97387 Kourou, France

Corresponding authors

Dr. Didier Stien, CNRS, UMR Ecofog, Institut d’Enseignement Supérieur de la Guyane, BP 792, 97337 Cayenne cedex, France. E-mail: didier.stien@guyane.cnrs.fr Phone: +594 594 29 75 17 Fax: +594 594 28 47 86.

23

24 **Abstract**

25

26 In effort to explain wood durability of Moraceae plants family, a phytochemical study was
27 undertaken on *Bagassa guianensis*. The phytochemical investigation of the ethyl acetate
28 extract obtained from the heartwood led to the isolation of 18 secondary metabolites,
29 including 6 moracins [the new 6-*O*-methyl-moracin M (**3**), 6- *O*-methyl-moracin N (**4**) and
30 moracin Z (**5**); the known moracin M (**1**), moracin N (**2**) and moracin P (**6**)], 8 phenolic
31 derivatives [the new (-)-epialboctalol (**12**), arachidin 4 (**10**) and the known alboctalol (**11**),
32 *trans*-resveratrol (**7**), arachidin 2 (**9**), *trans*-oxyresveratrol (**8**) and artogomezianol (**13**)], the 3
33 known flavonoids steppogenin (**14**), katuranin (**15**), dihydromorin (**16**), the β -sitosterol (**17**)
34 and the resorcinol (**18**). Comparison with literature data indicates that stilbenoids are
35 presumably responsible for the natural durability of the wood. In addition, chemical
36 composition points out that *B. guianensis* is closely related to *Morus* sp. in the phylogeny and
37 should be placed within the Moreae s. s. tribe in the Moraceae family.

38

39 **Keywords:** *Bagassa guianensis*, Moraceae, secondary metabolites, stilbenes, moracins,
40 flavonoids, natural durability

41

42

43

44 **1. Introduction**

45 Wood as a material is used extensively in construction and other applications where it can be
46 degraded by many different organisms, mainly fungi and insects. However, some trees have
47 specialized considerably long-lasting heartwoods. It has been demonstrated in the past that
48 wood natural durability can be ascribed to the presence of extractives (Smith et al., 1989;
49 Wang et al., 2005; Hsu et al., 2007), although structural components of the cell wall may also
50 contribute to its resistance to biodegradation (Silva et al., 2007). Heartwood natural durability
51 can also result from synergetic or additive effects of compounds with various modes of action
52 (toxic, hydrophobic, free radical scavengers and so on) (Suttie and Orsler, 1996; Okitani et al.,
53 1999; Schultz and Nicholas, 2000; Schultz et al. 2007; Binbuga et al., 2008). Future processes
54 to preserve wood constructions may involve returning to mankind's historical use of naturally
55 durable heartwood as well as discovering eco-friendly wood protection agents inspired from
56 long-lasting woods (Schultz et al., 2007).

57

58 *Bagassa guianensis* Aubl. (Moraceae) commercially known as tatajuba is a large rather
59 infrequent unbuttressed canopy tree naturally occurring in French Guiana. *Bagassa guianensis*
60 is a member of Moraceae family, which is divided in 5 unequal tribes when comparing the
61 number of species in these tribes (Mabberley, 2002). *Bagassa guianensis* (the only member of
62 its genus) was originally classified in the Artocarpeae tribe, but Weiblen genoma-based
63 classifications have suggested recently that this species would better be included in Moreae
64 tribe (Datweyler and Weiblen, 2004; Zerega et al, 2005).

65

66 Species in the Moraceae family have important economic and medicinal value. They are
67 widely acknowledged as a rich source of bioactive secondary metabolites such as flavonoids,

68 stilbenes, triterpenoids and xanthenes (Lee et al., 2009; Ngadjui et al., 2005; Han et al., 2006;
69 Jayasinghe et al., 2008). Also, some of them like *Maclura pomifera* and *B. guianensis* are
70 capable of specializing very long-lasting woods (Scheffer and Morrell, 1998; Schultz et al.,
71 1995), although in the latter case, the substances responsible for this high durability were
72 unknown. We therefore embarked upon identifying secondary metabolites of tatajuba wood
73 that may responsible for its natural durability. In addition, our secondary goal here was to
74 confirm (or refute) botanical classification of the *Bagassa* genus by chemotaxonomy.

75

76 2. Results and Discussion

77 The dried heartwood of *Bagassa guianensis* was extracted with ethyl acetate. This extract was
78 fractionated by silicagel column chromatography to give 9 fractions. Subsequent preparative
79 HPLC purifications of these fractions allowed us to isolate compounds **1-18** (figure 1).

80

81 **Figure 1** Compounds **1-18** isolated from *Bagassa guianensis* (Moraceae). (a) New
82 compounds; (b) New names.

83

84 Compounds **1** to **6** shared several common spectral characteristics. The ^1H and ^{13}C NMR
85 spectral data (Table 1) indicate the presence of two independent aromatic systems with a 3,5-
86 dihydroxyphenyl and a substituted benzofuran. For example, **3** exhibited the 3,5-
87 dihydroxyphenyl with characteristic ^1H spectrum composed of one doublet at δ 6.78 for H-
88 2'/H-6' and a triplet at δ 6.25 for H-4'. These protons are coupled to each other with a 4J
89 coupling of 2.1 Hz. In addition, ^{13}C spectrum indicates the presence of two equivalent aryl
90 hydroxyl groups at δ 159.7. The 3,5-dihydroxyphenyl moiety was linked to C-2 by the
91 observation of a long range ^1H - ^{13}C correlation between H-2'/H-6' and C-2 at δ 156.5. The
92 second aromatic system appeared characteristic of a 6-monosubstituted benzofuran with

93 signals of protons H-4, H-5 and H-7 being a broad doublet at δ 7.43 ($J = 8.5$ Hz), a doublet of
94 doublet at δ 6.85 ($J = 8.5$ and 2.0 Hz) and a doublet at δ 7.09 ($J = 2.0$ Hz), respectively. On
95 the furan ring H-3 gives a doublet at δ 6.95 ($J = 0.6$ Hz) due to a long range 5J coupling with
96 H-7 (confirmed by the presence of crosspeak between H-3 and H-7 on COSY NMR
97 spectrum). When compared to moracin M (**1**), it became obvious from signal at δ 3.85 (3H, s)
98 and the presence of crosspeak at δ 56.2 in the ^1H - ^{13}C HSQC spectra that compound **3** was a
99 moracin M methyl ether. The ^1H - ^{13}C long-range HMBC spectra gave a crosspeak with C-6 at
100 δ 159.6 unambiguously placing the methoxy group on C-6. HREIMS of **3** allowed us to
101 ascertain molecular formula $\text{C}_{15}\text{H}_{12}\text{O}_4$ further confirming that we had isolated the new 6-*O*-
102 methyl-moracin M (**3**).

103

104 **Table 1** ^1H and ^{13}C NMR spectroscopic data for moracins **3-5** in CD_3OD

105

106 Compound **4** was isolated as yellowish amorphous powder. The HREIMS indicated a
107 molecular formula $\text{C}_{20}\text{H}_{20}\text{O}_4$ deduced from the ion peak at m/z 325.1437 [$\text{M} + \text{H}$] $^+$ (calcd
108 325.1434). The ^1H and ^{13}C NMR spectral data of **4** were closely related to those of moracin N
109 (**2**) (Lee et al., 2001) except for the replacement of hydroxyl group by a methoxy group as
110 described for the above compound **3**. Indeed, the ^1H NMR data of **4** (Table 1) demonstrated
111 the presence a methoxy group on C-6 in the benzofuran ring, with a signal at δ 3.88 (3H, s), a
112 crosspeak at δ 56.2 in the ^1H - ^{13}C HSQC experiment and a crosspeak with C-6 at δ 157.4 in
113 the ^1H - ^{13}C long-range HMBC spectra. This novel molecule was named 6-*O*-methyl-moracin
114 N.

115

116 Compound **5** was isolated as an amorphous brown powder. The molecular formula $C_{20}H_{22}O_5$
117 was deduced from the HREIMS at m/z 343.1542 $[M + H]^+$ (calcd 343.1540). The 1H - and ^{13}C -
118 NMR spectral data of **5** were closely related to those of 6-*O*-methyl-moracin N (**4**) (Table 1).
119 The main difference was observed in the prenyl moiety at C-5. The double bond is absent in **5**
120 and it was unambiguously established that side chain at C-5 is hydrated and is therefore a 3-
121 hydroxy-3-methylbutyl group, with the upfield shifts of methylene group H-1'' from δ 3.34 to
122 δ 2.73 and the apparition of a methylene H-2'' at δ 1.74 in place of the vinyl proton at δ 5.52;
123 in addition, the two methyl groups H-4'' and H-5'' became equivalent at δ 1.27 (Table 1).
124 The 1H - ^{13}C long-range HMBC spectra exhibited a crosspeak between the methylene group H-
125 1'' and H-2'' with C-5 at δ 128.6 proving the linkage C-1''/C-5 between the 3-hydroxy-3-
126 methylbutyl moiety and the benzofuran ring. This molecule is a hydrate of 6-*O*-methyl-
127 moracin N and was named moracin Z.

128
129 Spectral data along with HREIMS of **1**, **2** and **6** allowed us to determine and ascertain by
130 comparison with literature data that we had also isolated moracin M (**1**) (Basnet et al. 1993,
131 Zhou et al., 1999), moracin N (**2**) (Lee et al. 2001) and moracin P (**6**) (Dat et al., 2009).

132
133 Stilbenoids *trans*-resveratrol (**7**) (Lee et al. 2001; Su et al., 2002), *trans*-oxyresveratrol (**8**)
134 (Likhitwitayawuid and Sritularak, 2001; Lee et al., 2001; Su et al., 2002; Li et al., 2007),
135 arachidin 2 (**9**) (Orsini et al., 2004) and artogomezianol (**13**) (Likhitwitayawuid and
136 Sritularak, 2001) were identified by comparison of the respective spectral and chemical data
137 with those described in the literature (Figure 1).

138
139 Compound **10** was a colorless syrup with molecular formula $C_{19}H_{22}O_4$ as deduced from the
140 HREIMS at m/z 315.1592 $[M + H]^+$ (calcd 315.1591). The 1H spectral data of **10** were closely

141 related to those of arachidin 2 (**9**) (Table 2) and suggested a stilbenoid compound with a *para*-
142 disubstituted aromatic ring A, a *trans* double bond between the aromatic rings, and a
143 1',3',4',5'-tetrasubstituted aromatic ring B. Ring A is symmetrical, with 2 doublets at δ 7.32
144 ($J = 8.7$ Hz, H-2/H-6) and δ 6.75 ($J = 8.7$ Hz, H-3/H-5). The *trans* configuration of the double
145 bond can be ascertained by the very large coupling constant between the two protons at δ 6.90
146 ($J = 16.5$ Hz, H- α) and δ 6.74 ($J = 16.5$ Hz, H- β), and the B ring is symmetrical as well and
147 was characterized by a singlet at δ 6.46 (H-2'/H-6'). In the same way as we identified a
148 hydrated side chain in the moracins series, the main difference here between **9** and **10** is in the
149 side chain in position 4', the double bond of which is also hydrated. This has been established
150 by the observation of methylene group H-1'' at δ 2.66 instead of δ 3.28 and the apparition of
151 a second methylene H-2'' at δ 1.68. In addition, the two methyl groups H-4'' and H-5''
152 became equivalent at δ 1.25. The chromatography collected quantities was too low to observe
153 heteronuclear ^1H - ^{13}C HSQC / HMBC correlations and direct ^{13}C chemicals shifts by
154 ^{13}C /DEPTQ sequence. However, the above-described data in comparison with those of
155 arachidin 2 are sufficient to ascertain identification of compound **10** as *trans*-4'-(3-hydroxy-3-
156 methylbutyl)-oxyresveratrol. We named this new compound arachidin 4.

157

158 **Table 2** ^1H and ^{13}C NMR spectroscopic data for stilbenes **9** and **10** in CD_3OD

159

160 Compounds **11** and **12** both isolated as brownish syrups presented the ion peak at m/z
161 489.1540 $[\text{M} + \text{H}]^+$ in HREIMS indicating that they are isomers with molecular formulas
162 $\text{C}_{28}\text{H}_{24}\text{O}_8$ (calcd 489.1544). The ^1H -NMR allowed us to identify a 3,5-dihydroxyphenyl group
163 and two distinct 2,4-dihydroxyphenyl groups in both compounds. By comparison of the
164 respective spectral and chemical data with those described in the literature, compound **11** was

165 identified as alboctalol (Bates et al., 1997). Compound **12** has an $[\alpha]_D^{20}$ value of -7.4° (c
166 0.004, CH₃OH). It was clear that **12** was a diastereoisomer of **11** with equivalent H-18/H-22
167 protons at δ 6.01 (Table 3). In **11**, H-18/H-22 pair gives a doublet at a strong upfield shift of δ
168 5.77 typical of the π -stacking effect of the neighboring 2,4-dihydroxyphenyl groups. In
169 addition, on this aliphatic ring, the main differences with **11** are on methylene H-5 and
170 methines H-6, H-7 and H-8. H-5_{ax} at δ 3.19 exhibited a broad triplet with large couplings ($J =$
171 13.7 Hz) with the gem H-5_{eq} and the vicinal H-6 suggesting that the 6-aryl group should be
172 equatorial and proton H-6 axial. This observation was corroborated by the multiplicity of H-
173 5_{eq} signal at δ 2.72. This signal is a doublet of doublet with a large coupling constant $J = 15.6$
174 Hz with H-5_{ax} and a small coupling constant $J = 3.0$ Hz with H-6_{ax}. Signal of H-6_{ax} at δ 3.51
175 is a broad triplet of doublet with two large coupling constants $J = 11.6$ Hz with H-5_{ax} and H-
176 7 and a small coupling constant $J = 2.1$ Hz with H-5_{eq}. This pattern indicates that the 7-aryl
177 group is equatorial and H-7 axial. H-7_{ax} at δ 3.41 exhibited one doublet of doublet with one
178 large coupling constant ($J = 11.3$ Hz) with H-6_{ax} and a second rather large coupling constant
179 ($J = 8.2$ Hz) with H-8 indicating that the 8-aryl group might be equatorial and proton H-8
180 axial. These assumptions were confirmed by NOESY experiment with cross peaks observed
181 between H-5_{eq} and H-6_{ax}, H-6_{ax} and H-8_{ax}, H-6_{ax} and H-18, H-8_{ax} and H-22 and between H-5_{ax}
182 and H-16, H-7_{ax} and H-16, H-7_{ax} and H-28 (Figure 2). All data permitted to confirm that we
183 had isolated a new epimer of alboctalol (**11**) therefore named (-)-epialboctalol (**12**).

184

185 **Table 3** ¹H and ¹³C NMR spectroscopic data for distilbenes **11** and **12** in CD₃OD

186

187 **Figure 2** Pertinent NOE interactions observed for (-)-epialboctalol (**12**) from NOESY
188 experiment

189

190 In addition to these moracins and stilbenoids, we isolated flavanones steppogenin (**14**) (Lee et
191 al., 2001), katuranin (**15**) (Lee et al., 2001) and dihydromorin (**16**) (Su et al., 2002), together
192 with β -sitosterol (**17**) (Basnet et al., 2003, Aldrich Library of ^{13}C and ^1H FT NMR spectra,
193 1992) and resorcinol (**18**) (Aldrich Library of ^{13}C and ^1H FT NMR spectra, 1992). These
194 known compounds were identified by comparison of the respective spectral and chemical data
195 with those described in the literature.

196

197 Essentially three classes of compounds were isolated in this study: moracins, stilbenes and
198 flavanones. Only resorcinol **18** and β -sitosterol **17** do not belong to these classes. These two
199 compounds are widely distributed in nature and cannot be viewed as chemotaxonomic
200 markers.

201

202 Moracin N, M and P have been isolated before from *Morus alba*. In general, it was found
203 from the literature that *Morus* genus is purveyor of moracins (Tagasuki et al., 1979; Hirakura
204 et al., 1986; Basnet et al., 1993; Nguyen et al., 2009). The only one exception is the isolation
205 of moracin M from *Artocarpus dadah* (Su et al., 2002).

206

207 Among stilbenes, *trans*-oxyresveratrol was isolated from various plants including *Morus* sp.
208 and *Artocarpus* sp. (Hirakura et al, 1986; Su et al, 2002; Shimizu et al., 1998; Song et al,
209 2009). *Trans*-resveratrol was isolated from many sources including the Moraceae *Cudrania*
210 *javanensis* classified today as *Maclura cochinchinensis* (Murti et al., 1972, Chapman & Hall,
211 2006). The distylbene artogomezianol **13** is a constituent of *Artocarpus gomezianus* roots and
212 albolactol **11** was isolated from heartwood of *Morus alba* (Likhitwitayawuid and Sritularak
213 2001, Ferlinahayati et al., 2008).

214

215 Regarding flavonoids, it has been described that many Moraceae can produce steppogenin
216 (El-Sohly et al, 1999; Su et al, 2002; Sheu et al., 2005). Katuranin was also isolated from
217 various biological sources in *Morus* and *Maclura* genera (El-sohly et al., 1999, Lee at al.,
218 2009) and dihydromorin was isolated from *Morus*, *Artocarpus*, and *Maclura* genera (Shimizy
219 et al., 1998, El-Sohly et al, 1999, Su et al., 2002).

220

221 It has been hypothesized before that stilbenes are the major types of compounds isolated from
222 Moraceae and may be useful chemotaxonomic markers (Rowe and Conner, 1979). Also,
223 Schultz has shown that stilbenoids play an important role in the high natural durability of
224 *Maclura pomifera* wood (Schultz et al., 1990). Stilbenes are known as fungicide, termicides
225 and bactericide (Hart and Shrimpton, 1979; Likhitwitayawui and Sritularak, 2001; Javasinghe
226 et al., 2004), and may also exhibit antioxidant properties (Dani et al., 2008; Iacopini et al.,
227 2008; Luo et al., 2005). If it is reasonable to believe that stilbenes are responsible for *Bagassa*
228 *guianensis* heartwood natural durability based on literature precedents, stilbenes can be
229 considered as a secondary chemotaxonomic marker here indicating that *Bagassa* is related to
230 *Morus*, *Artocarpus*, and *Maclura* genera. In Weiblen classification, *Artocarpus* belongs to the
231 Artocarpeae tribe and *Maclura* belongs to the Moreae sensu largo tribe, and both Moreae and
232 Artocarpeae tribes are rather closely related genetically.

233 The peculiarity of *B. guianensis* in comparison with other Moraceae is the very high
234 proportion of moracins. In this matter, it can be hypothesized that *Bagassa* genus is closely
235 related to *Morus* and that moracins are specific to these two genera. These findings are in
236 agreement with Weiblen genetic-based classification where both *Bagassa* and *Morus* belong
237 to the Moreae sensu stricto tribe. It should be mentioned that the *Sorocea* genus, which also
238 belongs to the Moreae s. s. tribe, has been investigated before in the literature and apparently

239 does not contain moracins (see for example Ferrari et al., 2003; Ross et al., 2008). This
240 observation speaks in favor of a very close relationship between *Bagassa* and *Morus*.

241

242 3. Concluding remarks

243 Studies of defensive wood chemicals in *Bagassa guianensis* allowed us to identify large
244 amount of diversely functionalized stilbenes presumably responsible for wood natural
245 durability. In addition, it was found based on the presence of moracins that *Bagassa* is very
246 closely related to *Morus* genus, therefore corroborating Weiblen phylogenetic classification
247 where *B. guianensis* belongs to the Moreae s. s. tribe rather than to the Artocarpeae tribe.

248

249 4. Experimental

250 4.1 General experimental procedure

251 The ^1H and ^{13}C -NMR spectra were recorded on a Bruker Avance DRX500 spectrometer (^1H -
252 500.13 MHz) equipped with a 5 mm triple resonance inverse Cryoprobe TXI (^1H - ^{13}C - ^{15}N),
253 with z gradient. Spectra were recorded with 1.7 mm NMR capillary tube in 40 μL of 99.99%
254 CD_3OD solvent ($\delta_{1\text{H}}$ 3.31 ppm - $\delta_{13\text{C}}$ 49.00 ppm) at 300 K. The ^1H (500 MHz) and ^{13}C NMR
255 (125 MHz) data are reported in ppm downfield from tetramethylsilane. Coupling constants are
256 in Hz and s stands for singlet, d for doublet, t for triplet, q for quartet, m for multiplet and br
257 for broad. Hydrogen connectivity (C, CH, CH_2 , CH_3) information was obtained from edited
258 HSQC and/or DEPTQ-135 experiments. Proton and carbon peak assignments were based on
259 2D NMR analyses (COSY, NOESY, HSQC and HMBC). HREI-MS were performed using a
260 QStar Elite mass spectrometer (Applied Biosystems SCIEX, Concord, ON, Canada) equipped
261 with an ESI source operated in the positive ion mode. The capillary voltage was set at 5,500
262 V, the cone voltage at 20 V and air was used as the nebulizing gas (20 psi). In this hybrid
263 instrument, ions were measured using an orthogonal acceleration time-of-flight (oa-TOF)

264 mass analyzer. Analyst software version 2.1 was used for instrument control, data acquisition
265 and data processing. The accurate mass measurements were performed in triplicate with two
266 internal calibrations. Direct sample introduction was performed at a 5 $\mu\text{L}/\text{min}$ flow rate using
267 a syringe pump. The UV spectra were recorded on a Perkin-Elmer Lambda 5
268 spectrophotometer . Optical rotations were measured with a Perkin-Elmer 241 polarimeter
269 equipped with a sodium lamp (589 nm) and a 1 dm cell. The HPLC separations were
270 performed on a Supelco Discovery[®] HS PEG column (250 \times 21.1 mm, 5 μm) using a Waters
271 system equipped with a W600 pump and a W2996 photodiode array absorbance detector. The
272 samples were injected manually through a Rheodyne injector and the flow rate was 15
273 $\text{mL}\cdot\text{min}^{-1}$. Silica gel 60 (35-70 μm) and analytical TLC plates (Si gel 60 F 254) were
274 purchased from SDS (France). All other chemicals and solvents were analytical grade and
275 purchased from SDS (France).

276 *4.2 Plant Material*

277 *Bagassa guianensis* was collected in Régina, French Guiana. A voucher specimen is kept at
278 the herbarium of Cayenne (CAY-RA13), French Guiana.

279 *4.3 Extraction and isolation*

280 The dried powdered heartwood of *Bagassa guianensis* (140 g) was extracted with ethyl
281 acetate (3 \times 500 mL) at room temperature to give a crude extract which was fractionated first
282 on a silica gel column chromatography with polarity gradient of hexane/ethyl acetate
283 mixtures: 80/20; 50/50; 20/80; 0/100. 9 fractions numbered F1 to F9 were obtained. Fractions
284 F1 to F5 were purified on HPLC with a linear gradient of hexane/isopropanol, by the
285 following method: 70:30 changing over 2 min to 60:40, then to 40:60 at 10 min and pure
286 isopropanol at 15 min and remaining as is for 5 min. The fractions F6 and F9 were analyzed
287 and purified with an isocratic method: 30:70 hexane/isopropanol. These methods allowed us
288 to isolate moracin M **1** (6.2 mg; w/w 0.019%), moracin N **2** (6.7 mg; w/w 0.020%), 6-*O*-

289 methyl-moracin M **3** (3.3 mg; w/w 0.010%), 6-*O*-methyl-moracin-N **4** (9.1 mg; w/w 0.027%),
290 moracin Z **5** (5.2 mg; w/w 0.016%), moracin P **6** (1.2 mg; w/w 0.003), *trans*-resveratrol **7**
291 (12.6 mg; w/w 0.038%), *trans*-oxyresveratrol **8** (112.3 mg; w/w 0.343%), arachidin 2 **9** (5.1
292 mg; w/w 0.015%), arachidin 4 **10** (0.4 mg; w/w 0.001%), alboctalol **11** (0.5 mg; w/w
293 0.001%), (-)-epialboctalol **12** (5.4 mg; w/w 0.016%), artogomezianol **13** (12.7 mg; w/w
294 0.038%), steppogenin **14** (11.5 mg; w/w 0.035%), katuranin **15** (1.5 mg; w/w 0.004%),
295 dihydromorin **16** (20.4 mg; w/w 0.062%), the β -sitosterol **17** (8.4 mg; w/w 0.025%) and the
296 resorcinol **18** (1.8 mg; w/w 0.005%). Compounds 1-6, 9-10 and 17-18 were obtained from
297 the purification of the fractions F1-F5 while compounds 7-8 and 11-16 were isolated from the
298 fractions F6-F9.

299 4.3.1 6-*O*-Methyl-moracin M (**3**)

300 Yellowish amorphous powder; HR-EIMS $[M + H]^+$ m/z 257.0805 $[M + H]^+$ (calcd 257.0808);
301 1H and ^{13}C NMR (500 MHz; CD₃OD) see table 1.

302 4.3.2 6-*O*-Methyl-moracin N (**4**)

303 Yellowish amorphous powder; HR-EIMS $[M + H]^+$ m/z 325.1437 $[M + H]^+$ (calcd 325.1434);
304 1H and ^{13}C NMR (500 MHz; CD₃OD) see table 1.

305 4.3.3 Moracin Z (**5**)

306 Yellowish amorphous powder; HR-EIMS $[M + H]^+$ m/z 343.1542 $[M + H]^+$ (calcd 343.1540);
307 1H and ^{13}C NMR (500 MHz; CD₃OD) see table 1.

308 4.3.4 Arachidin 4 (**10**)

309 Colorless syrup; HR-EIMS $[M + H]^+$ m/z 315.1592 $[M + H]^+$ (calcd 315.1591); 1H and ^{13}C
310 NMR (500 MHz; CD₃OD) see table 2.

311 4.3.5 (-)-Epialboctalol (**12**)

312 Brownish syrup; $[\alpha]_D^{20}$ -7.4° (c 0.004, CH₃OH); HR-EIMS $[M + H]^+$ m/z 489.1540 $[M + H]^+$
313 (calcd 489.1544); 1H and ^{13}C NMR (500 MHz; CD₃OD) see table 3.

314 The 3 known moracins M (**1**), N (**2**) and P (**6**) and the other known compounds 7-9, **11**, and
315 **13-18** were identified by comparison of their physical and spectral data with those reported in
316 the literature.

317

318 **Role of the funding source**

319 The authors gratefully acknowledge the *Programme Amazonie du CNRS* for financial support,
320 as well as the CNRS and the Région Guyane, France, for the research fellowship attributed to
321 MR.

322

323 **References**

324 Aldrich Library of ¹³C and ¹H FT NMR Spectra, 1992, 2, 207A; 243A (nmr).

325 Basnet P., Kadota S., Terashima S., Shimizu M., Namba T., 1993. Two new 2-arylbenzofuran
326 derivatives from hypoglycemic activity-bearing fractions of *Morus insignis*. Chemical
327 and Pharmaceutical Bulletin 41, 1238–1243.

328 Bates R.B., Caldera S., Deshpande V.H., Malik B.L., Paknikar S.K., 1997. Revised structure
329 of alboctalol. Journal of Natural Products 60, 1041–1042.

330 Binbuga N., Ruhs C., Hasty J.K., Henry W.P., Schultz T.P., 2008. Developing
331 environmentally benign and effective organic wood preservatives by understanding
332 the biocidal and non-biocidal properties of extractives in naturally durable heartwood.
333 Holzforschung 62, 264–269.

334 Chapman E., Hall W., 2006. Dictionary of Natural Products , CRC, Version 14:2.

335 Dani C., Bonnato D., Salvador M., Pereira M.D., Henriques J.A.P., Eleutherio E., 2008.
336 Antioxidant protection of resveratrol and catechin in *Saccharomyces cerevisiae*.
337 Journal of Agricultural and Food Chemistry 56, 4268–4272.

338 Dat N.T., Jin X., Lee K., Hong Y.-S., Kim Y.H., Lee J.J., 2009. Hypoxia-Inducible Factor-1
339 Inhibitory Benzofurans and Chalcone-Derived Diels-Alder Adducts from *Morus*
340 Species. *Journal of Natural Products* 72, 39–43.

341 Datwyler S.L., Weiblen G.D., 2004. On the origin of the fig: phylogenetic relationships of
342 Moraceae from *ndhF* sequence. *American Journal of Botany* 91, 767–777.

343 El-Sohly H.N., Joshi A., Li X-C., Ross S.A., 1999. Flavonoids from *Maclura tinctoria*.
344 *Phytochemistry* 52, 141–145.

345 Ferlinahayati S.Y.M., Juliawaty L.D., Achmad S.A., Hakim E.H., Takayama H., Said I.M.,
346 Latip J., 2008. Phenolic constituents from the wood of *Morus australis* with cytotoxic
347 activity. *Zeitschrift fur Naturforschung – Section C. Journal of Biosciences* 63, 35–39.

348 Ferrari F., Cechinel Filho V., Cabras T., Messana I., 2003. Sorocein L and sorocein M: two
349 Diels-Alder type adducts from *Sorocea ilicifolia*. *Journal of Natural Products* 66, 581–
350 582.

351 Han A-R., Kang Y-J., Windono T., Lee S.K., Seo E-K., 2006. Prenylated flavonoids from
352 heartwood of *Artocarpus communis* with inhibitory activity on lipopolysaccharide-
353 induced nitric oxide production. *Journal of Natural Products* 69, 719–721.

354 Hart J.H., Shrimpton D.M., 1979. Role of stilbenes in resistance of wood to decay.
355 *Phytopathology* 69, 1138–1143.

356 Hirakura K., Fujimoto Y., Fukai T., Nomura T., 1986. Two phenolic glycosides from the root
357 bark of the cultivated mulberry tree (*Morus lhou*). *Journal of Natural Products* 49,
358 218-224

359 Hsu, F.-L., Chang H.-T., Chang S.T., 2007. Evaluation of antifungal properties of octyl
360 gallate and its synergy with cinnamaldehyde. *Bioresource Technology* 98, 734–738.

361 Iacopini P., Baldi M., Storchi P., Sebastiani L., 2008. Catechin, epicatechin, quercetin, rutin
362 and resveratrol in red grape: content, *in vitro* antioxidant activity and interactions.
363 Journal of Food Composition and Analysis 21, 589–598.

364 Jahasinghe U.L.B., Samarakoon T.B., Kumarihamy B.M.M., Hara N., Fujimoto Y., 2008.
365 Four new prenylated flavonoids and xanthenes from the root bark of *Artocarpus*
366 *nobilis*. Fitoterapia 79, 37–41.

367 Jayasinghe U.L.B., Puvanendran S., Hara N., Fujimoto Y., 2004. Stilbene derivatives with
368 antifungal and radical scavenging properties from the stem bark of *Artocarpus nobilis*.
369 Natural Products Research 18, 571–574.

370 Lee D., Bhat K.P.L., Fong H.H.S., Farnsworth N.R., Pezzuto J.M., Kinghorn A.D., 2001.
371 Aromatase Inhibitors from *Broussonetia papyrifera*. Journal of Natural Products 64,
372 1286–1293.

373 Lee Y.J., Kim S., Lee S.J., Ham I., Whang W.K., 2009. Antioxydant activities of new
374 flavonoids from *Cudrania tricuspidata* root bark. Archives of Pharmacal Research 32,
375 195–200.

376 Li H., Cheng K.W., Cho C.H., He Z., Wang M., 2007. Oxyresveratrol as an antibrowning
377 agent for cloudy apple juices and fresh-cut apples. Journal of Agricultural and Food
378 Chemistry 55, 2604–2610.

379 Likhitwitayawuid K., Sritularak B., 2001. A new dimeric stilbene with tyrosinase inhibitory
380 activity from *Artocarpus gomezianus*. Journal of Natural Products 64, 1457–1459.

381 Luo M., Liang X.Q., Dang P., Holbrook C.C., Bausher M.G., Lee R.D., Guo B.Z., 2005.
382 Microarray-based screening of differentially expressed genes in peanut in response to
383 *Aspergillus parasiticus* infection and drought stress. Plant Science 169, 695–703.

384 Mabberley D.J. (2002) The Plant Book, second Ed, Cambridge University Press, 858 p.

385 Murti V.V.S., Seshadri T.R., Sivakumaran S., 1972. Cudriniaxanthon and butyrospermol
386 acetate from the roots of *Cudrania javanensis*. *Phytochemistry* 11, 2089–2092.

387 Ngadjui B.T., Watchueng J., Keumedjio F., Nagmeni B., Simo I.K., Abegaz B.M., 2005.
388 Prenylated chalcones, flavones and other constituents of the twigs of *Dorsteronia*
389 *angusticornis* and *Dorsteronia barteri* var. *subtriangularis*. *Phytochemistry* 66, 687–
390 692.

391 Nguyen T.D., Jin X., Lee K., Hog Y-S., Young H.K., Jung J.L., 2009. Hypoxia-inducible
392 factor-1 inhibitory benzofurans and chalcone-derived Diels-Alder adducts from *Morus*
393 species. *Journal of Natural Products* 72, 39–43.

394 Okitani T., Takabe K., Takahashi M., 1999. The role of extractives involved in the natural
395 durability of domestic softwood. *Wood Research* 86, 51–52.

396 Orsini F., Verotta L., Lecchi M., Restano R., Curia G., Redaelli E., Wanke E., 2004.
397 Resveratrol derivatives and their role as potassium channels modulators. *Journal of*
398 *Natural Products* 67, 412–426.

399 Ross S.A., Rodríguez-Guzmán R., Radwan M.M., Jacob M., Ding Y., Li X.-C., Ferreira D.,
400 Manly S.P., 2008. Sorocenols G and H, anti-MRSA oxygen heterocyclic Diels-Alder-
401 type adducts from *Sorocea muriculata* roots. *Journal of Natural Products* 71, 1764–
402 1767.

403 Rowe J.W., Conner A.H., 1979. Extractives in Eastern Hardwood. A review. USDA Forest
404 Service General Technical Report FPL 18. Forest Products Laboratory, Madison, WI,
405 66 pp.

406 Scheffer T.C., Morell J.J., 1998. Natural durability of wood: A worldwide checklist of
407 species. Forest Research Laboratory, Oregon State university; College of Forestry,
408 Research Contribution 22, 45 pp.

409 Schultz P., Hubbard J.T.F., Jin L., Fisher T.H., Nicholas D.D., 1990. Role of stilbenes in the
410 natural durability of wood: Fungicidal structure-activity relationships.
411 *Phytochemistry* 29, 1501–1507.

412 Schultz T.P., Harms W.B., Fischer T.H., McMurtrey K.D., Minn J., Nicholas D.D., 1995.
413 Durability of angiosperm heartwood: the importance of extractives. *Holzforschung* 49,
414 29–34.

415 Schultz P., Nicholas D., 2000. Naturally durable heartwood: evidence for a proposed dual
416 defensive function of the extractives. *Phytochemistry* 5, 47–52.

417 Schultz P., Nicholas D., Preston A.F., 2007. A brief review of the past, present and future of
418 wood preservation. *Pest Management Science* 63, 784–788.

419 Sheu Y-W., Chiang L-C., Chen I-C., Tsai I-L., 2005. Cytotoxic flavonoids and new
420 chromenes from *Ficus formosana* f. *formosana*. *Planta Medica* 71, 1165–1167.

421 Shimizu K., Kondo R., Sakai K., Lee S-H., Sato H., 1998. The inhibitory component from
422 *Artocarpus incisus* on melanin biosynthesis. *Planta Medica* 64, 408–412.

423 Silva C.A., Monteiro B.B., Brazolin S., Lopez A.C.G., Richter A., Braga M.R., 2007.
424 Biodeterioration of brazilwood *Caesalpinia echinata* Lam. (Leguminosae-
425 Caesalpinioideae) by rot fungi and termites. *International Biodeterioration and*
426 *Biodegradation* 60, 285–292.

427 Smith A.L., Campbell C.L., Diwakar M.P., Hanover J.W., Miller R.O., 1989. Extracts from
428 black locust as wood preservatives: A comparison of the methanol extract with
429 pentachlorophenol and chromated copper arsenate. *Holzforschung* 43, 293–296.

430 Song W., Wang H-H., Bucheli P., Zhang P-F., Wei D-Z., Lu Y-H., 2009. Phytochemical
431 profiles of different mulberry (*Morus* sp.) species from China. *Journal of Agricultural*
432 *and Food Chemistry* 57, 9133–9140.

433 Su B.N., Cuendet M., Hawthorne M.E., Kardono L.B.S., Riswan S.F., Harry H.S., Mehta
434 R.G., Pezzuto J.M., Kinghorn A.D., 2002. Constituents of the bark and twigs of
435 *Artocarpus dadah* with cyclooxygenase inhibitory activity. *Journal of Natural*
436 *Products* 65, 163–169.

437 Suttie E.D., Orsler R.J., 1996. The influence of the natural extractives of Opepe (*Nauclea*
438 *diderrichii*) and African Padauk (*Pterocarpus soyauxii*) timbers on their durability.
439 IRG/WP N° 96-30098, 1–15.

440 Takasugi M., Nagao S., Masamune T., 1979. Structure of moracins E, F, G and H, new
441 phytoalexins from diseased mulberry. *Tetrahedron Letters* 48, 4675–4678.

442 Wang S.Y., Chen P.F., Chang S.T., 2005. Antifungal activities of essential oils and their
443 constituents from indigenous cinnamon (*Cinnamomum osmophloeum*) leaves against
444 wood decay fungi. *Bioresource Technology* 96, 813–818.

445 Zerega N.J.C., Clement W.L., Datwyler S.L., Weiblen G.D., 2005. Biogeography and
446 divergence times in the mulberry family (Moraceae). *Molecular Phylogenetics and*
447 *Evolution* 37, 402–416.


448 Zhou C.X., Tanaka J., Cheng C.H.K., Higa T., Tan R.X., 1999. Steroidal alkaloids and
449 stilbenoids from *Veratrum taliense*. *Planta Medica* 65, 480–482.

450


451

452 **Figures and legends**


Moracins




Stilbenoids



Flavanones




Others



453

454 **Figure 1** Compounds **1-18** isolated from *Bagassa guianensis* (Moraceae). (a) New
455 compounds; (b) New names.

456



457
458

459 **Figure 2** Pertinent NOE interactions observed for (-)-epialboctalol (**12**) from NOESY
460 experiment

461

462

463 **Tables**464 **Table 1** ^1H and ^{13}C NMR spectroscopic data for moracins **3-5** in CD_3OD

Atom	3		4		5	
	δ_{C}	δ_{H} (<i>J</i> in Hz)	δ_{C}	δ_{H} (<i>J</i> in Hz)	δ_{C}	δ_{H} (<i>J</i> in Hz)
2	156.5	-	156.2	-	156.2	-
3	96.5	6.95, d (0.6)	102.1	6.90, s	102.0	6.91, d (0.6)
4	121.9	7.43, d (8.5)	121.3	7.25, s	121.6	7.30, s
5	112.9	6.85, dd (8.5, 2.1)	127.5	-	128.6	-
6	159.6	-	157.4	-	157.5	-
7	102.0	7.09, brd (2.0)	94.7	7.09, s	94.7	7.09, s
8	157.0	-	155.7	-	155.8	-
9	123.7	-	123.0	-	123.1	-
1'	133.6	-	133.9	-	133.6	-
2'/6'	104.0	6.78, d (2.1)	103.9	6.77, d (2.1)	103.5	6.77, d (2.1)
3'/5'	159.7	-	159.9	-	160.0	-
4'	103.5	6.25, t (2.1)	103.5	6.24, t (2.1)	103.4	6.25, t (2.1)
1''	-	-	29.7	3.34, brd (7.3)	26.7	2.73, m
2''	-	-	124.3	5.32, tm (7.3)	45.5	1.74, m
3''	-	-	132.7	-	71.5	-
4''	-	-	17.8	1.73 brs	28.9	1.27, s
5''	-	-	26.0	1.74 brs	28.9	1.27, s
MeO	56.0	3.85, s	56.2	3.88 s	56.0	3.89, s

465

466

467 **Table 2** ^1H and ^{13}C NMR spectroscopic data for stilbenes **9** and **10** in CD_3OD

Atom	9		10
	δ_{C}	δ_{H} (<i>J</i> in Hz)	δ_{H} (<i>J</i> in Hz)
1	130.6	-	-
2	128.6	7.31, d (8.6)	7.32, d (8.7)
3	116.5	6.75, d (8.6)	6.75, d (8.7)
4	158.1	-	-
5	116.5	6.75, d (8.6)	6.75, d (8.7)
6	128.6	7.31, d (8.6)	7.32, d (8.7)
α	128.3	6.88, d (16.3)	6.90, d (16.5)
β	127.2	6.74, d (16.3)	6.74, d (16.5)
1'	137.6	-	-
2'	105.7	6.46, s	6.46, s
3'	157.2	-	-
4'	116.0	-	-
5'	157.2	-	-
6'	105.7	6.46, s	6.46, s
1''	23.3	3.28, d (7.1)	2.66, m
2''	124.6	5.23, tm (7.1)	1.68, m
3''	131.4	-	-
4''	26.0	1.62, brs	1.25, s
5''	18.0	1.75, brs	1.25, s

468

469

Atom	12		11
	δ_{C}	δ_{H} (<i>J</i> in Hz)	δ_{H} (<i>J</i> in Hz)
1	156.7	-	-
2	101.8	6.10, d (2.2)	6.32, d (2.2)
3	156.2	-	-
4	107.3	6.19, d (2.2)	6.32, d (2.2)
5 _{ax}	40.1	3.19, brt (13.7)	2.98, dd (16, 14)
5 _{eq}		2.72, dd (15.6, 3.0)	2.53, dd (16.3, 4.3)
6 _{ax}	40.3	3.51, brtd (11.6, 2.1)	3.75, dt (14, 3.7)
7 _{ax}	56.2	3.41, dd (11.3, 8.2)	-
7 _{eq}	-	-	3.28, d (3.3)
8 _{ax}	44.1	4.42, d (8.2)	4.67, brs
9	119.6	-	-
10	142.2	-	-
11	123.7	-	-
12	156.4	-	-
13	103.3	6.16, d (2.2)	?
14	156.4	6.12, dd (8.4, 2.3)	6.13, dd (8.4, 2.3)
15	107.3	-	-
16	129.7	6.82, d (8.2)	6.44, d (8.2)
17	149.2	-	-
18	108.4	6.01, d (1.9)	5.77, d (1.9)
19	157.9	-	-
20	100.8	5.93, t (2.2)	6.02, t (2.2)
21	157.9	-	-
22	108.4	6.01, d (1.9)	5.77, d (1.9)
23	125.2	-	-
24	156.1	6.76, d (8.2)	6.25, d (8.2)
25	103.2	6.23, dd (8.2, 2.5)	6.04, dd (8.2, 2.5)
26	156.7	-	-
27	108.1	6.19, d (2.2)	6.25, d (2.2)
28	131.2	-	-