



HAL
open science

Evolution of a genetic polymorphism with climate change in a Mediterranean landscape

John D. Thompson, Anne Charpentier, Guillaume Bouguet, Faustine Charmasson, Stéphanie Roset, Bruno Buatois, Philippe Vernet, Pierre-Henri Gouyon

► To cite this version:

John D. Thompson, Anne Charpentier, Guillaume Bouguet, Faustine Charmasson, Stéphanie Roset, et al.. Evolution of a genetic polymorphism with climate change in a Mediterranean landscape. Proceedings of the National Academy of Sciences of the United States of America, 2013, 110 (8), pp.2893-2897. 10.1073/pnas.1215833110 . hal-00860487

HAL Id: hal-00860487

<https://hal.science/hal-00860487v1>

Submitted on 9 Dec 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



Evolution of a genetic polymorphism with climate change in a Mediterranean landscape

John Thompson^{a,1}, Anne Charpentier^b, Guillaume Bouguet^{b,c}, Faustine Charmasson^a, Stephanie Roset^b, Bruno Buatois^a, Philippe Vernet^d, and Pierre-Henri Gouyon^e

^aUnité Mixte de Recherche 5175, Centre d'Ecologie Fonctionnelle et Evolutive, Centre National de la Recherche Scientifique, 34293 Montpellier Cedex 5, France; ^bUnité Mixte de Recherche 5175, Centre d'Ecologie Fonctionnelle et Evolutive, Université de Montpellier 2, 34398 Montpellier, France; ^cFlore en Thym, 34725 Saint André de Sangonis, France; ^dLaboratoire de Génétique et Evolution des Populations Végétales, Unité Mixte de Recherche 8198, Centre National de la Recherche Scientifique–Université des Sciences et Technologies de Lille 1, 59655 Villeneuve d'Ascq Cedex, France; and ^eUnité Mixte de Recherche 7205, Origine, Structure et Évolution de la Biodiversité, Muséum National d'Histoire Naturelle, Centre National de la Recherche Scientifique, 75005 Paris, France

Edited by Johanna Schmitt, University of California, Davis, CA, and approved December 21, 2012 (received for review September 13, 2012)

Many species show changes in distribution and phenotypic trait variation in response to climatic warming. Evidence of genetically based trait responses to climate change is, however, less common. Here, we detected evolutionary variation in the landscape-scale distribution of a genetically based chemical polymorphism in Mediterranean wild thyme (*Thymus vulgaris*) in association with modified extreme winter freezing events. By comparing current data on morph distribution with that observed in the early 1970s, we detected a significant increase in the proportion of morphs that are sensitive to winter freezing. This increase in frequency was observed in 17 of the 24 populations in which, since the 1970s, annual extreme winter freezing temperatures have risen above the thresholds that cause mortality of freezing-sensitive morphs. Our results provide an original example of rapid ongoing evolutionary change associated with relaxed selection (less extreme freezing events) on a local landscape scale. In species whose distribution and genetic variability are shaped by strong selection gradients, there may be little time lag associated with their ecological and evolutionary response to long-term environmental change.

Ongoing changes in regional climates are pushing many species to shift their distribution toward higher latitudes and altitudes (1–7). Such changes in species distribution, with an expansion in previously hostile areas and contraction in areas becoming less favorable, can occur rapidly both in plants and animals (2, 3, 5, 6). As a result, major changes in community composition due to differential migration rates may occur (8). Indeed, habitat fragmentation may prevent many species from showing such a distributional response to climate change. As a result, only those species that can respond by phenotypic plasticity or genetically based local adaptation will persist (9). In animal and plant species, phenotypic plasticity of phenological traits can allow individuals to adjust to climate change (1, 10, 11). In addition to changes in distribution and plasticity, an evolutionary response to climate change may occur if species evolve a genetically based adaptation to climate change (12, 13). It is important to distinguish this genetic response from a plastic response of individuals if we are to fully understand the evolutionary potential of species to evolve with climate change (14). Adaptive trait variation in relation to climate change has been shown in the classic study of *Drosophila* (15, 16) and in experimental and natural populations of a small number of plant species (17–20). However, in some cases, the evolutionary response to climate change may be slow due to genetic constraints (21) causing a time lag between the environmental change and an observed evolutionary response. Understanding how species track climate change by genetically based adaptive trait variation and which traits facilitate the evolution of such adaptation is important; such issues determine which species may persist locally and which may shift their distribution (22, 23).

In this study, we test for an evolutionary response of a genetic polymorphism in essential oil composition in Mediterranean wild thyme, *Thymus vulgaris*, to reduced selection associated with a warming of extreme winter freezing events over a sharp spatial

climatic gradient. Our study was done in and around the Saint Martin-de-Londres basin (43°48'N, 03°46'E), which covers an area of ~80 km² and whose southern limits are ~20 km north of Montpellier in the Mediterranean climate region of southern France. The center of the basin (lowest altitude, 145 m) is surrounded by calcareous hills, ranging from 300 to 658 m. The study area has a Mediterranean climate with summer drought but also severe winter freezing temperatures within the basin as a result of a dramatic temperature inversion (Fig. 1). In this area, there are six different chemotypes that are the expression of a genetically controlled polymorphism in *T. vulgaris* (24). Two phenolic chemotypes (carvacrol and thymol) are largely dominant on the slopes outside of the basin on stony soils above 250-m elevation and four nonphenolic chemotypes (linalool, thuyanol-4, α -terpineol, and geraniol) occur within the basin below 200-m elevation on deeper, more humid soils (25–27), where they experience the winter temperature inversion. There is thus a sharp cline in chemotype frequency over only 3–5 km that goes from 100% of either phenolic or nonphenolic chemotypes to 100% of the other form, with a narrow transitional zone (Fig. 2). In short, nonphenolic chemotypes show marked adaptation to habitats, which in the past have frequently experienced extreme freezing temperatures in early winter, whereas phenolic chemotypes are sensitive to intense early-winter freezing and occur in habitats where extreme summer drought can exclude nonphenolic chemotypes (28, 29).

In the part of the study area where nonphenolic chemotypes dominated populations in the early 1970s, extreme winter temperatures have become less severe (Fig. 1) with a significant increase in temperature of extreme freezing events ($r = 0.36$, $n = 56$, $P < 0.01$). In the 20 y before and during the initial study, winter temperatures fell below levels (–15 °C in December) that would exclude phenolic chemotypes from sites dominated by nonphenolic chemotypes (29) on five occasions. In the 16 y following the initial study, three such events were recorded. In the last 20 y, no such extreme events have been recorded in the zone dominated by nonphenolic chemotypes. The last extreme freezing event that could impact the composition of thyme populations occurred 25 y ago.

Here, we test the hypothesis that phenolic chemotypes (thymol and carvacrol) now occur in sites where they were previously absent or have increased their frequency in transitional sites due to a relaxation of selection normally associated with extreme early-winter freezing temperatures. To do so, we compared the chemotype composition of populations observed in the early 1970s (26) to that in 2009–2010 for 36 populations sampled along six

Author contributions: J.T., A.C., P.V., and P.-H.G. designed research; J.T., A.C., G.B., F.C., S.R., B.B., P.V., and P.-H.G. performed research; B.B. contributed new reagents/analytic tools; J.T. and A.C. analyzed data; and J.T., A.C., and B.B. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

¹To whom correspondence should be addressed. E-mail: john.thompson@cefe.cnrs.fr.




Fig. 1. Coldest annual temperature from 1955 to 2010 at the weather station of Saint Martin-de-Londres (filled squares), which occurs in the zone dominated by freezing-tolerant nonphenolic chemotypes, and from 1970 to 2010 at the Centre d'Ecologie Fonctionnelle et Evolutive–Centre National de la Recherche Scientifique experimental gardens on the northern periphery of Montpellier (open circles), where natural thyme populations are dominated by freezing-sensitive phenolic chemotypes.

transects. Each transect is <10 km long, each containing six populations, with two “phenolic,” “mixed,” and “nonphenolic” populations (Fig. 2). To provide an indication of whether population-level changes are due to within-population adaptation or migration among populations, we also examine whether any increases in the abundance of phenolic chemotypes are primarily in nonphenolic or mixed populations that are spatially the closest to preexisting phenolic populations.

Results

We compared the contemporary composition of each of the 36 resampled populations with their composition in the early 1970s by geographically pairing samples, i.e., the two samples of each population, and performing a nonparametric Wilcoxon test across the paired samples of a given population. Nonparametric tests were used due to a severely skewed distribution associated with a consistent trend towards increasing abundance of phenolic chemotypes. Indeed, we found that the mean percentage of phenolic chemotypes in a population was significantly ($df = 35, S = 68.5, P < 0.01$) higher in the contemporary samples (overall value of 53.1%) than in those of the initial study (47.7%) (Table 1). The number of populations in which plants with the thymol chemotype were detected increased from 20 in the early 1970s to 30 in 2010 (the increase was from 16 to 22 populations for carvacrol, the other phenolic chemotype). Eight of the 13 exclusively nonphenolic populations in 1974 currently have a mixed composition, whereas only 1 of the 12 populations composed of exclusively phenolic plants in 1974 was observed to contain nonphenolic plants in 2010 (Table 1). In a goodness-of-fit G test, these changes produced a significant total ($G = 12.2, df = 3, P < 0.01$) and a significant heterogeneity among the two types of population ($G = 4.34, df = 1, P < 0.05$). The changes in composition of the initial nonphenolic populations were associated with the appearance of the thymol chemotype in all eight of the populations whose composition changed and the carvacrol chemotype in three of them. For the group of 11 populations that were mixed in 1974, the mean percentage of phenolic chemotypes significantly increased (Wilcoxon test for paired samples: $df = 10, S = 24, P < 0.05$) from 43.2 to 52.4% (Table 1). Nine of these populations showed an increase in the percentage of phenolic plants, primarily due to a 9% increase in the mean percentage of thymol plants (the carvacrol chemotype showed no mean percentage increase in mixed populations). In the eight populations that were mixed at both dates, five showed a 20–30% increase in percentage of phenolic plants, two showed no variation in composition, and one showed a 25% increase in nonphenolic plants.

Finally, we examined whether changes in chemotype frequency are associated with distance among populations. If migration is




Fig. 2. Spatial distribution and chemical composition of sampled thyme populations in the early 1970s and the 36 populations that were resampled in the present study along six transects. The chemical composition of resampled populations indicated on the map is that which was observed in the initial study. Black circles, phenolic populations; open circles, nonphenolic populations; gray circles, mixed populations. On each transect, the six populations are connected by a dashed line and are represented by a slightly larger circle than populations that were not resampled in the present study.

Table 1. Percentage of plants with a freezing-sensitive phenolic chemotype in 36 populations of *Thymus vulgaris* sampled in 1974 and 2010 along six transects (Fig. 2)

Population*	Transect											
	1		2		3		4		5		6	
1	100	100	100	100	100	100	100	100	100	100	100	90
2	100	100	100	100	100	100	100	100	100	100	100	100
3	82	100	91	100	40	67	70	44	60	80	23	24
4	19	47	78	87	16	17	26	46	14	0	0	17
5	0	0	0	0	0	7	0	6	0	14	0	3
6	0	0	0	0	0	3	0	44	0	17	0	0

*Initially phenolic (1 and 2), nonphenolic (5 and 6), and mixed (3 and 4) populations.

responsible for changes in chemotype frequency, increased frequency of phenolic chemotypes would be most likely to occur in nonphenolic or mixed populations that are spatially close to phenolic populations. For the 13 populations that were composed of only nonphenolic plants in 1974, the distance to the closest sampled population at that time ranged from 95 to 947 m. For the eight populations that became mixed in 2010, the mean distance to a known phenolic population was significantly (Student $T = 2.25$, $df = 11$, $P < 0.05$) smaller (mean \pm SE, 381 ± 119 m) than the distance between the five populations that remained composed of only nonphenolic individuals and a known phenolic population (mean \pm SE, 601 ± 200 m).

Discussion

Changes in the intensity and frequency of extreme climate events have fundamental significance for the response of natural organisms to climate change. Extreme climate events are well known to affect the sustainability of agricultural practices (30, 31), drive dramatic shifts in distribution ranges (2, 3, 5, 6), and impose strong directional selection pressures on natural populations (17–20). Here, we report a rapid and probably ongoing spatial reorganization of a genetic polymorphism that is closely associated with a warming of extreme winter freezing events on a highly localized spatial scale.

The changes we report involve a reduced intensity of freezing events and changes in frequency of freezing tolerant and non-tolerant phenotypes in natural populations of the Mediterranean aromatic plant, *Thymus vulgaris*. We observed both a significant appearance of freezing-sensitive phenolic chemotypes in sites where they were historically absent and an increase in their frequency in previously mixed populations. Such changes have occurred in 17 of the 24 populations where they could potentially occur. For more than 20 y, the freezing temperatures that have been shown to cause mortality of the phenolic chemotypes in early winter (28, 29) have not occurred in the study area (Fig. 1). Although our observations do not conclusively demonstrate a causal relationship, the highly sensitive nature of plants with a phenolic chemotype to the occurrence of intense early winter freezing and the reduced intensity of such extreme freezing events in the study area are indicative of an evolutionary change in response to relaxed selection. Such a decline in the frequency of extreme frosts are a repeated element of observed climate change in many areas (30); hence, there may be many species currently adapting to such relaxed selection. Our study illustrates how this can occur on a highly localized landscape scale.

Although most of the changes in population composition we detect are fairly small, some changes involve shifts of up to ~40% increase in percentage of phenolic plants in a population and several changes are related to the appearance or complete loss of morphs from a population. Also, the time period over which the

changes have occurred is of a very short duration. We sampled adult plants and thus may have minimized the chances of observing changes; it would now be interesting to sample a cohort of seedlings and small plants to see whether changes are even more apparent in the most recently established plants. The difficulty here is that, once thyme plants get through a very small seedling stage, they rapidly develop a woody stem and can often live >10 y, sometimes with little extra growth in size after 3–5 y. It is impossible to age such plants and some small woody plants are likely to have the same age as much larger plants. A study of seedlings in the wild and in offspring raised from seed collected in the field could, however, provide information on the ongoing nature of the observed changes.

The increased frequency of phenolic chemotypes in mixed populations and their appearance in populations where they were previously absent is primarily due to increased presence of the thymol chemotype. This chemotype has appeared in all eight of the historically exclusively nonphenolic populations that are now mixed populations. The carvacrol chemotype has appeared in only three of these populations. An important factor that may contribute to these differences concerns the relatively enhanced resistance to intense freezing temperatures in the thymol chemotype compared with the carvacrol chemotype. The latter is the chemotype that most suffers from extreme early winter freezing events in the field (28). In the wild, carvacrol plants occur in habitats that are the least subject to winter freezing, whereas thymol plants often occur in colder sites that are spatially closer to those of nonphenolic plants (25–27). The changes we present have occurred on a landscape scale and there is evidence that they may result from a combination of both local within-population adaptation and gene flow among populations.

First, the epistatic genetic control of the chemical polymorphism (24) may allow for fairly rapid local changes in chemotype frequency and within-population adaptive variation to a change in selection pressure. The chemical polymorphism is associated with a single biosynthetic chain of monoterpene production and five interacting loci. The genes for nonphenolic chemotypes precede those for the phenolic chemotypes in the epistatic chain of loci. In the presence of a dominant “G” allele (the first genetic locus in the chain), a plant will have the geraniol chemotype (the essential oil being primarily composed of the geraniol monoterpene) regardless of its allelic composition at subsequent loci. If a plant is recessive for the “G” allele, but has the dominant “A” allele at the second locus, it will produce the α -terpineol molecule and be that chemotype, again regardless of composition at subsequent loci. The series continues with the thuyanol-4 and then linalool (nonphenolic) chemotypes to the fifth and final locus where a plant that has only recessive alleles for the four “nonphenolic” loci and a dominant “C” allele will have the carvacrol phenotype. Finally, in the absence of any dominant genes in the chain, a plant will have the thymol phenotype. As a result, crosses among nonphenolic plants can produce other chemotypes including the phenolic chemotypes, whereas crosses among phenolic plants only produce phenolic chemotypes. Hence, the genes for phenolic chemotypes, particularly thymol, can be maintained in populations despite strong selection against them because their expression is masked by the presence of genes coding for nonphenolic chemotypes. Any change in selection regime could thus allow for a rapid evolution in the frequency of phenolic chemotypes, especially thymol. Hence, it is probable that at least some of the observed changes result from within-population genetic change and adaptation to relaxed selection on offspring produced by crosses among individuals heterozygous at the loci for nonphenolic chemotypes.

It is important to note here that there is no qualitative plasticity in chemotype expression: a given plant maintains its chemotype (i.e., phenotype) whatever the conditions of growth. The variations that can occur involve quantitative changes such as seasonal variation in the production of monoterpenes; it is well known that the best time to harvest thyme for its essential oil content is during the

flowering season when production is maximized and that production diminishes in winter (32). In samples of different individuals of a given chemotype, we have observed quantitative variation in the percentage composition of the dominant monoterpene; however, all individuals can be qualitatively assigned to the same chemotype (33). This recent study showed that carvacrol plants growing in marginal environments (i.e., toward the transition zone with nonphenolic chemotypes) produce a reduced percentage of carvacrol and a higher proportion of precursor molecules in their essential oil. They do not, however, change chemotype.

Second, the fact that historical nonphenolic populations that experienced the appearance of phenolic plants are closer to known phenolic populations than those populations that did not show such changes provides evidence that gene flow has at least in part contributed to the evolutionary change in chemotype frequency. This result is coherent with our previous work on the population genetics of *T. vulgaris* done in the study area during the 1990s. In a study of the population genetic structure of natural populations on three transects, each of which spanned nonphenolic, mixed-chemotype, and phenolic populations in the same manner as our current study (two of the transects involved several populations used in the present study), we found low spatial differentiation for allozymes despite strong spatial structure of the genes determining chemical phenotype (34). This study thus provided evidence of fairly high levels of gene flow among thyme populations in our study area. The fact that this study was done in the mid-1990s is indicative of a history of gene flow across the study landscape.

In *T. vulgaris*, chemotype migration can occur through the dissemination of seeds or pollen. Thyme seeds are not anemochorous and have no structure to facilitate animal dispersal and thus do not disperse more than few meters or tens of meters (35), whereas pollen is insect dispersed, potentially over much large distances. Hence, as the comparison of nuclear and chloroplast DNA genetic structure in thyme populations in our study area suggests (36), gene flow mainly occurs via pollen, whose movement is thus a likely source of gene flow for novel chemotype establishment. In our study, distances from nonphenolic populations to “source” phenolic chemotypes in mixed populations were around a few hundred of meters, a distance that is compatible with foraging distances estimated for bees (37), the most abundant contemporary pollinators of thyme (38). In addition, the study populations occur in a relatively nonfragmented garrigues landscape where there are no major barriers to pollinator movement other than small roads, oak woodland, or agricultural fields. In our study system, it thus appears that a combination of local adaptation and migration has contributed to the changes in the spatial organization of the secondary compound polymorphism. A detailed spatial analysis of population limits, neighboring population size, and distances among populations in relation to changes in chemotype composition is now necessary to more accurately identify the relative contributions of local adaptation and gene flow to observed changes in chemotype composition. Untangling their relative contributions to ongoing evolution of the polymorphism provides a fascinating perspective.

In thyme, chemotypes may be directly adapted to their climatic environment or alternatively to changes in other features of the biotic or abiotic environment that occur as a result of climate warming. In addition to the fact that phenolic chemotypes have been repeatedly shown to grow faster than nonphenolic chemotypes in the absence of early-winter freezing (28, 39), trichomes (glands where the oil is stocked on the outer surface of leaves) on nonphenolic adult plants are more fragile with respect to freezing temperatures than those on phenolic plants (29). The latter result indicates that nonphenolic plants may be able to liberate monoterpenes outside of the plant rather than into the leaf upon freezing. Nonphenolic chemotypes may thereby escape any negative effects of freezing (40) by releasing their essential oil into the atmosphere following severe freezing (which phenolic chemotypes are less able to do).

However, in *T. vulgaris* as in other plants that produce secondary compounds, the chemical variation cannot be treated in isolation from the range of biotic interactions that chemotypes may have with co-occurring species. First, changes in herbivore pressure may facilitate increased abundance of phenolic chemotypes due to their overall enhanced resistance to generalist herbivores (41, 42). If such herbivores show a response to climate change, then they may facilitate the colonization of nonphenolic populations by phenolic chemotypes. In addition, *T. vulgaris* shows close interactions with co-occurring grasses (43, 44), which can mediate the nature of plant/parasite/herbivore interactions entertained by the different chemotypes. The nonadditive interactions that can arise between thyme chemotypes in this multitrophic setting make it clear that, although the changes we observed in chemotype composition of natural populations are closely associated with climate warming, the precise processes mediating the response of chemotypes to climate change may lie in a highly complex combination of abiotic and biotic challenges.

Other than at one of our study sites, nonphenolic chemotypes, which are less tolerant of summer drought than phenolic chemotypes, have not colonized phenolic populations located around the basin on rocky and shallow soils more prone to extreme Mediterranean summer drought stress (27, 28). In such sites, there have been ongoing episodes of strong selection against nonphenolic chemotypes, e.g., the extreme summer conditions in 2003, which caused mortality of all chemotypes, but to which the phenolic chemotypes, particularly carvacrol, showed enhanced resistance (28). However, if this were the cause of increased frequency of phenolic chemotypes in our study populations, one would have predicted a significant change primarily caused by an increase in frequency of the carvacrol chemotype, which is not the case.

Climate-induced changes in the distribution of species have been increasingly modeled and documented. We thus now benefit from a corpus of knowledge on the nature of species' responses in terms of distribution change (1–7), how modified distribution patterns may lead to changes in community composition (8) and also potentially cause genetic erosion in areas of range contraction (45). Here, we provide an original illustration of a change in spatial organization of a genetic polymorphism in response to reduced extreme freezing events, which illustrates how, in the absence of any visible shifts in a species range, species' responses to climate change may be guided by gene flow and local adaptation. It is often thought that constraints associated with life history traits and genetic variability (21–23) and time lags (46, 47) may delay the evolutionary response of organisms, particularly plants, to long-term environmental change. Our study, done over a similar time span as work reporting ecological changes in plant species distribution (3, 5), illustrates that a rapid evolutionary response to temperature modifications can occur where genetic variation is combined with a change in a previously strong selection pressure, even for a perennial woody plant.

Materials and Methods

Six transects (Fig. 2), each <10 km long and containing six populations, with two “phenolic,” “mixed,” and “nonphenolic” populations, were sampled across the gradient in chemotype frequency. Transects 2, 3, and 6 were sampled in April 2009, and transects 1, 4, and 5 were sampled in April 2010. Due to changes in land use, one transect had three mixed populations. To be as objective as possible in the choice of the microsite to resample populations once at a given site, we placed the point indication (from an original map of field surveys in the early 1970s) of each original population on a digitalized 1/25,000 scale map of the region and selected its precise latitude and longitude coordinates. With these coordinates, we returned to each site with a global positioning system, and once at the precise point in the field we collected leaves on 30 adult plants (as in the original 1970s sampling scheme), sampled at least 2 m apart within a sampling area of up to 14 × 16 m.

For each plant, essential oils were extracted by putting leaves in 1 mL of methanol for 24 h. Analytical gas chromatography was carried out on

a Shimadzu GCMS-QP21010 fitted with a flame ionization detector and a fused silica capillary column SLB-MS (30 × 0.25 mm; 0.25- μ m thickness). Helium was used as a carrier gas. The injector was heated at 160 °C. The oven temperature was programmed from 50 °C to 180 °C at ~25 min, and increase to 270 °C for a total time of analysis of ~38 min. Compounds were identified by comparison of their retention times and mass spectra with those of standards published by Mass Spectral Library (National Institute of Standards and Technology, 2005)

- Parmesan C (2006) Ecological and evolutionary responses to recent climate change. *Annu Rev Ecol Syst Evol* 37:637–669.
- Chen I-C, Hill JK, Ohlemüller R, Roy DB, Thomas CD (2011) Rapid range shifts of species associated with high levels of climate warming. *Science* 333(6045):1024–1026.
- Walther GR, Beißner S, Burga CA (2005) Trends in the upward shift of alpine plants. *J Veg Sci* 16:541–548.
- Walther GR, et al. (2002) Ecological responses to recent climate change. *Nature* 416 (6879):389–395.
- Kelly AE, Goulden ML (2008) Rapid shifts in plant distribution with recent climate change. *Proc Natl Acad Sci USA* 105(33):11823–11826.
- Hickling R, Roy DB, Hill JK, Fox R, Thomas CD (2006) The distributions of a wide range of taxonomic groups are expanding polewards. *Glob Change Biol* 12:450–455.
- Parmesan C, Yohe G (2003) A globally coherent fingerprint of climate change impacts across natural systems. *Nature* 421(6918):37–42.
- Devictor V, et al. (2012) Differences in the climatic debts of birds and butterflies at a continental scale. *Nat Clim Chang* 2:121–124.
- Jump AS, Peñuelas J (2005) Running to stand still: Adaptation and the response of plants to rapid climate change. *Ecol Lett* 8:1010–1020.
- Bradshaw WE, Holzapfel CM (2008) Genetic response to rapid climate change: It's seasonal timing that matters. *Mol Ecol* 17(1):157–166.
- Cleland EE, Chuine I, Menzel A, Mooney HA, Schwartz MD (2007) Shifting plant phenology in response to global change. *Trends Ecol Evol* 22(7):357–365.
- Chevin L-M, Lande R, Mace GM (2010) Adaptation, plasticity, and extinction in a changing environment: Towards a predictive theory. *PLoS Biol* 8(4):e1000357.
- Hoffmann AA, Sgrò CM (2011) Climate change and evolutionary adaptation. *Nature* 470(7335):479–485.
- Gienapp P, Teplitsky C, Alho JS, Mills JA, Merilä J (2008) Climate change and evolution: Disentangling environmental and genetic responses. *Mol Ecol* 17(1):167–178.
- Umina PA, Weeks AR, Kearney MR, McKechnie SW, Hoffmann AA (2005) A rapid shift in a classic clinal pattern in *Drosophila* reflecting climate change. *Science* 308(5722):691–693.
- Balanyá J, Oller JM, Huey RB, Gilchrist GW, Serra L (2006) Global genetic change tracks global climate warming in *Drosophila subobscura*. *Science* 313(5794):1773–1775.
- Bradshaw WE, Holzapfel CM (2001) Genetic shift in photoperiodic response correlated with global warming. *Proc Natl Acad Sci USA* 98(25):14509–14511.
- Franks SJ, Sim S, Weis AE (2007) Rapid evolution of flowering time by an annual plant in response to a climate fluctuation. *Proc Natl Acad Sci USA* 104(4):1278–1282.
- Etterson JR (2004) Evolutionary potential of *Chamaecrista fasciculata* in relation to climate change. II. Genetic architecture of three populations reciprocally planted along an environmental gradient in the great plains. *Evolution* 58(7):1459–1471.
- Jump AS, et al. (2008) Simulated climate change provokes rapid genetic change in the Mediterranean shrub *Fumana thymifolia*. *Glob Change Biol* 14:637–643.
- Etterson JR, Shaw RG (2001) Constraint to adaptive evolution in response to global warming. *Science* 294(5540):151–154.
- Davis MB, Shaw RG, Etterson JR (2005) Evolutionary responses to changing climate. *Ecology* 86(7):1704–1714.
- Aitken SN, Yeaman S, Holliday JA, Wang T, Curtis-McClane S (2008) Adaptation, migration or extirpation: Climate change outcomes for tree populations. *Evol App* 1: 95–111.
- Vernet P, Gouyon PH, Valdeyron G (1986) Genetic control of the oil content in *Thymus vulgaris* L.: A case of polymorphism in a biosynthetic chain. *Genetica* 69:227–231.
- Thompson JD (2002) *Thyme: The Genus Thymus*, eds Stahl-Biskup E, Sáez F (Taylor & Francis, London), pp 44–74.
- Vernet P, Guillerme JL, Gouyon PH (1977) Le polymorphisme chimique de *Thymus vulgaris* L. (Labiée) I. Répartition des formes chimiques en relation avec certains facteurs écologiques. *Oecol Plant* 12(2):159–179.
- Gouyon PH, Vernet P, Guillerme JL, Valdeyron G (1986) Polymorphisms and environment: The adaptive value of the oil polymorphisms in *Thymus vulgaris* L. *Heredity* 57: 59–66.
- Thompson JD, et al. (2007) Ongoing adaptation to Mediterranean climate extremes in a chemically polymorphic plant. *Ecol Monogr* 77(3):421–439.
- Amiot J, Salmon Y, Collin C, Thompson JD (2005) Differential resistance to freezing and spatial distribution in a chemically polymorphic plant. *Thymus vulgaris*. *Ecol Lett* 8:370–377.
- Easterling DR, et al. (2000) Climate extremes: Observations, modeling, and impacts. *Science* 289(5487):2068–2074.
- Wigley TML (1985) Impact of extreme events. *Nature* 316:106–107.
- Passet J (1971) *Thymus vulgaris* L.: Chémotaxonomie et biogénèse monoterpénique. PhD thesis (Université de Montpellier 1, Montpellier, France).
- Thompson JD, Chalchat J-C, Michet A, Linhart YB, Ehlers B (2003) Qualitative and quantitative variation in monoterpene co-occurrence and composition in the essential oil of *Thymus vulgaris* chemotypes. *J Chem Ecol* 29(4):859–880.
- Tarayre M, Thompson JD (1997) Population genetic structure of the gynodioecious *Thymus vulgaris* (Labiatae) in southern France. *J Evol Biol* 10:157–174.
- Gouyon P-H, King EB, Bonnet JM, Valdeyron G, Vernet P (1987) Seed migration and the structure of plant populations: An experimental study on *Thymus vulgaris* L. *Oecologia* 72:92–94.
- Tarayre M, Saumitou-Laprade P, Cuguen J, Couvet D, Thompson JD (1997) The spatial genetic structure of cytoplasmic (cpDNA) and nuclear (allozyme) markers within and among populations of the gynodioecious *Thymus vulgaris* (Labiatae) in southern France. *Am J Bot* 84(12):1675–1684.
- Beekman M, Ratnieks FLW (2000) Long-range foraging by the honey-bee, *Apis mellifera* L. *Funct Ecol* 14:490–496.
- Brabant P, Gouyon PH, Lefort G, Valdeyron G, Vernet P (1980) Pollination studies in *Thymus vulgaris* L. (Labiatae). *Oecol Plantarum* 15(1):37–44.
- Thompson JD, Tarayre M, Gauthier P, Litrico I, Linhart YB (2004) Multiple genetic contributions to plant performance in *Thymus vulgaris*. *J Ecol* 92:45–56.
- Daday H (1965) Gene frequencies in wild populations of *Trifolium repens* L. IV. Mechanisms of natural selection. *Heredity* 20:355–365.
- Linhart YB, Thompson JD (1999) Thyme is of the essence: Biochemical polymorphism and multi-species deterrence. *Evol Ecol Res* 1:151–171.
- Linhart YB, Thompson JD (1995) Terpene-based selective herbivory by *Helix aspersa* (Mollusca) on *Thymus vulgaris* (Labiatae). *Oecologia* 102:126–132.
- Ehlers BK, Thompson JD (2004) Do co-occurring plant species adapt to one another? The response of *Bromus erectus* to the presence of different *Thymus vulgaris* chemotypes. *Oecologia* 141(3):511–518.
- Linhart YB, Keefeover-Ring K, Mooney KA, Breland B, Thompson JD (2005) A chemical polymorphism in a multitrophic setting: Thyme monoterpene composition and food web structure. *Am Nat* 166(4):517–529.
- Alsos IG, et al. (2012) Genetic consequences of climate change for northern plants. *Proc Biol Sci* 279(1735):2042–2051.
- Lindborg R, Eriksson O (2004) Historical landscape connectivity affects present plant species diversity. *Ecology* 85:1840–1845.
- Brooks TM, Pimm SL, Oyugi JO (1999) Time lag between deforestation and bird extinctions in tropical forest fragments. *Conserv Biol* 13(5):1140–1150.
- Adams RP (2007) *Identification of Essential Oil Components by Gas Chromatography/Mass Spectroscopy* (Allured Publishing, Carol Stream, IL), 4th Ed.