



HAL
open science

Tracking atmospheric and riverine terrigenous supplies variability during the last glacial and the Holocene in central Mediterranean

V. Bout-Roumazielles, N. Combourieu-Nebout, S. Desprat, G. Siani, J.-L. Turon, L. Essallami

► To cite this version:

V. Bout-Roumazielles, N. Combourieu-Nebout, S. Desprat, G. Siani, J.-L. Turon, et al.. Tracking atmospheric and riverine terrigenous supplies variability during the last glacial and the Holocene in central Mediterranean. *Climate of the Past*, 2013, 9 (3), pp.1065-1087. 10.5194/cp-9-1065-2013 . hal-00861322

HAL Id: hal-00861322

<https://hal.science/hal-00861322>

Submitted on 7 Jun 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Tracking atmospheric and riverine terrigenous supplies variability during the last glacial and the Holocene in central Mediterranean

V. Bout-Roumazeilles¹, N. Combourieu-Nebout², S. Desprat³, G. Siani⁴, J.-L. Turon⁵, and L. Essallami⁶

¹Geosystemes, CNRS – UMR8217, Université Lille 1, 59655 Villeneuve d'Ascq, France

²Laboratoire des Sciences du Climat et de l'Environnement, laboratoire Mixte CNRS-CEA-UVSQ, Avenue de la Terrasse, 91198 Gif-sur-Yvette Cedex, France

³Ecole Pratique des Hautes Etudes, Environnements et Paléoenvironnements Océaniques, CNRS – UMR5808 EPOC, Université Bordeaux I, 33405 Talence, France

⁴Laboratoire des Interactions et Dynamique des Environnements de Surface (IDES), UMR8148, CNRS-Université de Paris-Sud, Bât 504, 91405 Orsay Cedex, France

⁵Environnements et Paléoenvironnements Océaniques, CNRS – UMR5808 EPOC, Université Bordeaux I, 33405 Talence, France

⁶GEOGLOB, Sfax Faculty of Sciences, 3038 Sfax, Tunisia

Correspondence to: V. Bout-Roumazeilles (viviane.bout@univ-lille1.fr)

Received: 12 July 2012 – Published in Clim. Past Discuss.: 27 July 2012

Revised: 5 April 2013 – Accepted: 8 April 2013 – Published: 15 May 2013

Abstract. A multiproxy study – coupling mineralogical, grain size and geochemical approaches – was used to tentatively retrace eolian and fluvial contributions to sedimentation in the Sicilian–Tunisian Strait since the last glacial. The eolian supply is dominant over the whole interval, excepted during the sapropel S1 when riverine contribution apparently became significant. Saharan contribution increased during the Bølling–Allerød, evidencing the persistence of aridity over North Africa although the northern Mediterranean already experienced moister and warmer conditions. The Younger Dryas is marked by proximal dust inputs, highlighting intense regional eolian activity. A southward migration of dust provenance toward Sahel occurred at the onset of the Holocene, likely resulting from a southward position of the Inter Tropical Convergence Zone that was probably associated with a large-scale atmospheric reorganization. Finally, a peculiar high terrigenous flux associated with drastic modifications of the mineralogical and geochemical sediment signature occurred during the sapropel S1, suggesting the propagation of fine particles derived from major floodings of the Nile River – resulting from enhanced rainfall on northeastern Africa – and their transportation across the Sicilian–Tunisian Strait by intermediate water masses.

1 Introduction

The Mediterranean is a transitional area where northern and southern climatic influences tightly interact (e.g. Magny et al., 2009). Previous studies revealed that moist conditions developed in the Mediterranean during the early Holocene while a progressive orbitally-driven trend to aridification characterized the mid- and late Holocene. This climatic evolution displays a contrasting regional pattern, with an abrupt transition at 5.7 kyr BP on the western Mediterranean and a more gradual transition in the eastern Mediterranean (Cheddadi et al., 1991; Ariztegui et al., 2000; Magny et al., 2002, 2007; Frigola et al., 2007; Tzedakis, 2007; Sadori et al., 2008; Roberts et al., 2008, 2011; Jalut et al., 2009; Peyron et al., 2011). Precipitation estimations based on lake levels, fire and pollen association, and on speleothem and isotope records also provide evidences of contrasting seasonality across the Mediterranean during the Holocene (Magny et al., 2003, 2007, 2009; Zanchetta et al., 2007; Tzedakis, 2007; Roberts et al., 2008; Peyron et al., 2011; Vannièrè et al., 2011), but estimating the respective atmospheric and oceanic control on Mediterranean climatic evolution through their impact on eolian and fluvial systems is still complex.

A multiproxy study of the terrigenous supply would help in retracing the variability of both eolian and fluvial systems.

The nature and provenance of fine-grained terrigenous particles in the Mediterranean is mainly controlled by the balance between riverine supplies driven by precipitation regime of the surrounding continents and eolian supplies from the Sahara (Loÿe-Pilot et al., 1986; Martin et al., 1989; Bergametti et al., 1989b; Matthewson et al., 1995; Guerzoni and Chester, 1996; Foucault et Mélières, 2000; Goudie et Middleton, 2001). Several studies used deep-sea clay mineral associations as tracers of source regions and as indicators of water mass fluctuations (Chamley, 1975; Tomadin and Lenaz, 1989; Petschick et al., 1996; Fagel et al., 1997; Gingele et al., 2001; Liu et al., 2003; Boulay et al., 2005; Fagel and Hillaire-Marcel, 2006; Colin et al., 2010). Indeed, the mineralogical nature of sediments, which depends on the petrographic characteristics of their source areas (e.g. Bout-Roumzeilles et al., 1999; Sionneau et al., 2008), has been used to retrace detrital particles provenance in the Mediterranean and thus to assess the respective eolian and riverine contributions to deep-sea sedimentation and provide valuable information on the sediment propagation pathways (Caquineau et al., 1998, 2002; Bout-Roumzeilles et al., 2007; Ehrmann et al., 2007a; Hamann et al., 2009; Kandler et al., 2009; Formenti et al., 2011a,b). The clay mineral fraction also provides information on climatic conditions, such as precipitation and runoff patterns over the adjacent continents (Chamley, 1989; Montero-Serrano et al., 2009, 2010), as well as on the dynamics of river inputs (Pinsak et Murray, 1960). The geochemical signature of clays would complementary help in characterizing the main sources and evidencing specific transportation patterns or transfer processes (Haug et al., 2001; Kandler et al., 2009; Formenti et al., 2011a,b). Grain size distribution, which is primarily driven by physical processes (erosion, deflation, transportation, settling), will help in constraining the main transportation modes (Ehrmann et al., 2007b; Montero-Serrano et al., 2009; Sionneau et al., 2010). Within this framework, combining clay mineralogy with grain size analyses and geochemical tracers would allow for retracing significant variations of detrital supply in the Sicilian-Tunisian Strait and inferring any major modifications of both atmospheric and oceanic terrigenous transfers patterns since the last deglaciation.

2 Geographical settings

2.1 Oceanic circulation

The Mediterranean Sea is a concentration basin where evaporation exceeds precipitation plus freshwater discharge. The surface Atlantic water entering from the Gibraltar Strait transforms into Modified Atlantic Water (MAW) as it flows eastward (Fig. 1). The Levantine Intermediate Water (LIW) flows westward towards the western Mediterranean (Fig. 1)

via the relatively shallow Siculo–Tunisian Strait. The dense Eastern Mediterranean Deep Water (EMDW) fills the deep basin (> 800 m depth) (Wüst, 1961; Pickard et Emery, 1982; Malanotte-Rizzoli and Hecht, 1988; Klein et al., 1999). This general pattern is highly dependent on environmental conditions, including eolian activity and precipitations distribution, and recent alteration of the ocean/atmosphere coupling has resulted in enhanced deep-water formation in the Aegean Sea (Klein et al., 1999; Malanotte-Rizzoli et al., 1999).

2.2 Present-day river supplies

At present day, a major part of detrital clays is supplied to the Mediterranean via rivers (Fig. 1), the most important being the Nile River (from 120 to 230×10^6 t yr⁻¹) discharging in the eastern Mediterranean. The Po River, discharging into the Adriatic Sea (17×10^6 t yr⁻¹), and southeastern European rivers associated with Turkish rivers provide about, respectively, 30×10^6 t yr⁻¹ and 17×10^6 t yr⁻¹ to the Aegean Sea and are major contributors to sedimentation into the central Mediterranean (Holeman, 1968; Milliman and Syvitski, 1992; Stanley et al., 1992; Ehrmann et al., 2007b; Garzanti et al., 2006; Hamann et al., 2009). The detrital supply through the Dardanelles Strait is reduced (0.9×10^6 t yr⁻¹) because most sediment is trapped within the Black Sea and Marmara Sea (Ehrmann et al., 2007a). Central Mediterranean is not affected by detrital supply from the Rhône River nor from the Ebro River, which discharge into the western Mediterranean. Finally, riverine contribution to the central Mediterranean from areas bordering the southern shores is reduced due to the narrow drainage basin and sparse rainfall (Martin and Milliman, 1997). Nevertheless, the Medjerda and Miliane oueds, discharging into the Gulf of Tunis (Fig. 1), may influence the sedimentation in the study area.

2.3 Eolian supply

The estimation of the present-day eolian contribution to deep-sea sedimentation varies from 3.9×10^6 t yr⁻¹ to 120×10^6 t yr⁻¹ (Bergametti et al., 1989c; Matthewson et al., 1995; Guernozi and Chester, 1996; Goudie and Middleton, 2001). This wide range of estimation reflects intermittent and seasonal variations of dust outbreaks toward Europe (D'Almeida, 1986; Guerzoni et al., 1997). Satellite imagery, back trajectories and observations indicate that fine particles originating from North Africa (Fig. 1) are transported by southerly/southwesterly winds (Scirocco, Ghibbli) toward the central Mediterranean (Ganor et al., 2000; Goudie and Middleton, 2001; Washington and Todd, 2005; Engelstaeder et al., 2006; Israelevich et al., 2012). Arid and semi-arid regions of North Africa (Fig. 1) are the main areas of production of dust all year long (Coudé-Gaussen, 1982; Pye, 1987): Tunisia and northern Algeria, Morocco and western Sahara, the South Algeria–Mali region, the Bodélé depression and the southern Egypt–northern Sudan (Brooks and




Fig. 1. Geographical settings: (a) clay mineralogy of peri-Mediterranean river particles (green circles), sediments/soils (black circles) and dust particles (red circles) modified from Bout-Roumzeilles et al. (2007); additional data from Cyprus and Levantine Sea: e.g. Hamann et al. (2009); data from northwest Aegean province and West Turkey province: e.g. Ehrmann et al. (2007a); data from Marmara Sea: Arminot du Châtelet et al. (2013); North and South Libya dust from O'Hara et al. (2006). Position of the sediment cores mentioned in this study. Major, secondary and minor dust sources modified from D'Almeida (1986); Brooks and Legrand (2000); Caquineau et al. (2002); Prospero et al. (2002); Israelevich et al. (2002); Goudie (2003); Formenti et al. (2011a). (b) Main low level (red arrows) and altitude winds (black arrows), rivers supply (green arrows, thickness proportional to annual suspended supply) in the Mediterranean Sea. SAL: Saharan Air Layer, NSAL: northern branch of the Saharan Air Layer. The limit between Sahara and Sahel is reported. Main surface (MAW), intermediate (Levantine Intermediate Water, LIW) and deep water (WMDW and EMDW) masses (Salot, 2005). The inset shows the detailed bathymetry and oceanic circulation of the study area.

Legrand, 2000; Caquineau et al., 2002; Israelevich et al., 2002; Prospero et al., 2002; Goudie, 2003; Formenti et al., 2011a). Most of the northward aerosols transportation across the Mediterranean is linked to the seasonal displacement of cyclones over the Mediterranean (Folger, 1970; Guerzoni et al., 1997; Moulin et al., 1997; Rodriguez et al., 2001). Maximum aerosol is observed over central and eastern Mediterranean in spring (Luck and Ben Othman, 2002; O'Hara et al., 2006) and summer when anticyclonic conditions initiate drought over the area (Prospero et al., 2002; Koren et al., 2006; Roberts et al., 2008; Israelevich et al., 2012). Although Saharan fine particles are mainly produced during warm and hydrolysis period, it is transported from paleosols and small consolidated formations during a dry intervals when vegetation is reduced and eolian erosion becomes more efficient (Brooks and Legrand, 2000). During summer, the Saharan depression favors the transport of dust over the western basin, progressively moving toward central Mediterranean (toward Corsica and Italy) at the end of summer (Bergametti et al., 1989a; Moulin et al., 1997; Barry and Chorley, 1998).

3 Materials and methods

3.1 Materials

The core MD04-2797CQ (36°57' N–11°40' E) was taken during the PRIVILEGE-PRIMAROSA cruise in 2004 at 771 m water depth in the Sicilian–Tunisian Strait (Fig. 1). The Siculo–Tunisian Strait separates the Tunisian and Sicilian continental shelves. Quaternary sedimentation on the southern part of the strait is controlled by marine and eolian supplies. The morphology of the Strait is characterized by two troughs separated by a relatively shallow sill, which likely constrains the westward flow of the deeper part of the saline Levantine Intermediate Water (Astraldi et al., 2002). The upper 6 m of the core are composed of homogenous greenish-grey clays, with some silty clay intervals between 440 and 430 cm and between 155 and 140 cm. The uppermost part of the core is composed of orange-brown clay (Rouis-Zargouni, 2010; Rouis-Zargouni et al., 2010). A complete lithological description of the core is given in Rouis-Zargouni (2010). Smear slides of sediments were taken on 0.5 cm thick layers every 10 cm all along the 6 upper meters of the core. Microscopic observations did not reveal any evidence of volcanic material, but the occurrence of tephras cannot be ruled out in the absence of a detailed high-resolution investigation. According to its geographical setting, the sedimentation of the core is likely influenced by both marine and eolian supplies. The scarcity of rivers along the North African margin and their non-permanent characteristic prevents the area from any major and perennial fluvial contribution (O'Hara et al., 2006). By contrast, the area is subjected to an intense wind-driven supply of dust all year long, carried along by the dominant south-southwestern

low-altitude dusty winds (Scirocco) (Barry and Chorley, 1998) which reach the Adriatic Sea by crossing the Mediterranean (Fig. 1). According to the main wind system, the proximal source of dust for the Sicilian–Tunisian Strait is likely the Tunisia/northern Algeria area (Coudé-Gaussen, 1987). The Mali/Algeria frontier or the Moroccan Atlas (Brooks and Legrand, 2000; Caquineau et al., 2002; Prospero et al., 2002; Goudie, 2003) could also contribute to sedimentation either during intense dust episodes or when a specific and/or seasonal atmospheric configuration affect the main wind directions (Bergametti et al., 1989a,b; Ganor et al., 1991; Pye, 1992; Avila et al., 1997; Goudie and Middleton, 2001; Prospero et al., 2002; Washington and Todd, 2005). We compare our data with previously published $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ records and with the reconstructed salinity ($\Delta\delta_w$) from core MD04-2797 (Essallami et al., 2007). These data will help in constraining the hydrologic properties of the overlying water masses and evidencing alterations of freshwater input/evaporation budget in the central Mediterranean. The reliability of the $\Delta\delta_w$ reconstruction is high when sea surface temperature (SST) changes are small (Essallami et al., 2007).

3.2 Chronostratigraphy

The age model is based on 13 AMS ^{14}C age datings (Rouis-Zargouni, 2010; Rouis-Zargouni et al., 2012). The ages were corrected using reservoir ages of 400 yr during the Holocene, the Younger Dryas and the late glacial, 560 yr during the Bølling–Allerød (B–A) and 800 yr during the Heinrich event 1 and Older Dryas (H1-OD), following recommendations of Siani et al. (2001) (Table 1). The top of the core is dated at 668 yr calBP, suggesting that the sea–sediment interface was not preserved during coring. The terrigenous mass accumulation rates (MART) were calculated as follows: $\text{MART} (\text{g kyr}^{-1} \text{cm}^{-2}) = \text{linear sedimentation rates} (\text{cm kyr}^{-1}) \cdot \text{dry density} (\text{g cm}^{-3}) \cdot (100 - (\% \text{ opal} + \% \text{ CaCO}_3))$. The total CaCO_3 percentage was determined using a Bernard calcimeter and following standard procedures with a precision $< 5\%$. Total carbonate corresponds to the sum of biogenic carbonate plus detrital carbonates provided by eolian and riverine supply from Northern Africa. A recent review of the chemical composition of the main African mineral dust (Formenti et al., 2011b) indicates that dust originating from Tunisia and Algeria, and from the Mali–Algerian border, containing up to 50 % carbonates (Paquet et al., 1984; Coudé-Gaussen, 1991; Alastuey et al., 2005). A maximum carbonate content (5–70 %) is evidenced in western Saharan dust (Avila et al., 1997; Khiri et al., 2004; Kandler et al., 2009). By contrast, dust from central Libya (O'Hara et al., 2006) is characterized by intermediate carbonate content (1–25 %). Considering an eolian contribution toward the Mediterranean ranging from 10 to 50 % (Loÿe-Pilot et al., 1986; Loÿe-Pilot and Martin, 1996; Guerzoni et al., 1997),

Table 1. Radiocarbon ages performed on core MD04-2797. The ages were corrected using reservoir ages of 400 yr during the Holocene, the Younger Dryas and late glacial, 560 yr during the Bølling–Allerød and 800 yr during the Heinrich event H1 and the Older Dryas, following recommendations of Siani et al. (2001).

Depth (cm)	Species	¹⁴ C age (yr BP)	Error ±1σ	Corrected age (yr BP)	Error ±1σ	Calibrated age (cal yr BP)	Error ±1σ	Laboratory
0	<i>G. inflata</i>	1105*	20	705	20	668	9	ARTEMIS
80	foraminifera	5496	95	5093	95	5827	144	ARTEMIS
160	foraminifera	6700	85	6300	85	7241	120	ARTEMIS
199	foraminifera	7523	81	7123	81	7967	71	ARTEMIS
240	foraminifera	8113	81	7713	81	8488	98	ARTEMIS
330	<i>G. ruber</i>	8888	110	8488	110	9477	112	ARTEMIS
410	foraminifera	10 863	32	10 463	32	12 493	49	ARTEMIS
470	<i>G. inflata</i>	12 728	173	12 168	172	14 016	35	ARTEMIS
510	<i>G. ruber</i>	13 900	141	13 100	141	15 934	621	ARTEMIS
610	<i>G. ruber</i>	15 590	50	15 190	50	18 725	370	ARTEMIS
700	foraminifera	17 660	70	17 260	70	20 414	180	ARTEMIS
940	foraminifera	23 415	163	23 015	163	27 879	331	ARTEMIS
1030	foraminifera	26 095	7	25 695	7	30 490	156	ARTEMIS

* Data from Rouis-Zargouni et al. (2010).

carbonate of eolian origin may account for 2 to 35 % of the sediment, whereas total carbonate ranges from 30 to 50 % (Fig. 4). The importance of the eolian carbonate may thus lead to underestimation of MART, especially during the B–A when CaCO₃ represents 50 % of total sediment, but should be less significant over the Holocene as the carbonate content is rather stable.

3.3 Clay minerals

The analyses were performed according to the protocol described in Bout-Roumzeilles et al. (1999). XRD (X-ray diffraction) determinations were performed using a Bruker D4 Endeavor coupled with a Lynxeye detector. Each clay mineral was then characterized by its layer plus inter-layer interval as revealed by XRD analysis (Brindley and Brown, 1980). The illite crystallinity, or Kübler Index, is based on the expression of the width of the illite peak at 10 Å and allows for identifying the anchizone, the limit of diagenesis and the onset of the epizone. The presence of palygorskite has been confirmed by MET observations. Semi-quantitative estimation of clay mineral abundances, based on the pseudo-voigt deconvolution for the doublets illite-palygorskite (10–10.34 Å) and kaolinite-chlorite (3.57–3.53 Å), was performed using the software MacDiff developed by Petschick (2001). Quartz abundance is based on the intensity of diffraction measured at 4.25 Å weighted by the clay-size fraction deposited on the oriented glass.

3.4 Grain size

Grain size analyses were performed on the carbonate- and opal-free fraction of the sediment using a Malvern

Mastersizer 2000 laser (0.02–2000 μm) following protocols described in detail in Montero-Serrano et al. (2009). After deflocculation, an aliquot of the sample was measured using 10 % ultrasonication in the Hydro S dispersion cell, once beam obscuration ranged between 12 and 15 %. The main mode (i.e. most frequent grain size in μm), the percentage of clay (< 2 μm), cohesive silt (2–10 μm), sortable-silt (10–63 μm) and sand (> 63 μm) are reported.

3.5 XRF scanning

A series of element abundances – including K, Ti, Zr, Al – were measured on U-channels using the Avaatech core scanner from the EPOC laboratory at the University of Bordeaux. The sediment was protected with Ultralene during X-ray transmission foil in order to avoid contamination. The data were acquired at a 30 s count time, using 10 kV voltage and 400 mA intensity. Richter et al. (2006) give technical details on the XRF scanning technique. The results are expressed in counts per second, with a 2 % precision according to standard samples. The Zr/Al, Ti/Al and K/Al ratios as used in the text refer to Zr_{XRF}/Al_{XRF}, Ti_{XRF}/Al_{XRF}, and K_{XRF}/Al_{XRF} ratios respectively.

Trace elements are normalized to Al (Tribovillard et al., 2008), in order to avoid dilution effect by carbonates.

4 Results

4.1 Clay mineralogy

The clay mineral fraction is composed, on average, of kaolinite (45 %), smectite (25 %) and illite (15 %). Chlorite and




Fig. 2. Clay mineralogy of core MD04-2797: kaolinite (%), smectite (%), illite (%), chlorite (%), palygorskite (%), I/K: illite to kaolinite ratio, Kübler index (crystallinity), (MART) Terrigenous Mass Accumulation Rates ($\text{g kyr}^{-1} \text{cm}^{-2}$) and quartz content (counts/s) for the last 20 000 yr. B–A = Bølling–Allerød, YD = Younger Dryas, S1 = sapropel S1 (Meroni et al., 2000).

palygorskite are secondary components with 10 and 6% of the clay content, respectively (Table S1 in the Supplement). The composition of the clay mineral assemblage displays some variations over the studied interval (Fig. 2). The kaolinite content remains slightly below 45% between 18.5 and

11.7 ka. Afterward it increases and represents 50% between 10.5 and 8.5 ka. The percentage of kaolinite decreases at 8.5 ka and remains low between 8 and 5.8 ka. It then increases progressively between 5.8 and 2 ka and reaches its maximum ($> 50\%$) around 2 ka. The smectite remains

stable between 18.5 and 8.5 ka, excepted two maxima around 12 and 11 ka (Fig. 2). The smectite content increases up to 38 % of the clay mineral fraction between 8.5 and 5.8 ka, as the kaolinite content is minimum. This smectite-rich interval is interrupted by a slightly depleted level around 7.2 ka (Fig. 2). The smectite content decreases progressively starting at 5.8 ka down to 15 % at the top of the core. The illite content is maximum between 18.5 and 11.7 ka, then it drops down and remains low ($< 15\%$) between 11.7 and 5 ka. Then illite increases slightly and remains stable around 15 % over the more recent part of the record (Fig. 2). The Kübler Index is low ($< 0.4^\circ 2\theta$), indicating high structural order when the illite content is maximum. By contrast, the Kübler Index is higher ($< 0.5^\circ 2\theta$) – low ordering between 11.7 and 5 ka when the percentage of illite is reduced. The illite to kaolinite ratio (I/K) is characterized by a drastic change at 11.7 ka from values around 0.35 to values as low as 0.25. The chlorite content does not correlate with other clay species and shows slight variations, being higher than average between 18.5 and 15.5 ka. The palygorskite is less abundant between 8.5 and 5.8 ka, while smectite is maximum (Fig. 2).

4.2 Grain size

The grain size main mode varies between 4 and 10 μm , indicating the detrital fraction is mainly composed of cohesive silt-size particles. Cohesive particles indeed represent 50 to 70 % of the total terrigenous fraction, being slightly less abundant between 18.5 and 11.7 ka and between 7.5 and 5.5 ka (Fig. 3). By contrast, the clay-size fraction is generally low (around 11 %), excepted between 8.5 and 5.5 ka when it represents up to 25 % of the detrital material. The sortable silt (10 to 63 μm) composes 25 % of the detrital fraction, being more abundant between 13 and 12 ka where it represents 45 %. The sand-size fraction is rare, but shows two maxima between 14 and 12 ka and between 8.5 and 7.5 ka (Fig. 3).

4.3 Elemental geochemistry

Zr/Al and Ti/Al are used as eolian proxies (Boyle, 1983; Grousset et al., 1989; Martinez et al., 1999) because Ti and Zr are conservative. Ti oxides are present in the silt and fine-sand fractions from highly weathered bedrocks. Due to their density and size, Ti oxides are the main components of fine eolian dust together with quartz (e.g. Calvert and Pedersen, 2007). Due to its resistance to chemical weathering, and to its high specific gravity, zircon mainly belongs to the fine and medium sand fraction together with quartz. As a consequence, the Zr/Al ratio can be used as a grain size proxy, and may be useful to retrace coarse eolian input (Calvert and Fontugne, 2001). K/Al is used to reflect the balance between aluminium-rich supply (kaolinite) and potassium-rich supply (illite and feldspars) (Schneider et al., 1997; Yarin-cik et al., 2000). In the Mediterranean, these ratios have been successfully used in estimating the respective river and

eolian contribution to sedimentation (Jiménez-Espejo et al., 2007; Frigola et al., 2008; Rodrigo-Gamiz et al., 2011). The Al content displays a decreasing trend from 18.5 to 14 ka (Fig. 4). Some intervals are characterized by higher than average K/Al ratio, between 14 and 11.7, between 5.8 and 5.5 ka and between 2.5 and 1 ka. The two latter intervals are also characterized by the highest values of the Ti/Al ration (Fig. 4). The Zr/Al ratio displays a peculiar behavior, peaking at 13.4, 7, 6.4, 1.3 ka and being high between 12.8 and 12.3 ka.

5 Discussion

5.1 Mineralogical and geochemical characterization of particle provenance and transport patterns

Illite and chlorite are abundant in sediments from the Ionian Sea (Fig. 1), where they mainly derive from the southern European rivers flowing into the Adriatic (Po River) and Aegean Seas (southeastern European Rivers) (Venkatarathnam et Ryan, 1971; Dominik and Stoffers, 1978; Chamley, 1989; Alonso and Maldonado, 1990; Tomadin, 2000). Smectite predominates in the eastern basin, being abundant in the Marmara Sea (Ergin et al., 2012; Armynot et al., 2013), in the Levantine basin (Hamann et al., 2009) and in the southeastern Aegean Sea (Ehrmann et al., 2007a). Smectite characterizes the Nile River suspended loads, deriving from Cenozoic volcanic provinces of the Ethiopian Highlands (Ukstins et al., 2002; Hamann et al., 2009), and is further redistributed as suspended clay particles within the Mediterranean surface water (Venkatarathnam et Ryan, 1971; Foucault and Mélières, 2000; Hamann et al., 2009). Palygorskite is typical of arid and sub-arid environments from the Mediterranean characterized by chemically restricted conditions (Singer and Galan, 1984; Chamley, 1989). In these Mg-rich environments, alternating moist and drought periods promotes chemical concentration and favors the authigenic formation of palygorskite when evaporitic conditions prevailed (Singer and Galan, 1984). Saharan dust-blown particles contain palygorskite reworked from Neogene North African deposits, in addition to present-day neoformed palygorskite (e.g. Chamley, 1989; Coudé-Gaussen et al., 1982; Molinaroli, 1996; Regaya, 1984, 1992; Elloy and Thomas, 1981). The observed changes in clay mineralogy are rather abrupt, and are thus considered to primarily reflect varying terrigenous provenance/transportation patterns, because alteration/weathering processes are often slow (Thiry, 2000). But interaction between climate (rainfall/temperature) and the neoformation of clays has to be taken in account, because these processes are quite rapid in such evaporitic environments. However, an overview of the main sources of palygorskite related to their geological ages suggests that ancient formations constitute the main source of palygorskite, even if contribution from present-day neoformed palygorskite may




Fig. 3. Grain size analysis of core MD04-2797: main mode (μm), % of sand ($> 63 \mu\text{m}$), % of sortable silt ($10 < > 63 \mu\text{m}$), % of cohesive silt ($2 < > 10 \mu\text{m}$) and % of clay ($< 2 \mu\text{m}$). MART ($\text{g kyr}^{-1}\text{cm}^{-2}$) from Fig. 2. Sea surface temperatures (SST) for May–June and October–November based on foraminifera association (Essallami et al., 2007), $\delta^{18}\text{O}$ (‰) and $\delta^{13}\text{C}$ from planktonic foraminifera *G. bulloides* (Essallami et al., 2007). $\Delta\delta_w$ (‰) values are calculated using SSTs derived from MAT (blue) and alkenones (black) (Essallami et al., 2007). Mean analytical uncertainty is $\pm 0/3$ ‰.




Fig. 4. Geochemical data for core MD04-2797, Al_{XRF} content (counts/s), Ti_{XRF}/Al_{XRF} ratio, Zr_{XRF}/Al_{XRF} ratio (counts/s), $CaCO_3$ (%) and MART ($g\ kyr^{-1}\ cm^{-2}$) from Fig. 2.

be locally of importance (Verrecchia and Le Coustumer, 1996). Palygorskite has been shown to be abundant in Tunisian dunes and is used as a main eolian tracer, because it may be transported over long-range distance, crossing the Mediterranean through meridian transfers (Coudé-Gaussen et al., 1982; Robert et al., 1984; Molinaroli, 1996). Illite is also a dominant component of eolian dust originating from the Sahara when associated with palygorskite (Coudé-Gaussen et Blanc, 1985; Guerzoni et al., 1999; Foucault et Mélières, 2000; Goudie et Middleton, 2001), whereas illite associated with abundant kaolinite originates from south-eastern areas of North Africa (Chester et al., 1977; Stanley et Wingerath, 1996; Foucault et Mélières, 2000; Goudie et Middleton, 2001; O'Hara et al., 2006; Hamann et al., 2009). The illite to kaolinite ratio (I/K) is a useful tracer of Saharan versus Sahelian dust provenance (Schütz and Seibert, 1987; Caquineau et al., 1998, 2002; Formenti et al., 2011a), which is thought to be conservative after long-range transport

(Chester et al., 1972; Caquineau et al., 1998, 2002; Kandler et al., 2009). Highest ratio characterizes the northwestern areas (Fig. 1), whereas low ratio retraces the Sahelian belt: the lowest ratios are observed in dust collected in Sahel (0.1 to 0.3), intermediate values characterize south and central Sahara (0.3 to 0.7), moderate to high values are observed in Tunisia and northern Algeria (1 to 2), whereas highest ratios (1.3 to 2.6 up to 6) occur in northwestern Sahara (Fig. 1) and in Morocco (Chester et al., 1972; Glaccum and Prospero, 1980; Paquet et al., 1984; Coudé-Gaussen, 1991; Kiefert et al., 1996; Avila et al., 1997; Caquineau et al., 1998, 2002; Kandler et al., 2009). The I/K ratio also displays a longitudinal gradient (Fig. 1) with ratio decreasing from northwestern Africa ($I/K=2$ to 1.1) toward northeastern Africa ($I/K=0.2$ to 0.7) (Gomes et al., 1990; Ganor and Forer, 1996; Caquineau et al., 1998). The relation between enhanced supply of eolian particles and climate is complex. Indeed, Saharan dust is produced through deflation/erosion

processes that prevail during arid periods and/or associated with intense eolian activity, when scarce vegetation cover favors soils erosion. Production of Saharan dust particles is favored by the presence of small consolidated soils that mainly formed during a former humid period. Moreover, while arid conditions favor the erosion and transportation of dust, major deposition occurs as wet deposit with precipitation in the western Mediterranean (Bergametti et al., 1989c; Guerzoni et al., 1997; Loÿe-Pilot et al., 1986; Loÿe-Pilot and Martin, 1996). In that sense, enhanced eolian dust input is mainly related to wind activity, and indirectly related to aridity, which limits vegetation cover, promotes soils erosion and increases dust availability. The studied area is characterized by strong seasonal contrast, with warm and dry summers, and cool and wet winters when moisture being advected from the Atlantic Ocean. Increased aridity/summer-like conditions in the Mediterranean thus favor dust production and export toward the Mediterranean. Dust washing from the atmosphere is thus controlled by moisture advection from the Atlantic during winter. However, the relative contribution of wet fall-out versus dry deposits resulting from gravitational settling processes whose importance has been recently evidenced in the northeastern tropical Atlantic Ocean (Skonieczny et al., 2011, 2013) is still a matter of debate in the central Mediterranean.

The grain size of dust displays strong seasonal variations (Guerzoni et al., 1997), with a fine population (2 to 10 μm , main mode around 2.7 μm) characterizing long-distance and high latitude transport and a coarser population (5 to 50 μm , main mode around 20 μm) typical of dust storms carried over few km, being restricted to adjacent continental and marine areas (Prospero, 1981; Torres-Padron et al., 2002). Comparison between Ti/Al and Zr/Al ratios and grain size distribution are thus used to identify specific eolian vs. riverine transport processes, associated with the development of sand dunes during the Holocene, or with remote fine-grained eolian outbreaks or coarse-grained fluvial supply (Haug et al., 2001; Martinez-Ruiz et al., 2003; Ehrmann et al., 2007b; Hamann et al., 2009; Montero-Serrano et al., 2009).

5.2 Bølling–Allerød

The Bølling–Allerød (B–A) is marked by enhanced terrigenous supply starting at 14.6 ka, associated with increased contributions of both illite and palygorskite while kaolinite and chlorite decrease (Fig. 2). The relationship between illite content and sortable silt-size fraction (14 μm) indicates that illite is mostly transported through eolian processes. The increase of the I/K ratio suggests that the increased eolian supply is associated with a modification of the main detrital source (Fig. 5a). The increased proportion of illite may indicate a northward migration of the main clay provenance over Libya, as illite has been shown to be more abundant in dust collected over northern Libya (O’Hara et al., 2006). Nevertheless, the I/K ratio range – varying between 0.35

and 0.4 – during the B–A is rather low compared with observed values for Libya and does not further support this hypothesis (Fig. 5a). The observed sedimentological characteristics tend to indicate Tunisia/northern Algeria as the main dust provenance (Formenti et al., 2011a). According to previously published data, the mineralogical composition is indeed consistent with a source from northern or central Algeria (Fig. 7a), but the low content in palygorskite ruled out any major contribution from Tunisian loess (Bout-Roumazeilles et al., 2007). The development of humid conditions during the B–A is evidenced by an increased contribution of river vs. eolian supply in the Aegean Sea (core SL128, Fig. 1) (Hamann et al., 2008), by pollen association from the Alboran (ODP976, Fig. 1) and Adriatic Sea (Combourieu-Nebout et al., 1998, 2002; Fletcher and Sanchez-Goñi, 2008) and by a speleothem record from the eastern Mediterranean (Bar-Matthew et al., 2003), contrasting with the clay mineral record from the Sicilian–Tunisian Strait. The sea surface temperatures (SST, Fig. 3) – as reconstructed from planktonic foraminifera at site MD04-2797 – are high during the Bølling–Allerød (Ellassami et al., 2007), consistently with general warm oceanic conditions, relatively high humidity and temperatures compared with present-day in Sicily and central Mediterranean (Ramrath et al., 2000; Allen et al., 2002; Zielhofer et al., 2008). The dominance of semi-desert plants at site MD04-2797 (Desprat et al., 2013) confirms the persistence of dry continental conditions over Africa during the B–A. These results tend to confirm contrasting climatic evolution (e.g. Roberts et al., 2008), with humid conditions in the northeastern and eastern Mediterranean but aridity still persistent in the southern Mediterranean.

5.3 Younger Dryas

A major peak in the Zr/Al ratio is observed around 12.8 ka, associated with an increase in the abundance of both sortable silt and sand-size particles (Fig. 5b). The clay association does not exhibit major modification but smectite is more abundant at the expense of kaolinite, displaying a peak at 12 ka (Fig. 2). These characteristics indicate the supply of coarser particles that may either be associated with a volcanic supply, a river supply or an intense dust episode (Martinez-Ruiz et al., 2003; Ehrmann et al., 2007b; Hamann et al., 2009; Skonieczny et al., 2011). Smectite peaking at 12 ka supports the hypothesis of a tephra (Fig. 5d). Moreover, when compared with the chronology of tephtras (e.g. Zanchetta et al., 2011), the increase of Zr may be associated with the Agrano Pomici Principali (12.8–12.2 ka) tephra (DiVito et al., 1999; Siani et al., 2004). Nevertheless, optical observations of the sediment do not give any support to this hypothesis. The hypothesis of enhanced river supply is also unlikely, considering the general arid climatic conditions that prevailed during the Younger Dryas (YD). Fluvial archives evidence the development of arid conditions in Tunisia (Zielhofer et al., 2002, 2004, 2008; Faust et al.,




Fig. 5. Comparison of multiproxies data, (a) I/K ratio and its interpretation on particles provenance, increased aridity as indicated by Medjerda fluvial dynamic (Tunisia) modified from Zielhofer et al. (2008) and MART ($\text{g kyr}^{-1} \text{cm}^{-2}$) from Fig. 2, (b) grain size compared with $\text{Zr}_{\text{XRF}}/\text{Al}_{\text{XRF}}$ ratio; (c) quartz content (counts/s), $\text{Ti}_{\text{XRF}}/\text{Al}_{\text{XRF}}$ ratio, kaolinite (%) and % of cohesive-silt, (d) Smectite (%), characterizing the sapropel S1, % of clay, $\Delta\delta_w$ (‰) values calculated using SSTs derived alkenones (Essallami et al., 2007), and MART ($\text{g kyr}^{-1} \text{cm}^{-2}$) from Fig. 2.

2004) between 12.4 and 11.8 ka BP, attributed to the Younger Dryas. The Medjerda fluvial system was particularly active, characterized by aggradation, with sand deposits and high sedimentation rates in the mid-floodplain (Zielhofer et al., 2008). Such fluvial dynamic may also be responsible for the major increase in the Zr/Al ratio observed around 12.5 ka BP in the Sicilian–Tunisian Strait (Fig. 4). However, increase fluvial dynamic within the floodplain is not likely associated with enhanced terrigenous supply through the northern outlet, since the runoff was probably severely altered due to increased aridity and overall reduced precipitations. This specific level could thus alternatively reflect intense Sahara eolian activity that has previously been reported in sediment cores from the northeastern tropical Atlantic (deMenocal et al., 2000; Cole et al., 2009; Skonieczny et al., 2013).

Cold intervals are generally associated with intense dust deposition/eolian activity resulting from reduced vegetation, promoting the erosion of soils, low moisture availability and enhanced atmospheric circulation. The North Atlantic Oscillation (NAO) is likely responsible for some part of the Mediterranean climate variability, since its impact on the wind activity and precipitation balance (Pittalwala and Hameed, 1991; Moulin et al., 1997; Sanchez-Goñi et al.,

2002) has been shown to be positive during the YD (Kim et al., 2007). A positive phase of the NAO is thought to be responsible for the development of dry conditions over southern Europe and North Africa and for promoting northward intrusions of Saharan air masses toward the western Mediterranean (ODP976, Fig. 1) during cold climatic events of the last climatic cycle (Combourieu-Nebout et al., 2002; Bout-Roumzeilles et al., 2007). Salinity is low during the YD in central Mediterranean (Fig. 3), as during cold events like H1, suggesting enhanced contribution of less saline waters originating from the Atlantic Ocean (Essallami et al., 2007). These cold events were characterized by increased palygorskite supply associated with the dominance of arid steppe (*Artemisia*) and the occurrence of endemic pollen (*Argania*), pinpointing a dust source from western Morocco being transported along the North African coast (Bout-Roumzeilles et al., 2007). The absence of specific enrichment in palygorskite during YD in the Sicilian–Tunisian Strait indicates that the main long-range transportation pattern of fine Saharan dust is not responsible for the peculiar characteristic of sediments deposited in the Sicilian–Tunisian Strait. This is in agreement with grain size and geochemical data, which rather point to medium- to short-range transportation of coarse




Fig. 6. Comparison of the mineralogical and geochemical signatures before and after 11.7 kyr, **(a)** kaolinite versus illite (XRD counts/s) evidencing contrasting provenance between 18.5 and 11.7 ka (red squares) and between 11.7 ka and top of the core (blue squares); **(b)** illite to kaolinite ratio as a function of illite crystallinity (Kübler Index), diagenesis, epizone and anchizone limits.

particles from proximal continental areas (Pye and Tsoar, 1987; McTainsh et al., 1997; O'Hara et al., 2006). The occurrence of such a coarse-grain proximal supply during the YD, consistent with the presence of loess deposits and coastal dunes in Tunisia (Coudé-Gaussen et al., 1987; Grousset et al., 1992; Crouvi et al., 2010) and further supported by the model suggesting that Libyan desert and coastal areas are major coarse-grain dust sources for central and eastern Mediterranean (Callot et al., 2000), may thus reflect local eolian activity forced by ocean–atmosphere linkage at global scale (Fig. 7b).

5.4 Onset of the Holocene

The end of the Younger Dryas (11.7 ka) is marked by a drastic change in the clay assemblage (Figs. 5a, 6a and b) and a decrease of the sortable silt proportion. The kaolinite increases while illite diminishes, resulting in a progressive decrease of the I/K ratio. This shift, associated with a modification of the illite structural composition, evidences a major change of provenance of the clay fraction, interpreted as an increase contribution of the Sahelian area (Fig. 6). Low I/K indicates enhanced contribution from the southern Algeria–Mali border region (Paquet et al., 1984; Bergametti et al., 1989b; Alastuey et al., 2005; Bout-Roumazeilles et al., 2007), but the clay association suggests a provenance from southern Algeria rather than from northern Mali because kaolinite is more abundant southward (e.g. Bout-Roumazeilles et al., 2007), becoming dominant in southern Algeria whereas the Mali border is characterized by smectite-rich assemblages (Fig. 7c). The abundance of semi-desert pollen taxa exhibits a sharp decrease at 12.25 ka (Desprat et al., 2013), before the mineralogical transition, suggesting that the modification of vegetation, i.e. a major change in steppe composition and a development of scarce deciduous oak woodlands, might prevent soils from weathering,

displacing southward the areas of eolian erosion. The end of the YD is marked by a similar modification of the clay mineral fraction in a core from the northeastern tropical Atlantic off Senegal and interpreted as recording a southward shift of the Inter Tropical Convergence Zone (ITCZ), which strongly affects the transport of Saharan dust toward the Atlantic Ocean. Moreover, the onset of the Holocene is also marked by a decrease of the I/K ratio off Portugal (Stumpf et al., 2011). The synchronicity of the clay modification at these sites, subjected to distinct wind regimes, signifies that the end of the YD is marked by a large-scale atmospheric reorganization over the North African continent, with the development of dry conditions during the early Holocene, affecting both the high-latitude long-range transportation and the regional low-latitude wind regime over the Mediterranean.

5.5 Sapropel S1

The time interval between 8.6 and 5.5 ka is marked by the highest terrigenous flux and smectite content, with a significant grain size fining (Figs. 2 and 5d). The associated increase in particulate organic carbon (POC) during the whole interval suggests pre-sapropelic conditions (Essallami et al., 2007). The $\Delta\delta_w$ signal shows a decrease between 9 and 6.5 ka BP including the deposition of sapropel S1 (Fig. 5d). Both timing and peculiar characteristics suggest this interval to be related to the most recent organic-rich sapropel S1 (Kidd et al., 1978) deposited during the mid-Holocene in the eastern Mediterranean (Emeis et al., 1996, 2000; De Lange et al., 2008). During the sapropel, increased precipitation on the surrounding continents provided freshwater discharge and promoted strong stratification of the water column and enhanced nutrient supply, resulting in increased productivity and improved preservation due to oxygen depletion of the water column (Rossignol-Strick et al., 1982; Rohling, 1994; Abu-Zied et al., 2008). Increased precipitations are




Fig. 7. Main provenance (dashed areas) and potential eolian – low level (red arrows) or altitude winds (black arrows) – and riverine (green arrows) transportation patterns of clay-mineral particles for 5 time slices: **(a)** Bølling–Allerød; **(b)** Younger Dryas; **(c)** early Holocene; **(d)** Sapropel S1; **(e)** late Holocene.

thought to be linked to either orbitally-forced enhanced monsoon activity (Rossignol-Strick et al., 1982; Hilgen, 1991; Lourens et al., 1996; Emeis et al., 2000) or to increased rainfall over the Mediterranean region (Kallel et al., 1997; Magny et al., 2002). Indeed, although some data appear to be consistent over the Mediterranean basin (Sadori et al., 2011), rainfall records display seasonal modulation (Sadori et al., 2004, 2008; Frisia et al., 2006; Magny et al., 2007; Kotthoff et al., 2008; Jalut et al., 2009; Colonese et al., 2010; Giraudi et al., 2011; Peyron et al., 2011; Magny et al., 2012). Speleothems indicate increased moisture availability in both western (Corchia Cave, Zanchetta et al., 2007), and eastern Mediterranean (Soreq Cave, Bar-Matthews et al., 2000), consistent with lakes records from East Anatolia (Stevens et al., 2001; e.g. Roberts et al., 2008), and also give evidence for contrasting moisture originating either from the Atlantic (Bard et al., 2002) or from the Mediterranean (Arz et al., 2003). Moreover, rainfall patterns also display slightly different chronology in Italian lakes (Roberts et al., 2008) and stalagmite records (Frisia et al., 2006; Zanchetta et al., 2007). Consequently, most studies focused on the precipitation distribution and moisture availability that prevailed on the continent at the time of S1.

But investigations of the marine propagation of the sapropelic layer from eastern toward western Mediterranean evidenced the absence of true sapropels and/or synchronous events (Mercone et al., 2000) in the western basin, suggesting lower export of production fluxes into the basin (Martinez-Ruiz et al., 2003) or different hydrographic conditions (Weldeab et al., 2003). Smectite provenance during the sapropel in the Sicilian–Tunisian Strait would provide additional information on the modifications of oceanic environmental conditions. The enhanced supply of smectite may also be eolian, originating from Tunisian loesses, which are characterized by their high content in smectite and palygorskite (Fig. 1) (e.g. Bout-Roumazeilles et al., 2007). However, smectite and palygorskite display opposite variations during the sapropel, suggesting that they are not reworked from the same geological formation. Moreover, the overall evidences of increased moisture availability in the Mediterranean during the sapropel S1 rule out an eolian origin for smectite over that time interval. Considering the site location, a contribution from authigenic smectite should be discussed. Indeed, volcanic ashes deposited in deep-sea sediments are easily altered and may transform into smectite (Griffin et al., 1967). Formation of smectite at the expense of volcanic materials was evidenced near the Santorini archipelago in the eastern Mediterranean (Chamley, 1971). However, a volcanic origin of smectite from Sicilia is not supported by optical observations or by the chronology of the main tephra (Siani et al., 2004). The enhanced supply of smectite may originate from Tunisian loess (Fig. 1) (e.g. Bout-Roumazeilles et al., 2007), but the low content of palygorskite during the smectite-rich interval does not further support this interpretation. Moreover, the enhanced moisture availability over

that period, associated with pollen association – i.e. open oak forest with heath underbrush or maquis and Asteraceae–Poaceae–Cyperaceae steppe (Desprat et al., 2013) preventing soils from erosion – and the geochemical signal give little support to significant eolian activity. Considering the location of the core, near the Tunisian coast and directly under the MAV, the enhanced supply of smectite may result from an increase contribution of the Medjerda Oued, one of the main fluvial systems draining Algeria and Tunisia, as its main stream is transported southeastward via the Mediterranean surface layer (Astraldi et al., 2002). The hydrological regime of an Oued is typically irregular, being seasonally modulated by precipitation regimes, which may provoke major floods during heavy rainfall in winter and spring (Zahar et al., 2008). Even if the contribution of the Medjerda fluvial system to deep-sea sedimentation of the Siculo–Tunisian Strait is relatively low, such major flood events may transport one-year discharge in a few days (Claude et al., 1977). Moreover, the Medjerda suspended loads are enriched in smectite (up to 50%) (Claude et al., 1977), which is consistent with the observed mineralogy of the sapropel S1 deposit. In sub-humid to sub-arid environments, enhanced moisture availability likely promotes the development of vegetation that prevents soils from erosion, whereas enhanced fluvial dynamic occurs during dry conditions (Rohdenburg, 1989; Zielhofer et al., 2008). The Oued Medjerda experienced humid conditions between 11 and 7.8 ka cal BP with soil formation around 9.5 ka cal BP (Zielhofer et al., 2002). Such moist conditions generally do not support enhanced fluvial activity, unless rainfall is characterized by strong seasonality. Pollen assemblages show the expansion of temperate trees and shrubs between 10.1 and 6.6 ka – i.e. open oak forest with heath underbrush or maquis and Asteraceae, Poaceae, Cyperaceae steppe – while Mediterranean vegetation only developed after 6.6 ka cal BP, suggesting wetter conditions during sapropel S1 resulting from increased winter precipitation (Desprat et al., 2013). Strong winter precipitations may thus be responsible for moisture availability during the sapropel S1, sustaining the expansion of vegetation and triggering seasonal fluvial discharge, all of which may be responsible for the enhanced detrital supply observed in the Siculo–Tunisian Strait. However, the rather large sediment supply observed in the Siculo–Tunisian Strait makes the long-term contribution of the Medjerda Oued unrealistic over the whole sapropel S1. Indeed, sedimentation rate, smectite concentration, as well as the Zr/Al ratio and grain size distribution suggest that the terrigenous supply during S1 was multiphased (Fig. 5c and d). Especially the smectite contribution is maximum between 6.8 and 6 ka cal BP, with the Zr/Al ratio peaking at 7 and 6.5 ka cal BP, suggesting enhanced river supply. Enhanced fluvial activity has been evidenced in the Medjerda fluvial system between 6.6 and 6 ka BP (Zielhofer et al., 2008), suggesting that the Medjerda Oued may be responsible for the observed terrigenous supply over that interval. Smectite may alternatively

be supplied toward the Mediterranean by the Nile River and transported as suspended particles within the Mediterranean surface water (Ventakatarathnam and Ryan, 1971; Stanley and Wingerath, 1996; Foucault and Mélières, 2000; Hamann et al., 2009) and, to a lesser extent, by eolian processes since Saharan dust deposited in the eastern Mediterranean basin is composed of kaolinite associated with smectite (Chester et al., 1977). In that context, the LIW (Levantine Intermediate Water) likely redistributes smectite-rich fine-grain size particles as it flows westward in the sub-surface layers across the Siculo–Tunisian Strait (Fig. 1). This interpretation is consistent with an increased contribution of the Blue Nile draining the Cenozoic volcanic provinces of Ethiopian Highlands (Ukstins et al., 2002) during the African Humid period (Revel et al., 2010). In the Levantine basin (SL112, Fig. 1), the terrigenous fraction, deriving from the Nile River (Krom et al., 2002; Schilman et al., 2001; Hamann et al., 2008), reveals an increase in cohesive fine-grained particles during the sapropel S1 (Hamann et al., 2010), interpreted as enhanced pluvial period in Egypt (Revel et al., 2010) during the African Humid Period (AHP) (Cole et al., 2009; deMenocal et al., 2000). Alternatively, smectite may also be redistributed from the Adriatic Sea by the dense Eastern Mediterranean Deep Water (EMDW) flowing at depth across the Siculo–Tunisian Strait (Fig. 1). However, this hypothesis is not supported by data since the major low salinity episode that occurred around 7.7 ka BP likely resulted from a massive discharge of the Po River, which mainly carries illite and chlorite (Combourieu-Nebout et al., 2013; Tomadin, 2000).

Considering uncertainties on the age models, the enhanced contribution of fine particles is synchronous in the Levantine Sea and in the Sicilian–Tunisian Strait, supporting a provenance of the fine smectite-rich particles from the Eastern Mediterranean basin. The abundance of fine particles may thus result from slightly slower oceanic circulation during the sapropel S1, promoting the deposition of the finest fraction through settling processes (Cram and O’Sullivan, 1999). The circulation pattern is consistent with this hypothesis since the Levantine Intermediate Water (200–600 m water depth) overflows toward the western Mediterranean basin across the Sicilian–Tunisian Strait. Moreover, cold-water corals and benthic foraminifera assemblages evidenced the impact of sapropelic conditions in shallow water (600 m water depth) from the Ionian Sea (Fink et al., 2012), giving additional support to a westward propagation of sapropelic conditions through intermediate water masses (Fig. 7d). The occurrence of a smectite-rich interval in the Alboran Sea (Bout-Roumzeilles et al., 2007), and on the Atlantic side of the Gibraltar Strait during the AHP, suggest a potential transport of smectite through the Mediterranean Overflow Water (Stumpf et al., 2010).

5.6 Late Holocene

The upper part of the record is marked by an increase of the Ti/Al and Zr/Al ratios (Fig. 5c), associated with a high supply in quartz, both suggesting enhanced wind activity. The dominance of silt-size particles and the stable MART (Fig. 5c) suggest long-range transportation rather than enhanced eolian supply. The increases in kaolinite and palygorskite abundances (Fig. 5c) as well as the low I/K ratio indicate a provenance from the Sahelian area and/or the southwest Morocco (Fig. 7e). This event is synchronous with the occurrence of one of the cool/arid events in the Mediterranean (Rapid Climate Coolings-RCC; e.g. ~ 1.2–1.0 kyr BP), suggesting relationships with high-latitude climatic phenomena and altered runoff regime (Mayewski et al., 2004). The hypothesis of enhanced eolian activity during this interval is supported by an increase in palygorskite and quartz content, observed between 1.15 and 0.65 ka BP (starting at 1.7 ka BP) in the Alboran Sea (Rodrigo-Gamiz et al., 2011) and by several evidences of arid continental conditions over the Mediterranean North Africa, e.g. vegetation in Tunisia (Brun, 1992; Stevenson et al., 1993), terrestrial archives in Morocco (Zielhofer et al., 2010) and Tunisian fluvial archives (Faust et al., 2004; Zielhofer et al., 2002, 2004, 2008).

The sedimentation rate displays a sharp decrease at 5.5 ka BP, evidencing a major modification of environmental conditions. The late Holocene is characterized by increased palygorskite and quartz contents, while the kaolinite shows a progressive increase starting at 3 ka BP at the expense of smectite, associated with a high Ti/Al ratio. These observations suggest enhanced eolian supply that may be linked with the development of arid conditions on the North African continent. This interpretation is supported by the occurrence of arid periods (Fig. 4) in Tunisia around 4.7, 3.0, 1.6, 1.1 and 0.4 ka (Faust et al., 2004), evidenced by major modifications of fluvial dynamics in the Medjerda Oued and associated with late Holocene North Atlantic coolings (Faust et al., 2004; Zielhofer et al., 2008). Previous studies evidenced a climate change around 4.5 ka cal BP, with distinct regional late Holocene climatic evolution in the Mediterranean (Roberts et al., 2008; Sadori et al., 2008, 2011; Peyron et al., 2011; Vannière et al., 2011; Magny et al., 2012). Southern sites experienced a drastic decrease in summer precipitations (Magny et al., 2012) while northern sites were submitted to increased precipitation. According to present-day atmospheric patterns, the late Holocene mineralogical and geochemical records are consistent with a mean southward position of the ITCZ, which may also explain the low-latitude aridity associated to the RCC (Haug et al., 2001; Hodell et al., 2001). A southward position of the ITCZ would indeed explain the geographically contrasted Mediterranean climatic evolution by promoting the aridification of North Africa during which increased summer precipitation

occurred in northern Mediterranean (Vanni re et al., 2011) as did enhanced high-altitude export of dust (Skonieczny et al., 2011).

6 Conclusions

The clay mineral, grain size and geochemical studies of sediment deposited on and into the Sicilian–Tunisian Strait have helped retrace atmospheric vs. riverine terrigenous supplies variability since the last glacial in central Mediterranean. Although the eolian supply is dominant at the studied site – excepted during the sapropel S1 – both flux and the main provenance of particles display strong variations, related to aridity/moisture balance and vegetation cover, driven by large-scale atmospheric reorganization.

The B lling–Aller d is marked by increased terrigenous flux while both illite and palygorskite became main components of the clay mineral fraction. The concomitant dominance of silt-size particles indicates an eolian origin for this enhanced detrital supply. The geochemical and mineralogical data indicate a dominant Saharan contribution, pinpointing central and northern Algeria as a main provenance. This increased eolian contribution from the Sahara highlights contrasting regional climatic evolution, with the development of moist conditions over the north and eastern Mediterranean during the B lling–Aller d, while glacial aridity persisted in the southern Mediterranean.

A short-term coarse-grained and Zr-rich interval characterized the Younger Dryas, as arid and cold climatic conditions promoted soil erosion. Similar events characterize the western Mediterranean and the northeastern tropical Atlantic Ocean, although clay mineralogy clearly indicates different provenances. These results suggest local increase of wind activity driven by wide ocean–atmosphere interactions.

The onset of the Holocene is marked by a major change of clay mineralogy and cristallinity corresponding to a southward migration of the main clay provenance toward the Sahelian belt. This change is associated with a progressive development of the Mediterranean vegetation and the southward shift of the ITCZ. Similar signals are recorded in the northeastern tropical Atlantic Ocean and off Portugal, suggesting a large-scale atmospheric reorganization.

High terrigenous flux associated with the dominance of very fine-grained Pd-rich smectite characterized the early-to mid-Holocene in central Mediterranean, while the organic content indicates pre-sapropelic conditions. The sedimentological characteristics rule out any eolian supply but rather suggest enhanced riverine contribution from remote areas during the sapropel S1. The peculiar signature of this event in the central Mediterranean and in the Levantine basin suggests the propagation through intermediate water masses of fine-grained smectite originating from the Nile River during the sapropel S1, resulting from enhanced precipitations on northeast Africa, coincident with the late phase of the AHP.

Finally, an increase supply of dust originating from northern Sahel or southern Morocco characterizes the last 3 kyr, indicating intense eolian activity linked to modification of the rainfall regime.

Supplementary material related to this article is available online at: <http://www.clim-past.net/9/1065/2013/cp-9-1065-2013-supplement.pdf>.

Acknowledgements. This study was funded by the ANR LAMA. This manuscript benefits from thorough reviews from one anonymous referee and from Nejib Kallel. Thomas Richter is thanked for his advice on the use of presentation of XRF data. Philippe Martinez (EPOC – University of Bordeaux) is thanked for producing XRF analysis of the core. L ea-Marie Bernard Emaille, Deny Malengros and Romain Abraham are thanked for technical assistance for XRD analysis and laser grain size measurements. This is LSCE contribution no. 4586.

Edited by: D.-D. Rousseau



The publication of this article is financed by CNRS-INSU.

References

- Abu-Zied, R. H., Rohling, E. J., Jorissen, F. J., Fontanier, C., Casford, J. S. L., and Cooke, S.: Benthic foraminiferal response to changes in bottom-water oxygenation and organic carbon flux in the eastern Mediterranean during LGM to Recent times, *Mar. Micropal.*, 67, 46–68, 2008.
- Alastuey, A., Querol, X., Castillo, S., Escudero, M., Avila, A., Cuevas, E., Torres, C., Romero, P.-M., Exposito, F., Garcia, O., Pedro Diaz, J., Dingenen, R. V., and Putaud, J. P.: Characterization of TSP and PM_{2.5} at Izana and Sta. Cruz de Tenerife (Canary Islands, Spain) during a Saharan Dust Episode (July 2002), *Atmos. Environ.*, 39, 4715–4728, 2005.
- Allen, J. R. M., Watts, W. A., McGee, E., and Huntley, B.: Holocene environmental variability – The record from Lago Grande di Monticchio, Italy, *Quatern. Int.*, 88, 69–80, 2002.
- Alonso, B. and Maldonado, A.: Late Quaternary sedimentation patterns of the Ebro turbidite systems (northwestern Mediterranean): Two styles of deep-sea deposition, *Mar. Geol.*, 95, 353–377, 1990.
- Ariztegui, D., Asioli, A., Lowe, J. J., Trincardi, F., Vigliotti, L., Tamburini, F., Chondrogiani, C., Accorsi, C. A., Bandini Mazzanti, M., Mercuri, A. M., Van Der Kaars, S., Mckenzie, J. A., and Oldfield, F. Palaeoclimate and the formation of sapropel S1: inferences from Late Quaternary lacustrine and marine sequences in the central Mediterranean region, *Paleogeogr. Palaeoclimatol.*, 158, 215–240, 2000.

- Armynot du Châtelet, E., Bout-Roumazeilles, V., Coccioni, R., Frontalini, F., Guillot, F., Kaminski, M. A., Recourt, P., Riboulleau, A., Trentesaux, A., Tribouvillard, N., and Ventalon, S.: Environmental control on shell structure and composition of agglutinated foraminifera along a proximal-distal transect in the Mar-mara Sea, *Mar. Geol.*, 335, 114–128, 2013.
- Arz, H. W., Lamy, F., Pätzold, J., Müller, P. J., and Prins, M.: Mediterranean moisture source for an early-Holocene humid period in the Northern Red Sea, *Science*, 300, 118–121, 2003.
- Astraldi, M., Gasparini, G. P., Vetrano, A., and Vignudelli, S.: Hydrographic characteristics and interannual variability of water masses in the Central Mediterranean: a sensitivity test for long-term changes in the Mediterranean Sea, *Deep-Sea Res.-Pt. I*, 49, 661–680, doi:10.1016/s0967-0637(01)00059-0, 2002.
- Avila, A., Queralt-Mitjans, I., and Alarcón, M.: Mineralogical composition of African dust delivered by red rains over northeastern Spain, *J. Geophys. Res.*, 102, 21977–21996, 1997.
- Bard, E., Delaygue, G., Rostek, F., Antonioli, F., Silenzi, S., and Schrag, D.: Hydrological conditions in the western Mediterranean basin during the deposition of Sapropel 6 (ca 175 kyr), *Earth Planet. Sc. Lett.*, 202, 481–494, 2002.
- Bar-Matthews, M., Ayalon, A., and Kaufman, A.: Timing and hydrological conditions of Sapropel events in the eastern Mediterranean, as evident from speleothems, Soreq cave, Israel, *Chem. Geol.*, 169, 145–156, 2000.
- Bar-Matthews, M., Ayalon, A., Gilmour, M., Matthews, A., and Hawkesworth, C. J.: Sea-land oxygen isotopic relationship from planktonic foraminifera and speleothems in the Eastern Mediterranean region and their implication for paleorainfall during interglacial intervals, *Geochim. Cosmochim. Acta*, 67, 3181–3199, 2003.
- Barry, R. G. and Chorley, R. J.: *Atmosphere, Weather and Climate*, 7th Edn., Routledge, London, 409 pp. 1998.
- Bergametti, G., Dutot, A. L., Buat-Menard, P., Losno, R., and Rémoudaki, E.: Seasonal variability of the elemental composition of atmospheric aerosol particles over the northwestern Mediterranean, *Tellus*, 41, 353–361, 1989a.
- Bergametti, G., Gomes, L., Coudé-Gaussen, G., Rognon, P., and Le Coustumer, M. N.: African dust observed over Canary Islands: Source-regions identification and transport pattern for some summer situations *J. Geophys. Res.*, 94, 14855–14864, 1989b.
- Bergametti, G., Gomes, L., Remoudaki, E., Desbois, M., Martin, D., and Buat-Menard, P.: Present transport and deposition patterns of African dusts to the North-Western Mediterranean, in: *Paleoclimatology and Paleometeorology: Modern and past patterns of global atmospheric transport*, edited by: Leinen, M. and Sarnthein, M., Dordrecht, Kluwer Academic, 282, 227–251, 1989c.
- Boulay, S., Colin, C., Trentesaux, A., Frank, N., and Liu, Z.: Sediment sources and East Asian monsoon intensity over the last 450 ky, Mineralogical and geochemical investigations on South China Sea sediments, *Paleogeogr. Palaeocl.*, 228, 260–277, 2005.
- Bout-Roumazeilles, V., Cortijo, E., Labeyrie, L., and Debrabant, P.: Clay mineral evidence of nepheloid layer contribution to the Heinrich layers in the Northwest Atlantic, *Paleogeogr. Palaeocl.*, 146, 211–228, 1999.
- Bout-Roumazeilles, V., Nebout, N. C., Peyron, O., Cortijo, E., Landais, A., and Masson-Delmotte, V.: Connection between South Mediterranean climate and North African atmospheric circulation during the last 50,000 yr BP North Atlantic cold events, *Quaternary Sci. Rev.*, 26, 3197–3215, 2007.
- Boyle, E. A.: Chemical accumulation variations under the Peru Current during the past 130 000 years, *J. Geophys. Res.*, 88, 7667–7680, 1983.
- Brindley, G. W. and Brown, G.: *Crystal Structures of Clay Minerals and their X-ray identification*, Mineralogical Society, London, 1980.
- Brooks, N. and Legrand, M.: Dust variability over northern Africa and rainfall in the Sahel, in: *Linking Climate Change to Landsurface Change*, edited by: McLaren, S. J. and Kniveton, D., Kluwer Academic, Dordrecht, 1–25, 2000.
- Brun, A.: Pollens dans les séries marines du Golfe de Gabès et du plateau des Kerkennah (Tunisie): Signaux climatiques et anthropiques, *Quaternaire*, 3, 31–39, 1992.
- Callot, Y., Marticorena, B., and Bergametti, G.: Geomorphologic approach for modelling the surface features of arid environments in a model of dust emissions: application to the Sahara desert, *Geodin. Ac. Acta*, 13, 245–270, 2000.
- Calvert, S. E. and Fontugne, M. R.: On the late Pleistocene-Holocene sapropel record of climatic and oceanographic variability in the eastern Mediterranean, *Paleoceanography*, 16, 78–94, doi:10.1029/1999PA000488, 2001.
- Calvert, S. E. and Pedersen, T. F.: *Elemental Proxies for Palaeoclimatic and Palaeoceanographic Variability in Marine Sediments: Interpretation and Application*, in: *Proxies in Late Cenozoic Paleoclimatology*, edited by: Hillaire-Marcel, C. and De Vernal, A., Elsevier, Amsterdam, London, 567–644, 2007.
- Caquineau, S., Gaudichet, A., Gomes, L., Magonthier, M.-C., and Chatenet, B.: Saharan dust: clay ratio as a relevant tracer to assess the origin of soil-derived aerosols, *Geophys. Res. Lett.*, 25, 983–986, 1998.
- Caquineau, S., Gaudichet, A., Gomes, L., and Legrand, M.: Mineralogy of Saharan dust transported over northwestern tropical Atlantic Ocean in relation to source regions, *J. Geophys. Res.*, 107, 4251, doi:10.1029/2000JD000247, 2002.
- Chamley, H.: *Recherches sur la sédimentation argileuse actuelle en Méditerranée*, Sciences Géologiques de Strasbourg, Memoir, 35, 225 pp., 1971.
- Chamley, H.: *Sédimentation argileuse en mer Ionienne au Pliocène*, *B. Soc. Geol. Fr.*, 7, 1131–1143, 1975.
- Chamley, H.: *Clay sedimentology*, Berlin, Springer-Verlag, 623 pp., 1989.
- Cheddadi, R., Rossignol-Strick, M., and Fontugne, M.: Eastern Mediterranean paleoclimates from 26 to 5 ka BP by pollen and isotopic analysis of a core in the anoxic Bannock Basin, *Mar. Geol.*, 100, 53–66, 1991.
- Chester, R., Elderfield, H., Griffin, J. J., Johnson, L. R., and Padgham, R. C.: Eolian dust along the eastern margins of the Atlantic Ocean, *Mar. Geol.*, 13, 91–105, 1972.
- Chester, R., Baxter, G., Behairy, A. K. A., Connor, K., Cross, D., Elderfield, H., and Padgham, R. C.: Soil-sized eolian dusts from the lower troposphere of the eastern Mediterranean Sea, *Mar. Geol.*, 24, 201–217, 1977.
- Claude, J., Francillon, G., and Loyer, J. Y.: Les alluvions déposées par l'Oued Medjerda lors de la crue exceptionnelle de mars 1973, *Cah. ORSTOM, Hydrol.*, 14-1, 37–109, 1977.
- Cole, J. M., Goldstein, S. L., Demenocal, P. B., Hemming, S. R., and Grousset, F. E.: Contrasting compositions of Saharan dust in the

- eastern Atlantic Ocean during the last deglaciation and African Humid Period, *Earth Planet. Sc. Lett.*, 278, 257–266, 2009.
- Colin, C., Siani, G., Sicre, M.-A., and Liu, Z.: Impact of the east Asian monsoon rainfall changes on the erosion of the Mekong River basin over the past 25,000 yr, *Mar. Geol.*, 271, 84–92, 2010.
- Colonese, A. C., Zanchetta, G., Dotsika, E., Drysdale, R. N., Fallick, A. E., Grifoni Cremonesi, R., and Manganelli, G.: Early-Middle Holocene land snail shell stable isotope record from Grotta di Latronico 3 (Southern Italy), *J. Quaternary Sci.*, 25, 1347–1359, 2010.
- Comboureu-Nebout, N., Paterne, M., Turon, J. L., and Siani, G.: A high-resolution record of the last deglaciation on the central Mediterranean Sea: Palaeovegetation and palaeohydrological evolution, *Quaternary Sci. Rev.*, 17, 303–317, 1998.
- Comboureu Nebout, N., Turon, J. L., Zahn, R., Capotondi, L., Londeix, L., and Pahnke, K.: Enhanced aridity and atmospheric high-pressure stability over the western Mediterranean during the North Atlantic cold events of the past 50 k.y., *Geology*, 30, 863–866, 2002.
- Comboureu-Nebout, N., Peyron, O., Bout-Roumazeilles, V., Goring, S., Dormoy, I., Joannin, S., Sadori, L., Siani, G., and Magny, M.: Holocene vegetation and climate changes in central Mediterranean inferred from a high-resolution marine pollen record (Adriatic Sea), *Clim. Past Discuss.*, 9, 1969–2014, doi:10.5194/cpd-9-1969-2013, 2013.
- Coudé-Gaussen, G.: Les poussières éoliennes sahariennes, *Mise au point, Revue de géomorphologie dynamique*, 31, 49–69, 1982.
- Coudé-Gaussen, G.: Saharan dust, *J. Geophys. Res.*, 92, 9753–9771, 1987.
- Coudé-Gaussen, G.: Les poussières sahariennes, *John Libbey Euro Text*, 390 pp., 1991.
- Coudé-Gaussen, G. and Blanc, P.: Présence de grains éolisés de palygorskite dans les poussières actuelles et les sédiments récents d'origine désertique, *B. Soc. Geol. F.*, 1, 5 71–579, 1985.
- Coudé-Gaussen, G., Hillaire-Marcel, C., and Rognon, P.: Origine et évolution pédologique des fractions carbonatées dans les loess des Matmata (Sud-Tunisien) d'après leurs teneurs en ^{13}C et ^{18}O , *C. R. Acad. Sc. Paris*, 295, 939–942, 1982.
- Coudé-Gaussen, G., Rognon, P., Rapp, A., and Nihlén, T.: Dating of peridesert loess in Matmata, south Tunisia, by radiocarbon and thermoluminescence methods, *Z. Geomorph. N. F.*, 31, 129–144, 1987.
- Cram, A. and O'Sullivan, G.: Neogene sapropels in the Mediterranean: a review, *Mar. Geol.*, 153, 11–28, 1999.
- Crouvi, O., Amit, R., Enzel, Y., and Gillespie, A. R.: Active sand seas and the formation of desert loess, *Quaternary Sci. Rev.*, 29, 2087–2098, 2010.
- D'Almeida, G. A.: A model for Saharan dust transport, *J. Clim. Appl. Meteorol.*, 25, 903–916, 1986.
- De Lange, G. J., Thomson, J., Reitz, A., Slomp, C. P., Principato, M. S., Erba, E., and Corselli, C.: Synchronous basin-wide formation and redox-controlled preservation of a Mediterranean sapropel, *Nat. Geosci.*, 1, 606–610, 2008.
- deMenocal, P. B., Ortiz, J., Guilderson, T., Adkins, J., Sarnthein, M., Baker, L., and Yarusinsky, M.: Abrupt onset and termination of the African Humid Period: rapid climatic responses to gradual insolation forcing, *Quaternary Sci. Rev.*, 19, 347–361, 2000.
- Desprat, S., Comboureu-Nebout, N., Essallami, L., Sicre, M. A., Dormoy, I., Peyron, O., Siani, G., Bout Roumazeilles, V., and Turon, J. L.: Deglacial and Holocene vegetation and climatic changes in the southern Central Mediterranean from a direct land-sea correlation, *Clim. Past*, 9, 767–787, doi:10.5194/cp-9-767-2013, 2013.
- Di Vito, M. A., Isaia, R., Orsi, G., Southon, J., de Vita, S., D'Antonio, M., Pappalardo, L., and Piochi, M.: Volcanism and deformation since 12,000 years at the Campi Flegrei caldera (Italy), *J. Volcanol. Geoth. Res.*, 91, 221–246, 1999.
- Dominik, J. and Stoffers, P.: The influence of Late Quaternary stagnations on clay sedimentation in the eastern Mediterranean Sea, *Geol. Rund.*, 68, 302–317, 1978.
- Ehrmann, W., Schmiedl, G., Hamann, Y., and Kuhnt, T.: Distribution of clay minerals in surface sediments of the Aegean Sea: a compilation, *Int. J. Earth Sci.*, 96, 769–780, 2007a.
- Ehrmann, W., Schmiedl, G., Hamann, Y., Kuhnt, T., Hemleben, C., and Siebel, W.: Clay minerals in late glacial and Holocene sediments of the northern and southern Aegean Sea, *Paleogeogr. Palaeocl.*, 249, 36–57, 2007b.
- Elloy, R. and Thomas, G.: Dynamique de la g n se des croutes calcaires (calcretes) d velopp es sur series rouges pl istoc enes en Alg rie N. Occ., *Contexte g omorphologique et climatique, P trographie et g ochimie*, *Bull. Centre Rech. Explo.-Prod. Elf-Aquitaine*, 5, 53–112, 1981.
- Emeis, K.-C., Robertson, A. E. S., Richter, C., and Shipboard Scientific Party (Eds.): *Proc. ODP Init. Repts.*, 160, Ocean Drilling Program, College Station, TX, doi:10.2973/odp.proc.ir.160.1996, 1996.
- Emeis, K.-C., Struck, U., Schulz, H.-M., Bernasconi, S., Sakamoto, T., and Martinez-Ruiz, F.: Temperature and salinity of Mediterranean Sea surface waters over the last 16,000 years: constraints on the physical environment of S1 sapropel formation based on stable oxygen isotopes and alkenone unsaturation ratios, *Paleogeogr. Palaeocl.*, 158, 259–280, 2000.
- Engelstaedter, S., Tegen, I., and Washington, R.: North African dust emissions and transport, *Earth Sci. Rev.*, 79, 73–100, doi:10.1016/j.earscirev.2006.06.004, 2006.
- Ergin, M., Bayhan, E., and Temel, A.: Clay mineral distribution in last Glacial-Holocene sediment cores from the eastern Marmara Sea (Cinarcik Basin-Irzmir Gulf Transition), NW-Turkey: Multisources and transport paths in a two-way flow system, *Quatern. Int.*, 261, 53–74, 2012.
- Essallami, I., Sicre, M. A., Kallel, N., Labeyrie, L., and Siani, G.: Hydrological changes in the Mediterranean Sea over the last 30,000 years, *Geochem. Geophys. Geosy.*, 8, 73–100, doi:10.1029/2007GC001587, 2007.
- Fagel, N. and Hillaire-Marcel, C.: Glacial/interglacial instabilities of the Western Boundary Under Current during the last 365 kyr from Sm/Nd ratios of the sedimentary clay-size fractions at ODP site 646 (Labrador Sea), *Mar. Geol.*, 232, 87–99, 2006.
- Fagel, N., Hillaire-Marcel, C., and Robert, C.: Changes in the Western Boundary Undercurrent outflow since the Last Glacial Maximum, from smectite/illite ratios in deep Labrador Sea sediments, *Paleoceanography*, 12, 79–96, 1997.
- Faust, D., Zielhofer, C., Baena, R., and Diaz del Olmo, F.: High resolution fluvial record of late Holocene geomorphic change in northern Tunisia: climatic or human impact?, *Quaternary Sci. Rev.*, 23, 1757–1775, 2004.

- Fink, H. G., Wienberg, C., Hebbeln, D., McGregor, H. V., Schmiedl, G., Taviani, M., and Freiwald, A.: Oxygen control on Holocene cold-water coral development in the eastern Mediterranean Sea, *Deep-Sea Res. Pt. I*, 62, 89–96, 2012.
- Fletcher, W. I. and Sanchez-Goni, M. F.: Orbital- and sub-orbital-scale climate impacts on vegetation of the western Mediterranean basin over the last 48,000 yr, *Quaternary Res.*, 70, 451–464, 2008.
- Folger, D. W.: Wind transport of land derived mineral, biogenic and industrial matter in the Atlantic, *Deep-Sea Res.*, 17, 433–459, 1970.
- Formenti, P., Rajot, J. L., Desboeufs, K., Saïd, F., Grand, N., Chevaillier, S., and Schmechtig, C.: Airborne observations of mineral dust over western Africa in the summer Monsoon season: spatial and vertical variability of physico-chemical and optical properties, *Atmos. Chem. Phys.*, 11, 6387–6410, doi:10.5194/acp-11-6387-2011, 2011a.
- Formenti, P., Schütz, L., Balkanski, Y., Desboeufs, K., Ebert, M., Kandler, K., Petzold, A., Scheuvens, D., Weinbruch, S., and Zhang, D.: Recent progress in understanding physical and chemical properties of African and Asian mineral dust, *Atmos. Chem. Phys.*, 11, 8231–8256, doi:10.5194/acp-11-8231-2011, 2011b.
- Foucault, A. and Mélières, F.: Palaeoclimatic cyclicity in central Mediterranean Pliocene sediments: the mineralogical signal, *Paleogeogr. Palaeoclimatol.*, 158, 311–323, 2000.
- Frigola, J., Moreno, A., Cacho, I., Canals, M., Sierro, F. J., Flores, J. A., Grimalt, J. O., Hodell, D. A., and Curtis, J. H.: Holocene climate variability in the western Mediterranean region from a deepwater sediment record, *Paleoceanography*, 22, PA2209, doi:10.1029/2006PA001307, 2007.
- Frigola, J., Moreno, A., Cacho, I., Canals, M., Sierro, F. J., Flores, J. A., and Grimalt, J. O.: Evidence of abrupt changes in Western Mediterranean Deep Water circulation during the last 50 kyr: a high-resolution marine record from the Balearic Sea, *Quaternary Int.*, 181, 88–104, doi:10.1016/j.quaint.2007.06.016, 2008.
- Frisia, S., Borsato, A., Mangini, A., Spötl, C., and Madonna, G.: Holocene climate variability from a discontinuous stalagmite record and the Mesolithic to Neolithic transition, *Quaternary Res.*, 66, 388–400, 2006.
- Ganor, E. and Foner, H. A.: The mineralogical and chemical properties and behaviour of aeolian Sahara dust over Israel, in: *The Impact of Desert Dust Across the Mediterranean*, edited by: Guerzoni, S. and Chester, R., Kluwer Academic, Norwell, Mass., 163–172, 1996.
- Ganor, E., Foner, H., Brenner, S., Neeman, E., and Lavi, N.: Chemical composition of aerosols settling in Israel following dust storms, *Atmos. Environ.*, 25, 2665–2670, 1991.
- Ganor, E., Deutsch, Y., and Foner, H. A.: Mineralogical composition and sources of airborne settling particles on Lake Kinneret (the sea of Galilee) in Israel, *Water Air Soil Poll.*, 118, 245–262, doi:10.1023/A:1005167230795, 2000.
- Garzanti, E., Ando, S., Vezzoli, G., Abdel Megid, A. A., and El Kammar, A.: Petrology of Nile River sands (Ethiopia and Sudan): sediment budgets and erosion patterns, *Earth Planet. Sc. Lett.*, 252, 327–341, 2006.
- Gingele, F., De Decker, P., and Hillenbrand, C.-D.: Clay mineral distribution in surface sediments between Indonesia and NW Australia – source and transport by ocean currents, *Mar. Geol.*, 179, 135–146, 2001.
- Giraudi, C., Magny, M., Zanchetta, G., and Drysdale, R. N.: The Holocene climatic evolution of the Mediterranean Italy: A review of the continental geological data, *Holocene*, 21, 105–115, 2011.
- Glaccum, R. A. and Prospero, J. M.: Saharan aerosols over the tropical north Atlantic – Mineralogy, *Mar. Geol.*, 37, 295–321, 1980.
- Gomes, L., Bergametti, G., Coudé-Gaussen, G., and Rognon, P.: Submicron desert dust: A sandblasting process, *J. Geophys. Res.*, 95, 927–935, 1990.
- Goudie, A. S.: *Great warm deserts of the world*, Oxford University Press, Oxford, 2003.
- Goudie, A. S. and Middleton, N. J.: Saharan dust storms: nature and consequences, *Earth Sci. Rev.*, 56, 179–204, 2001.
- Griffin, J. J., Koide, M., Höhndorf, A., Hawkins, J. W., and Goldberg, E. D.: Sediments of the Lau Basin, rapidly accumulating volcanic deposits, *Deep-Sea Res.*, 19, 139–148, 1967.
- Grousset, F., Buat-Menard, P., Boust, D., Tian, R.-C., Baudel, S., Pujol, C., and Vergnaud-Grazzini, C.: Temporal changes of Aeolian Saharan input in the Cape Verde abyssal plain since the last Glacial period, *Oceanol. Acta*, 12, 177–185, 1989.
- Grousset, F. E., Rognon, P., Coudé-Gaussen, G., and Pédemay, P.: Origins of peri-Saharan dust deposits traced by their Nd and Sr isotopic composition, *Paleogeogr. Palaeoclimatol.*, 93, 203–212, 1992.
- Guerzoni, S. and Chester, R.: *The impact of Desert Dust across the Mediterranean*, Dordrecht, Kluwer Academic, 1996.
- Guerzoni, S., Molinaroli, E., and Chester, R.: Saharan dust inputs to the Western Mediterranean Sea: depositional patterns, geochemistry and sedimentological implications, *Deep-Sea Res. Pt. II*, 44, 631–654, 1997.
- Guerzoni, S., Chester, R., Dulac, F., Herut, B., Loyé-Pilot, M.-D., Measures, C., Migon, C., Molinaroli, E., Moulin, C., Rossini, P., Saydam, C., Soudine, A., and Ziveri, P.: The role of atmospheric deposition in the biogeochemistry of the Mediterranean Sea, *Prog. Oceanogr.*, 44, 147–190, 1999.
- Hamann, Y., Ehrmann, W., Schmiedl, G., Krüger, S., Stuut, J.-B., and Kuhnt, T.: Sedimentation processes in the Eastern Mediterranean Sea during the Late Glacial and Holocene revealed by end-member modelling of the terrigenous fraction in marine sediments, *Mar. Geol.*, 248, 97–114, 2008.
- Hamann, Y., Ehrmann, W., Schmiedl, G., and Kuhnt, T.: Modern and late Quaternary clay mineral distribution in the area of the SE Mediterranean Sea, *Quaternary Res.*, 71, 453–464, 2009.
- Hamann, Y., Bowen, S. W., Ersoy, O., Ehrmann, W., and Aydar, E.: First evidence of a distal early Holocene ash layer in Eastern Mediterranean deep-sea sediments derived from the Anatolian volcanic province, *Quaternary Res.*, 73, 497–506, 2010.
- Haug, G., Hughen, K. A., Sigman, D. M., Peterson, L. C., and Röhl, U.: Southward migration of the Intertropical Convergence Zone through the Holocene, *Science*, 293, 1304–1308, 2001.
- Hilgen, H. J.: Astronomical calibration of Gauss to Matuyama sapropels in the Mediterranean and implication for the geomagnetic polarity scale, *Earth Planet. Sc. Lett.*, 104, 226–244, 1991.
- Hodell, D. A., Brenner, M., Curtis, J. H., and Guilderson, T.: Solar forcing of drought frequency in the Maya lowlands, *Science*, 292, 1367–1370, doi:10.1126/science.1057759, 2001.
- Holeman, J. N.: The sediment yield of major rivers of the world, *Water Resour. Res.*, 4, 737–744, 1968.
- Israelevich, P. L., Levin, Z., Joseph, J. H., and Ganor, E.: Desert aerosol transport in the Mediterranean region as in-

- ferred from TOMS aerosol index, *J. Geophys. Res.*, 107, 4572, doi:10.1029/2001jd002011, 2002.
- Israeleovich, P., Ganor, E., Alpert, P., Kishcha, P., and Stupp, A.: Predominant transport paths of Saharan dust over the Mediterranean Sea to Europe, *J. Geophys. Res.*, 117, D02205, doi:10.1029/2011JD016482, 2012.
- Jalut, G., Deboubat, J. J., Fontugne, M., and Otto, T.: Holocene circum-Mediterranean vegetation changes: climate forcing and human impact, *Quatern. Int.*, 200, 4–18, 2009.
- Jiménez-Espejo, F. J., Martínez-Ruiz, F., Sakamoto, T., Iijima, K., Gallego-Torres, D., and Harada, N.: Paleoenvironmental changes in the western Mediterranean since the last glacial maximum: high resolution multiproxy record from the Algero-Balearic basin, *Palaeogeogr. Palaeoclimatol.*, 246, 292–306, doi:10.1016/j.palaeo.2006.10.005, 2007.
- Kallel, N., Paterne, J. C., Duplessy, C., Vergnaud-Grazzini, C., Labeyrie, L., Arnold, M., Fontugne, M., and Pierre, C.: Enhanced rainfall in the Mediterranean region during the last sapropel event, *Oceanol. Acta*, 20, 697–712, 1997.
- Kandler, K., Schütz, L., Deutscher, C., Ebert, M., Hofmann, H., Jäckel, S., Jaenicke, R., Knippertz, P., Lieke, K., Massling, A., Petzold, A., Schladitz, A., Weinzierl, B., Wiedensohler, A., Zorn, S., and Weinbruch, S.: Size distribution, mass concentration, chemical and mineralogical composition and derived optical parameters of the boundary layer aerosol at Tinfou, Morocco, during SAMUN 2006, *Tellus B*, 61, 32–50, 2009.
- Khiri, F., Ezaidi, A., and Kabbachi, K.: Dust deposits in Souss-Massa basin, South-West of Morocco: granulometrical, mineralogical and geochemical characterization, *J. Afr. Earth Sci.*, 39, 459–464, 2004.
- Kidd, R. B., Cita, M. B., and Ryan, W. B. F.: Stratigraphy of the Eastern Mediterranean Sapropel Sequences Recovered during DSDP LEG 42A and Their Paleoenvironmental Significance, edited by: Hsü, K. J., Montadert, L., Bernoulli, D., Bizon, G., Cita, M., Erickson, A., Fabricius, F., Garrison, R. E., Kidd, R. B., Mélières, F., Müller, C., and Wright, R. C., Initial Rep. Deep Sea 42, US Govt. Printing Office, Washington, DC, 421–443, 1978.
- Kiefert, L., McTainsh, G. H., and Nickling, W. G.: Sedimentological characteristics of Saharan and Australian Dusts, in: *The Impact of Desert Dust Across the Mediterranean*, edited by: Guerzoni, S. and Chester, R., Kluwer Academic Publishers, Dordrecht, 183–190, 1996.
- Kim, J., Jung, C. H., Choi, B.-C., Oh, S.-N., Brechtel, F. J., Yoon, S.-C., and Kim, S.-W.: Number size distribution of atmospheric aerosols during ACE-Asia dust and precipitation events, *Atmos. Environ.*, 41, 4841–4855, 2007.
- Klein, B., Roether, W., Manca, B. B., Bregant, D., Beitzel, V., Kovacevic, V., and Luchetta, A.: The large deep water transient in the Eastern Mediterranean, *Deep-Sea Res. Pt. I*, 46, 371–414, 1999.
- Koren, I., Kaufman, Y. J., Washington, R., Todd, M. C., Rudich, Y., Vanderlei Martins, J., and Rosenfeld, D.: The Bodélé depression: A single spot in the Sahara that provides most mineral dust to the Amazon forest, *Environ. Res. Lett.*, 1, 014005, doi:10.1088/1748-9326/1/1/014005, 2006.
- Kotthoff, U., Pross, J., Müller, U. C., Peyron, O., Schmiedl, G., Schulz, H., and Bordon, A.: Clim. Dynam. in the borderlands of the Aegean Sea during formation of sapropel S1 deduced from a marine pollen record, *Quaternary Sci. Rev.* 27, 832–845, 2008.
- Krom, M. D., Stanley, J. D., Cliff, R. A., and Woodwad, J. C.: Nile River sediment fluctuations over the past 7000 yr and their key role in sapropel development, *Geology*, 30, 71–74, 2002.
- Liu, Z., Trentesaux, A., Clemens, S. C., Colin, C., Wang, P., Huang, B., and Boulay, S.: Clay mineral assemblages in the northern South China Sea: implications for East Asian monsoon evolution over the past 2 million years. *Mar. Geol.*, 201, 133–146, 2003.
- Lourens, L. J., Antonarakou, F. J., Hilgen, F. J., Van Hoof, A. A. M., Vergnaud-Grazzini, C., and Zachariasse, W. J.: Evaluation of the Plio-Pleistocene astronomical time-scale, *Paleoceanography*, 11, 391–431, 1996.
- Loÿe-Pilot, M.-D. and Martin, J. M.: Saharan dust input to the Western Mediterranean; an eleven years record in Corsica, in: *The impact of desert dust across the Mediterranean*, edited by: Guerzoni, S. and Chester, R., Kluwer Academic, Dordrecht, 191–199, 1996.
- Loÿe-Pilot, M.-D., Martin, J.-M., and Morelli, J.: Influence of Saharan dust on the rainfall acidity and atmospheric input to the Mediterranean, *Nature*, 321, 427–428, 1986.
- Luck, J. M. and Ben Othman, D.: Trace element and Pb isotope variability during rainy events in the NW Mediterranean: constraints on anthropogenic and natural sources, *Chem. Geol.*, 182, 443–460, 2002.
- Magny, M., Miramont, C., and Sivan, O.: Assessment of the impact of climate and anthropogenic factors on Holocene Mediterranean vegetation in Europe on the basis of palaeohydrological records, *Palaeogeogr. Palaeoclimatol.*, 186, 47–59, 2002.
- Magny, M., Bégeot, C., Guiot, J., and Peyron, O.: Contrasting patterns of hydrological changes in Europe in response to Holocene climate cooling phases, *Quaternary Sci. Rev.*, 22, 1589–1596, 2003.
- Magny, M., Vannièrè, B., de Beaulieu, J.-L., Bégeot, C., Heiri, O., Millet, L., Peyron, O., and Walter-Simonnet, A. V.: Early-Holocene climatic oscillations recorded by lake-level fluctuations in west-central Europe and in central Italy, *Quaternary Sci. Rev.*, 26, 1951–1964, 2007.
- Magny, M., Vannièrè, B., Zanchetta, G., Fouache, E., Touchais, G., Petrika, L., Coussot, C., Walter-Simonnet, A. V., and Arnaud, F.: Possible complexity of the climatic event around 4300–3800 cal BP in the central and western Mediterranean, *Holocene*, 19, 823–833, 2009.
- Magny, M., Peyron, O., Sadori, L., Ortu, E., Zanchetta, G., Vannièrè, B., and Tinner, W.: Contrasting patterns of precipitation seasonality during the Holocene in the south- and north-central Mediterranean, *J. Quaternary Sci.*, 27, 290–296, 2012.
- Malanotte-Rizzoli, P. and Hecht, A.: Large-scale properties of the Eastern Mediterranean: a review, *Oceanol. Acta*, 11, 323–335, 1988.
- Malanotte-Rizzoli, P., Manca, B. B., Ribera, D., Alcalá, M., Theocharis, A., Brenner, S., Budillon, G., and Ozsoy, E.: The Eastern Mediterranean in the 80s and in the 90s: the big transition in the intermediate and deep circulations, *Dyn. Atmos. Ocean.*, 29, 365–395, 1999.
- Martin, J. M. and Milliman, J. D.: EROS 2000 (European Rivers Ocean System). The western Mediterranean Sea: an introduction, *Deep-Sea Res.*, 44, 521–529, 1997.
- Martin, J. M., Elbaz-Poulichet, F., Guieu, C., Loÿe-Pilot, M. D., and Han, G.: River versus atmospheric input of material to

- the Mediterranean Sea: an overview, *Mar. Chem.*, 28, 159–182, 1989.
- Martinez, P., Bertrand, P., Shimmield, G. B., Cochrane, K., Jorissen, F. J., Foster, J., and Dignan, M.: Upwelling intensity and ocean productivity changes off Cape Blanc (northwest Africa) during the last 70,000 years: Geochemical and micropaleontological evidence, *Mar. Geol.*, 158, 57–74, 1999.
- Martinez-Ruiz, F., Paytan, A., Kastner, M., Gonzalez-Donoso, J. M., Linares, D., Bernasconi, S., and Jimenez-Espejo, F. J.: A comparative study of the geochemical and mineralogical characteristics of the S1 sapropel in the western and eastern Mediterranean. *Palaeogeogr. Palaeoclimatol.*, 190, 23–37, 2003.
- Matthewson, A. P., Shimmield, G., Kroon, D., and Fallick, A. E.: A 300 kyr high-resolution aridity record of the North African continent, *Paleoceanography*, 10, 677–692, 1995.
- Mayewski, P. A., Rohling, E. J., Stager, J. C., Karlen, W., Maasch, K. A., Meeker, L. D., Meyerson, E. A., Gasse, F., Van Krevelend, S., Holmgren, K., Lee-Thorp, J., Rosqvist, G., Rack, F., Staubwasser, M., Schneider, R. R., and Steig, E. J.: Holocene climate variability, *Quaternary Res.*, 62, 243–255, 2004.
- McTainsh, G. H., Nickling, W. G., and Lynch, A. W.: Dust deposition and particle size in Mali, West Africa, *Catena*, 29, 307–322, 1997.
- Mercone, D., Thomson, J., Croudace, I. W., Siani, G., Paterne, M., and Troelstra, S.: Duration of S1, the most recent sapropel in the eastern Mediterranean Sea, as indicated by accelerator mass spectrometry radiocarbon and geochemical evidence, *Paleoceanography*, 15, 336–347, 2000.
- Milliman, J. D. and Syvitski, J. P. M.: Geomorphic/Tectonic control of sediment discharge to the ocean: The importance of small mountainous rivers, *J. Geol.*, 100, 520–540, 1992.
- Molinari, E.: Mineralogical characterization of Saharan dust with a review to its final destination in Mediterranean sediments, in: *The Impact of Desert Dust Across the Mediterranean*, edited by: Guerzoni, S. and Chester, R., Kluwer Academic, 153–162, 1996.
- Montero-Serrano, J. C., Bout-Roumazeilles, V., Tribouillard, N., Sionneau, T., Riboulleau, A., Bory, A., and Flower, B.: Sedimentary evidence of deglacial megafloods in the northern Gulf of Mexico (Pigmy Basin), *Quaternary Sci. Rev.*, 28, 3333–3347, 2009.
- Montero-Serrano, J. C., Bout-Roumazeilles, V., Sionneau, T., Tribouillard, N., Bory, A., Flower, B. P., Riboulleau, A., Martinez, P., and Billy, I.: Changes in precipitation regimes over North America during the Holocene as recorded by mineralogy and geochemistry of Gulf of Mexico sediments, *Global Planet. Change*, 74, 132–143, 2010.
- Moulin, C., Lambert, C. E., Dulac, F., and Dayan, U.: Control of atmospheric export of dust from North Africa by the North Atlantic Oscillation, *Nature*, 387, 691–694, 1997.
- O'Hara, S. L., Clarke, M. L., and Elatrash, M. S.: Field measurements of desert dust deposition in Libya, *Atmos. Environ.*, 40, 3881–3897, 2006.
- Paquet, H., Coudé-Gaussens, G., and Rognon, P.: Etude minéralogique de poussières sahariennes le long d'un itinéraire entre 19° et 35° de latitude nord, *Rev. Géol. Dyn. Géogr. Phys.*, 25, 257–265, 1984.
- Petschick, R.: MacDiff v4.2.5 (Free Geological Software), available at: <http://www.geol-pal.uni-frankfurt.de/Staff/Homepages/Petschick/classicsoftware.html#MacDiff> (last access: 2 July 2012), Geologisch-Paläontologisches Institut, Universität Frankfurt/Main, 2001.
- Petschick, R., Kuhn, G., and Gingele, F.: Clay mineral distribution in surface sediments of the South Atlantic: Sources, transport, and relation to oceanography, *Mar. Geol.*, 130, 203–230, 1996.
- Peyron, O., Goring, S., Dormoy, I., Kothoff, U., Porss, J., de Beaulieu, J. L., La Mantia, T., Schneider, R., and Magny, M.: Holocene seasonality changes in the central Mediterranean region reconstructed from the pollen sequences of Lake Accesa (Italy) and Tenaghi Philippon (Greece), *Holocene*, 21, 131–146, 2011.
- Pickard, G. L. and Emery, W. J.: *Descriptive Physical Oceanography – An Introduction*, Pergamon Press, San Diego, 249 pp., 1982.
- Pinsak, A. P. and Murray, H. H.: Regional clay mineral patterns in the Gulf of Mexico, *Clays and Clay Minerals*, 7, 162–177, 1960.
- Pittalwala, I. and Hameed, I.: Simulation of the North Atlantic Oscillation in a general circulation model, *Geophys. Res. Lett.*, 18, 841–844, 1991.
- Prospero, J. M.: Arid regions as sources of minerals aerosols in the marine atmosphere, *Geol. S. Am. S.*, 186, 71–86, 1981.
- Prospero, J. M., Ginoux, P., Torres, O., Nicholson, S. E. and Gill, T. E.: Environmental characterization of global sources of atmospheric soil dust identified with the nimbus 7 total ozone mapping spectrometer (TOMS) absorbing aerosol product, *Rev. Geophys.*, 40, 1002, doi:10.1029/2000RG000095, 2002.
- Pye, K.: *Aeolian dust and dust deposits*, Academic Press, New York, London, 334 pp., 1987.
- Pye, K.: Aeolian dust transport and deposition over Crete and adjacent parts of the Mediterranean Sea, *Earth Surf. Proc. Land.*, 17, 271–288, 1992.
- Pye, K. and Tsoar, H.: The mechanics and geological implications of dust transport and deposition in deserts with particular reference to loess formation and dune sand diagenesis in the northern Negev, Israel, in: *Desert Sediments Ancient and Modern*, edited by: Frostick, L. E. and Reid, I., *Geol. Soc. London, Spec. Publ.*, 35, 139–156, 1987.
- Ramrath, A., Sadori, L., and Negendank, J. F. W.: Sediments from Lago di Mezzano, central Italy: A record of Lateglacial/Holocene climatic variations and anthropogenic impact, *Holocene*, 10, 87–95, 2000.
- Regaya, K.: Les accumulations calcaires dans les limons de Matmata de la région de Gabès en Tunisie, *B. Soc. Geol. Fr.*, 37, 387–398, 1984.
- Regaya, K.: Les croutes calcaires de Tunisie orientale aux environs de Sousse, Signification climatique et historique, *B. Soc. Geol. Fr.*, 45, 99–107, 1992.
- Revel, M., Ducassou, E., Grousset, F. E., Bernasconi, S. M., Migeon, S., Revillon, S., Mascle, J., Murat, A., Zaragosi, S., and Bosch, D.: 100,000 years of African monsoon variability recorded in sediments of the Nile margin, *Quaternary Sci. Rev.*, 29, 1342–1362, 2010.
- Richter, T. O., van der Gaast, S., Koster, B., Vaasrs, A., Gieles, R., de Stigter, H. C., de Haas, H., van Weering, T. C. E., and Rothwell, R. G.: The Avaatech XRF Core Scanner: technical description and applications to NE Atlantic sediments, in: *Techniques in Sediment Core Analysis*, edited by: Rothwell, R. G., *New Geological Society of London, Special Publication*, 267, 39–50, 2006.

- Robert, C., Gauthier, A., and Chamley, H.: Origine autochtone et allochtone des argiles récentes de haute altitude en Corse, *Géologie de la Méditerranée*, XI, 243–253, 1984.
- Roberts, N., Jones, M. D., Benkaddour, A., Eastwood, W. J., Filippi, M. L., Frogley, M. R., Lamb, H. F., Leng, M. J., Reed, J. M., Stein, M., Stevens, L., Valero-Garcés, B., and Zanchetta, G.: Stable isotope records of Late Quaternary climate and hydrology from Mediterranean lakes: the ISOMED synthesis, *Quaternary Sci. Rev.*, 27, 2426–2441, 2008.
- Roberts, N., Brayshaw, D., Kuzucuoglu, C., Perez, R., and Sadori, L.: The mid-Holocene climatic transition in the Mediterranean: causes and consequences, *Holocene*, 21, 3–14, 2011.
- Rodrigo-Gamiz, M., Martínez-Ruiz, F., Jiménez-Espejo, F. J., Gallego-Torres, D., Nieto-Moreno, V., Romero, O., and Ariztegui, D.: Impact of climate variability in the western Mediterranean during the last 20,000 years: oceanic and atmospheric responses, *Quaternary Sci. Rev.*, 30, 2018–2034, 2011.
- Rodriguez, S., Querol, X., Alastuey, A., Kallos, G., and Kakaliagou, O.: Saharan dust contributions to PM₁₀ and TSP levels in Southern and Eastern Spain, *Atmos. Environ.*, 35, 2433–2447, 2001.
- Rohdenburg, H.: Landscape ecology – Geomorphology, Catena, Reiskirchen, 177 pp., 1989.
- Rohling, E. J.: Review and new aspects concerning the formation of eastern Mediterranean sapropels, *Mar. Geol.*, 122, 1–28, 1994.
- Rosignol-Strick, M., Nesteroff, W., Olive, P., and Vergnaud-Grazzini, C.: After the deluge: Mediterranean stagnation and sapropel formation, *Nature*, 295, 105–110, 1982.
- Rouis-Zargouni, I.: Evolution paléoclimatique et paléohydrologique de la Méditerranée Occidentale au cours des derniers 30 000 ans; contribution des dinokystes et des foraminifères planctoniques, Ph.D. Thesis, Bordeaux I-Sfax Univ., 210 pp., 2010.
- Rouis-Zargouni, I., Turon, J.-L., Londeix, L., Essallami, L., Kallel, N., and Sicre, M.-A.: Environmental and climatic changes in the central Mediterranean Sea (Siculo-Tunisian Strait) during the last 30 ka based on dinoflagellate cyst and planktonic foraminifera assemblages, *Paleogeogr. Palaeocl.*, 285, 17–29, 2010.
- Rouis-Zargouni, I., Turon, J. L., Londeix, L., Kallel, N., and Essallami, L.: The last glacial-interglacial transition and dinoflagellate cysts in the western Mediterranean sea, *C. R. Geoscience*, 344, 99–109, doi:10.1016/j.crte.2012.01.002, 2012.
- Sadori, L., Giraudi, C., Pettitti, P., and Ramrath, A.: Human impact at Lago di Mezzano (central Italy) during the Bronze Age: a multidisciplinary approach, *Quatern. Int.*, 113, 5–17, 2004.
- Sadori, L., Zanchetta, G. and Giardini, M.: Last Glacial to Holocene palaeoenvironmental evolution at Lago di Pergusa (Sicily, Southern Italy) as inferred by pollen, microcharcoal, and stable isotopes, *Quatern. Int.*, 181, 4–14, 2008.
- Sadori, L., Jahns, S., and Peyron, O.: Mid-Holocene vegetation history of the central Mediterranean, *Holocene*, 21, 117–129, 2011.
- Salot, A.: The Mediterranean Sea, *The Handbook of Environmental Chemistry*, v. 5, Water Pollution, Springer-Verlag, Berlin, 408 pp., 2005.
- Sánchez-Goñi, M. F., Cacho, I., Turon, J. L., Guiot, J., Sierro, F. J., Peyrouquet, J.-P., Grimalt, J. O., and Shackleton, N.: Synchronicity between marine and terrestrial responses to millennial scale climatic variability during the last glacial period in the Mediterranean region, *Clim. Dynam.*, 19, 95–105, 2002.
- Schilman, B., Bar-Matthews, M., Almogi-Labin, A., and Luz, B.: Global climate instability reflected by Eastern Mediterranean marine records during the late Holocene, *Paleogeogr. Palaeocl.*, 176, 157–176, 2001.
- Schneider, R. R., Price, B., Müller, P. J., Kroon, D., and Alexander, I.: Monsoon related variations in Zaire (Congo) sediment load and influence of fluvial silicate supply on marine productivity in the east equatorial Atlantic during the last 200,000 years, *Paleoceanography*, 12, 463–481, 1997.
- Schütz, L. and Seibert, M.: Mineral aerosols and source identification, *J. Aerosol Sci.*, 18, 1–10, 1987.
- Siani, G., Paterne, M., Michel, E., Sulpizio, R., Sbrana, A., Arnold, M., and Haddad, G.: Mediterranean sea-surface radiocarbon reservoir age changes since the last glacial maximum, *Science*, 294, 1917–1920, 2001.
- Siani, G., Sulpizio, R., Paterne, M., and Sbrana, A.: Tephrostratigraphy study for the last 18,000 ¹⁴C years in a deep-sea sediment sequence for the South Adriatic, *Quaternary Sci. Rev.*, 23, 2485–2500, 2004.
- Singer, A. and Galan, E.: Palygorskite-sepiolite: occurrences, genesis and uses, *Developments in Sedimentology*, Elsevier, Amsterdam, 352 pp., 1984.
- Sionneau, T., Bout-Roumazeilles, V., Biscaye, P. E., Van Vliet-Lanoe, B., and Bory, A.: Clay mineral distributions in and around the Mississippi River watershed and Northern Gulf of Mexico: sources and transport patterns, *Quaternary Sci. Rev.*, 27, 1740–1751, 2008.
- Sionneau, T., Bout-Roumazeilles, V., Flower, B. P., Bory, A., Tribouillard, N., Kissel, C., Van Vliet-Lanoe, B., and Montero Serrano, J. C.: Provenance of freshwater pulses in the Gulf of Mexico during the last deglaciation, *Quaternary Res.*, 74, 235–245, 2010.
- Skonieczny, C., Bory, A., Bout-Roumazeilles, V., Abouchami, W., Galer, S. J. G., Crosta, X., Stuut, J.-B., Meyer, I., Chiapello, I., Podvin, T., Chatenet, B., Diallo, A., and Ndiaye, T.: The 7–13 March 2006 major Saharan outbreak: Multi-proxy characterization of mineral dust deposited on the West African margin, *J. Geophys. Res.*, 116, D18210, doi:10.1029/2011JD016173, 2011.
- Skonieczny, C., Bory, A., Bout-Roumazeilles, V., Abouchami, W., Galer, S. J. G., Crosta, X., Diallo, A., Ndiaye, T.: A three-year time series of mineral dust deposits on the West African margin: Sedimentological and geochemical signatures and implications for interpretation of marine paleo-dust records, *Earth Planet. Sc. Lett.*, 364, 145–156, 2013.
- Stanley, D. J. and Wingerath, J. G.: Clay mineral distributions to interpret Nile cell provenance and dispersal, I. Lower River Nile to delta sector, *J. Coast. Res.*, 12, 911–929, 1996.
- Stanley, D. J., Warne, A. G., Davis, H. R., Bernasconi, M. P., and Chen, Z.: Nile Delta, *Nat. Geogr. Res. Expl.*, 8, 22–51, 1992.
- Stevens, L. R., Wright Jr., H. E., and Ito, E.: Proposed changes in seasonality of climate during the late-glacial and Holocene at Lake Zeribar, Iran, *Holocene*, 11, 747–756, 2001.
- Stevenson, A. C., Phethean, S. J., and Robinson, J. E.: The palaeosalinity and vegetational history of Garaet el Ichkeul, northwest Tunisia, *Holocene*, 3, 201–210, 1993.
- Stumpf, R., Frank, M., Schönfeld, J., and Haley, B. A.: Late Quaternary variability of Mediterranean Outflow Water from radiogenic Nd and Pb isotopes, *Quaternary Sci. Rev.*, 29, 2462–2572, doi:10.1016/j.quascirev.2010.06.021, 2010.

- Stumpf, R., Frank, M., Schönfeld, J., and Haley, B. A.: Climatically driven changes in sediment supply on the SW Iberian shelf since the last Glacial Maximum, *Earth Planet. Sc. Lett.*, 312, 80–90, 2011.
- Thiry, M.: Paleoclimatic interpretation of clay minerals in marine deposits: an outlook from the continental origin, *Earth Sci. Rev.*, 49, 201–221, 2000.
- Tomadin, L.: Sedimentary fluxes and different dispersion mechanisms of the clay sediments in the Adriatic Basin, *Rend. Fis. Acc. Lincei*, 9, 161–174, 2000.
- Tomadin, L. and Lenaz, R.: Eolian dust over the Mediterranean and their contribution to the present sedimentation, in: *Paleoclimatology and Paleometeorology: Modern and past patterns of global atmospheric transport*, edited by: Leinen, M. and Sarnthein, M., Kluwer Academic, Dordrecht, 282, 267–281, 1989.
- Torres-Padrón, M. E., Gelado-Caballero, M. D., Collado-Sánchez, C., Siruela-Matos, V. F., Cardona-Castellano, P. J., and Hernández-Brito, J. J.: Variability of dust inputs to the CANIGO zone, *Deep-Sea Res. Pt. II*, 49, 3455–3464, 2002.
- Tribovillard, N., Bout-Roumzeilles, V., Algeo, T., Lyons, T. W., Sionneau, T., Montero-Serrano, J. C., Riboulleau, A., and Baudin, F.: Paleodepositional conditions in the Orca Basin as inferred from organic matter and trace metal contents, *Mar. Geol.*, 254, 62–72, 2008.
- Tzedakis, P. C.: Seven ambiguities in the Mediterranean palaeoenvironmental narrative, *Quaternary Sci. Rev.*, 26, 2042–2066, 2007.
- Ukstins, I. A., Renne, P. R., Wolfenden, E., Baker, J. A., Ayalew, D., and Menzeis, M.: Matching conjugate volcanic rifted margins: $^{40}\text{Ar}/^{39}\text{Ar}$ chrono-stratigraphy of pre- and syn-rift bimodal flood volcanism in Ethiopia and Yemen, *Earth Planet. Sc. Lett.*, 198, 289–306, 2002.
- Vannièrè, B., Power, M. J., Roberts, N., Tinner, W., Carrion, J., Magny, M., and Bartlein, P.: Circum-Mediterranean fire activity and climate changes during the mid Holocene environmental transition (8500–2500 cal yr BP), *Holocene*, 21, 53–73, 2011.
- Venkatarathnam, K. and Ryan, W. B. F.: Dispersal patterns of clay minerals in the sediments of the Eastern Mediterranean Sea, *Mar. Geol.*, 11, 261–282, 1971.
- Verrecchia, E. P. and Le Coustumer, M.-N.: Occurrence and genesis of palygorskite and associated clay minerals in a Pleistocene calcrete complex, Sde Boqer, Negev Desert, Israel, *Clay Min.*, 31, 183–202, 1996.
- Washington, R. and Todd, M. C.: Atmospheric controls on mineral dust emission from the Bodélé depression, Chad: the role of the low level jet, *Geophys. Res. Lett.*, 32, L17701, doi:10.1029/2005GL023597, 2005.
- Weldeab, S., Siebel, W., Wehausen, R., Emeis, K.-C., Schmiedl, G., and Hemleben, C.: Late Pleistocene sedimentation in the Western Mediterranean Sea: implications for productivity changes and climatic conditions in the catchment areas, *Paleogeogr. Palaeocl.*, 190, 121–137, 2003.
- Wüst, G.: On the vertical circulation on the Mediterranean Sea, *J. Geophys. Res.*, 66, 3261–3271, 1961.
- Yarincik, K. M., Murray, R. W., and Peterson, L. C.: Climatically sensitive eolian and hemipelagic deposition in the Cariaco Basin, Venezuela, over the past 578,000 years: Results from Al/Ti and K/Al, *Paleoceanography*, 15, 210–228, 2000.
- Zahar, Y., Ghorbel, A., and Albergel, J.: Impacts of large dams on downstream flow conditions of rivers: aggradation and reduction of the Medjerda channel capacity downstream of the Sidi Salem dam (Tunisia), *J. Hydrol.*, 351, 318–330, 2008.
- Zanchetta, G., Drysdale, R. N., Hellstrom, J. C., Fallick, A. E., Isola, I., Gagan, M. K., and Pareschi, M. T.: Enhanced rainfall in the Western Mediterranean during deposition of sapropel S1: Stalagmite evidence from Corchia cave (Central Italy), *Quaternary Sci. Rev.*, 26, 279–286, 2007.
- Zanchetta, G., Sulpizio, R., Roberts, N., Cioni, R., Eastwood, W. J., Siani, G., Caon, B., Paterne, M., and Santacroce, R.: Tephrostratigraphy, chronology and climatic events of the Mediterranean basin during the Holocene: An overview, *Holocene*, 21, 33–52, 2011.
- Zielhofer, C., Faust, D., Diaz del Olmo, F., and Baena, R.: Sedimentation and soil formation phases in the Ghardimaou Basin (Northern Tunisia) during the Holocene, *Quatern. Int.*, 93–94, 109–125, 2002.
- Zielhofer, C., Faust, D., Baena, R., Diaz del Olmo, F., Kadereit, A., Moldenhauer, K.-M., and Porras, A.: Centennial-scale late Pleistocene to mid-Holocene synthetic profile of the Medjerda floodplain (Northern Tunisia), *Holocene*, 14, 851–861, 2004.
- Zielhofer, C., Faust, D., and Linstaedter, J.: Late Pleistocene and Holocene alluvial archives in the Southwestern Mediterranean: changes in fluvial dynamics and past human response, *Quatern. Int.*, 181, 39–54, doi:10.1016/j.quaint.2007.09.016, 2008.
- Zielhofer, C., Bussmann, J., Ibouhouten, H., and Fenech, K.: Flood frequencies reveal Holocene rapid climate changes (Lower Moulouya River, northeastern Morocco), *Quatern. Int.*, 25, 700–714, doi:10.1002/jqs.1347, 2010.