

In vitro photophysical and photobiological properties of Ce6-based dendrimer nanoparticles.

Estelle Bastien, Raphaël Schneider, Lina Bezdetnaya, Beate Röder, Steffen Hackbarth, Annegret Preuss, Jordane Jasniewski, Vadim Reshetov, Henri-Pierre Lassalle

▶ To cite this version:

Estelle Bastien, Raphaël Schneider, Lina Bezdetnaya, Beate Röder, Steffen Hackbarth, et al.. In vitro photophysical and photobiological properties of Ce6-based dendrimer nanoparticles.. 15th Congress of the European Society for Photobiology, ESP 2013, Sep 2013, Liège, Belgium. pp.CDROM. hal-00866009

HAL Id: hal-00866009

https://hal.science/hal-00866009

Submitted on 25 Sep 2013

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

In vitro photophysical and photobiological properties of Ce6-based dendrimer nanoparticles

Estelle Bastien^{1,2}, Raphael Schneider³, Lina Bezdetnaya^{1,2}, Beate Röder⁴, Steffen Hackbarth⁴, Annegret Preuß⁴, Jordane Jasniewski⁵, Vadzim Reshetov¹, Henri-Pierre Lassalle^{1,2}

Context: The photosensitizers vectorization enables better tumor localization and therefore a better tumoricidal PDT efficacy. Dendrimers are hyperbranched macromolecules that are considered as an attractive class of biocompatible nanovectors for drug targeting in tissues. In this study we report a novel PDT drug-carrier system, composed of chlorin e6 grafted to the periphery groups of poly(amido amine) dendrimers (PAMAM).

Materials & Methods: The size and zeta potential of our nanoparticles were determined by dynamic light scattering (DLS). The nanoparticle singlet oxygen generation in pharynx carcinoma (FaDu) cells was assessed by luminescence at 1268 nm. Phototoxicity was evaluated by MTT assay. Uptake was assessed by chemical extraction associated with fluorescence spectroscopy measurements. Finally, the uptake mechanism was determined by the use of endocytosis specific inhibitors.

Results: Ce6-dendrimers nanoparticles displayed a 40 times greater uptake along with a 25 fold higher photocytotoxic activity in FaDu cells compared to free Ce6. Also, a much longer singlet oxygen lifetime was demonstrated for Ce6-dendrimers compared to Ce6. Cellular incorporation of nanoparticles was completely inhibited at 4°C, demonstrating that uptake was mediated via endocytosis. Further investigations using specific inhibitors revealed that incorporation was partially mediated by macropinocytosis pathway.

Conclusion: Dendrimeric based nanoparticles showed a good photodynamic efficacy towards cancer cells *in vitro*, suggesting them as a promising vehicle for PDT. In the future we shall test dendrimer-based nanoparticles *in vivo*, in pre-clinical models.

¹Centre de recherché en Automatique de Nancy, Université de Lorraine, CNRS UMR 7039, Vandœuvre-lès-Nancy, France

²Institut de Cancérologie de Lorraine, Vandœuvre-lès-Nancy, France

³Laboratoire Réactions et Génie des Procédés, Ecole National Supérieure des Industries Chimiques, Université de Lorraine, Nancy, France

⁴Institut of physic, Humboldt University, Berlin, Germany

⁵Laboratoire d'ingénierie des biomolécules, Université de Lorraine, Vandœuvre-lès-Nancy, France