



HAL
open science

Hawking effect for a toy model of interacting fermions

Patrick Bouvier, Christian Gérard

► **To cite this version:**

Patrick Bouvier, Christian Gérard. Hawking effect for a toy model of interacting fermions. 2013. ⟨hal-00869929⟩

HAL Id: hal-00869929

<https://hal.science/hal-00869929v1>

Preprint submitted on 4 Oct 2013

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

HAWKING EFFECT FOR A TOY MODEL OF INTERACTING FERMIONS

P. BOUVIER AND C. GÉRARD

ABSTRACT. We consider a toy model of interacting Dirac fermions in a $1 + 1$ dimensional space time describing the exterior of a star collapsing to a black-hole. In this situation we give a rigorous proof of the *Hawking effect*, namely that under the associated quantum evolution, an initial vacuum state will converge when $t \rightarrow +\infty$ to a thermal state at Hawking temperature. We establish this result both for observables falling into the blackhole along null characteristics, and for static observables. We also consider the case of an interaction localized near the star boundary, obtaining similar results. We hence extend to an interacting model previous results of Bachelot and Melnyk, obtained for free Dirac fields.

1. INTRODUCTION

1.1. Introduction. The *Hawking effect*, see Hawking [Ha], predicts that in a space-time describing the collapse of a spherically symmetric star to a Schwarzschild black hole, an initial Boulware vacuum state will become an Unruh state at the future horizon: a static observer at infinity sees the Unruh state as a thermal state at Hawking temperature.

Despite the vast physical literature on the Hawking effect, there are few mathematically rigorous justifications of the Hawking effect. Dimock and Kay [DK1, DK2] gave a construction of the Unruh state in the Schwarzschild space-time and on its Kruskal extension, using scattering theory for Klein-Gordon fields.

The first mathematical proof of the Hawking effect, in the original setting of Hawking, is due to Bachelot [Ba1]. Bachelot considered a linear Klein-Gordon field in the exterior of a spherically symmetric star, collapsing to a Schwarzschild black hole. This result was extended to linear Dirac fields in the same situation, first by Bachelot [Ba2], and then by Melnyk [Me]. The only proof to date in a non-spherically symmetric situation is due to Häfner [H], who gave a rigorous proof of the Hawking effect for Dirac fields for a star collapsing to a Kerr black hole.

The common theme of all the above mentioned results is that they deal with *linear* quantum fields: the time evolution of observables is implemented by a group of linear (symplectic or unitary) transformations on the phase space, and all the states are quasi-free.

This means that the problem can be reduced to a question about linear partial differential equations, with boundary conditions on the star boundary. The Hawking effect emerges from the fact that the star boundary becomes asymptotically characteristic for large times. This leads to an exponentially fast concentration of Klein-Gordon or Dirac wave packets reflected by the star, which ultimately implies the Hawking effect.

Date: October 4, 2013.

2010 Mathematics Subject Classification. 81T10, 81T20.

Key words and phrases. Hawking effect, interacting fermions, $1 - d$ Dirac equations.

In this paper we investigate the Hawking effect for a toy model of *interacting Dirac fermions* in 1 + 1 space-time dimensions. A mathematical discussion of interacting quantum fields is of course difficult, because there are few rigorous constructions of interacting quantum fields, even on Minkowski space.

For Klein-Gordon fields, there are the well-known constructions of the $P(\varphi)_2$ and φ_3^4 models due to Glimm and Jaffe, which were the main successes of the constructive program from the seventies. We are not aware of any similar construction on a space-time which describes the exterior of a collapsing star, even when the interaction contains an ultraviolet and space cutoff.

For Dirac fields, the situation looks better, since fermionic fields are bounded, which in some situations allows to construct the interacting dynamics in a purely algebraic setting, independently of the choice of a representation. This is particularly convenient in the situation that we consider, since, even for free Dirac fields, two Fock representations in the exterior of the star at different times are inequivalent.

1.2. A toy model. To concentrate on the possibly new features introduced by the non-linear interactions and to keep the situation simple and manageable, we restrict ourselves to a toy model of Dirac fermions in 1 + 1 space-time dimensions:

we consider only 2 components spinors, and the effect of the metric is modeled by a vector potential. Note that if we forget about the non-linear interaction, our model is essentially identical to the one considered by Bachelot in [Ba2], after introduction of polar coordinates and suitable spin spherical harmonics.

Let us now briefly describe the model: the space-time is the region:

$$\mathcal{M} = \{(t, x) \in \mathbb{R}^2 : x > z(t)\},$$

where $x = z(t)$ is the star boundary. We assume that $z(t) \equiv z(0)$ for $t \leq 0$, i.e. the star is stationary in the past, the collapse starting at $t = 0$. As in [Ba2] we assume that $z(t) \sim -t - Ae^{-2\kappa t}$ for $t \rightarrow +\infty$, i.e. the star boundary becomes asymptotically characteristic for large positive times.

The Dirac fields are two-components spinors $\psi(t, x) \in \mathbb{C}^2$, solving (in absence of interaction) the Dirac equation:

$$(1.1) \quad \begin{cases} \partial_t \psi(t, x) + L \partial_x \psi(t, x) + iV(x)\psi(t, x) = 0, & \text{in } z > z(t), \\ \psi_1(t, z(t)) = \lambda(t)\psi_2(t, z(t)), \end{cases}$$

where $L = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$ and $V(x) = V(x)^* \in M_2(\mathbb{C})$ is a matrix-valued potential representing the influence of the metric, with

$$V(x) \rightarrow 0 \text{ at } -\infty, \quad V(x) \rightarrow m\Gamma \text{ at } +\infty,$$

$m > 0$ is the mass of the field, and $\Gamma \in M_2(\mathbb{C})$ satisfies

$$\Gamma = \Gamma^*, \quad \Gamma^2 = \mathbf{1}, \quad \Gamma L + L\Gamma = 0.$$

The reflection coefficient $\lambda(t)$ equals $\left(\frac{1+z(t)}{1-z(t)}\right)^{1/2}$, so that the L^2 norm

$$\int_{z(t)}^{+\infty} \|\psi(t, x)\|_{\mathbb{C}^2}^2 dx$$

is conserved. This implies that if $\mathfrak{h}_t := L^2(]z(t), +\infty[; \mathbb{C}^2)$, the evolution group $u^V(s, t) : \mathfrak{h}_t \rightarrow \mathfrak{h}_s$ (see Subsect. 2.2) associated to (1.1) is unitary, and hence generates a fermionic dynamics

$$\tau^V(s, t) : \text{CAR}(\mathfrak{h}_t) \rightarrow \text{CAR}(\mathfrak{h}_s),$$

where $\text{CAR}(\mathfrak{h})$ is the CAR C^* -algebra associated to a Hilbert space \mathfrak{h} .

The self-interaction of the Dirac field is described by a perturbation of the form

$$I = (\psi^*(g)M\psi(g))^n,$$

where $n \geq 2$, $M \in M_2(\mathbb{C})$ is a selfadjoint matrix, $g \in L^2_{\text{comp}}(\mathbb{R})$ is a compactly supported function. The associated interacting Dirac fields $\psi^{\text{int}}(t, x)$ formally solve the following non-linear Dirac equation:

$$(1.2) \quad \begin{cases} \partial_t \psi^{\text{int}}(t, x) + L\partial_x \psi^{\text{int}}(t, x) + iV(x)\psi^{\text{int}}(t, x) \\ - \text{in}(\overline{\psi^{\text{int}}(t, g)}|M\psi^{\text{int}}(t, g))_{\mathbb{C}^2}^{n-1} M\psi^{\text{int}}(t, g)g(x) = 0, \text{ in } x > z(t) \\ \psi_1(t, z(t)) = \lambda(t)\psi_2(t, z(t)), \end{cases}$$

where $\psi^{\text{int}}(t, g) := \int \psi^{\text{int}}(t, x)\bar{g}(x)dx \in \mathbb{C}^2$. The properties of the interaction which are essential for our analysis are the following:

- (1) I is bounded, which allows for a purely algebraic construction of the interacting dynamics $\tau^{V, \text{int}}(s, t)$;
- (2) I is even, which is the standard assumption needed to ensure locality,
- (3) I is localized in a (space) compact region.

1.3. Results. Let us now describe the results of the paper.

The first step is to construct interacting Dirac fields, i.e. to quantize the non-linear Dirac equation (1.2).

Since we deal with fermions, the interaction term I above is bounded, and one can work in a purely algebraic setting: one can introduce C^* -algebras $\mathfrak{A}_t = \text{CAR}(\mathfrak{h}_t)$ of observables at time t , and it is easy to construct the *interacting dynamics* $\tau^{V, \text{int}}(s, t)$ (see Sect. 4), which is a two parameter group of $*$ -isomorphisms from \mathfrak{A}_t to \mathfrak{A}_s describing the time evolution.

We investigate the Hawking effect in three different situations.

1.3.1. Hawking effect I. In the first situation we take an observable at time t , localized near the star boundary $x = z(t)$, i.e. of the form $\alpha^t(A)$ for some $A \in \mathfrak{A}_0$, where α^t is the group of *left space translations*. In terms of interacting space-time fields ψ^{int} , a typical observable would be $\psi^{\text{int}}(t, x - t)$, i.e. a field falling into the black hole along null characteristics. This is the analog for interacting fields of the situation in [Ba2].

To evaluate the time-evolved state at time t acting on $\alpha^t(A)$ we have to evolve $\alpha^t(A)$ back to time 0, which yields

$$\omega_{0, \text{vac}}^V(\tau^{V, \text{int}}(0, t) \circ \alpha^t(A)),$$

where $\omega_{0, \text{vac}}^V$ is the vacuum state at time $t = 0$, $\tau^{V, \text{int}}(t, 0)$ is the interacting dynamics. Our goal is to compute the limit of the above quantity when $t \rightarrow +\infty$. We prove in Thm. 5.6 that the limit

$$(1.3) \quad \lim_{t \rightarrow +\infty} \omega_{0, \text{vac}}^V(\tau^{V, \text{int}}(0, t) \circ \alpha^t(A)) = \omega_{\text{H, I}}(A) \text{ exists,}$$

for any A in the C^* -algebra \mathfrak{A}_0 . Let us describe the limiting state $\omega_{\text{H, I}}$, which is close to the one obtained by Bachelot in [Ba2]:

the algebra \mathfrak{A}_0 splits into the (\mathbb{Z}_2 -graded) tensor product $\mathfrak{A}_0^{\text{left}} \widehat{\otimes} \mathfrak{A}_0^{\text{right}}$ (see Subsect. A.2) of the *left/right moving observables*.

The limit state $\omega_{\text{H, I}}$ acts on right moving observables as a vacuum state (composed with an appropriate wave morphism), while on left moving observables it acts as the thermal state $\omega_{\infty, \beta}^0$ at inverse Hawking temperature $\beta = 2\pi\kappa^{-1}$, for the eternal black hole without interaction.

We also prove a similar result if the initial state $\omega_{0, \text{vac}}^V$ is replaced by another state $\tilde{\omega}$ which is *even* and belongs to the *folium* of $\omega_{0, \text{vac}}^V$ (see Corollary 5.8). As example

of such a state, one can choose an *interacting vacuum state*, whose existence is shown in Subsect. 5.5.

The first situation is graphically summarized in Figure 1 below: the grey region is the support of the non-linear self-interaction. The curve $x = z(t)$ is the star boundary. The dashed lines are the (backwards) characteristics for the Dirac equation, starting from the support of an observable at time T : left moving characteristics are reflected on the star boundary and asymptotically concentrated when $T \rightarrow +\infty$.




FIGURE 1. Hawking effect I

1.3.2. *Hawking effect II.* In the second situation the observable A at time t is localized near the origin. In terms of space-time fields, a typical example would be simply $\psi^{\text{int}}(t, x)$. This is the analog for interacting fields of the situation considered by Melnik in [Me].

The situation is now more complicated: one has to be sure that the observable A , under backwards propagation, will split into left and right moving parts. One way to formulate this property is to introduce the (future) *wave morphism* $\gamma_{\infty}^{\text{int}}$ between the dynamics on the eternal black hole $\tau_{\infty}^{V, \text{int}}$ and τ_{∞}^V (see Thm. 6.5). Then we have to require that A belongs to $\gamma_{\infty}^{\text{int}}\mathfrak{A}_{\infty}$. Observables outside this $*$ -subalgebra will not see the Hawking effect.

It is easier to formulate our result if we assume the *asymptotic completeness* of $\gamma_{\infty}^{\text{int}}$, i.e. that $\gamma_{\infty}^{\text{int}}\mathfrak{A}_{\infty} = \mathfrak{A}_{\infty}$: then we prove in Thm. 6.18 that the limit

$$(1.4) \quad \lim_{t \rightarrow +\infty} \omega_{0, \text{vac}}^V(\tau^{V, \text{int}}(0, t)(A)) = \omega_{\text{H}, \text{II}}(A) \text{ exists,}$$

for A a *local element* of \mathfrak{A}_{∞} (i.e. $A \in \mathfrak{A}_J$ for some interval $J \in \mathbb{R}$).

Without assuming asymptotic completeness, we have to restrict ourselves to observables $A \in \gamma_{\infty}^{\text{int}}\mathfrak{A}_{\infty}$. Such observables do not necessarily belong to \mathfrak{A}_t for t large, i.e. the expression $\tau^{V, \text{int}}(s, t)(A)$ may have no meaning. Therefore we replace A by $E_t A \in \mathfrak{A}_t$, where E_t is the natural projection $\mathfrak{A}_{\infty} \rightarrow \mathfrak{A}_t$ (see Remark 6.7).

Let us now describe the limiting state $\omega_{\text{H}, \text{II}}$. Again the algebra \mathfrak{A}_{∞} splits into a tensor product $\text{CAR}(P^l \mathfrak{h}_{\infty}) \widehat{\otimes} \text{CAR}(P^r \mathfrak{h}_{\infty})$ of left/right moving observables (see Subsect. 6.1). In this case elements of $\text{CAR}(P^{l/r} \mathfrak{h}_{\infty})$ are left/right moving only asymptotically for large times.

On right moving observables the limit state $\omega_{\text{H}, \text{II}}$ acts again as a vacuum state, composed with a wave morphism. On left moving observables it acts as the thermal state $\omega_{\infty, \beta}^V$. In contrast to case I, the potential term V is present in the thermal state.

A similar result holds if we replace the initial state by another even, state $\tilde{\omega}$ belonging to the folium of $\omega_{0,\text{vac}}^V$, see Corollary 6.19. However we have now to assume that $\tilde{\omega}$ is invariant under the interacting *stationary* dynamics, $\tau_0^{V,\text{int}}$, describing the interacting Dirac field in the past.

Fig. 2 summarizes the second situation, with the same conventions as in Fig. 1: note that left moving characteristics starting at time T from close to the origin, reach the star boundary at time close to $T/2$: after time $T/2$ the situation for left moving observables is similar to case I.




FIGURE 2. Hawking effect II

1.3.3. *Hawking effect III.* In the two previous situations, the interaction region is far away from the star boundary: the effect of the self-interaction is decoupled from the effect of the asymptotically characteristic boundary, which is essential in the Hawking effect.

For an initial observable starting at time T close to the star boundary $z = z(T)$, the Hawking effect (in the free situation), is essentially due to what happens between the times T and $T - 1$, i.e. to the reflection on the asymptotically characteristic star boundary. Therefore we consider a third situation where the interaction is localized near the star boundary for times $t \in [T - 1, T]$. We consider the following time-dependent interaction

$$I_T(t) = \mathbb{1}_{[T-1, T]}(t) \alpha^t(I),$$

which is at time t localized near the star boundary $x = z(t)$, and vanishes for $t \notin [T - 1, T]$. We denote by $\tilde{\tau}_T^{V,\text{int}}(s, t)$ the dynamics obtained as before by adding to the free dynamics $\tau^V(s, t)$ the time-dependent interaction $I_T(t)$. We obtain a dynamics depending on the parameter T , which differs from the free dynamics $\tau^V(s, t)$ only for $T - 1 \leq s \leq t \leq T$. We show in Thm. 7.7 that the limit

$$\lim_{T \rightarrow \infty} \omega_{0,\text{vac}}^V(\tilde{\tau}_T^{V,\text{int}}(0, T) \circ \alpha^t(A)) = \omega_{\text{H,III}}(A) \text{ exists}$$

for $A \in \mathfrak{A}_0$. The limiting state $\omega_{\text{H,III}}$ is actually quite explicit, being the pullback of the (free) limiting state $\omega_{\text{H}}^{\text{free}}$ obtained by Bachelot in [Ba2] by a simple effective interacting dynamics $\hat{\tau}_{\infty}^{0,\text{int}}(0, 1)$. The dynamics $\hat{\tau}_{\infty}^{0,\text{int}}(s, t)$ describes the combined effect of interaction and reflection on the star boundary between times $T + t$ and $T + s$, in the limit $T \rightarrow +\infty$. The situation is summarized in Fig. 3 below.




FIGURE 3. Hawking effect III

1.4. Plan of the paper. Let us now briefly describe the plan of our paper. In Sect. 2 we describe our geometrical setup and recall some results of [Ba2] about the linear case. The corresponding results for quantum dynamics are recalled in Sect. 3.

In Sect. 4 we construct the interacting dynamics in the algebraic, i.e. representation independent setting, by adapting standard perturbation arguments.

Sect. 5 resp. Sect. 6, Sect. 7 are devoted to the proof of the Hawking effect in the first, resp. second and third setup. In Appendix A we recall some standard facts about CAR algebras, the fermionic exponential law and perturbations of C^* -dynamics.

1.5. Notations. If \mathfrak{h}_i are Hilbert spaces $i = 1, 2$ we write $T : \mathfrak{h}_1 \xrightarrow{\sim} \mathfrak{h}_2$ if $T \in B(\mathfrak{h}_1, \mathfrak{h}_2)$ is bijective with bounded inverse. We will use the same notation if \mathfrak{A}_i are C^* -algebras and $T : \mathfrak{A}_1 \rightarrow \mathfrak{A}_2$ is a $*$ -isomorphism.

Various objects in the text, like Hilbert spaces, selfadjoint operators, C^* -algebras, $*$ -morphisms or states, are decorated with sub- and superscripts. As a rule subscripts are used to label a time or a time interval, while superscripts are used to label the various interaction terms, like 0 for no interaction, V for interaction potential, or int for the non-linear interaction. Superscripts l/r are also used to denote left/right moving observables. Subscripts vac and β in states are used to denote vacuum or thermal states, at temperature β^{-1} .

2. CLASSICAL FREE DYNAMICS

In this section we describe our setup and recall some results of [Ba2] about the free classical dynamics. We also collect some additional results which will be important in later sections.

2.1. Notations and hypotheses.

2.1.1. Collapsing star. We first recall the framework of Bachelot [Ba2], describing a star collapsing to a black hole, in a $1 + 1$ dimensional space-time.

The space-time is

$$\mathcal{M} := \{(t, x) \in \mathbb{R}^2 : x > z(t)\}$$

where the *star boundary* is $x = z(t)$ with:

$$(2.1) \quad \begin{aligned} z &\in C^2(\mathbb{R}), \\ z(t) &= z(0), \quad t \leq 0, \\ z(t) &= -t - Ae^{-2\kappa t} + \zeta(t), \quad t \geq 0, \\ -1 &\leq \dot{z}(t) \leq 0, \quad t \geq 0, \end{aligned}$$

for $A, \kappa > 0$ and

$$(2.2) \quad |\zeta(t)| + |\dot{\zeta}(t)| \leq Ce^{-4\kappa t}, \quad t \in \mathbb{R}, \quad C > 0.$$

The reflection coefficient on the star boundary is:

$$\lambda(t) = \left(\frac{1 + \dot{z}(t)}{1 - \dot{z}(t)} \right)^{1/2}.$$

Without loss of generality we can assume that $z(0) = 0$. The second condition in (2.1) means that the collapse start at $t = 0$, the star being stationary in the past.

2.1.2. *Dirac operators.* We now define various one dimensional Dirac operators. We set

$$\begin{aligned} \mathfrak{h}_t &:= L^2(]z(t), +\infty[, \mathbb{C}^2), \quad t \in \mathbb{R}, \\ \mathfrak{h}_\infty &:= L^2(\mathbb{R}, \mathbb{C}^2), \\ \mathfrak{h}_J &:= L^2(J, \mathbb{C}^2), \quad J \in \mathbb{R} \text{ interval.} \end{aligned}$$

We set

$$L := \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

and fix a matrix-valued potential (representing the influence of the metric):

$$\mathbb{R} \ni x \mapsto V(x) \in M_2(\mathbb{C}), \quad \text{with } V = V^*, \quad V \in C^1(\mathbb{R}),$$

and:

$$(2.3) \quad \begin{aligned} |V(x) - V(\infty)| + \langle x | V'(x) | &\in O(\langle x \rangle^{-1-\epsilon}), \quad x \rightarrow +\infty, \\ |V(x)| + \langle x | V'(x) | &\in O(\langle x \rangle^{-2-\epsilon}), \quad x \rightarrow -\infty, \end{aligned}$$

for some $\epsilon > 0$. We assume that

$$V_\infty = m\Gamma, \quad \Gamma \in M_2(\mathbb{C}),$$

where $m > 0$ is the mass of the field and

$$\Gamma = \Gamma^*, \quad \Gamma^2 = \mathbb{1}, \quad \Gamma L + L\Gamma = 0.$$

Let us now introduce Dirac operators; We set:

$$(2.4) \quad b_t^V := iL\partial_x - V(x) \text{ acting on } \mathfrak{h}_t,$$

with domain

$$\text{Dom } b_t^V = \{u \in H^1(]z(t), +\infty[, \mathbb{C}^2) : u_1(z(t)) = \lambda(t)u_2(z(t))\},$$

and:

$$(2.5) \quad b_\infty^V := iL\partial_x - V(x) \text{ acting on } \mathfrak{h}_\infty$$

with domain

$$\text{Dom } b_\infty^V = H^1(\mathbb{R}, \mathbb{C}^2).$$

2.2. Classical free dynamics. The classical free dynamics is generated by the following Dirac equation:

$$(2.6) \quad \begin{cases} \partial_s \psi(s, x) + L \partial_x \psi(s, x) + iV(x)\psi(s, x) = 0, & \text{in } x > z(s), \quad s \in \mathbb{R}, \\ \psi_1(s, z(s)) = \lambda(s)\psi_2(s, z(s)), & s \in \mathbb{R}, \\ \psi(t, x) = \psi(x), & \text{in } x > z(t). \end{cases}$$

In this subsection we recall some results of [Ba2], about the existence and properties of solutions of (2.6).

Definition 2.1. A $\{u(s, t)\}_{s, t \in \mathbb{R}}$ with values in $B(\mathfrak{h}_t, \mathfrak{h}_s)$ is called a (two-parameter) propagator if:

- i) $u(s, t) \in U(\mathfrak{h}_t, \mathfrak{h}_s)$,
- ii) $u(t, t) = \mathbb{1}$, $t \in \mathbb{R}$,
- iii) $u(s, t')u(t', t) = u(s, t)$, $s, t', t \in \mathbb{R}$,
- iv) $\forall (s_0, t_0) \in \mathbb{R}^2$, $\forall J \Subset]z(t_0), +\infty[$ $\forall f \in \mathfrak{h}_J$ the map $(s, t) \mapsto u(s, t)f \in \mathfrak{h}_\infty$ is continuous at (s_0, t_0) .

In the above definition we denoted by $U(\mathfrak{h}_t, \mathfrak{h}_s)$ the group of unitary operators from \mathfrak{h}_t to \mathfrak{h}_s .

Note that condition iv) is the appropriate replacement for the strong continuity of $(s, t) \mapsto u(s, t)$ in the case $\mathfrak{h}_t \equiv \mathfrak{h}$.

The following result can be found in [Ba2].

Theorem 2.2. Assume the hypotheses in Subsect. 2.1. Then there exists a unique propagator $u^V(s, t) \in B(\mathfrak{h}_t, \mathfrak{h}_s)$ such that:

$$\begin{aligned} u^V(s, t) &: \text{Dom } b_t^V \rightarrow \text{Dom } b_s^V, \quad s, t \in \mathbb{R}, \\ \partial_s u^V(s, t) &= i b_s^V u^V(s, t) \text{ on } \text{Dom } b_t^V, \\ \partial_t u^V(s, t) &= -i u^V(s, t) b_t^V \text{ on } \text{Dom } b_t^V. \end{aligned}$$

It follows that if $\psi \in \text{Dom } b_t^V$, then $\psi(s, x) = u^V(s, t)\psi(x)$ solves (2.6) in the strong sense. For the Dirac equation without boundary condition we will set accordingly:

$$u_\infty^V(s, t) := e^{i(s-t)b_\infty^V} \in U(\mathfrak{h}_\infty, \mathfrak{h}_\infty).$$

2.3. Additional results. In this subsection we collect some known results from Bachelot [Ba2] about the classical dynamics $u^V(s, t)$. For *free* Dirac fields outside of a collapsing star, they are sufficient to obtain a proof of the Hawking effect, as done in [Ba2]. In the toy model of interacting Dirac fields that we consider, they will also be important.

We first define the *left translations*:

Definition 2.3. If $f \in \mathfrak{h}_\infty$, we set $f^t(\cdot) := f(\cdot + t) \in \mathfrak{h}_\infty$.

2.3.1. Finite propagation speed. We first collect some properties of finite propagation speed for $u^V(s, t)$ and $u_\infty^V(s, t)$.

Proposition 2.4. (1) if $\text{supp } f \subset [R, +\infty[$ then $\text{supp } u^V(s, t)f \subset [R + |t - s|, +\infty[$;
(2) if $\text{supp } f \subset [a, b]$ then $\text{supp } u_\infty^V(s, t)f \subset [a - |t - s|, b + |t - s|]$;
(3) if $\text{supp } f \subset [0, R]$ then $\text{supp } u^V(s, t)f^t \subset [z(s), R - s]$ for all $s \leq t$.

Proof. the proof of (1) can be found in [Ba2]. (2) follows from classical arguments, see e.g. [CP]. (3) is shown in [Ba2, Proof of Thm. VI.5]. \square

Statement (2) of Prop. 2.4 and the uniqueness in Thm. 2.2 imply the following fact:

Proposition 2.5. *Let $J \in \mathbb{R}$ an interval. Then there exists $c \geq 0$ such that*

$$u^V(s, t)f = u_\infty^V(s, t)f, \quad \forall f \in \mathfrak{h}_J, \quad c + t/2 \leq s \leq t.$$

2.3.2. *Scattering results.* One can split \mathfrak{h}_t as direct sum:

$$\mathfrak{h}_t = \mathfrak{h}_t^l \oplus \mathfrak{h}_t^r,$$

for

$$(2.7) \quad \mathfrak{h}_t^l := \{f = (f_1, f_2) \in \mathfrak{h}_t : f_2 = 0\}, \quad \mathfrak{h}_t^r := \{f = (f_1, f_2) \in \mathfrak{h}_t : f_1 = 0\}.$$

If $f \in \mathfrak{h}_t$, we denote by $f^{l/r}$ its orthogonal projection on $\mathfrak{h}_t^{l/r}$.

If $V \equiv 0$ we easily see that:

$$(2.8) \quad u_\infty^0(0, t)f = f^t, \quad f \in \mathfrak{h}_\infty^l, \quad u_\infty^0(t, 0)f = f^t, \quad f \in \mathfrak{h}_\infty^r.$$

Proposition 2.6. *The strong limit*

$$w^r := s\text{-}\lim_{t \rightarrow +\infty} u^V(0, t)u_\infty^0(t, 0)$$

exists on \mathfrak{h}_∞^d .

Proof. See [Ba2, Prop. VI.4]. \square

Proposition 2.7.

$$w\text{-}\lim_{t \rightarrow +\infty} u^V(0, t)f^t = 0, \quad \forall f \in \mathfrak{h}_0^l.$$

Proof. We follow some arguments in [Ba2]. By density we can assume that $f \in \mathfrak{h}_0^l$ is compactly supported. We write for $0 \leq T \leq t$:

$$\begin{aligned} \|u^V(T, t)f^t - u^0(T, t)f^t\| &= \|u^0(t, T)u^V(T, t)f^t - f^t\| \\ &= \left\| \int_T^t u^0(t, \sigma)Vu^V(\sigma, t)f^t d\sigma \right\| \leq \int_T^t \|Vu^V(\sigma, t)f^t\| d\sigma. \end{aligned}$$

By Prop. 2.4 (3) we know that $\text{supp}u^V(\sigma, t)f^t \subset [z(\sigma), R - \sigma]$ for some $R \geq 0$, hence by hypothesis (2.3) we have $\|Vu^V(\sigma, t)f^t\| \in O(\langle \sigma \rangle^{-2-\epsilon})$. It follows that

$$(2.9) \quad \lim_{T \rightarrow +\infty} \sup_{T \leq t} \|u^V(T, t)f^t - u^0(T, t)f^t\| = 0.$$

Next we write

$$\begin{aligned} u^V(0, t)f^t &= u^V(0, T)u^0(T, t)f^t + u^V(0, T)(u^V(T, t)f^t - U^0(T, t)f^t) \\ &= u^V(0, T)u^0(T, 0)u^0(0, t)f^t + u^V(0, T)(u^V(T, t)f^t - U^0(T, t)f^t). \end{aligned}$$

We know from [Ba2, Lemma VI.8] that $w\text{-}\lim_{t \rightarrow +\infty} u^0(0, t)f^t = 0$. Using (2.9) and an $\epsilon/2$ argument we obtain the proposition. \square

2.3.3. *Limits of quasi-free states.* The following theorem is the key result of [Ba2].

Theorem 2.8. *For $f \in \mathfrak{h}_0^l$ one has:*

$$\lim_{t \rightarrow +\infty} (u^V(0, t)f^t | \mathbb{1}_{\mathbb{R}^+}(b_0^V)u^V(0, t)f^t) = (f | (\mathbb{1} + e^{-2\pi\kappa^{-1}b_\infty^0})^{-1}f).$$

The analogous result for $f \in \mathfrak{h}_0^r$ follows immediately from Prop. 2.6 and (2.8).

Proposition 2.9. *For $f \in \mathfrak{h}_0^r$ one has:*

$$\lim_{t \rightarrow +\infty} (u^V(0, t)f^t | \mathbb{1}_{\mathbb{R}^+}(b_0^V)u^V(0, t)f^t) = (w^r f | \mathbb{1}_{\mathbb{R}^+}(b_0^V)w^r f).$$

We recall that $(f | \mathbb{1}_{\mathbb{R}^+}(b_0^V)f)$ is the covariance of the quasi-free vacuum state for the Dirac field in the exterior of the star at $t = 0$, while $(f | (\mathbb{1} + e^{-2\pi\kappa^{-1}b_\infty^0})^{-1}f)$ is the covariance of the thermal state at Hawking temperature $\kappa/2\pi$ near the black hole horizon.

3. FREE QUANTUM DYNAMICS

In this section we define the free quantum dynamics corresponding to the classical dynamics constructed in Subsect. 2.2.

Let us first introduce some notation. For $t \geq 0$ we set $\mathfrak{A}_t := \text{CAR}(\mathfrak{h}_t)$, $\mathfrak{A}_\infty := \text{CAR}(\mathfrak{h}_\infty)$ and for an interval $J \Subset \mathbb{R}$, $\mathfrak{A}_J := \text{CAR}(\mathfrak{h}_J)$ (see Subsect. A.1). Note that $\mathfrak{A}_t, \mathfrak{A}_J \subset \mathfrak{A}_\infty$ isometrically.

We start by a definition analogous to Def. 2.1.

Definition 3.1. *A family $\{\tau(s, t)\}_{s, t \in \mathbb{R}}$ is a (two-parameter) quantum dynamics if:*

- i) $\tau(s, t) : \mathfrak{A}_t \xrightarrow{\sim} \mathfrak{A}_s$,
- ii) $\tau(t, t) = \mathbb{1}, t \in \mathbb{R}$,
- iii) $\tau(s, t')\tau(t', t) = \tau(s, t), s, t', t \in \mathbb{R}$,
- iv) $\forall (s_0, t_0) \in \mathbb{R}^2, \forall J \Subset]z(t_0), +\infty[\forall A \in \mathfrak{A}_J$ the map $(s, t) \mapsto \tau(s, t)A \in \mathfrak{A}_\infty$ is continuous at (s_0, t_0) .

Since $u^V(t, s)$ is a propagator, it generates a (free) quantum dynamics $\tau^V(s, t)$.

Definition 3.2. *We denote by $\tau^V(s, t)$ the quantum dynamics defined by:*

$$\tau^V(s, t)(\psi^{(*)}(f)) := \psi^{(*)}(u^V(s, t)f), f \in \mathfrak{h}_t.$$

Similarly we define the quantum dynamics $\tau^0(s, t), \tau_\infty^V(s, t)$ associated to $u^0(s, t)$ and $u_\infty^V(s, t)$.

Note that $\tau_\infty^V(s, t)$ is a *stationary* quantum dynamics on \mathfrak{A}_∞ , i.e. $\tau_\infty^V(s, t) = \tau_\infty^V(s + t', t + t')$, for all $s, t', t \in \mathbb{R}$.

We also define the (one-parameter) dynamics α^t on \mathfrak{A}_∞ defined by

$$(3.1) \quad \alpha^t(\psi^{(*)}(f)) := \psi^{(*)}(f^t), f \in \mathfrak{h}_\infty.$$

The properties of propagators recalled in Subsect. 2.3 immediately carry over to quantum dynamics. For example the following fact follows from Prop. 2.5.

Lemma 3.3. *Let $J \Subset \mathbb{R}$ an interval. Then there exists $c \geq 0$ such that*

$$\tau^V(s, t)A = \tau_\infty^V(s, t)A, \forall A \in \mathfrak{A}_J, c + t/2 \leq s \leq t.$$

4. INTERACTING QUANTUM DYNAMICS

In this section we construct the interacting dynamics $\tau^{V, \text{int}}(s, t)$ that we will consider in the sequel. It will be obtained by perturbing the free dynamics $\tau^V(s, t)$ by a bounded interaction term I localized in a bounded region of space. As usual, since we consider fermionic fields, interacting dynamics can be constructed at the algebraic level.

Formally the construction of the interacting dynamics $\tau^{V, \text{int}}(s, t)$ defined in Def. 4.4 corresponds to the quantization of the following non-linear Dirac equation:

$$(4.1) \quad \left\{ \begin{array}{l} \partial_s \psi(s, x) + L \partial_x \psi(s, x) + iV(x)\psi(s, x) \\ - \text{in}(\overline{\psi(s, g)}) | M \psi(s, g) \rangle_{\mathbb{C}^2}^{-1} M \psi(s, g) g(x) = 0, \\ \psi_1(s, z(s)) = \lambda(s) \psi_2(s, z(s)), s \in \mathbb{R}, \\ \psi(t, x) = \psi(x), \text{ in } x > z(t), \end{array} \right.$$

where $\psi(s, g) := \int \psi(s, x) \overline{g(x)} dx \in \mathbb{C}^2$, $M \in M_2(\mathbb{C})$ is a selfadjoint matrix and $g \in L^2(J)$ for some $J \Subset \mathbb{R}$ is a compactly supported function.

4.1. Construction of the interacting dynamics.

Definition 4.1. Let $M \in M_2(\mathbb{C})$ with $M = M^*$. We set for $g \in L^2(\mathbb{R})$:

$$\psi^*(g)M\psi(g) := \sum_{i=1}^2 \psi^*(g \otimes e_i) \lambda_i \psi(g \otimes e_i) \in \mathfrak{A}_\infty,$$

where $M = \sum_{i=1}^2 \lambda_i |e_i\rangle\langle e_i|$ (i.e. (e_1, e_2) is a basis of eigenvectors of M).

We fix $g \in L^2(J)$ for an interval $J \Subset]z(0), +\infty[$, $2 \leq n \in \mathbb{N}$ and set:

$$(4.2) \quad I := (\psi^*(g)M\psi(g))^n \in \text{CAR}_0(\mathfrak{h}_0).$$

The interaction term I represent a localized, even, self-interaction of the Dirac field in \mathcal{M} .

Remark 4.2. All the results below extend immediately to the case when I is replaced by a finite sum of I_k , associated to matrices M_k and compactly supported space-cutoffs g_k . The only important properties of I is that it should be even and localized.

For later use we state the following fact, which follows immediately from the CAR and the fact that $I \in \text{CAR}_0(\mathfrak{h}_\infty)$.

Lemma 4.3. Let $B = \prod_{i=1}^n \psi^{(*)}(f_i)$. Then there exists C_n such that

$$(4.3) \quad \|[I, B]\| \leq C_n \prod_{i=1}^n \|f_i\| \sum_{i=1}^n |(g|f_i)|.$$

Using the results of Subsect. A.6 we can now construct the *interacting dynamics* $\{\tau^{V,\text{int}}(s, t)\}_{s, t \in \mathbb{R}}$.

Definition 4.4. Let I be as in (4.2), $I(s, t) := \tau^V(s, t)I$ and $R(s, t) := R_s(s, t) \in U(\mathfrak{A}_s)$ be obtained as in Prop. A.11. We set:

$$(4.4) \quad \tau^{V,\text{int}}(s, t)(A) := R(s, t)\tau^V(s, t)(A)R(s, t)^*, \quad A \in \mathfrak{A}_t, \quad s, t \in \mathbb{R}$$

Then by Prop. A.11 $\{\tau^{V,\text{int}}(s, t)\}_{s, t \in \mathbb{R}}$ is a dynamics, called the interacting quantum dynamics.

For the convenience of the reader, we recall that $R(s, t) \in \mathfrak{A}_s$ solves:

$$(4.5) \quad \begin{cases} \partial_\sigma R(s, \sigma) = -iR(s, \sigma)I(s, \sigma), \\ R(s, s) = \mathbb{1}. \end{cases}$$

We can also define the corresponding interacting dynamics without boundary conditions, acting on \mathfrak{A}_∞ . We set $I_\infty(s, t) := \tau_\infty^V(s, t)(I)$ and define $R_\infty(s, t) \in U(\mathfrak{A}_\infty)$ as above and:

$$(4.6) \quad \tau_\infty^{V,\text{int}}(s, t)(A) := R_\infty(s, t)\tau_\infty^V(s, t)(A)R_\infty(s, t)^*, \quad A \in \mathfrak{A}_\infty, \quad s, t \in \mathbb{R}.$$

Again $\tau_\infty^{V,\text{int}}(s, t)$ is stationary.

Remark 4.5. Let us faithfully represent $\mathfrak{A}_\infty = \text{CAR}(\mathfrak{h}_\infty)$ in the fermionic Fock space $\Gamma_a(\mathfrak{h}_\infty)$ (see Subsect. A.1) by the Fock representation π_F . Then $\tau_\infty^V(s, t)$ is implemented in the Fock representation by the unitary group $e^{i(s-t)H_\infty^V}$, where $H_\infty^V = d\Gamma(b_\infty^V)$ is the second quantization of b_∞^V . The dynamics $\tau_\infty^{V,\text{int}}(s, t)$ is implemented by $e^{i(s-t)H_\infty^{V,\text{int}}}$ for $H_\infty^{V,\text{int}} = H_\infty^V + \pi_F(I)$.

4.2. Properties of $\tau^{V,\text{int}}(s, t)$.

Lemma 4.6. *There exists $c \geq 0$ such that:*

$$R_\infty(s, t) = R(s, t), \quad c + t/2 \leq s \leq t.$$

Proof. The interaction I defined in (4.2) belongs to \mathfrak{A}_J for some interval $J \in \mathbb{R}$. We apply then Lemma 3.3 to each term in the series defining $R(s, t)$, see Lemma A.10. \square

Lemma 4.7. *Let $J \in \mathbb{R}$ an interval. Then there exists $c \geq 0$ such that*

$$\tau^{V,\text{int}}(s, t)(A) = \tau_\infty^{V,\text{int}}(s, t)(A), \quad \forall A \in \mathfrak{A}_J, \quad c + t/2 \leq s \leq t.$$

Proof. It suffices to apply Lemmas 4.6 and 3.3 to the definition of $\tau^{V,\text{int}}, \tau_\infty^{V,\text{int}}$. \square

5. HAWKING EFFECT I

In this section we study the Hawking effect in the situation referred to as case I in the introduction (see Subsect. 1.3). For $t \in \mathbb{R}$ we set $\mathfrak{A}_t^{1/r} := \text{CAR}(\mathfrak{h}_t^{1/r}) \subset \mathfrak{A}_t$, called the *left/right moving observables*.

The algebra \mathfrak{A}_0 splits into a twisted tensor product of the left/right moving CAR algebras $\mathfrak{A}_0^{1/r}$. The first step consists in studying the evolution $\tau^{V,\text{int}}(0, t) \circ \alpha^t$ on left/right moving observables.

5.1. Left propagation.

Proposition 5.1. *Let $A \in \mathfrak{A}_0^1$. Then*

$$\lim_{t \rightarrow +\infty} \tau^{V,\text{int}}(0, t) \circ \alpha^t(A) - \tau^V(0, t) \circ \alpha^t(A) = 0.$$

To prove Prop. 5.1, we will need the following lemma.

Lemma 5.2. *For any $\epsilon > 0$ and $A \in \mathfrak{A}_0$, there exists T such that*

$$\sup_{t \geq T} \|\tau^{V,\text{int}}(T, t) \circ \alpha^t(A) - \tau^V(T, t) \circ \alpha^t(A)\| \leq \epsilon.$$

Proof. Let us set $A(s, t) = \tau^V(s, t) \circ \alpha^t(A)$ to simplify notation. We first claim that

$$(5.1) \quad \|\tau^{V,\text{int}}(s, t) \circ \alpha^t(A) - A(s, t)\| \leq \int_s^t \| [I, A(\sigma, t)] \| d\sigma.$$

Let us prove (5.1). By Def. 4.4 we have:

$$\begin{aligned} \tau^{V,\text{int}}(s, t) \circ \alpha^t(A) - A(s, t) &= R(s, t)A(s, t)R(s, t)^* - A(s, t) \\ &= [R(s, t), A(s, t)]R^*(s, t), \end{aligned}$$

using that $R(s, t)$ is unitary. Set :

$$F_{s,t}(\sigma) := [R(s, \sigma), A(s, t)], \quad G_{s,t}(\sigma) := [I(s, \sigma), A(s, t)].$$

We note first that

$$G_{s,t}(\sigma) = [\tau^V(s, \sigma)(I), \tau^V(s, t) \circ \alpha^t(A)] = \tau^V(s, \sigma)([I, A(\sigma, t)]),$$

using that τ^V is an homomorphism. Since τ^V is isometric, we have

$$(5.2) \quad \|G_{s,t}(\sigma)\| = \|[I, A(\sigma, t)]\|.$$

Recalling that $R(s, \sigma)$ solves

$$\begin{cases} \partial_\sigma R(s, \sigma) = -iR(s, \sigma)I(s, \sigma), \\ R(s, s) = \mathbb{1}, \end{cases}$$

we see next that $F_{s,t}(\cdot)$ solves the equation:

$$(5.3) \quad \begin{cases} \partial_\sigma F_{s,t}(\sigma) = -iF_{s,t}(\sigma)I(s,\sigma) - iR(s,\sigma)G_{s,t}(\sigma), \\ F_{s,t}(s) = 0, \end{cases}$$

which clearly has a unique solution. We look for $F_{s,t}(\sigma)$ of the form $F_{s,t}(\sigma) = H_{s,t}(\sigma)R(s,\sigma)$. We obtain the equation:

$$(5.4) \quad \begin{cases} \partial_\sigma H_{s,t}(\sigma) = -iR(s,\sigma)G_{s,t}(\sigma)R(s,\sigma)^*, \\ H_{s,t}(s) = 0. \end{cases}$$

Since $R(s,\sigma)$ is unitary we obtain

$$\begin{aligned} \|\tau^{V,\text{int}}(s,t) \circ \alpha^t(A) - A(s,t)\| &= \|F_{s,t}(t)\| = \|H_{s,t}(t)\| \\ &\leq \int_s^t \|G_{s,t}(\sigma)\| d\sigma = \int_s^t \| [I, A(\sigma,t)] \| d\sigma, \end{aligned}$$

which proves (5.1).

We can now complete the proof of the lemma. Assume first that A belongs to $\text{CAR}_{\text{alg}}(\mathfrak{h}_J)$ for some interval $J \subset [0, R]$ (recall that $z(0) = 0$). By linearity we may assume that $A = \prod_{i=1}^n \psi^{(*)}(f_i)$ with $\text{supp} f_i \subset [0, R]$. By Prop. 2.4 (3) we know that $\text{supp} u^V(\sigma, t) f_i^t \subset [z(\sigma), -\sigma + R]$ hence for $\sigma \geq \sigma(J)$ we have $[I, A(\sigma, t)] = 0$ by Lemma 4.3, hence

$$(5.5) \quad \tau^{V,\text{int}}(s,t)(A) - A(s,t) = 0, \quad \sigma(J) \leq s \leq t, \quad A \in \text{CAR}_{\text{alg}}(\mathfrak{h}_J).$$

Let now $A \in \mathfrak{A}_0$ and $\epsilon > 0$. By density we can choose J as above and $\tilde{A} \in \text{CAR}_{\text{alg}}(\mathfrak{h}_J)$ such that $\|A - \tilde{A}\| \leq \epsilon/2$. Applying (5.5) to \tilde{A} we obtain $T = \sigma(J)$ such that

$$\sup_{T \leq t} \|\tau^{V,\text{int}}(T,t) \circ \alpha^t(A) - A(T,t)\| \leq \epsilon.$$

This completes the proof of the lemma. \square

Proof of Prop. 5.1. Let $A \in \mathfrak{A}_0^1$. Again let us set $A(s,t) = \tau^V(s,t) \circ \alpha^t(A)$, so that we need to show that

$$\lim_{t \rightarrow +\infty} \tau^{V,\text{int}}(0,t) \circ \alpha^t(A) - A(0,t) = 0.$$

We fix $\epsilon > 0$ and T as in Lemma 5.2. We have:

$$\begin{aligned} \tau^{V,\text{int}}(0,t) \circ \alpha^t(A) &= \tau^{V,\text{int}}(0,T) \circ \tau^{V,\text{int}}(T,t) \circ \alpha^t(A) \\ &= R(0,T) \tau^V(0,T) \circ \tau^V(T,t) \circ \alpha^t(A) R(0,T)^* \\ &= R(0,T) A(0,t) R(0,T) + O(\epsilon), \end{aligned}$$

by Lemma 5.2. By (4.5) we have:

$$\partial_\sigma (R(0,\sigma) B R(0,\sigma)^*) = -iR(0,\sigma) [I(0,\sigma), B] R(0,\sigma)^*, \quad B \in \mathfrak{A}_0$$

hence:

$$R(0,T) A(t) R(0,T)^* - A(0,t) = -i \int_0^T R(0,\sigma) [I(0,\sigma), A(0,t)] R(0,\sigma)^* d\sigma.$$

By (5.2) for $s = 0$ we have $\| [I(0,\sigma), A(0,t)] \| = \| [I, A(\sigma,t)] \|$. To complete the proof of the proposition, it suffices to show that

$$(5.6) \quad \lim_{t \rightarrow +\infty} [I, A(\sigma,t)] = 0, \quad \forall \sigma \geq 0.$$

Since $\|A(\sigma,t)\| = \|A\|$, it suffices by density and linearity to prove (5.6) if $A = \prod_{i=1}^n \psi^{(*)}(f_i)$ for $f_i \in \mathfrak{h}_0^1$ with compact support. By Lemma 4.3 it suffices hence to prove that

$$\text{w-} \lim_{t \rightarrow +\infty} u^V(0,t) f_i^t = 0.$$

But this follows from Prop. 2.7. This completes the proof of the proposition. \square

5.2. Right propagation.

Proposition 5.3. *The strong limit*

$$s\text{-}\lim_{t \rightarrow +\infty} \tau^{V,\text{int}}(0,t) \circ \alpha^t =: \gamma^{r,\text{int}}$$

exists on \mathfrak{A}_0^r .

Before proving the proposition, let us note that $\gamma^{r,\text{int}}$ is an even homomorphism (see Subsect. A.1).

Lemma 5.4. *The homomorphism $\gamma^{r,\text{int}}$ is even i.e. $P \circ \gamma^{r,\text{int}} = \gamma^{r,\text{int}} \circ P$.*

Proof. α^t is even, so it suffices to prove that $\tau^{V,\text{int}}(0,t)$ is even. This follows if we prove that $R(s,t) \in \text{CAR}_0(\mathfrak{h}_s)$. We note that $R(s,\sigma)$ and $PR(s,\sigma)$ solve the same differential equation, using that I is even. \square

Proof of Prop. 5.3. Let $A \in \mathfrak{A}_0^r$. By (2.8) we have $\alpha^t = \tau_\infty^0(t,0)$ on \mathfrak{A}_0^r . Therefore we will be able to prove the proposition by the Cook argument. We will first prove that

$$(5.7) \quad \lim_{t \rightarrow +\infty} \tau^V(0,t) \circ \tau_\infty^0(t,0)(A) =: \gamma_0^r(A), \quad A \in \mathfrak{A}_0^r.$$

exists, and then that

$$(5.8) \quad s\text{-}\lim_{t \rightarrow +\infty} \tau^{V,\text{int}}(0,t) \circ \tau^V(t,0)(A), \quad A \in \gamma_0^r \mathfrak{A}_0^r.$$

exists. Let us first prove (5.7). Since $\tau^V(0,t)$ and $\tau_\infty^0(t,0)$ are free dynamics, this follows from Prop. 2.6 which states that:

$$(5.9) \quad \lim_{t \rightarrow +\infty} u^V(0,t)u_\infty^0(t,0)f = w_0^r f$$

exists for $f \in \mathfrak{h}_0^r$. It follows that

$$\gamma_0^r(\psi^{(*)}(f)) = \psi^{(*)}(w_0^r f), \quad f \in \mathfrak{h}_0^r.$$

To prove (5.8) we will need some estimates on the speed of convergence in (5.9), for well chosen initial data.

Assume that $f \in \mathfrak{h}_0^r$ is smooth with compact support. Then $u_\infty^0(t,0)f = f^t \equiv 0$ near $x = z(t)$ hence $u_\infty^0(t,0)f \in \text{Dom } b_t^V$. It follows that:

$$\partial_t u^V(0,t)u_\infty^0(t,0)f = iu^V(0,t)(b_\infty^0 - b_t^V)u_\infty^0(t,0)f = iu^V(0,t)Vf^t.$$

From hypothesis (2.3) we obtain that $\|Vf^t\| \in O(t^{-2-\epsilon})$ hence by integrating from t to $+\infty$, we obtain:

$$(5.10) \quad w_0^r f - u^V(0,t)u_\infty^0(t,0)f \in O(t^{-1-\epsilon}).$$

Let us now prove (5.8). By linearity, density and using that $\tau^{V,\text{int}}(0,t)$ and $\tau^V(t,0)$ are isomorphisms, we can assume that $A = \psi^{(*)}(w_0^r f)$ for $f \in \mathfrak{h}_0^r$ smooth with compact support. We have

$$\begin{aligned} \tau^{V,\text{int}}(0,t) \circ \tau^V(t,0)(A) &= R(0,t)\tau^V(0,t) \circ \tau^V(t,0)(A)R(0,t)^* \\ &= R(0,t)AR(0,t)^*. \end{aligned}$$

We apply once more the Cook argument and compute

$$\partial_t R(0,t)\gamma_0^r(A)R(0,t)^* = -iR(0,t)[I(0,t), A]R(0,t)^*.$$

As before

$$\begin{aligned} \|[I(0,t), A]\| &= \|[I, \tau^V(t,0)(A)]\| = \|[I, \psi^{(*)}(u^V(t,0)w_0^r f)]\| \\ &= \|[I, \psi^{(*)}(u^0(t,0)f)]\| + O(t^{-1-\epsilon}), \end{aligned}$$

by (5.10). Since f has compact support, and $u^0(t, 0)f = f^t$, we obtain that $[I, u^0(t, 0)f] = 0$ for t large enough. Therefore $\|\partial_t R(0, t)\gamma_0^r(A)R(0, t)^*\| \in O(t^{-1-\epsilon})$, which proves (5.8) by the Cook argument. \square

5.3. Hawking effect I.

5.3.1. *The limit state.* In the rest of the paper we denote by $\beta = 2\pi\kappa^{-1}$ the inverse Hawking temperature.

Let us denote by $\omega_{\infty, \beta}^0$ the gauge-invariant quasi-free thermal state on \mathfrak{A}_∞ with covariance:

$$\omega_{\infty, \beta}^0(\psi^*(f)\psi(g)) = (f | (\mathbb{1} + e^{-\beta b_\infty^0})^{-1} g), \quad f, g \in \mathfrak{h}_\infty.$$

This state restricts to a quasi-free state on \mathfrak{A}_0^1 , still denoted by $\omega_{\infty, \beta}^0$.

We denote by $\omega_{0, \text{vac}}^V$ the gauge-invariant quasi-free vacuum state on \mathfrak{A}_0 with covariance:

$$\omega_{0, \text{vac}}^V(\psi^*(f)\psi(g)) = (f | \mathbb{1}_{\mathbb{R}^+}(b_0^V)g), \quad f, g \in \mathfrak{h}_0.$$

The state $\omega_{0, \text{vac}}^V \circ \gamma^{r, \text{int}}$ is a gauge-invariant state on \mathfrak{A}_0^r , which is even by Lemma 5.4.

Since $\mathfrak{h}_0 = \mathfrak{h}_0^1 \oplus \mathfrak{h}_0^r$, we can by Def. A.7 define the following state on \mathfrak{A}_0 :

Definition 5.5. *We set*

$$\omega_{\text{H,I}} := \omega_{\infty, \beta}^0 \widehat{\otimes} (\omega_{0, \text{vac}}^V \circ \gamma^{r, \text{int}}),$$

which is a state on \mathfrak{A}_0 .

5.3.2. *Main result I.* The following theorem is the main result of this section.

Theorem 5.6.

$$\lim_{t \rightarrow +\infty} \omega_{0, \text{vac}}^V \circ \tau^{V, \text{int}}(0, t) \circ \alpha^t(A) = \omega_{\text{H,I}}(A), \quad A \in \mathfrak{A}_0.$$

Proof. By linearity and density we may assume that $A = A_1 \times A_2$, $A_1 \in \mathfrak{A}_0^1$, $A_2 \in \mathfrak{A}_0^r$. By Prop. 5.3 we have $\tau^{V, \text{int}} \circ \alpha^t(A_2) = \gamma^{d, \text{int}}(A_2) + o(t^0)$. Applying Lemma 5.7 we obtain that

$$\lim_{t \rightarrow +\infty} \omega_{0, \text{vac}}^V(A_1 A_2) = \omega_{\infty, \beta}^0 \widehat{\otimes} \omega_{0, \text{vac}}^V \circ \gamma^{r, \text{int}}(A_1 A_2),$$

which completes the proof of the theorem. \square

Lemma 5.7. *Let $A_1 \in \mathfrak{A}_0^1$, $A_2 \in \mathfrak{A}_0$. Then*

$$\lim_{t \rightarrow +\infty} \omega_{0, \text{vac}}^V(\tau^{V, \text{int}}(0, t) \circ \alpha^t(A_1) A_2) = \omega_{\infty, \beta}^0 \widehat{\otimes} \omega_{0, \text{vac}}^V(A_1 A_2).$$

Proof. By linearity and density we can assume that

$$A_1 = \prod_{i=1}^{n_1} \psi^*(f_i) \prod_{i=1}^{p_1} \psi(g_i), \quad A_2 = \prod_{i=1}^{n_2} \psi^*(f_{n_1+i}) \prod_{i=1}^{p_2} \psi(g_{p_1+i}),$$

where

$$f_i, g_j \in \mathfrak{h}_0^1, \quad \text{for } 1 \leq i \leq n_1, \quad 1 \leq j \leq p_1,$$

$$f_{n_1+i}, g_{p_1+j} \in \mathfrak{h}_\infty, \quad \text{for } 1 \leq i \leq n_2, \quad 1 \leq j \leq p_2.$$

To simplify notation we set

$$\gamma_{\text{int}}^t := \tau^{V, \text{int}}(0, t) \circ \alpha^t, \quad \gamma^t := \tau^V(0, t) \circ \alpha^t,$$

so that from Prop. 5.1 we have $\gamma_{\text{int}}^t(A_1) = \gamma^t(A_1) + o(t^0)$. It follows that

$$\gamma_{\text{int}}^t(A_1) = \prod_{i=1}^{n_1} \psi^*(u^V(0, t) f_{1i}^t) \prod_{i=1}^{p_1} \psi(g_{1i}^t) + o(t^0).$$

Using the CAR and Prop. 2.7, we obtain that:

$$\begin{aligned} & \gamma_{\text{int}}^t(A_1)A_2 \\ &= (-1)^{n_2(n_1+p_1)} \prod_{i=1}^{n_2} \psi^*(f_{2i}) \prod_{i=1}^{n_1} \psi^*(u^V(0,t)f_{1i}^t) \prod_{i=1}^{p_1} \psi(g_{1i}^t) \prod_{i=1}^{p_2} \psi(g_{2i}) + o(t^0). \end{aligned}$$

Since $\omega_{0,\text{vac}}^V$ is a gauge invariant quasi-free state (see Subsect. A.3), we see that $\omega_{0,\text{vac}}^V(\gamma_{\text{int}}^t(A_1)A_2) = o(t^0)$ if $n_1 + n_2 \neq p_1 + p_2$, and if $n_1 + n_2 = p_1 + p_2 = n$ we have:

$$(5.11) \quad \begin{aligned} & \omega_{0,\text{vac}}^V(\gamma_{\text{int}}^t(A_1)A_2) \\ &= (-1)^{n_2(n_1+p_1)} \sum_{\sigma \in S_n} \epsilon(\sigma) \prod_{k=1}^n \omega_{0,\text{vac}}^V(\psi^*(F_k^t)\psi(G_{\sigma(k)}^t)) + o(t^0), \end{aligned}$$

where:

$$F_k^t = \begin{cases} u^V(0,t)f_k^t & \text{for } 1 \leq k \leq n_1, \\ f_k & \text{for } n_1 + 1 \leq k \leq n, \end{cases}, \quad G_k^t = \begin{cases} u^V(0,t)g_k^t & \text{for } 1 \leq k \leq p_1, \\ g_k & \text{for } p_1 + 1 \leq k \leq n. \end{cases}$$

Recall that $\omega_{0,\text{vac}}^V(\psi(f)\psi(g)) = (f|\mathbb{1}_{\mathbb{R}^+}(b_0^V)g)_{\mathfrak{h}_0}$ and $w\text{-}\lim u^V(0,t)f^t = 0$ for $f \in \mathfrak{h}_0^1$ by Prop. 2.7. We see the sum on the r.h.s. is $o(t^0)$ unless $n_1 = p_1$ and $n_2 = p_2$. If this is the case the only permutations σ contributing to the sum are of the form $\sigma_1 \times \sigma_2$ where $\sigma_i \in S_{n_i}$. Collecting these terms we obtain that:

$$\omega_{0,\text{vac}}^V(\gamma_{\text{int}}^t(A_1)A_2) = \omega_{0,\text{vac}}^V(\gamma^t(A_1))\omega_{0,\text{vac}}^V(A_2) + o(t^0).$$

By the result of Bachelot [Ba2] recalled in Thm. 2.8 we know that

$$\lim_{t \rightarrow +\infty} \omega_{0,\text{vac}}^V(\gamma^t(A_1)) = \omega_{\infty,\beta}^0(A_1).$$

Now we use Remark A.4 and the definition of the \mathbb{Z}_2 -graded tensor product of two states (see Lemma A.6) to see that

$$\lim_{t \rightarrow +\infty} \omega_{0,\text{vac}}^V(\gamma_{\text{int}}^t(A_1)A_2) = \omega_{\infty,\beta}^0 \tilde{\otimes} \omega_{0,\text{vac}}^V(A_1 A_2),$$

which completes the proof of the lemma. \square

5.4. Change of initial state. We assumed in Subsect. 2.1 that the star was stationary in $t \leq 0$. It is hence natural to take as dynamics in the past the stationary *interacting* dynamics $\tau_0^{V,\text{int}}(s,t)$ on \mathfrak{A}_0 defined as follows: we first define the stationary analog of $\tau^V(s,t)$, acting on \mathfrak{A}_0 by

$$\tau_0^V(s,t)\psi^{(*)}(f) := \psi^{(*)}(e^{itb_0^V}f), \quad f \in \mathfrak{h}_0.$$

We can then define the stationary interacting dynamics $\tau_0^{V,\text{int}}(s,t)$ associated to I in Def. 4.1. It suffices to repeat the construction in Subsect. 4.1 with $\tau_0^V(s,t)$ instead of $\tau^V(s,t)$.

An adapted choice of the initial state in Thm. 5.6 would be an even state $\tilde{\omega}$ on \mathfrak{A}_0 , *invariant* under $\tau_0^{V,\text{int}}(s,t)$. The following easy result shows that Thm. 5.6 will extend to $\tilde{\omega}$, provided that $\tilde{\omega}$ belongs to the *folium* of $\omega_{0,\text{vac}}^V$, i.e. is represented by a density matrix in the GNS representation of $\omega_{0,\text{vac}}^V$. Recall that such states are physically interpreted as local perturbations of $\omega_{0,\text{vac}}^V$.

Corollary 5.8. *Let $\tilde{\omega}$ a state on \mathfrak{A}_0 which is even and belongs to the folium of $\omega_{0,\text{vac}}^V$. Then*

$$\lim_{t \rightarrow +\infty} \tilde{\omega} \circ \tau^{V,\text{int}}(0,t) \circ \alpha^t(A) = \tilde{\omega}_{\text{H,I}}(A), \quad A \in \mathfrak{A}_0,$$

where:

$$\tilde{\omega}_{\text{H,I}} = \omega_{\infty,\beta}^0 \tilde{\otimes} (\tilde{\omega} \circ \gamma^{\text{r,int}}).$$

Proof. Since $\tilde{\omega}$ belongs to the folium of $\omega_{0,\text{vac}}^V$, we are, by linearity and density, reduced to compute the limit:

$$\lim_{t \rightarrow +\infty} \omega_{0,\text{vac}}^V(P^*(\psi, \psi^*)\gamma^t(A)_1 A_2 P(\psi, \psi^*)),$$

where $A_1 \in \mathfrak{A}_0^1$, $A_2 \in \mathfrak{A}_0$ and $P(\psi, \psi^*)$ is a polynomial in $\text{CAR}_{\text{alg}}(\mathfrak{h}_0)$. Moreover since $\tilde{\omega}$ is even, we see that $P(\psi, \psi^*) \in \text{CAR}_{\text{alg},0}(\mathfrak{h}_0)$. By the same argument as in the proof of Lemma 5.7, we see that

$$P^*(\psi, \psi^*)\gamma^t(A)_1 A_2 P(\psi, \psi^*) = \gamma^t(A)_1 P^*(\psi, \psi^*) A_2 P(\psi, \psi^*) + o(t^0),$$

hence as in Lemma 5.7 we have:

$$\begin{aligned} & \lim_{t \rightarrow +\infty} \omega_{0,\text{vac}}^V(P^*(\psi, \psi^*)\gamma^t(A)_1 A_2 P(\psi, \psi^*)) \\ &= \omega_\beta^0(A_1) \omega_{0,\text{vac}}^V(P^*(\psi, \psi^*) A_2 P(\psi, \psi^*)) = \omega_\beta^0(A_1) \tilde{\omega}(A_2). \end{aligned}$$

We can then complete the proof as in Thm. 5.6. \square

5.5. Existence of interacting initial vacua. It remains to construct even states $\tilde{\omega}$ which belong to the folium of $\omega_{0,\text{vac}}^V$ and are invariant under $\tau_0^{V,\text{int}}(s, t)$. To do this it is convenient to work in the GNS representation of the vacuum state $\omega_{0,\text{vac}}^V$, i.e. the *Fock representation*. We refer the reader to Subsect. A.4.

Recall that b_0^V is defined in Subsect. 2.1.2. It is easy to show that

$$(5.12) \quad \sigma_{\text{ess}}(b_0^V) =] - \infty, -m] \cup [m, +\infty[.$$

We assume that $\text{Ker} b_0^V = \{0\}$ and equip \mathfrak{h}_0 with the complex structure $j := i \text{sgn}(b_0^V)$, and denote by \mathcal{Z} the associated one-particle space. If π_F is the corresponding Fock representation we have:

$$\omega_{0,\text{vac}}^V(A) = (\Omega | \pi_F(A) \Omega),$$

where $\Omega \in \Gamma_a(\mathcal{Z})$ is the vacuum vector. In other words $(\Gamma_a(\mathcal{Z}), \pi_F, \Omega)$ is the GNS triple associated to $\omega_{0,\text{vac}}^V$.

From Subsect. A.4 we know that:

$$\pi_F(\tau_0^V(s, t)A) = e^{i(s-t)H_0} \pi_F(A) e^{i(t-s)H_0}, \quad A \in \mathfrak{A}_0,$$

for $H_0 = d\Gamma(|b_0^V|)$, and if $Q = d\Gamma(\text{sgn}(b_0^V))$, then:

$$\psi_F^{(*)}(e^{i\theta} f) = e^{i\theta Q} \psi_F^{(*)}(f) e^{-i\theta Q}, \quad f \in \mathfrak{h}_0, \quad \theta \in \mathbb{R}.$$

It is also well known that if

$$H := H_0 + \pi_F(I),$$

then

$$\pi_F(\tau_0^{V,\text{int}}(s, t)(A)) = e^{i(s-t)H} \pi_F(A) e^{i(t-s)H}, \quad A \in \mathfrak{A}_0.$$

Since $\tau_0^{V,\text{int}}(s, t)$ is implemented by $e^{i(s-t)H}$ in the Fock representation, eigenvectors of H will yield invariant states for $\tau_0^{V,\text{int}}(s, t)$, which obviously belong to the folium of $\omega_{0,\text{vac}}^V$.

Existence of eigenvectors is ensured by the following theorem, whose proof follows by adapting arguments in [A, DG].

Theorem 5.9. *On has*

$$\sigma_{\text{ess}}(H) = [\inf \sigma(H) + m, +\infty[.$$

Therefore $\inf \sigma(H)$ is an eigenvalue of H .

To be able to apply Corollary 5.8, we need however the existence of an *even* eigenstate ψ of H , i.e. such that $Q\psi = 2n\psi$ for some $n \in \mathbb{Z}$. Note that since I is even we have $[H, Q] = 0$, which does not imply the existence of even eigenstates of H . However this is clearly true for small interactions. In fact setting $H(\lambda) = H_0 + \lambda I$ and $E(\lambda) = \inf \sigma(H(\lambda))$ we have

Lemma 5.10. *Assume $|\lambda|$ is small enough. Then $\mathbb{1}_{\{E(\lambda)\}}(H(\lambda))$ is rank one and $Q\mathbb{1}_{\{E(\lambda)\}}(H(\lambda)) = 0$.*

Therefore for λ small enough, $H(\lambda)$ has a unique ground state $\Omega(\lambda)$ of zero charge and the associated state satisfies the hypotheses of Corollary 5.8.

6. HAWKING EFFECT II

In this section we study the Hawking effect in case II (see Subsect. 1.3). Compared to Sect. 5, the observable is not translated to the left, therefore the influence of the potential V and of the non-linear self-interaction has to be taken into account. To this end we use tools from scattering theory, both for classical and quantum dynamics.

6.1. Asymptotic velocity for Dirac equations. In this subsection we state some results of Daudé [Da] on the existence of the *asymptotic velocity observable* for stationary Dirac equations. The asymptotic velocity provides a convenient way to separate left and right propagating initial states. More details can be found in [Da].

Theorem 6.1. *Let $\chi \in C_0^\infty(\mathbb{R}, \mathbb{C}^2)$. Then*

(1) *the limits*

$$\chi^\pm := s\text{-}\lim_{t \rightarrow \pm\infty} e^{-itb_\infty^V} \chi\left(\frac{x}{t}\right) e^{itb_\infty^V} \text{ exist.}$$

(2) *there exist bounded selfadjoint operators P^\pm on \mathfrak{h}_∞ such that $\chi^\pm = \chi(P^\pm)$ for $\chi \in C_0^\infty(\mathbb{R}, \mathbb{C}^2)$.*

(3) *one has*

$$[P^\pm, b_\infty^V] = 0, \quad \sigma(P^\pm) = [-1, 1], \quad \mathbb{1}_{\{0\}}(P^\pm) = \mathbb{1}_{\text{pp}}(b_\infty^V).$$

Remark 6.2. *Since it is known (see e.g. [Ba2, Lemma III.1] that b_∞^V has no eigenvalues, we have actually $\mathbb{1}_{\{0\}}(P^\pm) = 0$, i.e. any initial state has a non-vanishing asymptotic velocity.*

We will only use the future asymptotic velocity P^+ which we will denote simply by P . Moreover we will set

$$P^{l/r} := \mathbb{1}_{\mathbb{R}^\mp}(P),$$

so that by Remark 6.2 we have

$$P^l + P^r = \mathbb{1}.$$

We set now $V^{l/r} := \lim_{x \rightarrow \mp\infty} V(x)$, so that $V^l = 0$, $V^r = V_\infty$, see Subsect. 2.1. We set also

$$b_\infty^{l/r} := LD_x + V^{l/r}, \text{ with domain } H^1(\mathbb{R}, \mathbb{C}^2), \text{ acting on } \mathfrak{h}_\infty.$$

From Thm. 6.1 and the short-range nature of the interaction V (see (2.3)), we obtain by standard arguments the existence of wave operators:

Proposition 6.3. *The limit*

$$s\text{-}\lim_{t \rightarrow +\infty} e^{-itb_\infty^l} e^{itb_\infty^V} =: w_\infty^l$$

exists on $P^l \mathfrak{h}_\infty$.

Prop. 6.3 yields the following result for free quantum dynamics.

Proposition 6.4. *The limit*

$$\gamma_\infty^1 := s\text{-}\lim_{t \rightarrow +\infty} \tau_\infty^{V^1}(t, 0) \circ \tau_\infty^V(0, t) \text{ exists on } \text{CAR}(P^1 \mathfrak{h}_\infty)$$

The map $\gamma_\infty^1 : \text{CAR}(P^1 \mathfrak{h}_\infty) \rightarrow \text{CAR}(\mathfrak{h}_\infty) = \mathfrak{A}_\infty$ are $*$ -morphisms with

$$\gamma_\infty^1(\psi^{(*)}(f)) = \psi^{(*)}(w^1 f), \quad f \in P^1 \mathfrak{h}_\infty.$$

The similar limit γ_∞^r with V^1 replaced by V^r and P^1 replaced by P^r exists on $\text{CAR}(P^r \mathfrak{h}_\infty)$ also exists, but will play no role in the sequel. The 'right' analog of γ_∞^1 is the wave morphism γ_0^r introduced below in Prop. 6.13.

6.2. Wave morphisms. We now prove an analog of Prop. 6.4 for interacting dynamics.

Theorem 6.5. *The limit*

$$\gamma_\infty^{\text{int}} := s\text{-}\lim_{t \rightarrow +\infty} \tau_\infty^{V, \text{int}}(t, 0) \circ \tau_\infty^V(0, t) \text{ exists on } \mathfrak{A}_\infty$$

and is a $*$ -morphism of \mathfrak{A}_∞ .

The morphism $\gamma_\infty^{\text{int}}$ is an example of a *wave morphism*.

Proof of Thm. 6.5. The proof relies once again on the Cook argument, combined with minimal velocity estimates for the Dirac equation. For more details on minimal velocity estimates see [Da]. Since b_∞^V has no point spectrum we see that the space \mathcal{D} of vectors in $\cap_{n \in \mathbb{N}} \text{Dom} \langle x \rangle^n$ such that $f = \chi(b_\infty^V) f$ for some $\chi \in C_0^\infty(\mathbb{R} \setminus [-m, m])$ is dense in \mathfrak{h}_∞ . The strong minimal velocity estimates (see [Da]) give

$$(6.1) \quad \forall f \in \mathcal{D}, \exists 0 < c_0 < 1 \text{ such that } \|\mathbb{1}_{[0, c_0]} \left(\frac{|x|}{t} \right) e^{itb_\infty^V} f\| \in O(t^{-N}), \quad \forall N \in \mathbb{N}.$$

We can now argue as in the proof of Prop. 5.3, since by (6.1) $t \mapsto (g|e^{itb_\infty^V} f)$ is integrable for $f \in \mathcal{D}$. \square

We denote by $E_t : \mathfrak{A}_\infty \rightarrow \mathfrak{A}_t$ the $*$ -homomorphism defined in A.1.3, associated to the inclusion $\mathfrak{h}_t \subset \mathfrak{h}_\infty$. Since $\cup_{t \geq 0} \mathfrak{h}_t$ is dense in \mathfrak{h}_∞ , we have:

$$(6.2) \quad s\text{-}\lim_{t \rightarrow +\infty} E_t = \mathbb{1}, \text{ in } \mathfrak{A}_\infty.$$

We now combine Thm. 6.5 with Lemma 4.7 to obtain the following result.

Proposition 6.6. *Let $A = \gamma_\infty^{\text{int}}(B) \in \mathfrak{A}_\infty$. Then for any $\epsilon > 0$ there exist $C_\epsilon, T_\epsilon > 0$ such that*

$$\sup_{t/2 + C_\epsilon \leq s \leq t/2 + 2C_\epsilon, t \geq T_\epsilon} \|\tau_\infty^{V, \text{int}}(s, t) \circ E_t(A) - \tau_\infty^V(s, t)(B)\| \leq \epsilon.$$

Remark 6.7. *We do not know if $A \in \gamma_\infty^{\text{int}} \mathfrak{A}_\infty$ belongs to \mathfrak{A}_t for all t large enough, so a priori $\tau_\infty^{V, \text{int}}(s, t)(A)$ does not make sense. Replacing A by $E_t(A) \in \mathfrak{A}_t$ fixes this problem, at the price of an error $\|A - E_t(A)\|$ which is $o(t^0)$.*

Proof. Since $A = \gamma_\infty^{\text{int}}(B)$ we have:

$$\tau_\infty^V(0, t)(B) = \tau_\infty^{V, \text{int}}(0, t)(A) + o(t^0), \quad t \rightarrow +\infty.$$

Since τ_∞^V and $\tau_\infty^{V, \text{int}}$ are stationary dynamics, this implies that for any $c > 0$:

$$(6.3) \quad \lim_{t \rightarrow +\infty} \sup_{0 \leq s \leq t/2 + c} \|\tau_\infty^V(s, t)(B) - \tau_\infty^{V, \text{int}}(s, t)(A)\| = 0.$$

Since $\bigcup_{J \in \mathbb{R}} \mathfrak{A}(J)$ is dense in \mathfrak{A}_∞ , we can for any $\epsilon > 0$ find $J_\epsilon \in \mathbb{R}$ and $A_\epsilon \in \mathfrak{A}(J_\epsilon)$ such that $\|A - A_\epsilon\| \leq \epsilon/8$, hence:

$$(6.4) \quad \sup_{s,t} \|\tau_\infty^{V,\text{int}}(s,t)(A) - \tau_\infty^{V,\text{int}}(s,t)(A_\epsilon)\| \leq \epsilon/8.$$

By Lemma 4.7 there exists $C_\epsilon = C(J_\epsilon)$ such that:

$$(6.5) \quad \tau^{V,\text{int}}(s,t)(A_\epsilon) = \tau_\infty^{V,\text{int}}(s,t)(A_\epsilon), \quad \forall t/2 + C_\epsilon \leq s \leq t.$$

Next by (6.2), we can find T_ϵ such that $\sup_{t \geq T_\epsilon} \|A - E_t(A)\| \leq \epsilon/8$, hence

$$(6.6) \quad \sup_{s \leq t, t \geq T_\epsilon} \|\tau^{V,\text{int}}(s,t)(A_\epsilon) - \tau^{V,\text{int}}(s,t) \circ E_t(A)\| \leq \epsilon/4.$$

Combining (6.4), (6.5) and (6.6) we obtain:

$$\sup_{t/2 + C_\epsilon \leq s \leq t, t \geq T_\epsilon} \|\tau^{V,\text{int}}(s,t) \circ E_t(A) - \tau_\infty^{V,\text{int}}(s,t)(A)\| \leq \epsilon/2.$$

Using also (6.3) we obtain the proposition. \square

6.3. Left propagation. Recall that the subspaces $\mathfrak{h}_t^{1/r}$ were defined in (2.7).

Lemma 6.8. *We have $\text{Ran} w^1 = \mathfrak{h}_\infty^1$. It follows that $\gamma_\infty^1 : \text{CAR}(P^1 \mathfrak{h}_\infty) \xrightarrow{\sim} \text{CAR}(\mathfrak{h}_\infty^1)$ is a $*$ -isomorphism.*

Proof. Let us denote by \tilde{P} the asymptotic velocity for b_∞^1 . Since $V^1 = 0$ we have $b_\infty^1 = -LD_x$. If $x(t) = e^{-itb_\infty^1} x e^{itb_\infty^1}$, we see that $x(t) = -tL + x$, from which it follows that $\tilde{P} = -L$. It is well known (see e.g. [Da]) that the wave operator w^1 intertwines P and \tilde{P} , hence

$$\text{Ran} w^1 = \text{Ran} \mathbb{1}_{\mathbb{R}^-}(\tilde{P}) = \text{Ran} \mathbb{1}_{\mathbb{R}^+}(L) = \mathfrak{h}_\infty^1.$$

This completes the proof of the lemma. \square

Proposition 6.9. *Let $A = \gamma_\infty^{\text{int}}(B^1)$, $B^1 \in \text{CAR}(P^1 \mathfrak{h}_\infty)$. Then for any $\epsilon > 0$ there exist $C_\epsilon, T_\epsilon > 0$ such that*

$$\sup_{t \geq T_\epsilon} \|\tau^{V,\text{int}}(t/2 + C_\epsilon, t) \circ E_t(A) - \alpha^{t/2 - C_\epsilon} \circ \gamma_\infty^1(B^1)\| \leq \epsilon.$$

Proof. By Prop. 6.6 there exist $C_\epsilon, \tilde{T}_\epsilon$ such that

$$(6.7) \quad \sup_{t \geq \tilde{T}_\epsilon} \|\tau^{V,\text{int}}(t/2 + C_\epsilon, t) \circ E_t(A) - \tau_\infty^V(t/2 + C_\epsilon, t)(B^1)\| \leq \epsilon/2.$$

Moreover since $B^1 \in \text{CAR}(P^1 \mathfrak{h}_\infty)$, we have

$$\lim_{s \rightarrow +\infty} \tau_\infty^V(0, s)(B^1) - \tau_\infty^0(0, \sigma) \circ \gamma_\infty^1(B^1) = 0,$$

by Lemma 6.8. We now note that $\tau_\infty^V, \tau_\infty^0$ are stationary, $\gamma_\infty^1(B^1) \in \text{CAR}(\mathfrak{h}_\infty^1)$, and $\tau_\infty^0(0, s) = \alpha^s$ on $\text{CAR}(\mathfrak{h}_\infty^1)$, since $u_\infty^0(0, s)f = f^s$ for $s \in \mathfrak{h}_\infty^1$, by (2.8). It follows that we can find $T_\epsilon \geq \tilde{T}_\epsilon$ such that

$$\|\tau^{V,\text{int}}(t/2 + C_\epsilon, t) \circ E_t(A) - \alpha^{t/2 - C_\epsilon} \circ \gamma_\infty^1(B^1)\| \leq \epsilon,$$

for $t \geq T_\epsilon$. This completes the proof of the proposition. \square

6.4. Right propagation. The following lemma means that for an initial observable propagating to the right, the influence of the boundary condition on the star can be forgotten.

Lemma 6.10. *For any $c > 0$ and $B^r \in \text{CAR}(P^r \mathfrak{h}_\infty)$ one has:*

$$\tau^V(0, t/2 + c) \circ \tau_\infty^V(t/2 + c, t)(B^r) = \tau_\infty^V(0, t)(B^r) + o(t^0), \quad t \rightarrow +\infty.$$

Proof. By the usual arguments of linearity and density it suffices to prove that

$$(6.8) \quad u^V(0, t/2 + c) \circ u_\infty^V(t/2 + c, t)f = u_\infty^V(0, t)f + o(t^0), \quad f \in \text{Dom } b_\infty^V \cap P^r \mathfrak{h}_\infty, \quad t \rightarrow +\infty.$$

Using that $u_\infty^V(s, t) = e^{i(s-t)b_\infty^V}$ we can moreover assume that there exists $\epsilon > 0$ such that:

$$(6.9) \quad \begin{aligned} u_\infty^V(0, t)f &= \chi\left(\frac{x}{t} \geq \epsilon\right)u_\infty^V(0, t)f + o(t^0), \\ u_\infty^V(t/2 + c, t)f &= \chi\left(\frac{x}{t} \geq \epsilon\right)u_\infty^V(t/2 + c, t)f + o(t^0). \end{aligned}$$

To simplify notation let us set

$$f_{s,t} = \chi\left(\frac{x}{t} \geq \epsilon\right)u_\infty^V(s, t)f.$$

From (6.9) we obtain

$$\begin{aligned} & \|u^V(0, t/2 + c) \circ u_\infty^V(t/2 + c, t)f - u_\infty^V(0, t)f\| \\ &= \|u^V(0, t/2 + c)f_{t/2+c,t} - f_{0,t}\| + o(t^0) \\ &= \|f_{t/2+c,t} - u^V(t/2 + c, 0)f_{0,t}\| + o(t^0) \\ &\leq \left\| \int_0^{t/2+c} \partial_s u^V(t/2 + c, s)f_{s,t} ds \right\| + o(t^0). \end{aligned}$$

We have

$$\partial_s u^V(t/2 + c, s)f_{s,t} = -iu^V(t/2 + c, s) \left(b_s^V \chi\left(\frac{x}{t} \geq \epsilon\right) - \chi\left(\frac{x}{t} \geq \epsilon\right)b_\infty^V \right) u_\infty^V(s, t)f.$$

Since $f \in \text{Dom } b_\infty^V = H^1(\mathbb{R}, \mathbb{C}^2)$, we know that $f_{s,t} \in H^1(\mathbb{R}, \mathbb{C}^2)$. Since moreover $f_{s,t}$ is equal to 0 in $x \leq \epsilon t$, we have $f_{s,t} \in \text{Dom } b_s^V$ for $0 \leq s \leq t/2 + c$ and

$$\begin{aligned} & (b_s^V \chi\left(\frac{x}{t} \geq \epsilon\right) - \chi\left(\frac{x}{t} \geq \epsilon\right)b_\infty^V) u_\infty^V(s, t)f \\ &= [b_\infty^V, \chi\left(\frac{x}{t} \geq \epsilon\right)]u_\infty^V(0, t)f = \frac{1}{t}L\chi'\left(\frac{x}{t} \geq \epsilon\right)u_\infty^V(0, t)f. \end{aligned}$$

It follows that

$$(6.10) \quad \begin{aligned} & \left\| \int_0^{t/2+c} \partial_s u^V(t/2 + c, s)f_{s,t} ds \right\| \leq \frac{1}{t} \int_0^{t/2+c} \|\chi\left(\frac{x}{t} \simeq \epsilon\right)u_\infty^V(s, t)f\| ds \\ & \leq \frac{1}{t} \int_{t/2-c}^t \|\chi\left(\frac{x}{t} \simeq \epsilon\right)e^{-i\tau b_\infty^V} f\| d\tau, \end{aligned}$$

setting $\tau = t - s$. Denoting by $R_t(f)$ the r.h.s. in (6.10) we have $\|R_t(f)\| \leq C\|f\|$, $f \in \mathfrak{h}_\infty$ and from (6.1) $R_t(f) \in O(t^{-\infty})$ for $f \in \mathcal{D}$ hence $R_t(f) \in o(t^0)$ for any $f \in \mathfrak{h}_\infty$. This proves (6.8) and completes the proof of the lemma. \square

Proposition 6.11. *Let $A = \gamma_\infty^{\text{int}}(B^r)$, $B^r \in \text{CAR}(P^r \mathfrak{h}_\infty)$. Then for any $\epsilon > 0$ there exists $T_\epsilon > 0$ such that*

$$\sup_{t \geq T_\epsilon} \|\tau^{V,\text{int}}(0, t) \circ E_t(A) - \tau_\infty^V(0, t)(B^r)\| \leq \epsilon.$$

Proof. Let $\epsilon > 0$. By Prop. 6.6 there exist $C_\epsilon, \tilde{T}_\epsilon$ such that

$$(6.11) \quad \begin{aligned} \tau^{V,\text{int}}(0, t) \circ E_t(A) &= \tau^{V,\text{int}}(0, t/2 + C_\epsilon) \circ \tau^{V,\text{int}}(t/2 + C_\epsilon, t) \circ E_t(A) \\ &= \tau^{V,\text{int}}(0, t/2 + C_\epsilon) \circ \tau_\infty^V(t/2 + C_\epsilon, t)(B^r) + R_\epsilon(t), \end{aligned}$$

where $\|R_\epsilon(t)\| \leq \epsilon/4$ for $t \geq \tilde{T}_\epsilon$. Recall that the dense subspace $\mathcal{D} \subset \mathfrak{h}_\infty$ was introduced in the proof of Thm. 6.5. Since $B^r \in \text{CAR}(P^r \mathfrak{h}_\infty)$, we can by density

find $\tilde{B}^r \in \text{CAR}_{\text{alg}}(\mathcal{D} \cap P^r \mathfrak{h}_\infty)$ such that $\|B^r - \tilde{B}^r\| \leq \epsilon/4$. It follows then from (6.11) that

$$(6.12) \quad \sup_{t \geq \tilde{T}_\epsilon} \|\tau^{V,\text{int}}(0, t) \circ E_t(A) - \tau^{V,\text{int}}(0, t/2 + C_\epsilon) \circ \tau_\infty^V(t/2 + C_\epsilon, t)(\tilde{B}^r)\| \leq \epsilon/2.$$

Next we have

$$\begin{aligned} & \tau^{V,\text{int}}(0, t/2 + C_\epsilon) \circ \tau_\infty^V(t/2 + C_\epsilon, t)(\tilde{B}^r) \\ &= R(0, t/2 + C_\epsilon) \tau_\infty^V(0, t)(\tilde{B}^r) R(0, t/2 + C_\epsilon)^* + o(t^0), \end{aligned}$$

by Lemma 6.10 and the definition of $\tau^{V,\text{int}}$ in Def. 4.4. It follows that

$$\begin{aligned} & \tau^{V,\text{int}}(0, t/2 + C_\epsilon) \circ \tau_\infty^V(t/2 + C_\epsilon, t)(\tilde{B}^r) \\ &= \tau_\infty^V(0, t)(\tilde{B}^r) - \int_0^{t/2 + C_\epsilon} \partial_s \left(R(s, t/2 + C_\epsilon) \tau_\infty^V(0, t)(\tilde{B}^r) R(s, t/2 + C_\epsilon)^* \right) ds. \end{aligned}$$

Now

$$\begin{aligned} & \partial_s \left(R(s, t/2 + C_\epsilon) \tau_\infty^V(0, t)(\tilde{B}^r) R(s, t/2 + C_\epsilon)^* \right) \\ &= iR(s, t/2 + C_\epsilon) [I(s, t/2 + C_\epsilon), \tau_\infty^V(0, t)(\tilde{B}^r)] R(s, t/2 + C_\epsilon)^*. \end{aligned}$$

It remains to bound the norm of $[I(s, t/2 + C_\epsilon), \tau_\infty^V(0, t)(\tilde{B}^r)]$, since $R(\cdot, \cdot)$ is unitary. This amounts again to estimate scalar products of the form

$$(u^V(s, t/2 + C_\epsilon)g | u_\infty^V(0, t)f),$$

for g compactly supported, $f \in \mathcal{D} \cap P^r \mathfrak{h}_\infty$. We know that $u^V(s, t/2 + C_\epsilon)g$ is supported in $\{|x| \leq t/2 + C_\epsilon + C_0\}$ for $s \leq t/2 + C_\epsilon$ and $C_0 > 0$, since g has compact support. On the other hand if $f \in \mathcal{D} \cap P^r \mathfrak{h}_\infty$ we have by (6.1)

$$\|\mathbb{1}_{[0, c_0]} \left(\frac{|x|}{t} \right) u_\infty^V(0, t)f\| \in O(t^{-N}), \quad \forall N \in \mathbb{N}.$$

It follows that

$$\begin{aligned} & \|\tau^{V,\text{int}}(0, t/2 + C_\epsilon) \circ \tau_\infty^V(t/2 + C_\epsilon, t)(\tilde{B}^r) - \tau_\infty^V(0, t)(\tilde{B}^r)\| \\ & \leq D_\epsilon |t/2 + C_\epsilon| \langle t \rangle^{-N}, \end{aligned}$$

hence by (6.12) there exists $T_\epsilon \geq \tilde{T}_\epsilon$ such that

$$\sup_{t \geq T_\epsilon} \|\tau^{V,\text{int}}(0, t) \circ E_t(A) - \tau_\infty^V(0, t)(\tilde{B}^r)\| \leq 3\epsilon/4.$$

Since $\|B^r - \tilde{B}^r\| \leq \epsilon/4$, this completes the proof of the proposition. \square

We conclude this subsection by stating two easy scattering results for free dynamics with boundary conditions.

Lemma 6.12. *The limit*

$$s\text{-}\lim_{t \rightarrow +\infty} e^{itb_0^V} e^{-itb_\infty^V} \text{ exists on } P^r \mathfrak{h}_\infty.$$

Moreover the above limit is unitary from $P^r \mathfrak{h}_\infty$ to \mathfrak{h}_0 .

Proof. the proof follows by standard arguments (note that $e^{-itb_\infty^V} f$ propagates to the right when $f \in P^r \mathfrak{h}_\infty$, hence the boundary condition at $x = z(0)$ is irrelevant). \square

Lemma 6.12 immediately implies the following proposition:

Proposition 6.13. *The limit*

$$\gamma_0^r := s\text{-}\lim_{t \rightarrow +\infty} \tau_0^V(t, 0) \circ \tau_\infty^V(0, t) \text{ exists on } \text{CAR}(P^r \mathfrak{h}_\infty),$$

and $\gamma_0^r : \text{CAR}(P^r \mathfrak{h}_\infty) \xrightarrow{\sim} \text{CAR}(\mathfrak{h}_0)$ is a $*$ -isomorphism.

Combining Props. 6.11 and 6.13 we obtain the following proposition, which is the main result of this subsection.

Proposition 6.14. *Let $A = \gamma_\infty^{\text{int}}(B^r)$, $B^r \in \text{CAR}(P^r\mathfrak{h}_\infty)$. Then for any $\epsilon > 0$ there exists $T_\epsilon > 0$ such that*

$$\sup_{t \geq T_\epsilon} \|\tau^{V,\text{int}}(0, t) \circ E_t(A) - \tau_0^V(0, t) \circ \gamma_0^r(B^r)\| \leq \epsilon.$$

6.5. Hawking effect II.

6.5.1. *The limit state.* Before stating our main result on the Hawking effect, let us introduce some notation. Recall that $\gamma_\infty^1 : \text{CAR}(P^1\mathfrak{h}_\infty) \xrightarrow{\sim} \text{CAR}(\mathfrak{h}_\infty^1)$ defined in Prop. 6.4 is a $*$ -isomorphism. Similarly $\gamma_0^r : \text{CAR}(P^r\mathfrak{h}_\infty) \xrightarrow{\sim} \text{CAR}(\mathfrak{h}_0)$ defined in Prop. 6.13 is a $*$ -isomorphism.

Lemma 6.15. *Let us denote by $\omega_{\infty,\beta}^1$ the state on $\text{CAR}(P^1\mathfrak{h}_\infty)$ equal to*

$$\omega_{\infty,\beta}^1 := \omega_{\infty,\beta}^0 \circ \gamma_\infty^1,$$

and by $\omega_{\infty,\text{vac}}^r$ the state on $\text{CAR}(P^r\mathfrak{h}_\infty)$ equal to

$$\omega_{\infty,\text{vac}}^r := \omega_{0,\text{vac}}^V \circ \gamma_0^r.$$

Then

(1) $\omega_{\infty,\beta}^1$ is the restriction to $\text{CAR}(P^1\mathfrak{h}_\infty)$ of the quasi-free thermal state on $\text{CAR}(\mathfrak{h}_\infty)$ with covariance

$$(f | (\mathbb{1} + e^{-\beta b_\infty^V})^{-1} f), \quad f \in \mathfrak{h}_\infty.$$

(2) $\omega_{\infty,\text{vac}}^r$ is the restriction to $\text{CAR}(P^r\mathfrak{h}_\infty)$ of the quasi-free vacuum state on $\text{CAR}(\mathfrak{h}_\infty)$ with covariance

$$(f | \mathbb{1}_{\mathbb{R}^+}(b_\infty^V) f), \quad f \in \mathfrak{h}_\infty.$$

Proof. (1) follows from the fact that γ_∞^1 is implemented by the wave operator w^1 defined in Prop. 6.3, which intertwines b_∞^V and b_∞^0 . Similarly (2) follows from the intertwining properties of the wave operator constructed in Lemma 6.12. \square

Note that $\omega_{\infty,\beta}^1$ and $\omega_{\infty,\text{vac}}^r$ are even states. Since $\mathfrak{h}_\infty = P^1\mathfrak{h}_\infty \oplus P^r\mathfrak{h}_\infty$, we can define the following state, acting on $\gamma_\infty^{\text{int}} \text{CAR}(\mathfrak{h}_\infty)$:

Definition 6.16. *We set:*

$$\omega_{\text{H,II}} := (\omega_{\infty,\beta}^1 \widehat{\otimes} \omega_{\infty,\text{vac}}^r) \circ (\gamma_\infty^{\text{int}})^{-1},$$

which is a state on $\gamma_\infty^{\text{int}} \text{CAR}(\mathfrak{h}_\infty)$.

Remark 6.17. *Note that the limit state $\omega_{\text{H,II}}$ is a priori only defined on the C^* -algebra $\gamma_\infty^{\text{int}} \text{CAR}(\mathfrak{h}_\infty)$ and not on the whole of $\text{CAR}(\mathfrak{h}_\infty)$. One can of course assume the asymptotic completeness of the wave morphism $\gamma_\infty^{\text{int}}$:*

$$(AC) \quad \gamma_\infty^{\text{int}} \mathfrak{A}_\infty = \mathfrak{A}_\infty,$$

in which case the statement of Thm. 6.18 below simplifies.

To our knowledge, the asymptotic completeness is an essentially open question in the algebraic setting. For example we do not know of any argument which would ensure that the generator of the interacting dynamics $\tau_\infty^{V,\text{int}}(s, t)$ has no eigenvalues.

If we fix a state on \mathfrak{A}_∞ , like for example the vacuum state for τ_∞^V , and work in its GNS representation, replacing C^* -algebras by their weak closures, then by the same arguments as in Subsect. 5.5, the dynamics τ_∞^V and $\tau_\infty^{V,\text{int}}$ are implemented by unitary groups with selfadjoint generators H_0 and $H = H_0 + \pi_F(I)$. The (Hilbertian) scattering theory for the pair H_0, H is not trivial, but nevertheless completely understood, see e.g. [A]. In particular the Hilbertian version of asymptotic completeness was shown in [A]. However these Hilbertian results are of no use for the algebraic setting.

6.5.2. *Main result II.* The following theorem is the main result of this section.

Theorem 6.18. *The following holds:*

(1)

$$\lim_{t \rightarrow +\infty} \omega_{0,\text{vac}}^V \circ \tau^{V,\text{int}}(0,t) \circ E_t(A) = \omega_{\text{H,II}}(A), \quad A \in \gamma_\infty^{\text{int}} \mathfrak{A}_\infty.$$

(2) *Assume moreover that $\gamma_\infty^{\text{int}} \mathfrak{A}_\infty = \mathfrak{A}_\infty$. Then*

$$\lim_{t \rightarrow +\infty} \omega_{0,\text{vac}}^V \circ \tau^{V,\text{int}}(0,t)(A) = \omega_{\text{H,II}}(A), \quad A \in \mathfrak{A}_J, \quad \forall J \in \mathbb{R}.$$

Proof. Let us first prove (1). By linearity and density, it suffices to prove the theorem for

$$A = A^1 \times A^r, \quad A^{1/r} = \gamma_\infty^{\text{int}}(B^{1/r}), \quad B^{1/r} \in \text{CAR}(P^{1/r} \mathfrak{h}_\infty).$$

Let us fix $\epsilon > 0$. By Props. 6.9 and 6.14 there exist $C_\epsilon, T_\epsilon > 0$ such that

(6.13)

$$\sup_{t \geq T_\epsilon} \|\tau^{V,\text{int}}(0,t) \circ E_t(A^1) - \tau^{V,\text{int}}(0,t/2 + C_\epsilon) \circ \alpha^{t/2+C_\epsilon} \circ \alpha^{-2C_\epsilon} \circ \gamma_\infty^1(B^1)\| \leq \epsilon,$$

$$\sup_{t \geq T_\epsilon} \|\tau^{V,\text{int}}(0,t) \circ E_t(A^r) - \tau_0^V(0,t) \circ \gamma_0^r(B^r)\| \leq \epsilon.$$

We set $\tilde{B}_\epsilon^1 := \alpha^{-2C_\epsilon} \circ \gamma_\infty^1(B^1)$ and $\tilde{B}^r := \gamma_0^r(B^r)$. By Lemma 5.2 we can by increasing T_ϵ ensure that

$$\sup_{t \geq T_\epsilon} \|\tau^{V,\text{int}}(0,t/2 + C_\epsilon) \circ \alpha^{t/2+C_\epsilon}(\tilde{B}_\epsilon^1) - \tau^V(0,t/2 + C_\epsilon) \circ \alpha^{t/2+C_\epsilon}(\tilde{B}^1)\| \leq \epsilon.$$

Summarizing we have:

$$(6.14) \quad \sup_{t \geq T_\epsilon} \|\tau^{V,\text{int}}(0,t) \circ E_t(A) - \tau^V(0,t/2 + C_\epsilon) \circ \alpha^{t/2+C_\epsilon}(\tilde{B}_\epsilon^1) \times \tau_0^V(0,t)(\tilde{B}^r)\| \leq C\epsilon.$$

We now argue as in the proof of Lemma 5.7 to obtain that:

$$(6.15) \quad \lim_{s \rightarrow +\infty} \omega_{0,\text{vac}}^V \left(\tau^V(0,s) \circ \alpha^s(\tilde{B}_\epsilon^1) \times \tau_0^V(0,s)(\tilde{B}^r) \right) = \omega_{\infty,\beta}^0 \widehat{\otimes} \omega_{0,\text{vac}}^V(\tilde{B}_\epsilon^1 \times \tilde{B}^r).$$

To prove (6.15) we use that $\omega_{0,\text{vac}}^V$ is quasi-free, and the dynamics in (6.15) are free. The cross terms of the form:

$$(u^V(0,s) f_1^s | \mathbb{1}_{\mathbb{R}^+} (b_0^V) u_0^V(0,t) f_2) \text{ for } f_1 \in \mathfrak{h}_0^1, \quad f_2 \in \mathfrak{h}_0^r,$$

vanish when $s \rightarrow +\infty$. This is easy to see, since modulo errors which are $o(s^0)$ in norm, the vector $u^V(0,s) f_1^s$ is supported in $\{|x| \leq c_0\}$ for s large enough, while the vector $u_0^V(0,s) f_2$ is supported in $\{x \geq c_1 s\}$.

The rest of the proof is as in Lemma 5.7, using also that $\omega_{0,\text{vac}}^V$ is invariant under τ_0^V . A further observation is that the state $\omega_{\infty,\beta}^0$ is invariant under space translations. Since $\tilde{B}_\epsilon^1 = \alpha^{-2C_\epsilon} \circ \gamma_\infty^1(B^1)$, this implies that

$$\begin{aligned} \omega_{\infty,\beta}^0 \widehat{\otimes} \omega_{0,\text{vac}}^V(\tilde{B}_\epsilon^1 \times \tilde{B}^r) &= \omega_{\infty,\beta}^0 \widehat{\otimes} \omega_{0,\text{vac}}^V(\gamma_\infty^1(B^1) \times \gamma_0^r(B^r)) \\ &= \omega_{\infty,\beta}^1 \widehat{\otimes} \omega_{\infty,\text{vac}}^r(B^1 \times B^r). \end{aligned}$$

Therefore we can rewrite (6.15) as

$$\lim_{s \rightarrow +\infty} \omega_{0,\text{vac}}^V \left(\tau^V(0,s) \circ \alpha^s(\tilde{B}_\epsilon^1) \times \tau_0^V(0,s)(\tilde{B}^r) \right) = \omega_{\infty,\beta}^1 \widehat{\otimes} \omega_{\infty,\text{vac}}^r(B^1 \times B^r).$$

Using also (6.14) this completes the proof of (1). Statement (2) follows from (1), since if $A \in \mathfrak{A}_J$ for some $J \in \mathbb{R}$ then $A = E_t(A)$ for t large enough. \square

6.6. Change of initial state. As in Subsect. 5.4 one can try to replace the initial state $\omega_{0,\text{vac}}^V$ by another (even) state $\tilde{\omega}$ which belongs to the folium of $\omega_{0,\text{vac}}^V$.

There is however a difference with the situation considered in Sect. 5: in Sect. 5 we have to consider the evolution of a right-going observable $A^r \in \mathfrak{A}_0^r$ under $\tau^{V,\text{int}}(0,t) \circ \alpha^t$ when $t \rightarrow +\infty$: this converges to the limit observable $\gamma^{r,\text{int}}(A^r)$, which implies that Thm. 5.6 extends to any such state $\tilde{\omega}$, see Corollary 5.8.

In the present situation, we have to consider the evolution of an observable $B^r \in \text{CAR}(\mathfrak{h}_0)$ under $\tau_0^V(0,t)$ (note that all observables in $\text{CAR}(\mathfrak{h}_0)$ are right-going, since $\mathfrak{h}_0 = L^2(\]z(0), +\infty[, \mathbb{C}^2)$). This has obviously no limit in $\text{CAR}(\mathfrak{h}_0)$. We need to restrict ourselves to initial states $\tilde{\omega}$ on \mathfrak{A}_0 which have the property that

$$(6.16) \quad \text{w-} \lim_{t \rightarrow +\infty} \tilde{\omega} \circ \tau_0^V(0,t) \text{ exists.}$$

Examples of such states are states which are *invariant* under the stationary interacting dynamics $\tau_0^{V,\text{int}}$, which were considered in Subsect. 5.4.

Let us now explain this in more details. We first recall some facts about the algebraic scattering theory in $\text{CAR}(\mathfrak{h}_0)$. We will use the notation introduced in Subsect. 5.5. It is easy to prove that the limit:

$$\gamma_0^{\text{int}} := \text{s-} \lim_{t \rightarrow +\infty} \tau_0^{V,\text{int}}(t,0) \circ \tau_0^V(0,t)$$

exists on $\text{CAR}(\mathfrak{h}_0)$. From Prop. 6.13 and the chain rule for wave homomorphisms, we obtain the existence of the limit

$$\gamma_0^{r,\text{int}} := \gamma_0^{\text{int}} \circ \gamma_0^r = \text{s-} \lim_{t \rightarrow +\infty} \tau_0^{V,\text{int}}(t,0) \circ \tau_\infty^V(0,t) \text{ on } \text{CAR}(P^r \mathfrak{h}_\infty).$$

We obtain the following analog of Corollary 5.8.

Corollary 6.19. *Let $\tilde{\omega}$ be an even state on \mathfrak{A}_0 , which belongs to the folium of $\omega_{0,\text{vac}}^V$ and is invariant under $\tau_0^{V,\text{int}}$. Let:*

$$\tilde{\omega}_{\text{H,II}} = (\omega_{\infty,\beta}^1 \hat{\otimes} (\tilde{\omega} \circ \gamma_0^{r,\text{int}})) \circ (\gamma_\infty^{\text{int}})^{-1}.$$

Then:

(1)

$$\lim_{t \rightarrow +\infty} \tilde{\omega} \circ \tau^{V,\text{int}}(0,t) \circ E_t(A) = \tilde{\omega}_{\text{H,II}}(A), \quad A \in \gamma_\infty^{\text{int}} \mathfrak{A}_\infty.$$

(2) *Assume moreover that $\gamma_\infty^{\text{int}} \mathfrak{A}_\infty = \mathfrak{A}_\infty$. Then*

$$\lim_{t \rightarrow +\infty} \tilde{\omega} \circ \tau^{V,\text{int}}(0,t)(A) = \tilde{\omega}_{\text{H,II}}(A), \quad A \in \mathfrak{A}_J, \quad \forall J \in \mathbb{R}.$$

Note that we proved in Subsect. 5.5 that such states $\tilde{\omega}$ exist, at least for small interactions.

Proof. We will only sketch the proof, since it is an easy combination of the arguments in Thm. 6.18 and Corollary 5.8. From (6.14) we see that modulo an error of size ϵ , uniformly for $t \geq T_\epsilon$, we have to compute

$$\lim_{s \rightarrow +\infty} \tilde{\omega}(\tau^V(0,s) \circ \alpha^s(\tilde{B}_\epsilon^1) \times \tau_0^V(0,s)(\tilde{B}^r)).$$

We set $\tilde{B}^1(s) := \tau^V(0,s) \circ \alpha^s(\tilde{B}_\epsilon^1)$ and $\tilde{B}^r(s) := \tau_0^V(0,s)(\tilde{B}^r)$ to simplify notation. Since $\tilde{\omega}$ belongs to the folium of $\omega_{0,\text{vac}}^V$, we can find $P = P(\psi^*, \psi) \in \text{CAR}_{\text{alg}}(\mathfrak{h}_0)$ even, such that:

$$|\tilde{\omega}(B) - \omega_0^V(P^*BP)| \leq \epsilon \|B\|, \quad B \in \text{CAR}(\mathfrak{h}_0).$$

By the same argument as in the proof of Corollary 5.8 we have

$$\omega_0^V(P^* \tilde{B}^1(s) \tilde{B}^r(s) P) = \omega_0^V(\tilde{B}^1(s) P^* \tilde{B}^r(s) P) + o(s^0).$$

Again the cross terms vanish when $s \rightarrow +\infty$, the terms coming from $\tilde{B}^1(s)$ give in the limit $s \rightarrow +\infty$ the contribution $\omega_\infty^0(\tilde{B}^1)$ equal to $\omega_{\infty,\beta}^1(B^1)$.

The terms coming from $\tilde{B}^r(s)$ give modulo an error of size ϵ the contribution $\tilde{\omega}(\tau_0^V(0, s)(\tilde{B}^r))$. Now we use the hypothesis that $\tilde{\omega}$ is invariant under $\tau_0^{V, \text{int}}$ hence:

$$\begin{aligned} \tilde{\omega}(\tau_0^V(0, s)(\tilde{B}^r)) &= \tilde{\omega}(\tau_0^{V, \text{int}}(s, 0) \circ \tau_0^V(0, s)(\tilde{B}^r)) \\ &= \tilde{\omega}(\gamma_0^{\text{int}}(\tilde{B}^r)) + o(s^0) = \tilde{\omega}(\gamma_0^{r, \text{int}}(B^r)) + o(s^0). \end{aligned}$$

We can now complete the proof as in Thm. 6.18. \square

7. HAWKING EFFECT III

In this section we study the Hawking effect in case III (see Subsect. 1.3). As explained in the introduction, the interaction should now be localized in a region

$$\{(x, t) : z(t) < x < z(t) + C, T - 1 \leq t \leq T\},$$

and we will apply the interacting evolution to an observable $\alpha^{-z(T)}(A)$ for $A \in \mathfrak{A}_0$, letting eventually $T \rightarrow +\infty$. Let us now make this more precise.

7.1. Definition of the interacting dynamics. We fix $I \in \text{CAR}_0(\mathfrak{h}_0)$ as in (4.2) and set

$$I(t) := \alpha^t(I), \quad I_T(t) := \mathbb{1}_{[T-1, T]}(t)I(t),$$

where the group α^s of space translations is defined in (3.1), and $T \gg 1$ is a parameter which will eventually tend to $+\infty$. To be sure that $I(t) \in \mathfrak{A}_t$ we assume in this section that $z(t) \leq -t$ for all $t \geq 0$, which is not a restriction.

Definition 7.1. We denote by $\tilde{\tau}_T^{V, \text{int}}(s, t)$ the interacting dynamics (depending on the parameter T), constructed using Prop. A.11, with free dynamics $\tau^V(s, t)$ and time dependent interaction $\mathbb{R} \ni t \mapsto I_T(t)$.

Our goal in this section is to study the limit:

$$(7.1) \quad \lim_{T \rightarrow +\infty} \omega_{0, \text{vac}}^V(\tilde{\tau}_T^{V, \text{int}}(0, T) \circ \alpha^T(A)), \quad A \in \mathfrak{A}_0.$$

Since $I_T(t)$ vanishes for $0 \leq t \leq T - 1$ we have:

$$\tilde{\tau}_T^{V, \text{int}}(0, T - 1) = \tau^V(0, T - 1),$$

hence

$$(7.2) \quad \tilde{\tau}_T^{V, \text{int}}(0, T) \circ \alpha^T = \tau^V(0, T - 1) \circ \alpha^{T-1} \circ \alpha^{-(T-1)} \circ \tilde{\tau}_T^{V, \text{int}}(T - 1, T) \circ \alpha^T.$$

Applying Thm. 2.8, the existence of the limit (7.1) will follow from the existence of

$$(7.3) \quad \underset{T \rightarrow +\infty}{\text{s-}} \lim \alpha^{-(T-1)} \circ \tilde{\tau}_T^{V, \text{int}}(T - 1, T) \circ \alpha^T \text{ on } \mathfrak{A}_0.$$

Recall that

$$\gamma^t f(x) := f(x + t), \quad x, t \in \mathbb{R}, \quad f \in \mathfrak{h}.$$

Let us introduce the following notation:

$$(7.4) \quad \begin{aligned} \hat{u}_T^V(s, t) &:= \gamma^{-(T+s)} \circ u^V(T + s, T + t) \circ \gamma^{T+t} \in \mathcal{U}(\mathfrak{h}_0, \mathfrak{h}_0), \\ \hat{\tau}_T^V(s, t) &:= \alpha^{-(T+s)} \circ \tau^V(T + s, T + t) \circ \alpha^{T+t}, \quad \mathfrak{A}_0 \rightarrow \mathfrak{A}_0, \\ \hat{\tau}_T^{V, \text{int}}(s, t) &:= \alpha^{-(T+s)} \circ \tilde{\tau}_T^{V, \text{int}}(T + s, T + t) \circ \alpha^{T+t}, \quad \mathfrak{A}_0 \rightarrow \mathfrak{A}_0, \end{aligned}$$

so that the automorphism appearing in (7.3) equals $\hat{\tau}_T^{V, \text{int}}(-1, 0)$. Note that

$$(7.5) \quad \hat{\tau}_T^V(s, t)(\psi^{(*)}(f)) = \psi^{(*)}(\hat{u}_T^V(s, t)f), \quad f \in \mathfrak{h}_0,$$

and that $\{\hat{u}_T^V(s, t)\}_{s, t \in \mathbb{R}}$ is a two-parameter propagator, while $\{\hat{\tau}_T^V(s, t)\}_{s, t \in \mathbb{R}}$ and $\{\hat{\tau}_T^{V, \text{int}}(s, t)\}_{s, t \in \mathbb{R}}$ are two-parameter quantum dynamics.

7.2. Preparations. We start by considering the limit (7.3) for $I = 0$.

Lemma 7.2. *The strong limit*

$$\hat{u}_\infty^0(s, t) := s\text{-}\lim_{T \rightarrow +\infty} \hat{u}_T^0(s, t)$$

exists on \mathfrak{h}_0 and the convergence is uniform in $a \leq s \leq t \leq b$ for any $a \leq b$. Moreover $\{\hat{u}_\infty^0(s, t)\}_{s, t \in \mathbb{R}}$ is a two-parameter propagator given by

$$\hat{u}_\infty^0(s, t)f = \begin{pmatrix} \gamma^{2(t-s)} f_1 \\ f_2 \end{pmatrix}, \quad f \in \mathfrak{h}_0.$$

Remark 7.3. *The convergence above holds a priori only for $s \leq t$. Nevertheless the limit $\hat{u}_\infty^0(s, t)$ is defined for all $s, t \in \mathbb{R}$.*

Proof. It is easy to obtain an explicit expression for $\hat{u}_T^0(s, t)$. In fact from [Ba2, Lemme VI.3] we know that if $\psi(s, \cdot) = u^0(s, t)f(\cdot)$ for $f \in \mathfrak{h}_t$ then:

$$\begin{aligned} \psi_1(s, x) &= f_1(x - s + t) \\ \psi_2(s, x) &= \begin{cases} \lambda \circ \tau(x + s)^{-1} f_1(x + t + s - 2\tau(x + s)) & \text{for } z(s) < x < z(t) + t - s, \\ f_2(x - t + s) & \text{for } x > z(t) + t - s, \end{cases} \end{aligned}$$

where the reflection coefficient λ is defined in Subsect. 1.2, and the function $y \mapsto \tau(y)$ is the inverse of the function $s \mapsto s + z(s)$ (see [Ba2, Equ. VI.40]).

From this a routine computation gives that:

$$\begin{aligned} (\hat{u}_T^0(s, t)f)_1(x) &= f_1(x + 2(t - s)), \\ (\hat{u}_T^0(s, t)f)_2(x) &= \begin{cases} \lambda \circ \tau(x)^{-1} f_1(x + 2(T + t) - 2\tau(x)), \\ \text{for } \tilde{z}(T + s) < x < \tilde{z}(T + t), \\ f_2(x), & \text{for } x > \tilde{z}(T + t), \end{cases} \end{aligned}$$

where $\tilde{z}(\sigma) := \sigma + z(\sigma) \in o(\sigma^0)$, by (2.1). Using this fact and that f is compactly supported, we easily see that

$$\lim_{T \rightarrow +\infty} \hat{u}_T^0(s, t)f = \begin{pmatrix} \gamma^{2(t-s)} f_1 \\ f_2 \end{pmatrix},$$

uniformly for $a \leq s \leq t \leq b$. This completes the proof. \square

We now establish the same result for arbitrary V .

Lemma 7.4. *The limit*

$$s\text{-}\lim_{T \rightarrow +\infty} \hat{u}_T^V(s, t) = \hat{u}_\infty^0(s, t),$$

exists, and the convergence is uniform in $a \leq s \leq t \leq b$ for any $a \leq b$.

Proof. From Duhamel's formula we obtain that

$$\|u^V(T + s, T + t)f - u^0(T + s, T + t)f\| \leq \int_s^t \|Vu^0(T + \sigma, T + t)f\| d\sigma,$$

hence

$$\|\hat{u}_T^V(s, t)f - \hat{u}_T^0(s, t)f\| \leq \int_s^t \|Vu^0(T + \sigma, T + t)\gamma^{T+t}f\| d\sigma.$$

By the usual density argument we can assume that $\text{supp} f \subset [0, b]$. We deduce from this that $\text{supp} u^0(T + \sigma, T + t)\gamma^{T+t}f \subset [-T + o(T^0), b - T + o(T^0)]$, uniformly for $a \leq s \leq \sigma \leq t \leq b$. Using the decay property of V near $-\infty$, this implies that

$$\lim_{T \rightarrow +\infty} \sup_{-1 \leq s \leq t \leq 0} \|\hat{u}_T^V(s, t)f - \hat{u}_T^0(s, t)f\| = 0,$$

which completes the proof of the lemma. \square

By (7.5) we obtain

Proposition 7.5. *The strong limit*

$$\hat{\tau}_\infty^0(s, t) := s\text{-}\lim_{T \rightarrow +\infty} \hat{\tau}_T^V(s, t)$$

exists on \mathfrak{A}_0 and the convergence is uniform in $a \leq s \leq t \leq b$ for any $a \leq b$.

7.3. Hawking effect III.

Proposition 7.6. *Let $\hat{\tau}_\infty^{0, \text{int}}(s, t)$ be the interacting dynamics obtained from Prop. A.11 from the free dynamics $\hat{\tau}_\infty^0(s, t)$ and the interaction I . Then*

$$s\text{-}\lim_{T \rightarrow +\infty} \hat{\tau}_T^{V, \text{int}}(s, t) = \hat{\tau}_\infty^{0, \text{int}}(s, t),$$

and the convergence is uniform in $a \leq s \leq t \leq b$.

Proof. Let $R_T(s, t)$ the unitary operator in Prop. A.11 for the time-dependent interaction $I_T(\cdot)$. Then from (7.4), we see that:

$$\begin{aligned} & \hat{\tau}_T^{V, \text{int}}(s, t)(A) \\ &= \alpha^{-(T+s)} R_{T+s}(T+s, T+t) \times \hat{\tau}_T^V(s, t)(A) \times \alpha^{-(T+s)} R_{T+s}(T+s, T+t)^*, \quad A \in \mathfrak{A}_0. \end{aligned}$$

Using Prop. 7.5 it hence suffices to show that

$$R_\infty(s, t) := \lim_{T \rightarrow +\infty} \alpha^{-(T+s)} R_{T+s}(T+s, T+t) \text{ exists.}$$

By Lemma A.10, we have:

$$\begin{aligned} (7.6) \quad & \alpha^{-(T+s)} R_T(T+s, T+t) \\ &= \sum_{n \geq 0} (-i)^n \int_{T+s \leq t_n \leq \dots \leq t_1 \leq T+t} \alpha^{-(T+s)} I_T(T+s, t_n) \dots \alpha^{-(T+s)} I_T(T+s, t_1) dt_n \dots dt_1 \\ &= \sum_{n \geq 0} (-i)^n \int_{s \leq t_n \leq \dots \leq t_1 \leq t} \alpha^{-(T+s)} I_T(T+s, T+t_n) \dots \alpha^{-(T+s)} I_T(T+s, T+t_1) dt_n \dots dt_1 \end{aligned}$$

Note that:

$$\alpha^{-(T+s)} I_T(T+s, \sigma+T) = \mathbb{1}_{[-1, 0]}(s) \hat{\tau}_T^V(s, \sigma)(I).$$

By Prop. 7.5 we know that

$$\lim_{T \rightarrow +\infty} \hat{\tau}_T^V(s, \sigma)(I) = \hat{\tau}_\infty^0(s, \sigma)(I),$$

uniformly for $-1 \leq s \leq \sigma \leq 0$. Therefore using that the convergence of the series in (7.6) is uniform in T , we can pass to the limit under the sum and integrals, which range over compact regions. The limit

$$\hat{R}_\infty(s, t) = \lim_{T \rightarrow +\infty} \alpha^{-(T+s)} R_T(T+s, T+t)$$

equals the unitary operator obtained in Prop. A.11 from the free dynamics $\hat{\tau}_\infty^0(s, t)$ and the interaction I . Therefore

$$\lim_{T \rightarrow \infty} \hat{\tau}_T^{V, \text{int}}(T+s, T+t)(A) = \hat{R}_\infty(s, t) \times \hat{\tau}_\infty^0(s, t)(A) \times \hat{R}_\infty(s, t)^* = \hat{\tau}_\infty^{0, \text{int}}(s, t)(A).$$

This completes the proof of the proposition. \square

We can now formulate the main result of this section. We first define the limiting state. Let $\gamma^r = s\text{-}\lim_{t \rightarrow +\infty} \tau^V(0, t) \circ \alpha^t$ on \mathfrak{A}_0^r , obtained as in Prop. 5.3 if the interaction I vanishes. Note that γ^r is implemented by the classical wave operator $s\text{-}\lim_{t \rightarrow +\infty} u^V(0, t) \circ \gamma^t$ on \mathfrak{H}_0^r .

Let

$$\omega_{\text{H}}^{\text{free}} := \omega_{\infty, \beta}^0 \hat{\otimes} (\omega_{0, \text{vac}}^V \circ \gamma^r).$$

This is the limiting state obtained by Bachelot in [Ba2] in the case when the interaction vanishes.

Theorem 7.7.

$$\lim_{T \rightarrow \infty} \omega_{0, \text{vac}}^V \circ \tilde{\tau}_T^{V, \text{int}}(0, T) \circ \alpha^T(A) = \omega_{\mathbb{H}}^{\text{free}} \circ \hat{\tau}_{\infty}^{0, \text{int}}(-1, 0)(A), \quad A \in \mathfrak{A}_0.$$

Proof. The result follows from Prop. 7.6, formula (7.2) and Thm. 5.6 in the case $I = 0$, (which in this case was already obtained in [Ba2]). \square

A similar result can be obtained if we replace the initial state $\omega_{0, \text{vac}}^V$ by another state $\tilde{\omega}$ as in Corollary 5.8.

APPENDIX A

In this appendix we recall various well-known facts about CAR algebras, fermionic Fock spaces and groups of $*$ -isomorphisms on C^* -algebras.

A.1. CAR algebras.

A.1.1. *CAR algebras.* Let \mathfrak{h} be a (complex) Hilbert space, with scalar product denoted by $(\cdot | \cdot)$.

Definition A.1. The algebraic CAR algebra over \mathfrak{h} , denoted by $\text{CAR}_{\text{alg}}(\mathfrak{h})$ is the unital $*$ -algebra generated by the generators $\mathbb{1}$, $\psi(h)$, $h \in \mathfrak{h}$ and the relations:

$$\begin{aligned} \psi(h_1 + h_2) &= \psi(h_1) + \psi(h_2), \quad h_i \in \mathfrak{h}, \\ \psi(\lambda h) &= \bar{\lambda} \psi(h), \quad h \in \mathfrak{h}, \quad \lambda \in \mathbb{C}, \\ [\psi(h_1), \psi(h_2)]_+ &= 0, \quad [\psi(h_1), \psi^*(h_2)]_+ = (h_1 | h_2) \mathbb{1}. \end{aligned}$$

It is well known that $\text{CAR}_{\text{alg}}(\mathfrak{h})$ is simple, hence has a unique C^* -norm. A concrete expression for this norm can be obtained by taking the representation $\pi : \text{CAR}_{\text{alg}}(\mathfrak{h}) \rightarrow B(\Gamma_{\mathfrak{a}}(\mathfrak{h}))$ with $\pi(\psi^{(*)}(h)) := a^{(*)}(h)$ for $h \in \mathfrak{h}$, see Subsect. A.4 below. (This corresponds to the choice of $j = i$ as Kähler structure).

Definition A.2. The CAR C^* -algebra over \mathfrak{h} , denoted by $\text{CAR}(\mathfrak{h})$ is the completion of $\text{CAR}_{\text{alg}}(\mathfrak{h})$ for its unique C^* -norm.

An element of $\text{CAR}_{\text{alg}}(\mathfrak{h})$ can be written in a unique way in *normal ordered form*, i.e. with ψ^* 's to the left of ψ 's. This allows to define unambiguously the *monomials*. A monomial A has a degree denoted by $\text{deg } A$. Sometimes we will also use the *bi-degree* (n, p) , where n is the number of factors of ψ^* , p the number of factors of ψ .

A.1.2. *Parity.* $\text{CAR}(\mathfrak{h})$ is equipped with the *parity automorphism* P , defined by

$$P\psi^{(*)}(h) := \psi^{(*)}(-h), \quad h \in \mathfrak{h}.$$

We denote by $\text{CAR}_0(\mathfrak{h})$, resp. $\text{CAR}_1(\mathfrak{h})$ the subspace of even, resp. odd elements of $\text{CAR}(\mathfrak{h})$. $\text{CAR}_0(\mathfrak{h})$ is a C^* -sub-algebra of $\text{CAR}(\mathfrak{h})$.

A state ω on $\text{CAR}(\mathfrak{h})$ is *even* if $\omega \circ P = \omega$, or equivalently $\omega = 0$ on $\text{CAR}_1(\mathfrak{h})$.

A $*$ -automorphism α of $\text{CAR}(\mathfrak{h})$ is *even* if $P \circ \alpha = \alpha \circ P$.

A.1.3. *Conditional expectations.* If \mathfrak{h}_1 is a closed subspace of \mathfrak{h} , then $\text{CAR}(\mathfrak{h}_1)$ is a C^* -subalgebra of $\text{CAR}(\mathfrak{h})$. The converse construction is as follows: define $E_{\mathfrak{h}_1} : \text{CAR}_{\text{alg}}(\mathfrak{h}) \rightarrow \text{CAR}_{\text{alg}}(\mathfrak{h}_1)$ by

$$E_u \psi^{(*)}(f) := \psi^{(*)}(\pi f), \quad f \in \mathfrak{h},$$

where $\pi : \mathfrak{h} \rightarrow \mathfrak{h}_1$ is the orthogonal projection. Then $E_{\mathfrak{h}_1}$ extends as a $*$ -homomorphism from $\text{CAR}(\mathfrak{h})$ to $\text{CAR}(\mathfrak{h}_1)$. This can be easily checked by using the Fock representations of $\text{CAR}(\mathfrak{h})$ (resp. $\text{CAR}(\mathfrak{h}_1)$) on $\Gamma_{\mathfrak{a}}(\mathfrak{h})$ (resp. $\Gamma_{\mathfrak{a}}(\mathfrak{h}_1)$) and the second quantized

map $\Gamma(\pi)$. Moreover if $\{\mathfrak{h}_i\}_{i \in I}$ is an increasing net of closed subspaces of \mathfrak{h} with $\cup_{i \in I} \mathfrak{h}_i$ dense in \mathfrak{h} then

$$\text{s-}\lim_i E_{\mathfrak{h}_i} = \mathbb{1}, \text{ in } \text{CAR}(\mathfrak{h}).$$

A.2. Fermionic exponential law.

A.2.1. \mathbb{Z}_2 -graded tensor product. Let \mathfrak{h}_i , $i = 1, 2$ be two Hilbert spaces. We equip the vector space $\text{CAR}_{\text{alg}}(\mathfrak{h}_1) \otimes \text{CAR}_{\text{alg}}(\mathfrak{h}_2)$ with the $*$ -algebra structure defined by:

$$(A.1) \quad \begin{aligned} (a_1 \otimes a_2) \cdot (b_1 \otimes b_2) &:= (-1)^{\deg a_2 \deg b_1} a_1 b_1 \otimes a_2 b_2, \\ (a_1 \otimes a_2)^* &:= (-1)^{\deg a_1 \deg a_2} a_1^* \otimes a_2^*, \end{aligned}$$

for a_i, b_i monomials in $\text{CAR}_{\text{alg}}(\mathfrak{h}_i)$, and extended to $\text{CAR}_{\text{alg}}(\mathfrak{h}_1) \otimes \text{CAR}_{\text{alg}}(\mathfrak{h}_2)$ by linearity. The resulting $*$ -algebra is denoted by $\text{CAR}_{\text{alg}}(\mathfrak{h}_1) \widehat{\otimes} \text{CAR}_{\text{alg}}(\mathfrak{h}_2)$.

Lemma A.3. *The map*

$$\begin{aligned} I : \text{CAR}_{\text{alg}}(\mathfrak{h}_1 \oplus \mathfrak{h}_2) &\rightarrow \text{CAR}_{\text{alg}}(\mathfrak{h}_1) \widehat{\otimes} \text{CAR}_{\text{alg}}(\mathfrak{h}_2) \\ \psi^{(*)}(h_1 \oplus h_2) &\mapsto \psi^{(*)}(h_1) \otimes \mathbb{1} + \mathbb{1} \otimes \psi^{(*)}(h_2) \end{aligned}$$

extends as a $$ -isomorphism.*

Remark A.4. *If $u : \mathfrak{h} \rightarrow \tilde{\mathfrak{h}}$ is an isometry then the map $\psi^{(*)}(h) \mapsto \psi^*(uh)$ extends to an $*$ -homomorphism from $\text{CAR}(\mathfrak{h})$ to $\text{CAR}(\tilde{\mathfrak{h}})$. This allows to see $\text{CAR}(\mathfrak{h}_i)$ $i = 1, 2$ as $*$ -sub-algebras of $\text{CAR}(\mathfrak{h}_1 \oplus \mathfrak{h}_2)$. If $A_i \in \text{CAR}(\mathfrak{h}_i)$ let us still denote by A_i its image in $\text{CAR}(\mathfrak{h}_1 \oplus \mathfrak{h}_2)$. Then clearly we have*

$$I(A_1 A_2) = A_1 \otimes A_2, \text{ for } A_i \in \text{CAR}(\mathfrak{h}_i).$$

Definition A.5. *The \mathbb{Z}_2 -graded tensor product $\text{CAR}(\mathfrak{h}_1) \widehat{\otimes} \text{CAR}(\mathfrak{h}_2)$ is the completion of $\text{CAR}_{\text{alg}}(\mathfrak{h}_1) \widehat{\otimes} \text{CAR}_{\text{alg}}(\mathfrak{h}_2)$ for the norm $\|I^{-1} \cdot\|$.*

A.2.2. *Tensor product of states and automorphisms.*

Lemma A.6. *Let ω_i be a state on $\text{CAR}(\mathfrak{h}_i)$, $i = 1, 2$. Assume that ω_1 is even, i.e. $\omega_1 \circ P_1 = \omega_1$. Then $\omega_1 \otimes \omega_2$ is a state on $\text{CAR}(\mathfrak{h}_1) \widehat{\otimes} \text{CAR}(\mathfrak{h}_2)$.*

Proof. It suffices to check positivity. If $A = \sum_1^n \lambda_i a_{1i} \otimes a_{2i}$, where a_{ki} are monomials, $\lambda_i \in \mathbb{C}$, then from (A.1) we get that:

$$A^* A = \sum_{i,j} \bar{\lambda}_i \lambda_j (-1)^{d_{2i}(d_{1i}+d_{1j})} a_{1i}^* a_{1j} \otimes a_{2i}^* a_{2j},$$

for $d_{ki} = \deg a_{ki}$. Since ω_1 is even, we obtain that:

$$\omega(A^* A) = \sum_{i,j} \bar{\lambda}_i \lambda_j \omega_1(a_{1i}^* a_{1j}) \omega_2(a_{2i}^* a_{2j}).$$

The positivity follows from the well-known fact that the pointwise product of two positive selfadjoint matrices is positive selfadjoint. \square

Definition A.7. *Let ω_i be a state on $\text{CAR}(\mathfrak{h}_i)$, $i = 1, 2$ with ω_1 even. The \mathbb{Z}_2 -graded tensor product $\omega_1 \widehat{\otimes} \omega_2$ is the state on $\text{CAR}(\mathfrak{h}_1 \oplus \mathfrak{h}_2)$ equal to $\omega_1 \otimes \omega_2 \circ I$.*

Similarly one easily check that if α_i , $i = 1, 2$ are even $*$ -automorphisms of $\text{CAR}(\mathfrak{h}_i)$ then $\alpha_1 \otimes \alpha_2$ is a $*$ -automorphism of $\text{CAR}(\mathfrak{h}_1) \widehat{\otimes} \text{CAR}(\mathfrak{h}_2)$.

Definition A.8. *Let α_i $i = 1, 2$ be even $*$ -automorphisms of $\text{CAR}(\mathfrak{h}_i)$. The \mathbb{Z}_2 -graded tensor product $\alpha_1 \widehat{\otimes} \alpha_2$ is the $*$ -automorphism of $\text{CAR}(\mathfrak{h}_1 \oplus \mathfrak{h}_2)$ equal to $I^{-1} \circ (\alpha_1 \otimes \alpha_2) \circ I$.*

A.3. Quasi-free states. We now recall some well-known facts on quasi-free states.

Definition A.9. A state ω on $\text{CAR}(\mathfrak{h})$ is a (gauge-invariant) quasi-free state if

$$\begin{aligned} \omega\left(\prod_{i=1}^n \psi^*(f_i) \prod_{i=1}^p \psi(g_i)\right) &= 0, \text{ for } n \neq p, \\ \omega\left(\prod_{i=1}^n \psi^*(f_i) \prod_{i=1}^n \psi(g_i)\right) &= \sum_{\sigma \in S_n} \epsilon(\sigma) \prod_{i=1}^n \omega(\psi^*(f_i) \psi(g_{\sigma(i)})). \end{aligned}$$

The bounded selfadjoint operator c on \mathfrak{h} defined by

$$\omega(\psi^*(f) \psi(g)) =: (g|cf)_{\mathfrak{h}}, \quad f, g \in \mathfrak{h}$$

is called the covariance of ω .

It is well-known that a necessary and sufficient condition for a selfadjoint operator c to be the covariance of a quasi-free state is:

$$(A.2) \quad 0 \leq c \leq \mathbb{1}.$$

A.4. Fermionic Fock spaces. Let \mathcal{Z} be a complex Hilbert space. The *fermionic Fock space* over \mathcal{Z} is the Hilbert space

$$\Gamma_{\mathfrak{a}}(\mathcal{Z}) := \bigoplus_{n=0}^{\infty} \otimes_{\mathfrak{a}}^n \mathcal{Z},$$

where $\otimes_{\mathfrak{a}}^n \mathcal{Z}$ denotes the anti-symmetric n -th tensor power of \mathcal{Z} . On \mathcal{Z} one defines the *creation/annihilation operators* $a^*(u)$, $a(u)$ (see e.g. [RS]) satisfying the canonical anti-commutation relations:

$$[a(u), a(v)]_+ = [a^*(u), a^*(v)]_+ = 0, \quad [a(u), a^*(v)]_+ = (u|v)_{\mathcal{Z}} \mathbb{1},$$

where $[\cdot, \cdot]_+$ denotes the anti-commutator. If b is an operator acting on \mathcal{Z} , $d\Gamma(b)$ is its second quantization, acting on $\Gamma_{\mathfrak{a}}(\mathcal{Z})$ (see e.g. [RS]). The unit vector $\Omega = (1, 0, \dots) \in \Gamma_{\mathfrak{a}}(\mathcal{Z})$ is called the *vacuum*.

A.5. Fock representation associated to a Kähler structure. Let \mathfrak{h} be a complex Hilbert space. We denote its complex structure by i and its scalar product by $(\cdot|\cdot)$. The space \mathfrak{h} considered as a real vector space will be denoted by $\mathfrak{h}_{\mathbb{R}}$.

A *Kähler structure* on \mathfrak{h} is a unitary anti-involution j acting on \mathfrak{h} . Note that $\kappa := -ij$ is a selfadjoint involution. Therefore we can split \mathfrak{h} as $\mathfrak{h}^+ \oplus \mathfrak{h}^-$, where $\mathfrak{h}^{\pm} := \mathbb{1}_{\{\pm 1\}}(\kappa)\mathfrak{h}$. We also set $f^{\pm} := \mathbb{1}_{\{\pm 1\}}(\kappa)f$ for $f \in \mathfrak{h}$.

Let us denote by \mathcal{Z} the real vector space $\mathfrak{h}_{\mathbb{R}}$ equipped with the complex structure j . We can furthermore turn \mathcal{Z} into a Hilbert space by equipping it with the scalar product:

$$(u|v)_{\mathcal{Z}} := (u^+|v^+) + (v^-|u^-).$$

The Hilbert space \mathcal{Z} is called the *one-particle space* (associated to the Kähler structure j).

One can then define the *Fock representation* of $\text{CAR}(\mathfrak{h})$ in $\Gamma_{\mathfrak{a}}(\mathcal{Z})$ by setting

$$\psi_F(f) = \pi_F(\psi(f)) := \frac{1}{2}a^*(f^+) + \frac{1}{2}a(f^-), \quad f \in \mathfrak{h}.$$

The operator $Q := d\Gamma(\kappa)$ acting on $\Gamma_{\mathfrak{a}}(\mathcal{Z})$ is usually called the *charge operator*. One has:

$$(A.3) \quad e^{i\theta Q} \psi_F^{(*)}(f) e^{-i\theta Q} = \psi_F(e^{i\theta} f), \quad f \in \mathfrak{h}, \quad \theta \in \mathbb{R}.$$

If b is a selfadjoint operator on \mathfrak{h} which commutes with j , then $c = \kappa b$ is selfadjoint on \mathcal{Z} . The operator $H = d\Gamma(c)$ acting on $\Gamma_{\mathfrak{a}}(\mathcal{Z})$ is usually called the (quantum) *Hamiltonian*. Note that $H \geq 0$ and $\Omega \in \Gamma_{\mathfrak{a}}(\mathcal{Z})$ is its unique ground state. One has:

$$(A.4) \quad e^{itH} \psi_F(f) e^{-itH} = \phi_F(e^{itb} f), \quad f \in \mathfrak{h}, \quad t \in \mathbb{R},$$

i.e. the unitary group e^{itH} implements the dynamics generated by e^{itb} in the Fock representation. Note that to conform with the common usage, we denoted by i in (A.3) and (A.4) the complex structure on $\Gamma_{\mathfrak{a}}(\mathcal{Z})$.

A.6. Some auxiliary results. The following lemma is well known.

Lemma A.10. *Let \mathfrak{A} a C^* -algebra and $\mathbb{R} \ni t \mapsto H(t) \in \mathfrak{A}$ a continuous map with $H(t) = H^*(t)$. Then there exists a unique C^1 map*

$$\mathbb{R}^2 \ni (s, t) \mapsto U_{H(\cdot)}(s, t) \in \mathfrak{A},$$

such that:

- i) $\partial_t U_{H(\cdot)}(s, t) = -iU_{H(\cdot)}(s, t)H(t)$, $s, t \in \mathbb{R}$,
- ii) $\partial_s U_{H(\cdot)}(s, t) = iH(s)U_{H(\cdot)}(s, t)$, $s, t \in \mathbb{R}$,
- iii) $U_{H(\cdot)}(t, t) = \mathbb{1}$.

Moreover one has

- iv) $U_{H(\cdot)}(s, t) = \sum_{n=0}^{\infty} (-i)^n \int_{s \leq t_n \leq \dots \leq t_1 \leq t} H(t_n) \cdots H(t_1) dt_n \dots dt_1$,
- v) $U_{H(\cdot)}(s, t')U_{H(\cdot)}(t', t) = U_{H(\cdot)}(s, t)$, $s, t', t \in \mathbb{R}$,
- vi) $U_{H(\cdot)}(s, t)$ is unitary in \mathfrak{A} .

We now further study $U_{H(\cdot)}(s, t)$ if $t \mapsto H(t)$ is obtained from a quantum dynamics.

It is natural to generalize the framework of Def. 3.1. Instead of choosing $\mathfrak{A}_t = \text{CAR}(\mathfrak{h}_t)$ we can assume that \mathfrak{A}_t for $t \in \mathbb{R}$ are C^* -algebras with $\mathfrak{A}_t \subset \mathfrak{A}_s \subset \mathfrak{A}_\infty$ for $s \leq t$ for some C^* -algebra \mathfrak{A}_∞ . Moreover we can assume that for each $t_0 \in \mathbb{R}$, there exists a $*$ -sub-algebra $\tilde{\mathfrak{A}}_{t_0}$ dense in \mathfrak{A}_{t_0} such that an $A \in \tilde{\mathfrak{A}}_{t_0}$ belongs to \mathfrak{A}_t for t near t_0 . Then the obvious generalization of Def. 3.1 makes sense.

Proposition A.11. *Let \mathfrak{A}_t for $t \in \mathbb{R}$ be a family of C^* -algebras satisfying the above conditions and $\tau^0(s, t) : \mathfrak{A}_t \xrightarrow{\sim} \mathfrak{A}_s$ a quantum dynamics. Let $\mathbb{R} \ni t \mapsto I(t) \in \mathfrak{A}_\infty$ a continuous map with $I(t) = I^*(t)$ and $I(t) \in \mathfrak{A}_t$ for $t \in \mathbb{R}$. Set:*

$$I(s, t) := \tau^0(s, t)(I(t)) \in \mathfrak{A}_s, \quad R_s(t', t) := U_{I(s, \cdot)}(t', t) \in U(\mathfrak{A}_s).$$

Then

$$(1) \quad \tau^0(s, t')R_{t'}(t', t) = R_s(t', t), \quad s, t', t \in \mathbb{R};$$

(2) Set

$$\tau(s, t)(A) := R_s(s, t)\tau^0(s, t)(A)R_s(s, t)^*, \quad s, t \in \mathbb{R}.$$

Then $\tau(s, t) : \mathfrak{A}_t \xrightarrow{\sim} \mathfrak{A}_s$ is a quantum dynamics.

Proof. Statement (1) follows by differentiating both members w.r.t. t and using the uniqueness result in Lemma A.10. (2) follows from (1). \square

The following remark will hopefully clarify the meaning of $\tau(s, t)$ constructed in Prop. A.11.

Remark A.12. *Assume that $\mathfrak{A}_t \equiv \mathfrak{A}$ and let $H_0 = H_0^* \in \mathfrak{A}$ and $\mathbb{R} \ni t \mapsto I(t) \in \mathfrak{A}$ be continuous. Set $\tau^0(s, t)A = e^{i(s-t)H_0} A e^{-i(s-t)H_0}$ and let $\tau(s, t)$ be obtained from Prop. A.11. Then $\tau(s, t)A = U(s, t)AU(t, s)$ where $\{U(s, t)\}_{s, t \in \mathbb{R}}$ is the two-parameter propagator obtained from Lemma A.10 for $H(t) = H_0 + I(t)$.*

REFERENCES

- [A] Ammari, Z.: Scattering theory for a class of fermionic Pauli-Fierz models , J. Funct. Anal. **208** (2004), 302-359.
- [Ba1] Bachelot, A.: The Hawking effect, Ann. Inst. H. Poincaré Phys. Théor. **70** (1999), 41-99.
- [Ba2] Bachelot, A.: Creation of fermions at the charged black-hole horizon, Ann. Henri Poincaré, **1** (2000), 1043-1095.
- [CP] Chazarain, J., Piriou, A.: *Introduction to the Theory of Linear Partial Differential Equations*, Studies in Mathematics and Its Applications. Elsevier Science, (1982).

- [Da] Daudé, T.: Propagation estimates for Dirac operators and application to scattering theory, *Annales de l'Institut Fourier* **54** (2004), 2021-2083.
- [DG] Dereziński, J., Gérard, C.: Asymptotic Completeness in Quantum Field Theory: massive Pauli-Fierz Hamiltonians, *Rev. Math. Phys.* **11** (1999), 383-450.
- [DK1] Dimock, J., Kay, B.: Classical and Quantum scattering theory for linear scalar fields on the Schwarzschild metric I, *Ann. Phys.* **175** (1987), 366-426.
- [DK2] Dimock, J., Kay, B.: Classical and Quantum scattering theory for linear scalar fields on the Schwarzschild metric II, *J. Math. Phys.* **27** (1986), 2520-2525.
- [H] Häfner, D.: Creation of fermions by rotating charged black holes, *Mémoires de la S.M.F.* **117** (2009), 158 p.
- [Ha] Hawking, S.: Particle creation by black holes, *Comm. Math. Phys.* **43** (1975), 199-220.
- [Me] Melnyk, F.: The Hawking effect for spin 1/2 fields, *Comm. Math. Phys.* **244**, (2003), 483-55.
- [RS] Reed, M., Simon, B.: *Methods of Modern Mathematical Physics, Vol. II Fourier Analysis, Self-Adjointness*, Academic Press, (1975).

DÉPARTEMENT DE MATHÉMATIQUES, UNIVERSITÉ DE PARIS XI, 91405 ORSAY CEDEX FRANCE
E-mail address: `patrick.bouvier@math.u-psud.fr`

DÉPARTEMENT DE MATHÉMATIQUES, UNIVERSITÉ DE PARIS XI, 91405 ORSAY CEDEX FRANCE
E-mail address: `christian.gerard@math.u-psud.fr`