



HAL
open science

Dimensional contraction in Wasserstein distance for diffusion semigroups on a Riemannian manifold

Ivan Gentil

► **To cite this version:**

Ivan Gentil. Dimensional contraction in Wasserstein distance for diffusion semigroups on a Riemannian manifold. 2013. hal-00873097v2

HAL Id: hal-00873097

<https://hal.science/hal-00873097v2>

Preprint submitted on 22 May 2014 (v2), last revised 14 Dec 2014 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Dimensional contraction in Wasserstein distance for diffusion semigroups on a Riemannian manifold

Ivan Gentil*

May 22, 2014

Abstract

We prove a refined contraction inequality for diffusion semigroups with respect to the Wasserstein distance on a compact Riemannian manifold taking account of the dimension. The result generalizes in a Riemannian context, the dimensional contraction established in [BGG13] for the Euclidean heat equation. It is proved by using a dimensional coercive estimate for the Hodge-de Rham semigroup on 1-forms.

Key words: Diffusion equations, Wasserstein distance, Hodge-de Rham operator, Curvature-dimension bounds.

Mathematics Subject Classification (2000): 58J65, 58J35, 53B21

1 Introduction

The von Renesse-Sturm Theorem (c.f. [vRS05]) insures that the contraction of the heat equation on a Riemannian manifold with respect to the Wasserstein distance is equivalent to a uniform lower bound of the Ricci curvature. This result is one of the first equivalence theorem relating the Wasserstein distance and the Ricci curvature. Actually, there are many extensions of this result including the case of a heat equation on a metric measurable space and we would like to take account of the dimension into such contraction.

Let us explain the contraction inequality and its extensions with more details. For simplicity, we focus on the heat equation on a Riemannian manifold but all of these results have been proved for a general diffusion semigroup on a Riemannian manifold (or more general spaces). Let Δ be the Laplace-Beltrami operator on a smooth Riemannian manifold (M, g) and let $P_t f$ be the solution of the heat equation $\partial_t u = \Delta u$ with f as the initial condition.

The von Renesse-Sturm Theorem states that: Let $R \in \mathbb{R}$, the following assertions are equivalent (the Wasserstein distance is denoted W_2),

- (i) for any f, g probability densities with respect to the Riemannian measure dx

$$W_2^2(P_t f dx, P_t g dx) \leq e^{-2Rt} W_2^2(f dx, g dx), \quad \forall t \geq 0,$$

*Institut Camille Jordan, Umr Cnrs 5208, Université Claude Bernard Lyon 1, 43 boulevard du 11 novembre 1918, F-69622 Villeurbanne cedex. gentil@math.univ-lyon1.fr

- (ii) $Ricci_g \geq R$ (uniformly in M) where $Ricci_g$ is the Ricci tensor of (M, g) . The inequality has to be understood in the sense of inequality between symmetric tensors.

There are many proofs and extensions of this result, one can see for instance [Wan04, OW05, AGS08, Kuw10, Wan11, Sav14, GKO13, BGL14, BGL15].

Recently, many extensions have been given taking account of the dimension of the manifold. For instance in [Kuw13, BGL15] the authors prove that if M is a n -dimensional Riemannian manifold with a non-negative Ricci curvature, then for any f, g probability densities with respect to dx , and any $s, t \geq 0$,

$$W_2^2(P_s f dx, P_t g dx) \leq W_2^2(f dx, g dx) + 2n(\sqrt{s} - \sqrt{t})^2, \quad \forall s, t \geq 0, \quad (1)$$

Non-negative curvature condition has been removed in [Kuw13, EKS13]. If M is a n -dimensional Riemannian manifold, the main extensions are the followings:

- In [Kuw13], K. Kuwada proves that the Ricci curvature is bounded from below by $R \in \mathbb{R}$ if and only if for every $s, t \geq 0$,

$$W_2^2(P_t f dx, P_s g dx) \leq A(s, t, R)W_2^2(f dx, g dx) + B(s, t, n, R), \quad \forall s, t \geq 0, \quad (2)$$

for probability densities f, g with respect to dx , for appropriate functions $A, B \geq 0$.

- In [EKS13], M. Erbar, K. Kuwada and K.-T. Sturm prove that the Ricci curvature of the n -dimensional manifold M is bounded from below by a constant $R \in \mathbb{R}$ if and only if

$$s_{\frac{R}{n}} \left(\frac{1}{2} W_2(P_t f dx, P_s g dx) \right)^2 \leq e^{-R(t+s)} s_{\frac{R}{n}} \left(\frac{1}{2} W_2(f dx, g dx) \right)^2 + \frac{n}{R} (1 - e^{-R(s+t)}) \frac{(\sqrt{t} - \sqrt{s})^2}{2(t+s)}, \quad (3)$$

for any $s, t \geq 0$ and any probability densities f and g . Here $s_r(x) = \sin(\sqrt{r}x)/\sqrt{r}$ if $r > 0$, $s_r(x) = \sinh(\sqrt{-r}x)/\sqrt{-r}$ if $r < 0$ and $s_0(x) = x$, hence recovering (1) when $R = 0$.

- In [BGG13] we prove that the classical heat equation in the Euclidean space \mathbb{R}^n satisfies, for any f, g probability densities with respect to the Lebesgue measure λ ,

$$W_2^2(P_t f \lambda, P_t g \lambda) \leq W_2^2(f \lambda, g \lambda) - \frac{2}{n} \int_0^t (\text{Ent}_\lambda(P_u f) - \text{Ent}_\lambda(P_u g))^2 du, \quad \forall t \geq 0, \quad (4)$$

where Ent is the Entropy (it will be defined later).

- In the same way, again in [BGG13], we prove a more general result for a n -dimensional Riemannian manifold with a Ricci curvature bounded from below by R and for the Markov transportation distance T_2 (a new distance on measures). We obtain for any f and g ,

$$T_2^2(P_t f dx, P_t g dx) \leq e^{-2Rt} T_2^2(f dx, g dx) - \frac{2}{n} \int_0^t e^{-2R(t-u)} (\text{Ent}_\mu(P_u g) - \text{Ent}_\mu(P_u f))^2 du,$$

for any $t \geq 0$.

The goal of this paper is to prove the previous inequality for the Wasserstein distance instead the Markov transportation distance, in other words to extend inequality (4) on a n -dimensional Riemannian manifold with a lower bound on the Ricci curvature.

The main result of this paper can be stated as follow, let $d\mu = e^{-\Psi}dx$ be a probability measure with Ψ a smooth function on M (for the sequel the manifold will be compact). We denote by $(P_t)_{t \geq 0}$ the Markov semigroup associated to the generator $L = \Delta - \nabla\Psi \cdot \nabla$. Then under the curvature-dimension condition

$$\text{Ricci}_g + \text{Hess}(\Psi) \geq R + \frac{1}{m-n} \nabla\Psi \otimes \nabla\Psi, \quad (5)$$

for some $R \in \mathbb{R}$ and $m \geq n$ (when $m = n$ then $\Psi = 0$), then for any smooth probability densities f, g and any $t \geq 0$,

$$W_2^2(P_t f \mu, P_t g \mu) \leq e^{-2Rt} W_2^2(f \mu, g \mu) - \frac{2}{m} \int_0^t e^{-2R(t-u)} [\text{Ent}_\mu(P_u g) - \text{Ent}_\mu(P_u f)]^2 du.$$

The main advantage of such inequality with respect to (2) and (3) is to obtain a contraction inequality with the same time t instead two different times s and t . Moreover, the additional term is given with a minus sign which shows the improvement given by the dimension.

The method to get a dimensional contraction is radically different that one one used in [EKS13]. Here the strategy to prove such inequality is the Benamou-Brenier dynamical formulation (the Eulerian formulation) of the Wasserstein distance associated to a sharp dimensional estimate on the Hodge-de Rham semigroup. In [EKS13] the authors use in force the definition of the heat equation as a gradient flow of the entropy with respect to the Wasserstein distance.

The paper is organized as follow. First in Section 2, we recall the Riemannian setting and the Wasserstein distance through the Benamou-Brenier dynamical formulation. We need to introduce the Hodge-de Rham operator on forms and its associated semigroup. In Section 3, we improve the Bochner-Lichnerowicz-Weitzenböck identity for 1-forms to get a coercive inequality for the Hodge-de Rham semigroup. Finally in Section 4 the main theorem is proved.

For simplicity reasons, the result will be stated and proved in the context of a compact Riemannian manifold but its generalization in a metric space, including the equivalence with respect to the condition (5), is actually a project with F. Bolley, N. Gigli and A. Guillin.

2 Framework and main result

2.1 Geometrics tools

Conventions and notations. Let (M, g) be a n -dimensional, connected, compact and differentiable Riemannian manifold without boundary. We assume for simplicity that the manifold is \mathcal{C}^∞ . For each $x \in M$ we denote by $T_x M$ the tangent space to M and by TM the whole tangent bundle of M . Moreover g_x is a symmetric definite positive quadratic form on $T_x M$, in a local basis of $T_x M$ $(e_i)_{1 \leq i \leq n}$, $g_x = (g_{ij})$. (For simplicity, the x dependance of the metric g is omitted.) For every $x, y \in M$, $d(x, y)$ denotes the usual (Riemannian) distance and dx its measure. In the sequel we will use the Einstein's convention of summation over repeated indices: for instance $x_i y^i = \sum_{i=1}^n x_i y^i$, $x^i g_{ij} y^j = \sum_{i=1}^n x^i g_{ij} y^j$.

The Riemannian scalar product between two tensors X and Y is denoted $X \cdot Y$, its associated norm is noted $|X|$ (depending on g). For instance, locally in a basis $(e_i)_{1 \leq i \leq n}$ for two smooth functions $f, h : M \mapsto \mathbb{R}$, $\nabla f \cdot \nabla h = \partial_i f g^{ij} \partial_j h$ where $(g^{ij}) = (g_{ij})^{-1}$. As usual the covariant derivative in the direction e_i of a tensor X (vector field or form) is noted $\nabla_i X$. The geometric

musicology will be used in force, for instance $\nabla^i f = g^{ij} \nabla_j f = g^{ij} \partial_j f$ or $\nabla^i X = g^{ij} \nabla_j X$. If ω is a 1-form then ω^* is its dual representation as a vector field with components $\omega^i = g^{ij} \omega_j$. The Ricci tensor of (M, g) is noted $Ricci_g$.

The Laplace-Beltrami operator Δ is acting on smooth functions f ,

$$\Delta f = \nabla \cdot \nabla f,$$

where $\nabla \cdot$ is the divergence operator acting on vector fields, $\nabla \cdot X = \nabla_i X^i$ for every vector field X . (We use the analyst's convention with respect to the sign.) A smooth function (or form) is a C^∞ -function (or form). The divergence operator $\nabla \cdot$ satisfies for any smooth vector fields X and any smooth functions f ,

$$\int_M \nabla \cdot X f dx = - \int_M \delta df \cdot X dx.$$

where δ is the divergence operator on 1-form

$$\delta \omega = \nabla_i \omega^i = g^{ij} \nabla_i \omega_j = g^{ij} (\partial_i \omega_j - \Gamma_{ij}^p \omega_p),$$

where Γ_{ij}^p are Christoffel symbols. Operators δ and $\nabla \cdot$ are related by the formula $\delta \omega = \nabla \cdot \omega^*$, for any 1-forms ω .

The Markov (or heat) semigroup. Let $\Psi : M \mapsto \mathbb{R}$ be a fixed C^∞ -function. Let $\mu(dx) = e^{-\Psi} dx$ and since M is compact we can assume that μ is a probability measure. Let $Lf = \Delta f - \nabla \Psi \cdot \nabla f$, for any smooth functions f . Since the manifold is compact, the operator L defined a unique Markov semigroup $(P_t)_{t \geq 0}$ on $L^2(\mu)$ and it is called a Markov generator. This semigroup is symmetric in $L^2(\mu)$ and for any f , $P_t f$ is solution of the equation $\partial_t u = Lu$ with f as the initial condition. If f is a probability density (with respect to μ) then for all $t \geq 0$, $P_t f \mu$ remains a probability measure. Finally, for any smooth functions f and g on M ,

$$\int_M f Lg d\mu = \int_M g Lf \mu = - \int_M \nabla f \cdot \nabla g d\mu = - \int_M \Gamma(f, g) d\mu,$$

where Γ is Carré du champ operator defined on functions, $\Gamma(f, g) = \nabla f \cdot \nabla g$. If X is a vector field, we define $\nabla_\Psi \cdot X = \nabla \cdot X - \nabla \Psi \cdot X$, and the generator L takes then the form

$$L = \nabla_\Psi \cdot \nabla.$$

As before we note by δ_Ψ the divergence operator acting on forms:

$$\delta_\Psi \omega = \nabla_\Psi \cdot \omega^*. \tag{6}$$

It satisfies the integration by parts formula, for any smooth 1-forms ω and functions f ,

$$\int_M \delta_\Psi \omega f d\mu = - \int_M \omega \cdot df d\mu.$$

The triple (M, μ, Γ) is a compact Markov triple as defined in [BGL14, Chap. 3].

The Hodge-de Rham semigroup. Connected to the Markov semigroup (P_t) (associated to the generator L) one can define the Hodge-de Rham semigroup. As explained in this context in [Bak87] the (modified) Hodge-de Rham operator is acting on smooth 1-forms,

$$\vec{L} = (d^{(0)} \delta_\Psi^{(1)} + \delta_\Psi^{(2)} d^{(1)}), \tag{7}$$

where $d^{(i)}$ is the differential operator acting on i -forms and $\delta_{\Psi}^{(i)}$ is its adjoint operator in $L^2(\mu)$ (in the sequel, we omit the exponent (i)). For computations, we use the Hodge-de Rham operator toward the Weitzenböck formula, that is for any $1 \leq i \leq n$,

$$(\vec{L}\omega)_i = \nabla^k \nabla_k \omega_i - (\nabla_{\nabla \Psi} \omega)_i - \text{Ricci}(L)(\omega^*, e_i) = \nabla^k \nabla_k \omega_i - \nabla^k \Psi \nabla_k \omega_i - \text{Ricci}(L)(\omega^*, e_i), \quad (8)$$

where $\text{Ricci}(L) = \text{Ricci}_g + \text{Hess}(\Psi)$ is the so-called Bakry-Émery tensor (see for instance [Bak87, Prop. 1.5]). Again, we use the analyst's convention with respect to the sign. If $\Psi = 0$, then $L = \Delta$ and \vec{L} is the usual Hodge-de Rham operator noted $\vec{\Delta}$.

Since M is compact, the operator \vec{L} induces a semigroup $(R_t)_{t \geq 0}$ on 1-forms. It is also symmetric in $L^2(d\mu)$, for any smooth 1-forms ω and η ,

$$\int_M \omega \cdot \vec{L}\eta \, d\mu = \int_M \eta \cdot \vec{L}\omega \, d\mu.$$

Then for any smooth 1-forms ω , $R_t\omega$ is the solution of $\partial_t u = \vec{L}u$ where $u : [0, \infty) \times M \mapsto TM^*$ (TM^* is the cotangent bundle) with ω as the initial condition. The details of the construction of the Hodge-de Rham semigroup can be found in [Bak87] (see also the references therein).

The Hodge-de Rham semigroup is related to the Markov generator by the following commutation property: for any smooth 1-forms ω and $t \geq 0$,

$$P_t \delta_{\Psi} \omega = \delta_{\Psi} R_t \omega. \quad (9)$$

The easiest way to prove this fundamental identity is to use the definition (7) and the identity $\delta_{\Psi}^{(1)} \delta_{\Psi}^{(2)} = 0$.

2.2 The Wasserstein distance

Let $\mathcal{P}(M)$ be the set of probability measures in M . The Wasserstein distance between two probability measures $\nu_1, \nu_2 \in \mathcal{P}(M)$, is defined by

$$W_2(\nu_1, \nu_2) = \inf \left(\int_{M \times M} d^2(x, y) d\pi(x, y) \right)^{1/2},$$

where the infimum runs over all probability measures π in $M \times M$ with marginals ν_1 and ν_2 . We refer to the monumental work [Vil09] for a reference presentation of this distance, its interplay with the optimal transportation problem and many other issues.

The Wasserstein distance has a dynamical formulation: for any probabilities measure $\nu_1, \nu_2 \in \mathcal{P}(M)$,

$$W_2^2(\nu_1, \nu_2) = \inf \int_0^1 \int_M |\eta_s|^2 d\mu_s ds,$$

where the infimum is running over all paths of probabilities $(\mu_s)_{s \in [0,1]}$ and $\eta_s \in TM^*$ satisfying in the distributional sense

$$\begin{cases} \partial_s \mu_s + \delta(\mu_s \eta_s) = 0 \\ \mu_0 = \nu_1, \mu_1 = \nu_2. \end{cases}$$

The Euclidean case has been proved by Benamou-Brenier in [BB00] and the Riemannian case by F. Otto and M. Westdickenberg on the other hand by L. De Pascale, M. S. Gelli and L. Granieri in [OW05, DPGG06].

Let $\omega_s = \eta_s \rho_s$ where $\frac{d\mu_s}{d\mu} = \rho_s$, in [OW05] the authors state that for any f and g , smooth probability densities (with respect to μ),

$$W_2^2(f\mu, g\mu) = \inf \int_0^1 \int_M \frac{|\omega_s|^2}{\rho_s} d\mu ds, \quad (10)$$

where the infimum is running over smooth couples $(\rho_s, \omega_s)_{s \in [0,1]}$ where for any $s \in [0, 1]$, ρ_s is a positive probability density (with respect to μ) and ω_s is a 1-form, satisfying

$$\begin{cases} \partial_s \rho_s + \delta_\Psi \omega_s = 0 \\ \rho_0 = f, \quad \rho_1 = g. \end{cases}$$

This dynamical formulation of optimal transportation has been mainly used to get contraction result or Evolutional Variational Inequality (see e.g. [DNS09, DNS12, BGG13]).

2.3 Main result

We can now state the main result of this short paper.

Theorem 2.1 (Dimensional contraction in Wasserstein distance) *Let (M, g) be a C^∞ , n -dimensional, connected and compact Riemannian manifold. We assume that there exists $R \in \mathbb{R}$ and $m \geq n$ such that uniformly in M (if $m = n$ then we impose that $\Psi = 0$),*

$$\text{Ricci}(L) \geq R + \frac{1}{m-n} \nabla \Psi \otimes \nabla \Psi. \quad (11)$$

Then for any smooth probability densities f, g with respect to μ and any $t \geq 0$,

$$W_2^2(P_t f \mu, P_t g \mu) \leq e^{-2Rt} W_2^2(f\mu, g\mu) - \frac{2}{m} \int_0^t e^{-2R(t-u)} [\text{Ent}_\mu(P_u g) - \text{Ent}_\mu(P_u f)]^2 du, \quad (12)$$

where $\text{Ent}_\mu(h) = \int_M h \log h d\mu$ for every probability density h (with respect to μ).

Remark 2.2 *When $m \rightarrow \infty$ we recover the von Renesse-Sturm result, the exponential contraction of the heat equation with respect to the Wasserstein distance. When $\Psi = 0$, one can choose $m = n$, and then the Laplace-Beltrami operator satisfies the condition (11) under a lower bound on the Ricci curvature.*

As it will be explained in Remark 3.7, condition (11) is equivalent to the so-called Bakry-Émery curvature-dimension condition $CD(R, m)$.

3 Coercive inequality for the Hodge-de Rham semi-group

The next proposition states a refined Bochner-Lichnerowicz-Weitzenböck formula for 1-forms.

Proposition 3.1 (Refined Bochner-Lichnerowicz-Weitzenböck formula) *For any smooth 1-forms α, η and any $a \in \mathbb{R}$,*

$$L \frac{|\eta|^2}{2} - \eta \cdot \vec{L} \eta + 2a\alpha \cdot d|\eta|^2 + 4a^2 |\alpha|^2 |\eta|^2 = |\nabla \eta + 2a\alpha \otimes \eta|^2 + \text{Ricci}(L)(\eta^*, \eta^*), \quad (13)$$

where $|\nabla \eta + 2a\alpha \otimes \eta|$ has to be understood as the norm of the 2-tensor $\nabla_i \eta_j + 2a\alpha_i \eta_j$.

When $\Psi = 0$, $\overrightarrow{\Delta}$ be the usual Hodge-de Rham operator. The Weitzenböck (8) implies

$$\overrightarrow{L}\omega = \overrightarrow{\Delta}\omega - \nabla_{\nabla\Psi}\omega - \text{Hess}(\Psi)(\omega^*, \cdot). \quad (14)$$

According to Lemma 3.2, equation (13) is equivalent to the identity

$$\Delta \frac{|\eta|^2}{2} - \eta \cdot \overrightarrow{\Delta}\eta + 2a\alpha \cdot d|\eta|^2 + 4a^2|\alpha|^2|\eta|^2 = |\nabla\eta + 2a\alpha \otimes \eta|^2 + \text{Ricci}_g(\eta^*, \eta^*),$$

which will be proved in Proposition 3.3. The main difficulty is to prove it for all $a \in \mathbb{R}$ since when $a = 0$, equation (13) is classic.

Next Lemma can be found for instance in [Lic58, p. 3].

Lemma 3.2 *For any 1-forms η and any $1 \leq k \leq n$, $\left(d\frac{|\eta|^2}{2}\right)_k = \eta^i \nabla_k \eta_i$.*

Proposition 3.3 *For any smooth 1-forms α , η and any $a \in \mathbb{R}$,*

$$\Delta \frac{|\eta|^2}{2} - \eta \cdot \overrightarrow{\Delta}\eta + 2a\alpha \cdot d|\eta|^2 + 4a^2|\alpha|^2|\eta|^2 = |\nabla\eta + 2a\alpha \otimes \eta|^2 + \text{Ricci}_g(\eta^*, \eta^*). \quad (15)$$

Proof

◁ The Bochner-Lichnerowicz-Weitzenböck formula (c.f. [Lic58, P. 3]) insures that for any smooth 1-forms ω ,

$$\Delta \frac{|\omega|^2}{2} - \omega \cdot \overrightarrow{\Delta}\omega = |\nabla\omega|^2 + \text{Ricci}_g(\omega^*, \omega^*), \quad (16)$$

recall that $|\nabla\omega|^2 = \nabla_i \omega_j g^{il} g^{jk} \nabla_l \omega_k$ is the square of the norm of the 2-tensor $\nabla_i \omega_j$.

The idea is to change of variables into this formula. We would like to prove this result at some $x_0 \in M$ which is supposed to be fixed. Let

$$\omega = (ag + 1)\eta + af\alpha,$$

where η and α are 1-forms and f, g are actually smooth functions satisfying $f(x_0) = g(x_0) = 0$.

First we have

$$|\omega|^2 = (ag + 1)^2 |\eta|^2 + a^2 f^2 |\alpha|^2 + 2a(ag + 1)f\alpha \cdot \eta.$$

Since, for any smooth functions F and G we have $\Delta(FG) = 2\Gamma(F, G) + F\Delta G + G\Delta F$, where Γ is the carré du champ operator, a straight forward computation gives at the point x_0 ,

$$\begin{aligned} \Delta|\omega|^2 &= 4a\Gamma(g, |\eta|^2) + 2a|\eta|^2 \Delta g + 2a^2|\eta|^2 |\nabla g|^2 + \Delta|\eta|^2 + 2a^2|\alpha|^2 |\nabla f|^2 \\ &\quad + 4a\Gamma(f, \eta \cdot \alpha) + 4a^2(\eta \cdot \alpha)\Gamma(f, g) + 2a(\eta \cdot \alpha) \Delta f. \end{aligned}$$

Since $\omega(x_0) = \eta(x_0)$, we have in x_0 ,

$$2\omega \cdot \overrightarrow{\Delta}\omega = 2a\eta \overrightarrow{\Delta}(g\eta) + 2\eta \overrightarrow{\Delta}\eta + 2a\eta \overrightarrow{\Delta}(f\alpha).$$

Thanks to Lemma 3.5, at the point x_0 (recall that $f(x_0) = g(x_0) = 0$),

$$2\omega \cdot \overrightarrow{\Delta}\omega = 2a|\eta|^2 \Delta g + 4a\eta \cdot \nabla_{\nabla g} \eta + 2\eta \overrightarrow{\Delta}\eta + 2a\eta \cdot \alpha \Delta f + 4a\eta \cdot \nabla_{\nabla f} \alpha.$$

Moreover, at the point x_0 ,

$$|\nabla\omega|^2 = |\nabla_i\omega_j|^2 = a^2|\eta|^2\Gamma(g) + |\nabla_i\eta_j|^2 + a^2\Gamma(f)|\alpha|^2 + 2a\eta^j\nabla^i g\nabla_i\eta_j \\ + 2a\nabla^i f\alpha^j\nabla_i\eta_j + 2a^2\eta_j\nabla_i g\nabla^i f\alpha^j,$$

and $Ricci_g(\omega^*, \omega^*) = Ricci_g(\eta^*, \eta^*)$. At the point x_0 , the identity (16) applied to ω , becomes

$$\Delta\frac{|\eta|^2}{2} - \eta\vec{\Delta}\eta + 2a\Gamma(g, |\eta|^2) + a^2|\eta|^2\Gamma(g) + a^2|\alpha|^2\Gamma(f) + 2a\Gamma(f, \eta \cdot \alpha) \\ + 2a^2(\eta \cdot \alpha)\Gamma(f, g) - 2a\eta \cdot \nabla_{\nabla g}\eta - 2a\eta \cdot \nabla_{\nabla f}\alpha = a^2|\eta|^2\Gamma(g) + |\nabla_i\eta_j|^2 \\ + a^2\Gamma(f)|\alpha|^2 + 2a\eta^j\nabla^i g\nabla_i\eta_j + 2a\nabla^i f\alpha^j\nabla_i\eta_j + 2a^2\eta_j\nabla_i g\nabla^i f\alpha^j + Ricci_g(\eta^*, \eta^*)$$

Let now assume that f and g satisfied moreover $df(x_0) = \eta(x_0)$ and $dg(x_0) = \alpha(x_0)$ (which is allays possible), we obtain

$$\Delta\frac{|\eta|^2}{2} - \eta\vec{\Delta}\eta + 2a\alpha \cdot \nabla|\eta|^2 + 4a^2|\eta|^2|\alpha|^2 \\ = 4a^2|\eta|^2|\alpha|^2 - 2a\eta \cdot \nabla(\eta \cdot \alpha) + 2a\eta \cdot \nabla_\alpha\eta - 2a\eta \cdot \nabla_\eta\alpha + |\nabla_i\eta_j|^2 \\ + 2a\eta^j\alpha^i\nabla_i\eta_j + 2a\eta^i\alpha^j\nabla_i\eta_j + Ricci_g(\eta^*, \eta^*).$$

Then Lemma 3.4 implies that

$$4a^2|\eta|^2|\alpha|^2 - 2a\eta \cdot d(\eta \cdot \alpha) + 2a\eta \cdot \nabla_\alpha\eta - 2a\eta \cdot \nabla_\eta\alpha + |\nabla_i\eta_j|^2 + 2a\eta^j\alpha^i\nabla_i\eta_j + 2a\eta^i\alpha^j\nabla_i\eta_j \\ = 4a^2|\eta|^2|\alpha|^2 + 4a\eta^j\alpha^i\nabla_i\eta_j + \nabla_i\eta_j\nabla^i\eta^j = |\nabla\eta + 2a\alpha \otimes \eta|^2,$$

which implies the identity (15). \triangleright

The next lemma is classic in the Riemannian context and we skip the proof.

Lemma 3.4 *For any 1-forms η and α , $d(\eta \cdot \alpha) = \nabla\eta \cdot \alpha + \nabla\alpha \cdot \eta$.*

The next result can be seen as a kind of diffusion property as defined in [BGL14]. The result is probably classic but I didn't find it in the literature.

Lemma 3.5 *For any smooth functions f and 1-form ω ,*

$$\vec{\Delta}(f\omega) = f\vec{\Delta}(\omega) + \omega\Delta f + 2\nabla_{\nabla f}\eta. \quad (17)$$

In other words, for any i , $(\vec{\Delta}(f\omega))_i = f(\vec{\Delta}(\omega))_i + \omega_i\Delta f + 2\nabla^j f\nabla_j\eta_i$.

Proof

\triangleleft The Weitzenböck formula (8) insures that

$$(\vec{\Delta}(f\omega))_i = \nabla^k\nabla_k(f\omega)_i - Ricci_g(f\omega^*, e_i) = \nabla^k\nabla_k(f\omega)_i - fRicci_g(\omega^*, e_i).$$

Now, using in force the formula $\nabla_k(f\omega)_i = \omega_i\partial_k f + f\nabla_k\omega_i$, we get

$$\begin{aligned} \nabla^k\nabla_k(f\omega)_i &= g^{kl}\nabla_l\nabla_k(f\omega)_i \\ &= g^{kl}(\partial_l\nabla_k(f\omega)_i - \Gamma_{li}^p\nabla_k(f\omega)_p - \Gamma_{lk}^p\nabla_p(f\omega)_i) \\ &= g^{kl}(\partial_l\omega_i\partial_k f - \Gamma_{lk}^p\partial_p f)\omega_i + f g^{kl}(\partial_l(\nabla_k\omega)_i - \Gamma_{lk}^p(\nabla_p\omega)_i - \Gamma_{li}^p(\nabla_k\omega)_p) \\ &\quad + g^{kl}(\partial_l\omega_i\partial_k f - \partial_k f\Gamma_{li}^p\omega_p + \partial_l f(\nabla_k\omega)_i) \\ &= g^{kl}(\nabla_k\nabla_l f)\omega_i + f g^{kl}\nabla_l\nabla_k\omega_i + 2g^{kl}\partial_k f\nabla_l\omega_i \\ &= \omega_i\Delta f + f\nabla^j\nabla_j\omega_i + 2\nabla^j f\nabla_j\omega_i = \omega_i\Delta f + f\nabla^j\nabla_j\omega_i + 2\nabla_{\nabla f}\omega_i, \end{aligned}$$

which implies (17). \triangleright

Finally we can state our main estimation.

Corollary 3.6 *Assume that there exists $R \in \mathbb{R}$ and $m \geq n$ such that uniformly in M (if $m = n$ then we impose that $\Psi = 0$),*

$$\text{Ricci}(L) \geq R + \frac{1}{m-n} \nabla \Psi \otimes \nabla \Psi. \quad (18)$$

Then for any smooth 1-forms η , α and $a \in \mathbb{R}$, (with δ_Ψ defined in (6)),

$$L \frac{|\eta|^2}{2} - \eta \cdot \vec{L} \eta + 2a\alpha \cdot d|\eta|^2 + 4a^2 |\alpha|^2 |\eta|^2 \geq \frac{1}{m} (\delta_\Psi \eta + 2a\alpha \cdot \eta)^2 + R|\eta|^2. \quad (19)$$

Proof

\triangleleft From (13) we only have to verify

$$|\nabla \eta + 2a\alpha \otimes \eta|^2 + \text{Ricci}(L)(\eta^*, \eta^*) \geq \frac{1}{m} (\delta_\Psi \eta + 2a\alpha \cdot \eta)^2 + R|\eta|^2.$$

Let $x \in M$ and let assume that $(e_i)_{1 \leq i \leq n}$ is an orthonormal basis of $T_x M$. Then the left hand side can be written

$$|\nabla \eta + 2a\alpha \otimes \eta|^2 = \sum_{i,j=1}^n (\nabla_i \eta_j + 2a\alpha_i \eta_j)^2 \geq \frac{1}{n} \left(\sum_{i=1}^n \nabla_i \eta_i + 2a\alpha \cdot \eta \right)^2,$$

from the Cauchy-Schwartz inequality. So it remains to prove that for any $t \in \mathbb{R}$

$$\frac{1}{n} (t + 2a\alpha \cdot \eta)^2 + \text{Ricci}(L)(\eta^*, \eta^*) \geq \frac{1}{m} (t - d\Psi \cdot \eta + 2a\alpha \cdot \eta)^2 + R|\eta|^2,$$

where $t = \delta\eta = \sum_{i=1}^n \nabla_i \eta_i$ (recall that (e_i) is an orthonormal basis). The formula is valid since $m \geq n$ and from (18) the discriminant of this two order polynomial function in variable t is non-positive (it doesn't depend on the parameter a). \triangleright

Remark 3.7 (Link with the curvature-dimension condition) *The so-called Bakry-Émery curvature-dimension condition $CD(R, m)$ for an operator L is satisfied when for every smooth function f ,*

$$\Gamma_2(f) \geq R\Gamma(f) + \frac{1}{m} (Lf)^2,$$

where

$$\Gamma_2(f) = \Gamma_2(f, f) = \frac{1}{2} (L\Gamma(f) - 2\Gamma(f, Lf)).$$

The same procedure in the case of closed 1-forms has been stated in [Bak94] (see also [ABC⁺00, Chap. 5]) in the context of Γ_2 -calculus. It is proved that if $\eta = df$ and $\alpha = dg$, then under the curvature-dimension inequality $CD(R, m)$, and any $a \in \mathbb{R}$,

$$\Gamma_2(f) + 2a\Gamma(f, \Gamma(g)) + 4a^2 \Gamma(f)\Gamma(g) \geq \frac{1}{n} (Lf + 2a\Gamma(f, g))^2 + \rho\Gamma(f), \quad (20)$$

which is (19) for closed 1-form. Moreover, inequality (20) for every function f (with $a = 0$) is equivalent to $CD(R, m)$. Since inequality (19) is a generalization of (20), it is also equivalent to (18) and then to $CD(R, m)$.

We can now give the main estimation of our semigroups:

Theorem 3.8 (Coercive estimation) *Assume that there exists $R \in \mathbb{R}$, $m \geq n$ such that (if $m = n$ then we impose that $\Psi = 0$),*

$$\text{Ricci}(L) \geq R + \frac{1}{m-n} \nabla \Psi \otimes \nabla \Psi. \quad (21)$$

Then for any smooth 1-forms ω and smooth functions $g > 0$, for any $t \geq 0$,

$$\frac{|R_t \omega|^2}{P_t g} \leq e^{-2Rt} P_t \left(\frac{|\omega|^2}{g} \right) - \frac{2}{m} \int_0^t \frac{e^{-2Ru}}{P_t g} [P_t \delta_\Psi \omega - P_u (d(\log P_{t-u}) \cdot R_{t-u} \omega)]^2 du. \quad (22)$$

Proof

\triangleleft Since M is compact, one can assume that there exists $\varepsilon > 0$ such that $g = f + \varepsilon$ with $f > 0$ and then $\varepsilon \rightarrow 0$ in (22). Let ω be a smooth 1-form and $t \geq 0$. For any $s \in [0, t]$, we define

$$\Lambda(s) = P_s \left(\frac{|R_{t-s} \omega|^2}{P_{t-s} g} \right).$$

For any $s \in [0, 1]$,

$$\Lambda'(s) = P_s \left(L \left(\frac{|R_{t-s} \omega|^2}{P_{t-s} g} \right) - 2 \frac{R_{t-s} \omega \cdot \vec{L} R_{t-s} \omega}{P_{t-s} g} + L P_{t-s} g \frac{|R_{t-s} \omega|^2}{(P_{t-s} g)^2} \right).$$

Since for any smooth functions F, G , $L(FG) = 2\Gamma(F, G) + FLG + GLF$, the identity becomes, with $\eta = R_{t-s} \omega$ and $G = P_{t-s} g$,

$$\begin{aligned} \Lambda'(s) &= P_s \left(-\frac{2}{G^2} \Gamma(|\eta|^2, G) - \frac{2}{G} \eta \cdot \vec{L} \eta + \frac{2}{G^3} |\eta|^2 \Gamma(G) + \frac{1}{G} L |\eta|^2 \right) \\ &= P_s \left[\frac{2}{G} \left(L \frac{|\eta|^2}{2} - \eta \cdot \vec{L} \eta - \Gamma(|\eta|^2, \log G) + |\eta|^2 \Gamma(\log G) \right) \right]. \end{aligned}$$

Corollary 3.6 applied to $\alpha = d \log G$ and $a = -1/2$, implies

$$\Lambda'(s) \geq \frac{2}{m} P_s \left(\frac{1}{G} (\delta_\Psi \eta - d(\log G) \cdot \eta)^2 \right) + 2R P_s \left(\frac{|\eta|^2}{G} \right).$$

Thanks to the Cauchy-Schwarz inequality,

$$P_s \left(\frac{1}{G} (\delta_\Psi \eta - d(\log G) \cdot \eta)^2 \right) \geq \frac{1}{P_s G} \left[P_s (\delta_\Psi \eta - d(\log G) \cdot \eta) \right]^2.$$

Since $P_s G = P_s P_{t-s} g = P_t g$ and from (9), $P_s \delta_\Psi R_{t-s} \omega = P_t \delta_\Psi \omega$, the inequality becomes

$$(\Lambda(s) e^{-2Rs})' \geq \frac{2}{m} \frac{e^{-2Rs}}{P_t g} \left[P_t \delta_\Psi \omega - P_s (d(\log P_{t-s} g) \cdot R_{t-s} \omega) \right]^2.$$

The integration over $s \in [0, t]$ of the previous inequality implies (22). \triangleright

4 Proof of Theorem 2.1

◁ **Regularity assumption.** Let $f_\varepsilon = (P_\varepsilon f + \varepsilon)/(1 + \varepsilon)$ and $g_\varepsilon = (P_\varepsilon g + \varepsilon)/(1 + \varepsilon)$, for $\varepsilon > 0$. The probability measure $f_\varepsilon \mu$ (resp. $g_\varepsilon \mu$) converges weakly to $f \mu$ (resp. $g \mu$), when $\varepsilon \rightarrow 0$ and since M is compact $W_2^2(f_\varepsilon \mu, g_\varepsilon \mu)$ converges to $W_2(f \mu, g \mu)$. The same is also true for $W_2^2(P_t f_\varepsilon \mu, P_t g_\varepsilon \mu)$. Thus, one can assume that f and g are two smooth functions satisfying $f, g \geq \varepsilon$ for some $\varepsilon > 0$.

Contraction inequality. Let $f, g \geq \varepsilon$, be two smooth functions and let $(\rho_s, \omega_s)_{s \in [0,1]}$ be a smooth couple satisfying

$$\begin{cases} \partial_s \rho_s + \delta_\Psi \omega_s = 0 \\ \rho_0 = f, \quad \rho_1 = g. \end{cases} \quad (23)$$

For any $t \geq 0$, from the commutation property (9), the couple $(P_t \rho_s, R_t \omega_s)_{s \in [0,1]}$ satisfies

$$\begin{cases} \partial_s P_t \rho_s + \delta_\Psi R_t \omega_s = 0 \\ P_t \rho_0 = P_t f, \quad P_t \rho_1 = P_t g. \end{cases} \quad (24)$$

One can apply Theorem 3.8 to get

$$\begin{aligned} \int_0^1 \int_M \frac{|R_t \omega_s|^2}{P_t \rho_s} d\mu ds &\leq e^{-2Rt} \int_0^1 \int_M \frac{|\omega_s|^2}{\rho_s} d\mu ds \\ &\quad - \frac{2}{m} \int_0^1 \int_0^t e^{-2Ru} \int_M \frac{[P_t \delta_\Psi \omega_s - P_u(d(\log P_{t-u} \rho_s) \cdot R_{t-u} \omega_s)]^2}{P_t \rho_s} d\mu du ds. \end{aligned} \quad (25)$$

Cauchy-Schwarz inequality implies

$$\begin{aligned} &\int_M \frac{[P_t \delta_\Psi \omega_s - P_u(d(\log P_{t-u} \rho_s) \cdot R_{t-u} \omega_s)]^2}{P_t \rho_s} d\mu \\ &\geq \frac{1}{\int P_t \rho_s d\mu} \left[\int_M (P_t \delta_\Psi \omega_s - P_u(d(\log P_{t-u} \rho_s) \cdot R_{t-u} \omega_s)) d\mu \right]^2 = \left[\int_M d(\log P_{t-u} \rho_s) \cdot R_{t-u} \omega_s d\mu \right]^2, \end{aligned}$$

since $\int_M P_t \delta_\Psi \omega_s d\mu = \int_M \delta_\Psi \omega_s d\mu = 0$. Integrating over $s \in [0, 1]$, thanks again to the Cauchy-Schwartz inequality,

$$\begin{aligned} \int_0^1 \left[\int_M \nabla(\log P_{t-u} \rho_s) \cdot R_{t-u} \omega_s d\mu \right]^2 ds &\geq \left[\int_0^1 \int_M d(\log P_{t-u} \rho_s) \cdot R_{t-u} \omega_s d\mu ds \right]^2 \\ &= [\text{Ent}_\mu(P_{t-u} f) - \text{Ent}_\mu(P_{t-u} g)]^2, \end{aligned}$$

since coming from (24) we have (since M is compact, the next integration by parts is valid)

$$\begin{aligned} \frac{d}{ds} \text{Ent}_\mu(P_{t-u} \rho_s) &= \frac{d}{ds} \int_M P_{t-u} \rho_s \log P_{t-u} \rho_s d\mu = \int_M \partial_s P_{t-u} \rho_s \log P_{t-u} \rho_s d\mu \\ &= - \int_M \delta_\Psi R_{t-u} \omega_s \log P_{t-u} \rho_s d\mu = \int_M R_{t-u} \omega_s \cdot d(\log P_{t-u} \rho_s) d\mu. \end{aligned}$$

The inequality (25) becomes

$$\int_0^1 \int_M \frac{|R_t \omega_s|^2}{P_t \rho_s} d\mu ds \leq e^{-2Rt} \int_0^1 \int_M \frac{|\omega_s|^2}{\rho_s} d\mu ds - \frac{2}{m} \int_0^t e^{-2Ru} [\text{Ent}_\mu(P_{t-u} f) - \text{Ent}_\mu(P_{t-u} g)]^2 du$$

Thanks to the Brenier-Benamou formulation (10) with respect $P_t f \mu$ and $P_t g \mu$ and formula (24),

$$W_2^2(P_t f \mu, P_t g \mu) \leq e^{-2Rt} \int_0^1 \int_M \frac{|\omega_s|^2}{\rho_s} d\mu ds - \frac{2}{m} \int_0^t e^{-2Ru} [\text{Ent}_\mu(P_{t-u} f) - \text{Ent}_\mu(P_{t-u} g)]^2 du.$$

Taking now the infimum over all couples $(\rho_s, \omega_s)_{s \in [0,1]}$ satisfying (23),

$$W_2^2(P_t f \mu, P_t g \mu) \leq e^{-2Rt} W_2^2(f \mu, g \mu) - \frac{2}{m} \int_0^t e^{-2Ru} [\text{Ent}_\mu(P_{t-u} f) - \text{Ent}_\mu(P_{t-u} g)]^2 du,$$

which is the inequality desired changing $t - u$ by u . \triangleright

Acknowledgements. The author warmly thank F. Bolley and A. Guillin for stimulating discussions. This research was supported by the French ANR-12-BS01-0019 STAB project.

References

- [ABC⁺00] C. Ané, S. Blachère, D. Chafaï, P. Fougères, I. Gentil, F. Malrieu, C. Roberto, and G. Scheffer. *Sur les inégalités de Sobolev logarithmiques*, volume 10 of *Panoramas et Synthèses*. Société Math. de France, Paris, 2000.
- [AGS08] L. Ambrosio, N. Gigli, and G. Savaré. *Gradient flows in metric spaces and in the space of probability measures*. Lectures in Math. ETH Zürich. Birkhäuser, Basel, 2008.
- [Bak87] D. Bakry. Étude des transformations de Riesz dans les variétés riemanniennes à courbure de Ricci minorée. In *Séminaire de Probabilités, XXI*, volume 1247 of *Lecture Notes in Math.*, pages 137–172. Springer, Berlin, 1987.
- [Bak94] D. Bakry. L’hypercontractivité et son utilisation en théorie des semigroupes. In *Lectures on probability theory (Saint-Flour, 1992)*, volume 1581 of *Lecture Notes in Math.*, pages 1–114. Springer, Berlin, 1994.
- [BB00] J.-D. Benamou and Y. Brenier. A computational fluid mechanics solution to the Monge-Kantorovich mass transfer problem. *Numer. Math.*, 84(3):375–393, 2000.
- [BGG13] F. Bolley, I. Gentil, and A. Guillin. Dimensional contraction via Markov transportation distance. *To appear in Journal of the London Mathematical Society*, 2013.
- [BGL14] D. Bakry, I. Gentil, and M. Ledoux. *Analysis and geometry of Markov diffusion operators*, volume 348 of *Grund. Math. Wiss.* Springer, Berlin, 2014.
- [BGL15] D. Bakry, I. Gentil, and M. Ledoux. On Harnack inequality and optimal transportation. *To appear in Ann. Sc. Norm. Sup. Pisa*, 2015.
- [DNS09] J. Dolbeault, B. Nazaret, and G. Savaré. A new class of transport distances between measures. *Calc. Var. Part. Diff. Eq.*, 34(2):193–231, 2009.
- [DNS12] J. Dolbeault, B. Nazaret, and G. Savaré. From Poincaré to Logarithmic Sobolev inequalities: a gradient flow approach. *SIAM J. Math. Anal.*, 44(5):3186–3216, 2012.
- [DPGG06] L. De Pascale, M. S. Gelli, and L. Granieri. Minimal measures, one-dimensional currents and the Monge-Kantorovich problem. *Calc. Var. Partial Differential Equations*, 27(1):1–23, 2006.

- [EKS13] M. Erbar, K. Kuwada, and K.-T. Sturm. On the equivalence of the entropic curvature-dimension condition and Bochner’s inequality on metric measure spaces. *Preprint*, 2013.
- [GKO13] N. Gigli, K. Kuwada, and S.-I. Ohta. Heat flow on Alexandrov spaces. *Comm. Pure Appl. Math.*, 66(3):307–331, 2013.
- [Kuw10] K. Kuwada. Duality on gradient estimates and Wasserstein controls. *J. Funct. Anal.*, 258(11):3758–3774, 2010.
- [Kuw13] K. Kuwada. Space-time Wasserstein controls and Bakry-Ledoux type gradient estimates. *Preprint*, 2013.
- [Lic58] A. Lichnerowicz. *Géométrie des groupes de transformations*. Travaux et Recherches Mathématiques, III. Dunod, Paris, 1958.
- [OW05] F. Otto and M. Westdickenberg. Eulerian calculus for the contraction in the Wasserstein distance. *SIAM J. Math. Anal.*, 37(4):1227–1255, 2005.
- [Sav14] G. Savaré. Self-improvement of the Bakry-Émery condition and Wasserstein contraction of the heat flow in $\text{RCD}(K, \infty)$ metric measure spaces. *Discrete Contin. Dyn. Syst.*, 34(4):1641–1661, 2014.
- [Vil09] C. Villani. *Optimal transport, old and new*, volume 338 of *Grund. Math. Wiss.* Springer, Berlin, 2009.
- [vRS05] M.-K. von Renesse and K.-T. Sturm. Transport inequalities, gradient estimates, entropy and Ricci curvature. *Comm. Pure Appl. Math.*, 68:923–940, 2005.
- [Wan04] F.-Y. Wang. *Functional Inequalities, Markov Processes, and Spectral Theory*. Science Press, Beijing, 2004.
- [Wan11] F.-Y. Wang. Equivalent semigroup properties for the curvature-dimension condition. *Bull. Sci. Math.*, 135(6-7):803–815, 2011.