



HAL
open science

A numerical lower bound for the spectral radius of random walks on surface groups

Sebastien Gouezel

► **To cite this version:**

Sebastien Gouezel. A numerical lower bound for the spectral radius of random walks on surface groups. 2013. hal-00873167v1

HAL Id: hal-00873167

<https://hal.science/hal-00873167v1>

Preprint submitted on 15 Oct 2013 (v1), last revised 6 Jun 2014 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A NUMERICAL LOWER BOUND FOR THE SPECTRAL RADIUS OF RANDOM WALKS ON SURFACE GROUPS

SÉBASTIEN GOUËZEL

ABSTRACT. Estimating numerically the spectral radius of a random walk on a non-amenable graph is complicated, since the cardinality of balls grows exponentially fast with the radius. We propose an algorithm to get a bound for below for this spectral radius in Cayley graphs with finitely many cone types (including for instance hyperbolic groups). In the genus 2 surface group, it improves by an order of magnitude the previous best bound, due to Bartholdi.

1. MAIN ALGORITHM

Let Γ be a countable group, generated by a finite symmetric set S of cardinality $|S|$. The simple random walk X_0, X_1, \dots on Γ is defined by $X_0 = e$ the identity of Γ , and $X_{n+1} = X_n s$ with probability $1/|S|$ for any $s \in S$. A crucial numerical parameter of this random walk is its *spectral radius* $\rho = \lim P(X_{2n} = e)^{1/2n}$. Equivalently, denote by W_n the number of words of length n in the generators that represent e in Γ , then $P(X_n = e) = W_n/|S|^n$, so that $\rho = \lim W_{2n}^{1/2n}/|S|$. It is equivalent to study the spectral radius or the *cogrowth* $\lim W_{2n}^{1/2n}$.

The spectral radius is at most 1, and $\rho = 1$ if and only if Γ is amenable. In the free group with d generators, the generating function $\sum W_n z^n$ can be computed explicitly (it is algebraic), and the exact value of the spectral radius follows: $\rho = \sqrt{2d-1}/d$. Since words that reduce to the identity in the free group also reduce to the identity in any group with the same number of generators, one infers that in any group Γ , $\rho \geq 2\sqrt{|S|-1}/|S|$. Moreover equality holds if and only if the Cayley graph of Γ is a tree [Kes59].

In general, there are no explicit formulas for ρ , and even giving precise numerical estimates is a delicate question. In this short note, we will describe an algorithm giving bounds from below on ρ in some classes of groups, particularly for the fundamental group Γ_g of a compact surface of genus $g \geq 2$, given by its usual presentation

$$(1.1) \quad \Gamma_g = \langle a_1, \dots, a_g, b_1, \dots, b_g \mid [a_1, b_1] \cdots [a_g, b_g] = e \rangle.$$

Since there are 4 generators in Γ_2 , the above trivial bound obtained by comparison to the free group gives $\rho \geq 0.661437$. Our main estimate is the following result:

Theorem 1.1. *In the surface group Γ_2 , one has $\rho \geq 0.662772$.*

This improves on the previously best known result, due to Bartholdi [Bar04], giving $\rho \geq \rho_{\text{Bar}} = 0.662421$.¹ Bartholdi's method is to study a specific class of paths from the

Date: October 14, 2013.

Key words and phrases. spectral radius, cogrowth, numerical algorithm, surface groups.

¹Bartholdi claims that $\rho \geq 0.662418$, but implementing his algorithm in multiprecision one gets in fact the better bound $\rho \geq 0.662421$.

identity to itself (called cactus trees), for which he can compute the generating function. The radius of convergence of this generating function is a lower bound for ρ .

The best known upper bound for ρ in Γ_2 is $\rho \leq \rho_{\text{Nag}} = 0.662816$, due to Nagnibeda [Nag97]. Non-rigorous numerical estimates suggest that $\rho = 0.662812\dots$, so the upper bound is still sharper than our lower bound, although our lower bound is an order of magnitude better than the bound of [Bar04]: indeed, $\rho_{\text{Nag}} - \rho_{\text{Bar}} \sim 4.10^{-4}$ while our estimate ρ from Theorem 1.1 satisfies $\rho_{\text{Nag}} - \rho \sim 4.10^{-5}$.

Nagnibeda's upper bound does not rely on a counting argument for closed paths, but on another spectral interpretation of ρ . Indeed, ρ is also the spectral radius of the Markov operator Q on $\ell^2(\Gamma)$ corresponding to the random walk, i.e., the convolution with the probability measure μ which is uniformly distributed on S . It is also the norm of this operator, since it is symmetric. Nagnibeda gets the above upper bound by using a lemma of Gabber about norms of convolution operators on graphs and the precise geometry of Γ_2 .

Our approach to get Theorem 1.1 is very similar to Nagnibeda's. To bound from below the norm of the convolution operator Q , it is sufficient to exhibit one function u (which ought to be close to an hypothetical eigenfunction for the element ρ of the spectrum of Q) for which $\|Qu\| / \|u\|$ is large. This is exactly what we will do.

For any $\alpha < \rho$, the function $u_\alpha = \sum_{n=0}^{\infty} \alpha^n Q^n \delta_e$ is in ℓ^2 , and $\|Qu_\alpha\| / \|u_\alpha\|$ converges to ρ when α tends to ρ . Unfortunately, u_α is not explicit enough. To find estimates, one should rather find an ansatz for the function u , depending on finitely many parameters, and then optimize over these parameters.

A first strategy would be the following: take a very large ball B_n in the Cayley graph, and compute the function u supported in this ball such that $\|Qu\| / \|u\|$ is largest. This gives a lower bound ρ_n on ρ , and ρ_n converges to ρ when n tends to infinity. However, this strategy is computationally not efficient at all: one would need to take a very large n to obtain good estimates (since most mass of u_α is supported close to infinity if α is close to ρ), and the cardinality of B_n grows exponentially with n . On the other hand, it can be implemented in any finitely presented group for which the word problem is solvable (see for instance [ERW12] for examples in Baumslag-Solitar and Thompson groups). We will use a more efficient method, but which requires more assumptions on the group: it should have finitely many cone types.

To illustrate our method of construction of u , let us describe it quickly in the case of the free group \mathbb{F}_d with d generators. The sphere \mathbb{S}^n of radius $n \geq 1$ has cardinality $2d(2d-1)^{n-1}$. Fix some $\alpha < 1/\sqrt{2d-1}$, and define a function u_α by $u_\alpha(x) = \alpha^n$ for $x \in \mathbb{S}^n$, $n \geq 1$. This function belongs to $\ell^2(\mathbb{F}_d)$. We write $x \sim y$ if x and y are neighbors in the Cayley graph of Γ , and $x \rightarrow y$ if $x \sim y$ and $d(e, y) = d(e, x) + 1$. Then

$$\langle Qu_\alpha, u_\alpha \rangle = \frac{1}{2d} \sum_{x \sim y} u_\alpha(x) u_\alpha(y) = \frac{1}{d} \sum_{x \rightarrow y} u_\alpha(x) u_\alpha(y) = \frac{1}{d} \sum_{n=1}^{\infty} (2d-1) \alpha^{2n+1} |\mathbb{S}^n|,$$

since a point in \mathbb{S}^n has $2d-1$ successors in \mathbb{S}^{n+1} . Since $\langle u_\alpha, u_\alpha \rangle = \sum_{n=1}^{\infty} \alpha^{2n} |\mathbb{S}^n|$, we get

$$\langle Qu_\alpha, u_\alpha \rangle = \alpha \frac{2d-1}{d} \langle u_\alpha, u_\alpha \rangle.$$

Hence, $\rho = \|Q\| \geq \alpha(2d - 1)/d$. Letting α tend to $1/\sqrt{2d - 1}$, we finally obtain $\rho \geq \sqrt{2d - 1}/d$.

One can check that the spectral arguments of Nagnibeda give $\rho \leq \sqrt{2d - 1}/d$, hence we have a completely spectral proof of the formula for the spectral radius $\rho = \sqrt{2d - 1}/d$ that was previously obtained by word-counting arguments.

In the free group, it is natural to take a function u that is constant of the sphere \mathbb{S}^n of radius n , since all the points in such a sphere are equivalent: the automorphisms of the Cayley graph of \mathbb{F}_d fixing the identity act transitively on \mathbb{S}^n . In more general groups, for instance surface groups, this is not the case. Intuitively, we would like to take a function that decays exponentially as above, but with different values on different equivalence classes under the automorphism group. However, this automorphism group is finite in the case of surface groups, so instead of true equivalence classes (which are finite), we will consider larger classes, of points that “locally behave in the same way”, and we will construct functions that are constants on such classes of points (leaving only finitely many parameters which one can optimize using a computer).

This intuition is made precise with the notion of *type* of the elements of the group (as in [Nag97]). Let Γ be a countable group generated by a finite symmetric set S . Assume that there are no cycles of odd length, so that any edge can be oriented from the closer point from e to the farther point. Let $\mathcal{S}(x)$ be the set of successors of x , i.e., the points y which are neighbors of x with $d(e, y) = d(e, x) + 1$.

Definition 1.2. Let T be a finite set, let t be a function from Γ to T and let M be a square matrix indexed by T . We say that (T, t, M) is a type system for (Γ, S) if, for all i and j in T , for all but finitely many $x \in \Gamma$ with $t(x) = j$, one has

$$\text{Card}\{y \in \mathcal{S}(x) : t(y) = i\} = M_{ij}.$$

We will often simply say that t is a type system, since it determines T and M .

In other words, if one knows the type of a point x , then one knows the number of successors of each type, thanks to the matrix M . For instance, in \mathbb{F}_d , one can use one single type, with $M_{11} = 2d - 1$: every point but the identity has $2d - 1$ successors.

Using a type system, we will be able to find a lower bound for the spectral radius of the simple random walk. While the argument works in general, it is more convenient to formulate using an additional assumption, which is satisfied for surface groups:

Definition 1.3. A type system (T, t, M) is Perron-Frobenius if the matrix M is Perron-Frobenius, i.e., some power M^n has only positive entries.

The algorithm to estimate the spectral radius follows.

Theorem 1.4. Let (Γ, S) be a countable group with a finite symmetric generating set, and let (T, t, M) be a Perron-Frobenius type system for (Γ, S) .

Define a new matrix \tilde{M} by $\tilde{M}_{ij} = M_{ij}/p_i$, where p_i is the number of predecessors of a point of type i (it is given by $p_j = |S| - \sum_i M_{ij}$). Since it is Perron-Frobenius, its dominating eigenvalue e^v is simple. Let (A_1, \dots, A_k) be a corresponding eigenvector, with positive entries, let D be the diagonal matrix with entries A_i , and let $M' = D^{-1/2}MD^{1/2}$.

Define

$$(1.2) \quad \lambda = \max_{|q|=1} \langle M'q, q \rangle.$$

Then

$$(1.3) \quad \rho \geq \frac{2e^{-v/2}\lambda}{|S|}.$$

Proof. Let $s_n(i) = \text{Card}\{x \in \mathbb{S}^n : t(x) = i\}$. By definition of a type system, if n is large enough (say $n \geq n_0$),

$$p_i s_{n+1}(i) = \sum_{y \in \mathbb{S}^n} \text{Card}\{x \in \mathcal{S}(y) : t(x) = i\} = \sum_j M_{ij} s_n(j).$$

This shows that $s_{n+1} = \tilde{M} s_n$. Therefore, the cardinality of \mathbb{S}^n grows like ce^{nv} for some $c > 0$. Moreover, $s_n(i) = c' A_i e^{nv} + O(e^{n(v-\varepsilon)})$ for some $\varepsilon > 0$.

Take some parameters $b_1, \dots, b_k > 0$ to be chosen later, and let $\alpha < e^{-v/2}$. We define a function u_α by $u_\alpha(x) = \alpha^n b_i$ if $x \in \mathbb{S}^n$ and $t(x) = i$ with $n \geq n_0$. For $n < n_0$, let $u_\alpha(x) = 0$. We have when α tends to $e^{-v/2}$

$$\begin{aligned} \langle Qu_\alpha, u_\alpha \rangle &= \frac{1}{|S|} \sum_{x \sim y} u_\alpha(x) u_\alpha(y) = \frac{2}{|S|} \sum_x \sum_{y \in \mathcal{S}(x)} u_\alpha(x) u_\alpha(y) \\ &= \frac{2}{|S|} \sum_{n \geq n_0} \sum_{i,j} s_n(j) b_j \alpha^n M_{ij} b_i \alpha^{n+1} = \frac{2\alpha}{|S|} \sum_{n \geq n_0} \sum_{i,j} c' A_j e^{nv} b_j \alpha^{2n} M_{ij} b_i + O(1) \\ &= \frac{2\alpha}{|S|} \sum_{i,j} c' A_j b_j M_{ij} b_i / (1 - \alpha^2 e^v) + O(1). \end{aligned}$$

On the other hand,

$$\langle u_\alpha, u_\alpha \rangle = \sum_{n \geq n_0} \sum_i s_n(i) b_i^2 \alpha^{2n} = \sum_{n \geq n_0} \sum_i c' A_i e^{nv} b_i^2 \alpha^{2n} + O(1) = \sum_i c' A_i b_i^2 / (1 - \alpha^2 e^v) + O(1).$$

We have $\rho \geq \langle Qu_\alpha, u_\alpha \rangle / \langle u_\alpha, u_\alpha \rangle$. Comparing the above two equations and letting α tend to $e^{-v/2}$, we get

$$\rho \geq \frac{2e^{-v/2} \sum_{i,j} A_j b_j M_{ij} b_i}{|S| \sum_i A_i b_i^2}.$$

To conclude, we need to optimize in b_i . Writing b_i as $A_i^{-1/2} c_i$, this lower bound becomes

$$\frac{2e^{-v/2} \sum A_j^{1/2} c_j M_{ij} A_i^{-1/2} c_i}{|S| \sum c_i^2}.$$

The last factor is the maximum on the unit sphere of the quadratic form with matrix $M' = D^{-1/2} M D^{1/2}$. This proves (1.3). \square

Remark 1.5. It follows from the formula (1.2) that λ is the maximum on the unit sphere of $\langle M''q, q \rangle$, where M'' is the symmetric matrix $(M' + M'^*)/2$. Since any symmetric matrix is diagonal in some orthogonal basis, it also follows that λ is the maximal eigenvalue of M'' . Hence, it is easy to compute using standard algorithms.

2. APPLICATION TO SURFACE GROUPS




FIGURE 1. Two examples of geodesics from e to x

2.1. Cannon’s types. Consider a countable group with the word distance coming from a finite generating set S . The *cone* of a point x is the set of points y for which there is a geodesic from e to y going through x . The *cone type* of x is the set $\{x^{-1}y\}$, for y in the cone of x . Note that knowing the cone type of a point determines the number of its successors, and the number of its successors having any given cone type. Cannon proved that, in any hyperbolic group, there are finitely many cone types. Therefore, such a group admits a type system in the sense of Definition 1.2. This is in particular the case of the surface groups Γ_g . However, the number of cone types is too large, and it is more convenient for practical purposes to reduce them using symmetries. We obtain Cannon’s canonical types for the surface groups, described in [Can83] or [FP87] as follows.

The hyperbolic plane can be tessellated by regular $4g$ -gons, with $4g$ of them around each vertex. The Cayley graph of Γ_g (with its usual presentation (1.1)) is dual to this (self-dual) tessellation, and is therefore isomorphic to it. Define the type of a point $x \in \Gamma_g$ as the maximal length along the last $4g$ -gon of a geodesic starting from e and ending at x . Beware that one really has to take the maximum: for instance, in Figure 1, the thick geodesic from e to x shares only one edge with the last octagon, while the wiggly one shares two edges. Hence, the type of x is 2.

The type can also be described combinatorially as follows: write $x = c_1 \cdots c_{|x|}$ as a product of minimal length in the generators a_1, \dots, b_g , look at the length n of its longest common suffix with a fundamental relator (i.e., a cyclic permutation of the basic relation $[a_1, b_1] \cdots [a_g, b_g]$ in (1.1) or its inverse: $c_{|x|-n+1} \cdots c_{|x|}$ should be a subword of the basic relation or its inverse, up to cyclic permutation), and take the maximum of all such n over all ways to write $x = c_1 \cdots c_{|x|}$. It is obvious that the geometric and combinatorial descriptions are equivalent, we will mostly rely on the geometric one.

The type of a group element x can be at most $2g$ (otherwise, taking the same path but going the other way around the last $4g$ -gon, one would get a strictly shorter path, contradicting the fact that the initial path is geodesic), and it is 0 only for the identity. Points x of type $i < 2g$ have only one predecessor, and $4g - 1$ successors. Among them, 2 are followers of x on the $4g$ -gon to its left and to its right, while the other ones correspond to newly created $4g$ -gons (whose closest point to e is x). It follows that those points have $4g - 3$ successors of type 1, one of type 2 and one of type $i + 1$. Points of type $2g$ are special, since they have two predecessors (one can reach them with a geodesic either from the left

or from the right of a single $4g$ -gon). They have 2 successors of type 2, corresponding to the extremal outgoing edges of x (they extend the two $4g$ -gons adjacent to both incoming edges to x), and the $4g - 4$ remaining successors are on newly created $4g$ -gons, and are of type 1. See Figure 2 for an illustration in genus 2 (of course, additional octagons should be drawn around all outgoing edges, but since this is notoriously difficult to do in a Euclidean drawing, we have to rely on the reader's imagination).




FIGURE 2. Types of the points in (part of) the Cayley graph of Γ_2

Keeping only the types from 1 to $2g$ (since type 0 only happens for the identity, while Definition 1.2 allows to discard finitely many points), we obtain a type system for Γ_g with $T = \{1, \dots, 2g\}$, where the matrix M has been described in the previous paragraph. For instance, in genus 2,

$$M = \begin{pmatrix} 5 & 5 & 5 & 4 \\ 2 & 1 & 1 & 2 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix}.$$

One can now apply the algorithm of Theorem 1.4 to this matrix to bound the spectral radius of the simple random walk from below. All points but points of type 4 have 1 predecessor, so the matrix \tilde{M} is

$$\tilde{M} = \begin{pmatrix} 5 & 5 & 5 & 4 \\ 2 & 1 & 1 & 2 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1/2 & 0 \end{pmatrix}.$$

The dominating eigenvalue of this matrix is $e^v = 6.979835\dots$ (this is also the growth of the group), while the corresponding eigenvector is

$$A = (0.715987\dots, 0.246211\dots, 0.035274\dots, 0.002526\dots).$$

One gets that the matrix M'' of Remark 1.5 is

$$M'' = \begin{pmatrix} 5 & 3.171316\dots & 0.554905\dots & 0.118814\dots \\ 3.171316\dots & 1 & 1.510223\dots & 0.101307\dots \\ 0.554905\dots & 1.510223\dots & 0 & 1.868132\dots \\ 0.118814\dots & 0.101307\dots & 1.868132\dots & 0 \end{pmatrix},$$

with dominating eigenvalue $\lambda = 7.000902\dots$. Finally,

$$(2.1) \quad \rho \geq \frac{2e^{-v/2}\lambda}{8} = 0.662477\dots$$

This is already slightly better than Bartholdi's estimate $\rho \geq 0.662421$, but much weaker than the estimate $\rho \geq 0.662772$ that we claimed in Theorem 1.1 (to be compared with the true value $\rho \sim 0.662812$).

In the next sections, we will explain how to get better estimates by using different type systems, that distinguish between more points (but, of course, give rise to larger matrices M and therefore to more computer-intensive computations).

Remark 2.1. Using cone types instead of Cannon's canonical types does not give rise to better estimates for ρ (although the number of types is much larger). Indeed, if some type system t' is obtained from some type system t by quotienting by some symmetries of t , then the dominating eigenvector of $M(t)$, being unique, is invariant by those symmetries and reduces to the dominating eigenvector of $M(t')$. It follows that the dominating eigenvector of $M''(t)$ is also invariant by those symmetries, and that the dominating eigenvalue of $M''(t)$ is the same as that of $M''(t')$. Hence, the estimates on ρ given by Theorem 1.1 for t and t' are the same.

2.2. Suffix types. There are many ways to define new type systems in surface groups, that separate more points. If a type system is finer than another one, then the estimate on the spectral radius coming from Theorem 1.4 is better, but the matrix involved in the computation is larger. To get manageable estimates, we should find the right balance.

In this paragraph, we describe a very simple extension of Cannon's canonical type systems in surface groups, that we call suffix types. Given a point $x \in \Gamma_g$, there can be several geodesics from e to x . Consider the longest ending that is common to all these geodesics, say x_{n-k+1}, \dots, x_n (with $x_n = x$), and define the suffix type of x to be

$$t_{\text{suff}}(x) = (t(x_n), t(x_{n-1}), \dots, t(x_{n-k+1})),$$

where t is the canonical type of Cannon.

For any x , $t_{\text{suff}}(x)$ is easy to compute inductively:

- If $t(x) = 0$, i.e., $x = e$, then $t_{\text{suff}}(x) = (0)$.
- If x is of type $2g$, it has two predecessors, so the common ending to all geodesics ending at x is simply x , and $t_{\text{suff}}(x) = (2g)$.
- If $t(x) \in \{1, \dots, 2g-1\}$, then x has a unique predecessor z . The common ending to all geodesics ending at x is the common ending to all geodesics ending at z , followed with x . Hence, $t_{\text{suff}}(x) = (t(x), t_{\text{suff}}(z))$.

It also follows from this description that, if one knows $t_{\text{suff}}(x)$, it is easy to determine $t_{\text{suff}}(y)$ for any successor y of x : if $t(y) = 2g$, then $t_{\text{suff}}(y) = (2g)$, otherwise x is the only predecessor of y and $t_{\text{suff}}(y) = (t(y), t_{\text{suff}}(x))$.

We have shown that t_{suff} shares most properties of type systems as described in Definition 1.2, except that it does not take its values in a finite set. To ensure this additional property, one should truncate the suffix type. For instance, one can fix some maximal length k , and define the k -truncated suffix type $t_{\text{suff}}^{(k)}(x)$ by keeping only the first k elements of $t_{\text{suff}}(x)$ if its length is $> k$.

The following proposition is obvious from the previous discussion:

Proposition 2.2. *For any $k \geq 1$, the k -truncated suffix type system $t_{\text{suff}}^{(k)}$ is a (Perron-Frobenius) type system in the sense of Definition 1.2.*

The matrix size increases with k , but the estimates on the spectral radius following from Theorem 1.4 get better. For instance, in Γ_2 , for $k = 5$, the matrix size is 148, and we get $\rho \geq 0.662694$.

A drawback of this truncation process is that it truncates uniformly, independently of the likeliness of the type, while it should be more efficient to extend mostly those types that are more likely to happen. This intuition leads to another truncation process: fix a system of weights $w = (w_0, \dots, w_{2g}) \in [0, +\infty)^{2g+1}$, a threshold k , and truncate a suffix type (t_0, t_1, \dots) at the smallest n such that $t_0 + \dots + t_n > k$. This gives another type system denoted by $t_{\text{suff}}^{(k,w)}$ ($t_{\text{suff}}^{(k)}$ corresponds to the weights $w = (1, \dots, 1)$ and the threshold $k - 1$). Define for instance a weight system \bar{w} by $\bar{w}_0 = 1$ and $\bar{w}_i = i$ for $i \geq 1$: the corresponding type system $t_{\text{suff}}^{(k,\bar{w})}$ truncates more quickly the suffix types involving a lot of large types, that happen less often in the group. Hence, it should give a smaller matrix than the naive truncation only according to length, while retaining a comparatively good estimate for the spectral radius.

This intuition is correct: for instance, in Γ_2 , using $t_{\text{suff}}^{(k,\bar{w})}$ with $k = 6$, one gets a matrix with size 109 and an estimate $\rho \geq 0.662697$: the matrix is smaller than for the naive truncation $t_{\text{suff}}^{(5)}$, while the estimate on the spectral radius is better.

We can now push the computations, to a larger matrix size: using in Γ_2 the weight \bar{w} and the truncation threshold $k = 25$, one obtains a type system where the matrix is of size 2,774,629, and the following estimate on the spectral radius:

Proposition 2.3. *In Γ_2 , one has $\rho \geq 0.662757$.*

This is definitely better than (2.1), but not yet as good as Theorem 1.1.

A few comments on the practical implementation. There are three main steps in the algorithm of Theorem 1.4:

- (1) Compute the matrix M corresponding to the type system.
- (2) Find the eigenvector A , to define the matrix M' .
- (3) Find the maximal expansion rate of M' .

Computing the matrix of the type system is a matter of simple combinatorics: we explained above all the transitions from one suffix type to the next ones. The resulting matrix M is very sparse: each type has at most $2g$ successors. However, it is extremely large, so that finding the eigenvector A and then the maximal expansion rate of M' might seem computationally expensive. This is not the case, as we explain now.

Let $A^{(0)}$ be the eigenvector for the original Cannon type, so that $\text{Card}\{x \in \mathbb{S}^n : t(x) = i\} \sim A_i^{(0)} e^{nv}$. Let also $M^{(0)}$ be the matrix for the original Cannon type. Given a new type $\bar{i} = (i_0, \dots, i_m)$, the entry $A_{\bar{i}}$ of the eigenvector A for the new type $t_{\text{suff}}^{(w,k)}$ is such that $\text{Card}\{x \in \mathbb{S}^n : t_{\text{suff}}^{(w,k)}(x) = \bar{i}\} \sim A_{\bar{i}} e^{nv}$. Such a point x can be obtained uniquely by starting from a point $y \in \mathbb{S}^{n-m}$ with type i_m , and then taking successors respectively of type i_{m-1}, \dots, i_0 . Hence,

(2.2)

$$\text{Card}\{x \in \mathbb{S}^n : t_{\text{suff}}^{(w,k)}(x) = (i_0, \dots, i_m)\} = M_{i_0 i_1}^{(0)} \cdots M_{i_{m-1} i_m}^{(0)} \text{Card}\{y \in \mathbb{S}^{n-m} : t(y) = i_m\}.$$

It follows that the new eigenvector is given by

$$A_{\bar{i}} = M_{i_0 i_1}^{(0)} \cdots M_{i_{m-1} i_m}^{(0)} A_{i_m}^{(0)} e^{-mv}.$$

This shows that A is very easy to compute.

By Remark 1.5, to determine the maximal expansion rate λ of M' , it suffices to find the maximal eigenvalue of $M'' = (M' + M'^*)/2$. This matrix is real, symmetric, with nonnegative coefficients, and it is Perron-Frobenius (i.e., it has one single maximal eigenvalue). It follows that, for any vector v with positive coefficients, $\lambda = \lim \|M''^n v\| / \|M''^{n-1} v\|$ (and moreover this sequence is non-decreasing, see for instance [Woe00, Corollary 10.2]). Hence, one can readily estimate λ from below, by starting from a fixed vector v and iterating M'' . Again, there is no issue of instability or complexity.

2.3. Essential types. To improve the suffix types, to separate even more points, one can for instance replace the canonical Cannon types with the true cone types. However, the matrix size increases so quickly that this is not usable in practice. Moreover, this does not solve the main problem of suffix types: they do not separate at all points with Cannon type $2g$, although such points are clearly not always equivalent. In this paragraph, we introduce a new type system that can separate such points, that we call the *essential type*.

The basic idea (that will not work directly) is to memorize not only the common ending of all geodesics ending at a point x , but all the parts that are common to such geodesics: i.e., the sequence $F_{\text{ess}}(x) = (x_0 = e, x_1, \dots, x_n = x)$ (with $n = |x|$) where $x_i = *$ if there are two geodesics from e to x that differ at position i , and x_i is the point that is common to all those geodesics at position i otherwise. We then associate to x the sequence $t_{\text{ess}}(x) = (t(x_n), t(x_{n-1}), \dots, t(x_0))$ where $t(x_i)$ is the Cannon type of x_i if $x_i \neq *$, and $t(*) = *$.

The problem with this notion is that $t_{\text{ess}}(x)$ does not determine $t_{\text{ess}}(y)$ for y a successor of x : in Figure 3, the points x and x' have the same essential type $(3, 2, 1, 3, 2, 1, 0)$, while their successors y and y' have respective essential types $(4, *, *, *, 3, 2, 1, 0)$ and $(4, *, *, *, *, *, *, 0)$ (this follows from the fact that the thick paths and wiggly paths are geodesics). This shows that, as we have defined it, t_{ess} can not be used to define a type system.

This problem can be solved if we do not use directly the Cannon types in the definition of t_{ess} , but a slightly refined notion, the Cannon modified type, taking values in $\{1, \dots, 2g, 1', 2'\}$. The modified type of a point is the same as its Cannon type, except for some points of Cannon types 1 and 2, that have modified types $1'$ and $2'$ respectively: considering any point y of type $2g - 1$, it has a unique successor z of type $2g$, and a unique




FIGURE 3. x and x' have the same essential type, contrary to y and y' .

successor x of type 1 that is on the same $2g$ -gon as z . We say that x is of modified type $1'$. Moreover, x has a unique successor of type 2 that is also on the same $2g$ -gon as z , we say that it is of modified type $2'$. See Figure 4 for an example in genus 2. By definition, the modified Cannon type t' is also a type system. The transition matrix is the same as for the usual Cannon type, except that

- A point of type $2g - 1$ has one successor of type $1'$, one successor of type $2g$, one successor of type 2, and $4g - 4$ successors of type 1.
- A point of type $1'$ has one successor of type 2, one successor of type $2'$, and $4g - 3$ successors of type 1.
- A point of type $2'$ has one successor of type 2, one successor of type 3, and $4g - 3$ successors of type 1.

We define $t_{\text{ess}}(x) = (t'(x_n), \dots, t'(x_0))$ where $(x_0, \dots, x_n) = F_{\text{ess}}(x)$ and $t'(*) = *$.

Proposition 2.4. *The essential type $t_{\text{ess}}(x)$ of a point x determines the essential type of its successors.*

Proof. We argue by induction on $|x| = d(x, e)$.

Consider a point x , and one of its successors y . If the type of y is not $2g$, then x is the unique predecessor of y , and $t_{\text{ess}}(y) = (t'(y), t_{\text{ess}}(x))$. Assume now that $t'(y) = 2g$ (so that $t'(x) = 2g - 1$). Let $z_{2g-1} = x$, and define inductively z_i as the unique predecessor of z_{i+1} for $i \geq 1$, so that $F(x) = (e, \dots, z_1, z_2, \dots, z_{2g-1} = x)$. Those points are on a common $2g$ -gon R , and $t(z_i) = i$ for $i \geq 2$, while $t(z_1)$ can be anything. In the same way, let $\tilde{z}_{2g-1}, \dots, \tilde{z}_1$ be the successive preimages of y going around R in the other direction. They also satisfy $t(\tilde{z}_i) = i$ for $i \geq 2$.

If $t(z_1) \neq 2g$, then z_1 has a unique preimage z_0 , which also belongs to R . Moreover, z_0 is the unique closest point to e on R . The path $P = (z_0, z_1, \dots, z_{2g-1}, y)$ is a geodesic path going around R in one direction, and $\tilde{P} = (z_0, \tilde{z}_1, \dots, \tilde{z}_{2g-1}, y)$ is also geodesic and goes around R in the other direction. If, along \tilde{P} , all points different from z_0, y have type $< 2g$




FIGURE 4. Modified Cannon types of the points in (part of) the Cayley graph of Γ_2




FIGURE 5. Determining the essential type of y from that of x

(so that they have a unique preimage), then any geodesic from e to y has to follow either P or \tilde{P} , so that $t_{\text{ess}}(y) = (2g, *, \dots, *, t_{\text{ess}}(z_0))$, with $2g - 1$ ambiguous points. See the first part of Figure 5 for an illustration.

The only way to have a point of type $2g$ along \tilde{P} is if $t(z_0) = 2g - 1$ and $t(\tilde{z}_1) = 2g$, since $t(\tilde{z}_i) = i$ for $i \geq 2$ (see the second part of Figure 5). By definition of the modified type, this happens exactly when $t'(z_1) = 1'$ and $t'(z_2) = 2'$. In this case, a geodesic from e to y can either go through z_0 and then follow P , or go through \tilde{z}_1 and then follow $(\tilde{z}_2, \dots, \tilde{z}_{2g-1}, y)$. In the first case, if one truncates the geodesic when it reaches z_0 and then adds the edge $[z_0 \tilde{z}_1]$,

one gets a geodesic from e to \tilde{z}_1 . It follows that the essential type of y is $(2g, *, \dots, *, \widehat{t_{\text{ess}}(\tilde{z}_1)})$, where $\widehat{t_{\text{ess}}(\tilde{z}_1)}$ is the essential type of \tilde{z}_1 minus its first entry (i.e., the type $2g$ of \tilde{z}_1), and where there are $2g - 1$ stars. Since \tilde{z}_1 is a successor of z_0 , the induction hypothesis ensures that its essential type can be determined from that of z_0 , which is given by that of x . When $t(z_1) \neq 2g$, we have shown that in all situations $t_{\text{ess}}(x)$ determines $t_{\text{ess}}(y)$ in an algorithmic way.

Assume now that $t(z_1) = 2g$ (see the third part of Figure 5). This case is very similar to the previous one. To reach y , one has to reach z_1 or its preimage on R , and then reach y by going around R in one direction or the other. It follows that $t_{\text{ess}}(y) = (2g, *, \dots, *, \widehat{t_{\text{ess}}(z_1)})$ as above. Since $t_{\text{ess}}(z_1)$ is obtained by removing the last $2g - 2$ entries of $t_{\text{ess}}(x)$, it follows again that $t_{\text{ess}}(x)$ determines $t_{\text{ess}}(y)$. \square

For $k > 0$, let $t_{\text{ess}}^{(k)}(x)$ be the truncated essential type, obtained by keeping the first k entries of the essential type of x . The above proof also shows that $t_{\text{ess}}^{(k)}(x)$ determines $t_{\text{ess}}^{(k+1)}(y)$ (and therefore $t_{\text{ess}}^{(k)}(y)$) for any successor y of x . In the same way, if one considers a truncation according to a weight $w = (w_0, w_1, \dots, w_{2g}, w_{1'}, w_{2'}, w_*)$ and a threshold k , then $t_{\text{ess}}^{(k,w)}(x)$ determines $t_{\text{ess}}^{(k,w)}(y)$ for any successor y of x , if the weight w_* is maximal among all weights. This last requirement is necessary for the following reason: the essential type of y can be obtained from that of x by replacing some entries with stars; if this could decrease the weight of the resulting sequence, one might need to look further to determine $t_{\text{ess}}^{(k,w)}(y)$, and $t_{\text{ess}}^{(k,w)}(x)$ might not be sufficient.

Under these conditions, it follows that $t_{\text{ess}}^{(k)}$ and $t_{\text{ess}}^{(k,w)}$ are (Perron-Frobenius) type systems, in the sense of Definition 1.2. Hence, we can use Theorem 1.4 to estimate the spectral radius of the corresponding random walk. Again, it turns out that it is more efficient to truncate using weights than length.

In genus 2, taking the weights $w = (1, 2, 3, 4, 1, 2, 4)$ and the threshold $k = 25$, we obtain a matrix of size 8,999,902. The corresponding bound on the spectral radius is $\rho \geq 0.662772$, proving Theorem 1.1. Those bounds were obtained on a personal computer with a memory of 12GB (memory is indeed the main limiting factor, since one should store all truncated essential types to create the matrix M). With more memory, one would get better estimates, but it is unlikely that those estimates converge to the true spectral radius when k tends to infinity: to recover it, it is probably necessary to distinguish even more points, for instance by using Cannon's cone types instead of the canonical types (but this would become totally impractical).

In higher genus, here are the bounds we obtain:

Theorem 2.5. *In genus 3, using $t_{\text{ess}}^{(k,w)}$ with $w = (1, 2, 3, 4, 5, 6, 1, 2, 6)$ and $k = 25$, we get a matrix of size 7,307,293 and the estimate $\rho \geq 0.5527735593$.*

In genus 4, using $t_{\text{suff}}^{(k,w)}$ $w = (1, 2, 3, 4, 5, 6, 7, 8)$ and $k = 24$, we get a matrix of size 4,120,495 and the estimate $\rho \geq 0.48412292068$.

When the genus increases, the groups look more and more like free groups. This means that the spectral radius is very close to that of the random walk on a tree with the corresponding number of generators (i.e., $\sqrt{4g - 1}/(2g)$ in general, specializing to 0.55277079 in genus 3 and 0.48412291827 in genus 4), and to get a significant improvement one needs to

take very large matrices. The paths counting arguments of Bartholdi [Bar04], on the other hand, are more and more precise when the groups looks more and more like a free group: in genus 3, he gets $\rho \geq 0.5527735401$, which is just slightly worse than our estimate, while requiring considerably less computer power. In genus 4, he gets $\rho \geq 0.48412292074$, which is already better than our estimate, and the situation is certainly the same in higher genus.

In genus 3, the upper bound of Nagnibeda [Nag97] is $\rho \leq 0.552792$, while our lower bound (or Bartholdi's) is much closer to the naive lower bound coming from the free group. It is unclear which one is closer to the real value of the spectral radius.

For the practical implementation, as in the end of Paragraph 2.2, it is important to know the asymptotics of the number of points on \mathbb{S}^n having a given truncated essential type. We illustrate how to compute such asymptotics in three significant examples, that can be combined to handle the general case.

- Assume first that the type (i_0, \dots, i_m) does not contain any ambiguous letter, i.e., $i_\ell \neq *$ for all ℓ . In this case, the formula (2.2) still holds.
- Assume now that the type is of the form $(i, *, \dots, *, j)$ for some types $i, j \neq *$, and some number N of stars. Let x be a point with the above truncated essential type, on a sphere \mathbb{S}^n , and let y be the point of modified type j , on the sphere \mathbb{S}^{n-N-1} , such that any geodesic from e to x goes through y . Since the type $2g$ is the only one to have several predecessors, necessarily $i = 2g$. On the other hand, j can be any type in $\{1, 1', 2, 2', 3, \dots, 2g\}$.

The discussion in the proof of Proposition 2.4 (see in particular the last 2 cases of Figure 5) implies that $N = (2g - 1)m$, for some integer m : a geodesic from e to x goes through y , and then it follows m $2g$ -gons.

Let us first study the case $m = 1$. Consider a point y of type j , and a $2g$ -gon R based at y (i.e., y is the closest point to e on this $2g$ -gon). There are $4g - 2$ such R if $j \neq 2g$ and $4g - 3$ if $j = 2g$ (since a point of type $2g$ has two incoming edges). On R , consider the point x that is the farthest from e : it has type $2g$, and one can reach it from y by going around R in one direction or the other. It follows that

$$\begin{aligned} \text{Card}\{x \in \mathbb{S}^n : t_{\text{suff}}^{(w,k)}(x) = (2g, *, \dots, *, j) \text{ with } N = 2g - 1 \text{ stars}\} \\ = a_j \text{Card}\{y \in \mathbb{S}^{n-N-1} : t'(y) = j\}, \end{aligned}$$

where $a_j = 4g - 3$ if $j \in \{2g - 1, 2g\}$, and $a_j = 4g - 2$ otherwise. The case of a point y of type $j = 2g - 1$ is special since, on the $2g$ -gon R containing the successors of y of types $1'$ and $2'$, one of the geodesic paths from y to the opposite point x goes through a vertex of type $2g$, giving rise to further ambiguities. Hence, this $2g$ -gon should be discarded from the above counting, leaving only $2g - 3$ suitable $2g$ -gons.

In the case of a general $m \geq 1$, the number of points x corresponding to a given point y of type j may be obtained first by choosing a suitable $2g$ -gon R_1 based at y (giving a_j choices). Further ambiguities can only be obtained by choosing one of the two predecessors (of type $2g - 1$) of the point that is opposite to y on R_1 , and then following the $2g$ -gon R_2 based at this point and containing its successors of type $1'$ and $2'$. This gives 2 choices for R_2 , then 2 more choices for the next $2g$ -gon R_3 , and

so on. In the end, we obtain

$$\begin{aligned} \text{Card}\{x \in \mathbb{S}^n : t_{\text{suff}}^{(w,k)}(x) = (2g, *, \dots, *, j) \text{ with } N = (2g-1)m \text{ stars}\} \\ = a_j 2^{m-1} \text{Card}\{y \in \mathbb{S}^{n-N-1} : t'(y) = j\}. \end{aligned}$$

- Finally, assume that the type is $(i, *, \dots, *)$ with some number N of stars (the situation is very similar to the previous one). Necessarily, as above, $i = 2g$. Write $N = (2g-1)m + k$ for some $k \in \{1, \dots, 2g-1\}$.

For $m = 0$, a point of type $2g$ has two predecessors, so there are always ambiguities regarding his first $2g-1$ ancestors. Hence, the set of points we are considering is simply the set of points of type $2g$ on the sphere \mathbb{S}^n , and there is nothing to do.

For $m = 1$, there can be further ambiguities only if x has an ancestor x' of generation $2g-1$ that has type $2g$, and x is at the tip of the $2g$ -gon R based at one of the two predecessors of x' (of type $2g-1$), and containing x' . There are two choices for R .

Proceeding inductively in the case of a general m , we get

$$\begin{aligned} \text{Card}\{x \in \mathbb{S}^n : t_{\text{suff}}^{(w,k)}(x) = (2g, *, \dots, *) \text{ with } N = (2g-1)m + k \text{ stars}\} \\ = 2^m \text{Card}\{y \in \mathbb{S}^{n-(2g-1)m} : t(y) = 2g\}. \end{aligned}$$

A general truncated essential type is the concatenation of successive types of the form just described in the example. We can thus count the number of points with a given type just by combining the above formulas.

REFERENCES

- [Bar04] Laurent Bartholdi, *Cactus trees and lower bounds on the spectral radius of vertex-transitive graphs*, Random walks and geometry, Walter de Gruyter GmbH & Co. KG, Berlin, 2004, pp. 349–361. MR2087788. Cited pages 1, 2, and 13.
- [Can83] James W. Cannon, *The growth of the closed surface groups and the compact hyperbolic Coxeter groups*, Preprint, 1983. Cited page 5.
- [ERW12] Murray Elder, Andrew Rechnitzer, and Thomas Wong, *On the cogrowth of Thompson’s group F* , Groups Complex. Cryptol. **4** (2012), 301–320. MR3043436. Cited page 2.
- [FP87] William J. Floyd and Steven P. Plotnick, *Growth functions on Fuchsian groups and the Euler characteristic*, Invent. Math. **88** (1987), 1–29. MR877003. Cited page 5.
- [Kes59] Harry Kesten, *Symmetric random walks on groups*, Trans. Amer. Math. Soc. **92** (1959), 336–354. MR0109367. Cited page 1.
- [Nag97] Tatyana Nagnibeda, *An upper bound for the spectral radius of a random walk on surface groups*, Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI) **240** (1997), 154–165, 293–294. MR1691645. Cited pages 2, 3, and 13.
- [Woe00] Wolfgang Woess, *Random walks on infinite graphs and groups*, Cambridge Tracts in Mathematics, vol. 138, Cambridge University Press, Cambridge, 2000. MR1743100. Cited page 9.

IRMAR, CNRS UMR 6625, UNIVERSITÉ DE RENNES 1, 35042 RENNES, FRANCE
E-mail address: `sebastien.gouezel@univ-rennes1.fr`