



The right angle to look at orthogonal sets

Frank Olaf Wagner

► To cite this version:

| Frank Olaf Wagner. The right angle to look at orthogonal sets. 2015. hal-00876033v3

HAL Id: hal-00876033

<https://hal.science/hal-00876033v3>

Preprint submitted on 1 Oct 2015 (v3), last revised 6 Jul 2016 (v5)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

THE RIGHT ANGLE TO LOOK AT ORTHOGONAL SETS

FRANK O. WAGNER

ABSTRACT. If X and Y are orthogonal hyperdefinable sets such that X is simple, then any group G interpretable in $X \cup Y$ has a normal hyperdefinable X -internal subgroup N such that G/N is Y -internal; N is unique up to commensurability. In order to make sense of this statement, local simplicity theory for hyperdefinable sets is developed.

INTRODUCTION

Two definable sets X and Y in some structure are said to be *orthogonal* if every definable subset of $X \times Y$ is a finite union of *rectangles*, i.e. of subsets of the form $U \times V$ with $U \subseteq X$ and $V \subseteq Y$ definable. It easily follows that if X and Y are orthogonal groups, every definable subgroup of $X \times Y$ is of the form $U \times V$ with $U \leq X$ and $V \leq Y$ subgroups. However, the situation is considerably more complicated for a group G definable, or more generally interpretable, in $X \cup Y$, as it need not be the sum of a group interpretable in X and a group interpretable in Y . In fact, an example by Berarducci and Mamino [2, Example 1.2] shows that G need not have *any* subgroup interpretable in either X or Y . However, they prove [2, Theorem 7.1] that if X is superstable of finite and definable Lascar rank, then any group G interpretable in $X \cup Y$ has a normal subgroup N interpretable in X , such that G/N is interpretable in Y .

In this paper we shall generalize their result to the case where X is merely simple. In this context, definability has to be replaced by type-definability, as even for a definable group the tools of simplicity theory in general only yield type-definable subgroups. In fact, we even have to study *hyperdefinable* groups, since the quotient G/N , for N

2000 *Mathematics Subject Classification.* 03C45.

Key words and phrases. orthogonality; simplicity; internality.

Partially supported by ANR-09-BLAN-0047 Modig and ANR-13-BS01-0006 ValCoMo.

type-definable, will be of that form. We therefore put ourselves in the hyperdefinable context and assume right from the start that our orthogonal sets X and Y are merely hyperdefinable. To this end, we shall include a quick development of hyperdefinability in section 1, and of local simplicity theory for hyperdefinable sets in section 5. Note that we do recover the theorem by Berarducci and Mamino even for general supersimple (and definable) X (Corollary 6.2).

Another problem is that of parameters. The usual hypothesis would be that of *stable embedding*, i.e. that every hyperdefinable subset of X is hyperdefinable with parameters in X . We shall circumvent this issue by only ever considering parameters from $X \cup Y$, as orthogonality automatically yields stable embeddedness of X and of Y in $X \cup Y$.

We shall work in a big κ -saturated and strongly κ -homogeneous monster model \mathfrak{M} , where κ is bigger than any cardinality we wish to consider. We shall not usually distinguish between elements and tuples.

1. HYPERIMAGINARIES

- Definition 1.1.** • A *countable equivalence relation* is an equivalence relation given by the conjunction of countably many formulas (and hence only using countably many parameters and variables); it is *over* a set A of parameters if the formulas only use parameters from A .
- A *hyperimaginary* (element) of *type* E is the class a_E of some tuple a (of the right length) modulo a countable equivalence relation E over \emptyset .
 - A hyperimaginary e is *definable* over some set B of hyperimaginaries if every automorphisms of the monster model which fixes B pointwise fixes also e ; it is *bounded* over B if its orbit under the group of automorphisms fixing B pointwise has bounded size (smaller than the saturation degree of the monster model). The *hyperimaginary definable closure* $\text{dcl}^{\text{heq}}(B)$ of B is the set of hyperimaginaries definable over B ; the *hyperimaginary bounded closure* $\text{bdd}(B)$ of B is the set of all hyperimaginaries bounded over B . Clearly, both dcl^{heq} and bdd are idempotent operators.¹
 - Two hyperimaginaries are *equivalent* if they are interdefinable.
 - If e is hyperimaginary, a *representative* for e is any real (or imaginary) tuple a with $e \in \text{dcl}^{\text{heq}}(a)$.

¹By compactness, an imaginary element in $\text{bdd}(B)$ is already in the algebraic closure $\text{acl}(B)$. So there is no need for a superscript bdd^{heq} .

Remark 1.2. (1) If E is an arbitrary type-definable equivalence relation over \emptyset (given by an intersection of arbitrary size, on tuples of arbitrary length), it is easy to see that E is equivalent to a conjunction of subintersections E_i , each one defining a countable equivalence relation on a countable subtuple x_i . So

$$aEb \Leftrightarrow \forall i \ a_i E_i b_i.$$

This means that an automorphism fixes a_E if and only iff it fixes $(a_i)_{E_i}$ for all i , and we can replace a_E by the sequence $((a_i)_{E_i})_i$ of hyperimaginaries.

- (2) If E_A is a countable equivalence relation over A , we consider the countable equivalence relation

$$(xX)F(yY) \Leftrightarrow (X = Y \models \text{tp}(A) \wedge xE_X y) \vee (xX) = (yY)$$

where we only consider the countable subset of A actually occurring in the definition of E_A . Then for any a the class a_{E_A} is fixed by an automorphism fixing A iff and only iff $(aA)_F$ is fixed, and we can use the hyperimaginary $(aA)_F$ instead of a_{E_A} .

- (3) If E is a countable partial type over A which defines an equivalence relation on some partial type π over A , then by compactness there is a countable subtype π_0 such that E defines an equivalence relation on π_0 . Then

$$xFy \Leftrightarrow (\pi_0(x) \wedge \pi_0(y) \wedge xEy) \vee x = y$$

is a countable equivalence relation over A extending E .

If $E = (E_i : i \in I)$ is a sequence of countable equivalence relations over \emptyset and $a = (a_i : i \in I)$ is a sequence of tuples of the right length, we put $a_E = ((a_i)_{E_i} : i \in I)$. Similarly, we write aEb if $a_i E_i b_i$ for all $i \in I$.

Definition 1.3. Let a_E and b_F be tuples of hyperimaginaries. The type $\text{tp}(a_E/b_F)$ is given by all partial types over B of the form

$$\exists yz [xEy \wedge zFb \wedge \varphi(y, z)]$$

true of a . It is easy to see that (in the monster model) two tuples of hyperimaginaries of type E have the same type over b_F if and only if they are conjugate by an automorphism fixing b_F .

Note that the type of a hyperimaginary over b_F is just a maximal E -invariant partial real type over b invariant under automorphisms fixing b_F . For any two representatives of b_F , any two such types are equivalent. We shall say that a partial type $\pi(y)$ is a partial E -type if $\pi(y)$ is E -invariant.

Definition 1.4. A set X is *hyperdefinable* over some parameters A if it is of the form Y/E , where Y is a type-definable set in countably many variables and E a countable equivalence relation on Y , both over A . We denote by X^{heq} the collection of all hyperimaginaries in the definable closure of A and some tuple from X .

For the rest of the paper, all tuples and parameter sets are hyperimaginary, unless stated otherwise. We shall not distinguish between elements and tuples of elements from a set.

2. ORTHOGONALITY

Definition 2.1. Let X, Y be A -hyperdefinable sets in some structure \mathfrak{M} . We say that X and Y are *orthogonal* over A , denoted $X \perp_A Y$, if for any tuples a from X and b from Y , the partial type $\text{tp}(a/A) \cup \text{tp}(b/A)$ determines $\text{tp}(ab/A)$.

Remark 2.2. Note that we do not require X (or Y) to be *stably embedded*, i.e. that every hyperdefinable subset of X be hyperdefinable with parameters in $A \cup X$. In a stable theory, every hyperdefinable subset is stably embedded, but this need not hold in general. We shall compensate for the lack of stable embeddedness by restricting our additional parameters to $X^{heq} \cup Y^{heq}$.²

Example 2.3. If \mathfrak{M}_1 and \mathfrak{M}_2 are two structures and $\mathfrak{N} = \mathfrak{M}_1 \times \mathfrak{M}_2$ with a predicate X for \mathfrak{M} and a predicate Y for \mathfrak{N} , then X and Y are orthogonal in \mathfrak{N} .

Remark 2.4. If X and Y are orthogonal type-definable sets over A and $Z \subseteq X^k \times Y^\ell$ is relatively A -definable, then Z is a finite union of rectangles $A_i \times B_i$, where $A_i \subseteq X^k$ and $B_i \subseteq Y^\ell$ are relatively A -definable.

Proof: For any $z = (x, y) \in Z$ we have that

$$\text{tp}(x/A) \cup \text{tp}(y/A) \vdash (x, y) \in Z.$$

By compactness there are relatively A -definable subsets $A_z \subseteq X^k$ in $\text{tp}(x/A)$ and $B_z \subseteq Y^\ell$ in $\text{tp}(y/A)$ with $A_z \times B_z \subseteq Z$. Again by compactness, finitely many of these rectangles suffice to cover Z . \square

Remark 2.5. $X \perp_A X$ if and only if $X \subseteq \text{dcl}^{heq}(A)$.

²See Proposition 2.6. In a stable theory, $X \perp_A Y$ implies $X \perp_B Y$ for any $B \supseteq A$ (full orthogonality).

Proof: For any $x, x' \in X$ we have

$$\text{tp}(x/A) \cup \text{tp}(x'/A) \vdash \text{tp}(x, x'/A).$$

If $x \notin \text{dcl}^{\text{heq}}(A)$ choose $x' \equiv_A x$ with $x' \neq x$. Then $xx' \equiv_A xx$, a contradiction. \square

For the rest of this section, X and Y will be orthogonal \emptyset -hyperdefinable sets. We note first that orthogonality is preserved under adding parameters from $X^{\text{heq}} \cup Y^{\text{heq}}$, and interpretation:

Proposition 2.6. *If $X' \subseteq X^{\text{heq}}$ and $Y' \subseteq Y^{\text{heq}}$ are hyperdefinable over some parameters $A \subseteq X^{\text{heq}} \cup Y^{\text{heq}}$, then $X' \perp_A Y'$.*

Proof: Suppose $A = (a, b)$ with $a \in X^{\text{heq}}$ and $b \in Y^{\text{heq}}$, and consider tuples $a' \in X'$ and $b' \in Y'$. Choose representatives $\bar{a}, \bar{a}' \in X$ of a, a' and $\bar{b}, \bar{b}' \in Y$ of b, b' . Then $\text{tp}(\bar{a}\bar{a}') \cup \text{tp}(\bar{b}\bar{b}') \vdash \text{tp}(\bar{a}\bar{a}'\bar{b}\bar{b}')$.

Now if $a'' \equiv_A a'$ and $b'' \equiv_A b'$, we can find A -conjugates $\tilde{a}\tilde{a}''$ of $\bar{a}\bar{a}'$ and $\tilde{b}\tilde{b}''$ of $\bar{b}\bar{b}'$ such that $a''\tilde{a}\tilde{a}'' \equiv_A a'\bar{a}\bar{a}'$ and $b''\tilde{b}\tilde{b}'' \equiv_A b'\bar{b}\bar{b}'$. By orthogonality of X and Y , we obtain $\tilde{a}''\tilde{a}\tilde{b}''\tilde{b} \equiv \tilde{a}'\bar{a}\bar{b}'\bar{b}$, whence $a''ab''b \equiv a'ab'b$, and thus $a''b'' \equiv_A a'b'$. \square

Proposition 2.7. *X is stably embedded in $X \cup Y$: For tuples $a \in X^{\text{heq}}$ and $b \in Y^{\text{heq}}$, every ab -hyperdefinable subset X' of X^{heq} is hyperdefinable over a .*

Proof: If $\Phi(x, a, b)$ hyperdefines X' and $\Psi(y) = \text{tp}(b)$, put

$$\Phi'(x, a) = \exists y [\Psi(y) \wedge \Phi(x, a, y)].$$

Clearly $\Phi(x, a, b) \vdash \Phi'(x, a)$. Conversely, suppose $a' \models \Phi'(x, a)$, and choose $b' \models \Psi$ with $a' \models \Phi(x, a, b')$. By orthogonality $a'ab \equiv a'ab'$, whence $a' \models \Phi(x, a, b)$, and $\Phi'(x, a)$ hyperdefines X' . \square

We put $\text{dcl}_X^{\text{heq}}(A) = \text{dcl}^{\text{heq}}(A) \cap X^{\text{heq}}$ and $\text{bdd}_X(A) = \text{bdd}(A) \cap X^{\text{heq}}$.

Corollary 2.8. *Suppose $a \in X^{\text{heq}}$ and $b \in Y^{\text{heq}}$. Then*

$$\text{dcl}_X^{\text{heq}}(a, b) = \text{dcl}_X^{\text{heq}}(a) \quad \text{and} \quad \text{bdd}_X(a, b) = \text{bdd}_X(a).$$

Proof: Immediate from Proposition 2.7: If X' is a singleton (resp. bounded) subset of X^{heq} hyperdefinable over ab containing some element $e \in \text{dcl}_X^{\text{heq}}(ab)$ (resp. $e \in \text{bdd}_X(ab)$), then X' is hyperdefinable already over a . \square

3. WEAK ELIMINATION OF HYPERIMAGINARIES

In this section, X and Y will be \emptyset -hyperdefinable sets.

Definition 3.1. Let Z be \emptyset -hyperdefinable. We say that Z has *weak elimination of hyperimaginaries with respect to X^{heq} and Y^{heq}* if for every $z \in Z^{heq}$ there is some $x \in \text{bdd}_X(z)$ and $y \in \text{bdd}_Y(z)$ with $z \in \text{dcl}^{heq}(xy)$.

Theorem 3.2. *If X and Y are orthogonal over \emptyset , then $X \cup Y$ has weak elimination of hyperimaginaries with respect to X^{heq} and Y^{heq} .*

Proof: Consider $z \in (X \cup Y)^{heq}$, say $z = (x, y)_E$ for some tuples $x \in X$, $y \in Y$ and countable equivalence relation E over \emptyset . For $x' \equiv x$ consider the hyperdefinable equivalence relation $E_{x'}$ on $\text{tp}(y)$ given by

$$yE_{x'}y' \Leftrightarrow (x', y)E(x', y').$$

Then $E_{x'}$ is \emptyset -hyperdefinable by Proposition 2.7, and does not depend on the choice of $x' \equiv x$. Similarly, for $y' \equiv y$ the equivalence relation

$$xE_{y'}x' \Leftrightarrow (x, y')E(x', y')$$

on $\text{tp}(x)$ does not depend on $y' \equiv y$. Clearly $z \in \text{dcl}^{heq}(x_{E_{y'}}, y_{E_{x'}})$.

We claim that $x_{E_{y'}}$ is bounded over z . If not, there is an indiscernible sequence $(x_i, y_i : i < \omega)$ in $\text{tp}(x, y/z)$ with $\neg x_i E_{y'} x_j$ for $i \neq j$. By orthogonality, for $i < j$,

$$\text{tp}(x_i, x_j) \cup \text{tp}(y_i, y_j) \vdash \text{tp}((x_i, y_i), (x_j, y_j)).$$

But $\text{tp}(x_i, x_j) = \text{tp}(x_i, x_k)$ for $i < k < j$, whence

$$\text{tp}((x_i, y_i), (x_j, y_j)) = \text{tp}((x_i, y_i), (x_k, y_j)).$$

Now $(x_i, y_i)E(x, y)E(x_j, y_j)$ holds since $z = (x, y)_E$. Hence

$$(x_k, y_j)E(x_i, y_i)E(x_j, y_j).$$

Thus $x_k E_{y'} x_j$, a contradiction.

Hence $x_{E_{y'}} \in \text{bdd}_X(z)$; similarly $y_{E_{x'}} \in \text{bdd}_Y(z)$. \square

Corollary 3.3. *If $X \perp Y$, then $\text{bdd}_{XY}(A)$ and $\text{bdd}_X(A) \cup \text{bdd}_Y(A)$ are interdefinable for any A . Moreover, for $aA \subset (X \cup Y)^{heq}$ we have $\text{tp}(a/\text{bdd}_{XY}(A)) \vdash \text{tp}(a/\text{bdd}(A))$.*

Proof: Clearly $\text{bdd}_X(A) \cup \text{bdd}_Y(A) \subseteq \text{bdd}_{XY}(A)$.

For the converse inclusion, let $z \in \text{bdd}_{XY}(A)$. By Theorem 3.2 there is $x \in \text{bdd}_X(z)$ and $y \in \text{bdd}_Y(z)$ with $z \in \text{dcl}^{eq}(xy)$. So

$$z \in \text{dcl}^{heq}(\text{bdd}_X(\text{bdd}(A)), \text{bdd}_Y(\text{bdd}(A))) = \text{dcl}^{heq}(\text{bdd}_X(A), \text{bdd}_Y(A)).$$

For the second assertion, note that equality of E -type over $\text{bdd}(A)$ is a bounded type-definable equivalence relation E_B over any set B of representatives for $\text{bdd}(A)$ with $B_F = \text{bdd}(A)$, given by

$$xE_By \Leftrightarrow \bigwedge_{\varphi \text{ a } B\text{-formula}} [(\varphi(x, B) \rightarrow \exists y'Z [yEy' \wedge BFZ \wedge \varphi(y', Z)]) \wedge (\varphi(y, B) \rightarrow \exists x'Z [xEx' \wedge BFZ \wedge \varphi(x', Z)])].$$

As E_B is invariant under any A -automorphism, it is in fact type-definable over A . By Remark 1.2 the class a_{E_B} is interdefinable with a tuple

$$((aA_i)_{E_i} : i \in I) \in \text{bdd}_{XY}(A),$$

where the $A_i \subseteq A$ are countable. Since the partial type $(xA_i)E_i(aA_i)$ is in $\text{tp}(a/\text{bdd}_{XY}(A))$ for all $i \in I$, we get the result. \square

Corollary 3.4. *If $X \perp Y$ and $X' \subset \text{bdd}(X)$ and $Y' \subset \text{bdd}(Y)$ are \emptyset -hyperdefinable, then $X' \perp_{\text{bdd}_{XY}(\emptyset)} Y'$.*

Proof: It is clearly sufficient to show $X \perp_{\text{bdd}_{XY}(\emptyset)} Y'$. Given $x \in X$ and $y \in Y'$, consider $y_0 \in Y$ with $y \in \text{bdd}(y_0)$, and put $\bar{y} = \text{bdd}_Y(y_0)$. Now if

$$x' \equiv_{\text{bdd}_{XY}(\emptyset)} x \quad \text{and} \quad y' \equiv_{\text{bdd}_{XY}(\emptyset)} y,$$

choose \bar{y}' with $\bar{y}'y' \equiv_{\text{bdd}_{XY}(\emptyset)} \bar{y}y$. As $X \perp_{\text{bdd}_{XY}(\emptyset)} Y$ by Lemma 2.6, we have $x\bar{y} \equiv_{\text{bdd}_{XY}(\emptyset)} x'\bar{y}'$. Since

$$\text{bdd}_{XY}(\bar{y}) \in \text{dcl}^{heq}(\text{bdd}_X(\bar{y}), \text{bdd}_Y(\bar{y})) = \text{dcl}^{heq}(\text{bdd}_X(\emptyset), \bar{y})$$

by Corollary 2.8, we obtain

$$x\text{bdd}_{XY}(\bar{y}) \equiv x'\text{bdd}_{XY}(\bar{y}') \quad \text{and} \quad \text{bdd}_{XY}(\bar{y})y \equiv \text{bdd}_{XY}(\bar{y}')y'.$$

Choose x'' with $x\text{bdd}_{XY}(\bar{y})y \equiv x''\text{bdd}_{XY}(\bar{y}')y'$. Then

$$x''\text{bdd}_{XY}(\bar{y}') \equiv x\text{bdd}_{XY}(\bar{y}) \equiv x'\text{bdd}_{XY}(\bar{y}'),$$

so $\text{tp}(x''/\text{bdd}_{XY}(\bar{y}')) = \text{tp}(x'/\text{bdd}_{XY}(\bar{y}'))$. By Corollary 3.3 we obtain $\text{tp}(x''/\text{bdd}(\bar{y}')) = \text{tp}(x'/\text{bdd}(\bar{y}'))$. As $y \in \text{bdd}(\bar{y})$, we get in particular

$$x'y' \equiv x''y' \equiv xy.$$

The result follows. \square

Example 3.5. We do need $\text{bdd}_{XY}(\emptyset)$ in Corollary 3.4, as we might take $X' = X \times \text{bdd}_Y(\emptyset)$. Then $X' \not\perp Y$ unless $\text{bdd}_Y(\emptyset) = \text{dcl}_Y^{heq}(\emptyset)$.

4. INTERNALITY AND ANALYSABILITY

Definition 4.1. Let X and Y be hyperdefinable sets over A . We say that X is (almost) Y -internal if there is some parameter set B such that for every $a \in X$ there is a tuple $b \in Y$ with $a \in \text{dcl}^{\text{heq}}(Bb)$ (or $a \in \text{bdd}(Bb)$, respectively).³

If the parameters B can be chosen in some set Z , we say that X is (almost) Y -internal *within* Z .

We say that X is Y -analysable (within Z) if for all $a \in X$ there is a sequence $(a_i : i < \alpha)$ such that $\text{tp}(a_i/A, a_j : j < i)$ is Y -internal (within Z) for every $i < \alpha$, and $a \in \text{bdd}(A, a_i : i < \alpha)$.

For the rest of the section, X and Y will be hyperdefinable orthogonal sets over \emptyset .

Proposition 4.2. *If an \emptyset -hyperdefinable set X' is X -analysable within $X \cup Y$, then X' is almost X -internal within $\text{bdd}_Y(\emptyset)$; if X' is X -internal within $X \cup Y$, then X' is X -internal within $\text{bdd}_Y(\emptyset)$.*

Proof: We first show the assertion for (almost) X -internal X' . So suppose $\bar{a} \in X$ and $\bar{b} \in Y$ are such that for every $c_E \in X'$ there is a tuple $a \in X$ with $c_E \in \text{bdd}(\bar{a}\bar{b}a)$. Let $\Phi(x, \bar{a}\bar{b}a)$ be the E -type $\text{tp}(c_E/\bar{a}\bar{b}a)$. Then for every symmetric formula $\psi(x, y) \in E$ there is $n_\psi < \omega$ such that a maximal ψ -antichain in Φ has size n_ψ , and a formula $\phi_\psi(x, \bar{a}\bar{b}a) \in \Phi$ such that every ψ -antichain in ϕ_ψ has size at most n_ψ . Consider the type-definable relation F on $\text{tp}(\bar{a}\bar{b}a)$ given by

$$(\bar{a}'\bar{b}'a')F(\bar{a}''\bar{b}''a'') \Leftrightarrow \bigwedge_{\psi \in E} [\forall x(\phi_\psi(x, \bar{a}'\bar{b}'a') \rightarrow \exists x'[\Phi(x', \bar{a}''\bar{b}''a'') \wedge x\psi^2x']) \\ \wedge \forall x'(\phi_\psi(x', \bar{a}''\bar{b}''a'') \rightarrow \exists x[\Phi(x, \bar{a}'\bar{b}'a') \wedge x\psi^2x'])],$$

where $x\psi^2x'$ means $\exists x''[\psi(x, x'') \wedge \psi(x'', x')]$. Then $(\bar{a}'\bar{b}'a')F(\bar{a}''\bar{b}''a'')$ holds if and only if $\Phi(x, \bar{a}'\bar{b}'a')$ and $\Phi(x, \bar{a}''\bar{b}''a'')$ contain the same points modulo E : If they contain the same points modulo E , for every ψ in E let $(x_i : i < n_\psi)$ be a ψ -antichain in $\Phi(x, \bar{a}'\bar{b}'a')$, and choose $(x'_i : i < n_\psi)$ in $\Phi(x, \bar{a}''\bar{b}''a'')$ with $x_i E x'_i$ for all $i < n_\psi$. Then whenever x satisfies $\phi_\psi(x, \bar{a}'\bar{b}'a')$ there is $i < n_\psi$ with $\psi(x, x_i)$, whence $x\psi^2x'_i$. By symmetry the converse also holds, so $(\bar{a}'\bar{b}'a')F(\bar{a}''\bar{b}''a'')$. On the other hand, if there is x such that $\Phi(x, \bar{a}'\bar{b}'a')$ but $\neg x E x'$ for all x' with

³In simplicity theory, this is called *finite generation*; for *internality* we would require for every $a \in X$ the existence of some $B \perp_A a$ and tuple $b \in Y$ with $a \in \text{dcl}^{\text{heq}}(Bb)$.

$\Phi(x', \bar{a}''\bar{b}''a'')$, by compactness there is $\psi' \in E$ such that $\neg\psi'(x, x')$ for all x' with $\Phi(x', \bar{a}''\bar{b}''a'')$. Then any $\psi \in E$ with $\psi^2 \vdash \psi'$ witnesses $\neg(\bar{a}'\bar{b}'a')F(\bar{a}''\bar{b}''a'')$.

It follows that F is an equivalence relation, and any automorphism fixes $(\bar{a}\bar{b}a)_F$ if and only if it permutes the set $C \subset X'$ of E -classes in $\Phi(x, \bar{a}\bar{b}a)$. In particular $(\bar{a}\bar{b}a)_F \in \text{dcl}^{heq}(C)$ and $C \subseteq \text{bdd}((\bar{a}\bar{b}a)_F)$, whence in particular $c_E \in \text{bdd}((\bar{a}\bar{b}a)_F)$. Moreover, if X' is X -internal, then $C = \{c_E\}$ and $c_E \in \text{dcl}^{heq}(\bar{a}\bar{b}a)_F$.

By weak elimination of hyperimaginaries, there is $\tilde{a} \in \text{bdd}_X((\bar{a}\bar{b}a)_F)$ and $\tilde{b} \in \text{bdd}_Y((\bar{a}\bar{b}a)_F)$ with $(\bar{a}\bar{b}a)_F \in \text{dcl}^{heq}(\tilde{a}\tilde{b})$. Thus we are done if we can show $\tilde{b} \in \text{bdd}_Y(\emptyset)$.

Suppose $\tilde{b} \notin \text{bdd}_Y(\emptyset)$. Then there is an \emptyset -conjugate \tilde{b}' of \tilde{b} outside $\text{bdd}(\bar{a}\bar{b})$; if σ is an automorphism mapping \tilde{b}' to \tilde{b} , put $\bar{a}'\bar{b}' = \sigma(\bar{a}\bar{b})$. Then $\tilde{b} \notin \text{bdd}(\bar{a}'\bar{b}')$. On the other hand, since $\bar{a}'\bar{b}' \equiv \bar{a}\bar{b}$, for every $e \in C$ there is $a_e \in X$ with $e \in \text{bdd}(\bar{a}'\bar{b}'a_e)$. Therefore

$$\tilde{b} \in \text{bdd}_Y((\bar{a}\bar{b}a)_F) \subseteq \text{bdd}_Y(C) \subseteq \text{bdd}_Y(\bar{a}', \bar{b}', a_e : e \in C),$$

whence $\tilde{b} \in \text{bdd}_Y(\bar{b}')$ by Corollary 2.8, a contradiction.

Now assume that $x \in X'$ and $(x_i : i < \alpha)$ is an X -analysis of x within $X \cup Y$. We show inductively on $i \leq \alpha$ that $\text{tp}(x_j : j < i)$ is X -internal within $\text{bdd}_Y(\emptyset)$. So suppose $\text{tp}(x_j : j < k)$ is X -internal within $\text{bdd}_Y(\emptyset)$ for all $k < i$. If i is limit, then clearly $\text{tp}(x_j : j < i)$ is X -internal within $\text{bdd}_Y(\emptyset)$. If $i = k + 1$, then by the result for internality $\text{tp}(x_k/x_j : j < k)$ is X -internal within $\text{bdd}_Y(x_j : j < k)$ and there is $a \in X$ with

$$x_k \in \text{dcl}^{heq}(a, \text{bdd}_Y(x_j : j < k), x_j : j < k).$$

Or, by X -internality of $\text{tp}(x_j : j < k)$ within $\text{bdd}_Y(\emptyset)$ there is $a' \in X$ with $(x_j : j < k) \in \text{dcl}^{heq}(a', \text{bdd}_Y(\emptyset))$. Then by Corollary 2.8

$$\text{bdd}_Y(x_j : j < k) \subseteq \text{bdd}_Y(a', \text{bdd}_Y(\emptyset)) = \text{bdd}_Y(\emptyset),$$

and $x_k \in \text{bdd}(a, a', \text{bdd}_Y(\emptyset))$. So $\text{tp}(x_j : j < i)$ is X -internal within $\text{bdd}_Y(\emptyset)$, and $\text{tp}(x)$ is almost X -internal within $\text{bdd}_Y(\emptyset)$. \square

Corollary 4.3. *Let X' and Y' be \emptyset -hyperdefinable. If X' is almost X -internal within $X \cup Y$ and Y' is almost Y -internal within $X \cup Y$, then $X' \perp_{\text{bdd}_{XY}(\emptyset)} Y'$.*

Proof: Proposition 2.6 and Corollary 3.3 yield $X \perp_{\text{bdd}_{XY}(\emptyset)} Y$. By Proposition 4.2 we have

$$X' \subset \text{bdd}(X, \text{bdd}_{XY}(\emptyset)) \quad \text{and} \quad Y' \subset \text{bdd}(Y, \text{bdd}_{XY}(\emptyset)).$$

Hence $X' \perp_{\text{bdd}_{XY}(\emptyset)} Y'$ by Corollary 3.4. \square

Corollary 4.4. *If an \emptyset -hyperdefinable set Z is almost X - and almost Y -internal within $X \cup Y$, then it is bounded.*

Proof: We have $Z \perp_{\text{bdd}_{XY}(\emptyset)} Z$ by Corollary 4.3, so Z is bounded by Remark 2.5. \square

Corollary 4.5. *If $Z \subseteq (X \cup Y)^{\text{heq}}$ is \emptyset -hyperdefinable and almost X -internal within $X \cup Y$, then it is X -internal within $\text{bdd}_Y(\emptyset)$.*

Proof: Let $z \in Z$. By weak elimination of hyperimaginaries there is $x \in \text{bdd}_X(z)$ and $y \in \text{bdd}_Y(z)$ with $z \in \text{dcl}^{\text{heq}}(xy)$. Then $\text{tp}(y)$ is Y -internal since $y \in Y^{\text{heq}}$, but also almost X -internal, as $y \in \text{bdd}(z)$ and $\text{tp}(z)$ is almost X -internal. So $y \in \text{bdd}_Y(\emptyset)$ by Corollary 4.4. \square

Again Example 3.5 shows that we need $\text{bdd}_Y(\emptyset)$ in Corollaries 4.3 and 4.5.

5. LOCAL SIMPLICITY

Definition 5.1. Let $A \subseteq B$, and $\pi(x, B)$ be a partial type over B . We say that $\pi(x, B)$ does not divide over A if for any indiscernible sequence $(B_i : i < \omega)$ in $\text{tp}(B/A)$ the partial type

$$\bigcup_{i < \omega} \pi(x, B_i)$$

is consistent. Clearly, $\text{tp}(a/B)$ divides over A if and only if $\text{tp}(a_0/B)$ does so for some finite subtuple $a_0 \subseteq a$.

Example 5.2. If $\text{tp}(a) \perp \text{tp}(b)$, then $\text{tp}(a/b)$ does not divide over \emptyset .

We now define the appropriate version of local rank. We follow Ben Yaacov's terminology [1, Definition 1.4], more general than [5, Definition 4.3.5].

Definition 5.3. Let $\pi(x)$, $\Phi(x, y)$ and $\Psi(y_1, \dots, y_k)$ be partial types in (at most) countably many variables.

- (1) Ψ is a k -inconsistency witness for Φ if

$$\models \forall y_1 \dots y_k \neg \exists x [\Psi(y_1, \dots, y_k) \wedge \bigwedge_{i=1}^k \Phi(x, y_i)].$$

- (2) Let Ψ be a k -inconsistency witness for Φ . The *local (Φ, Ψ) -rank* $D(., \Phi, \Psi)$ is defined on partial types in x as follows:

- $D(\pi(x), \Phi, \Psi) \geq 0$ if $\pi(x)$ is consistent.
- $D(\pi(x), \Phi, \Psi) \geq n + 1$ if there is a sequence $(a_i : i < \omega)$ such that $\models \Psi(\bar{a})$ for any k -tuple $\bar{a} \subset (a_i : i < \omega)$, and $D(\pi(x) \wedge \Phi(x, a_i), \Phi, \Psi) \geq n$ for all $i < \omega$.

If $D(\pi, \Phi, \Psi) \geq n$ for all $n < \omega$, we put $D(\pi, \Phi, \Psi) = \infty$.

Remark 5.4. Note that $D(\pi(x, a), \Phi, \Psi) \geq n$ is a closed condition on a , and $D(\text{tp}(x/a), \Phi, \Psi) \geq n$ is a closed condition on x over a . By compactness and Ramsey's theorem, we may require $(a_i : i < \omega)$ to be indiscernible in Definition 5.3 (2).

Lemma 5.5. *Let Ψ be an inconsistency witness for Φ , and π a partial type over A . Then $D(\pi, \Phi, \Psi)$ is infinite if and only if for every linear order I there are elements $(b_i, a_i^j : i \in I, j < \omega)$ such that $\models \Psi(\bar{a})$ for all $\bar{a} \subset (a_i^j : j < \omega)$ of the right length, $b_i \models \pi \wedge \bigwedge_{k \leq i} \Phi(x, a_k^0)$, and $(a_i^j : j < \omega)$ is indiscernible over $A \cup \{b_k a_k^0 : k < i\}$, for all $i \in I$. Moreover, we may require $(b_i a_i^0 : i \in I)$ to be indiscernible.*

Proof: If the condition is satisfied, we can take $I = \omega$. Then for all $n \in \omega$ the partial type $\pi \wedge \bigwedge_{i \leq n} \Phi(x, a_i^0)$ is satisfied by b_n and hence non-empty. So

$$\begin{aligned} D(\pi, \Phi, \Psi) &> D(\pi \wedge \Phi(x, a_0^0), \Phi, \Psi) > D(\pi \wedge \Phi(x, a_0^0) \wedge \Phi(x, a_1^0), \Phi, \Psi) \\ &> \cdots > D(\pi \wedge \Phi(x, a_0^0) \wedge \cdots \wedge \Phi(x, a_n^0), \Phi, \Psi) \geq 0. \end{aligned}$$

Hence $D(\pi, \Phi, \Psi) > n$ for all $n < \omega$, and $D(\pi, \Phi, \Psi) = \infty$.

For the converse, by compactness it is sufficient to consider finite I . We show by induction that if $D(\pi, \Phi, \Psi) \geq n$, then the condition is satisfied for I of size n . For $n = 0$ there is nothing to show. Suppose $D(\pi, \Phi, \Psi) \geq n + 1$. Then by definition there is a sequence $(a_0^j : j < \omega)$ whose subsequences satisfy Ψ , and such that $D(\pi \wedge \Phi(x, a_0^0), \Phi, \Psi) \geq n$ for all $j < \omega$. By Remark 5.4 we may assume that $(a_0^j : j < \omega)$ is indiscernible over A . Choose $b_0 \models \pi \wedge \Phi(x, a_0^0)$. By inductive hypothesis for the partial type $\pi \wedge \Phi(x, a_0^0)$ over $A \cup \{b_0, a_0^0\}$, there are $(b_i, a_i^j : 1 \leq i \leq n, j < \omega)$ such that $\models \Psi(\bar{a})$ for all $\bar{a} \subset (a_i^j : j < \omega)$ of the right length,

$$b_i \models \pi \wedge \Phi(x, a_0^0) \wedge \bigwedge_{1 \leq k \leq i} \Phi(x, a_k^0),$$

and $(a_i^j : j < \omega)$ is indiscernible over $A \cup \{b_0, a_0^0\} \cup \{b_k, a_k^0 : 1 \leq k < i\}$, for all $1 \leq i \leq n$, as required.

The final assertion follows by compactness and Ramsey's theorem. \square

Definition 5.6. Let I be an ordered set. A sequence $I = (a_i : i \in I)$ is *independent over A* , or *A -independent*, if $\text{tp}(a_i/A, a_j : j < i)$ does not divide over A for all $i \in I$. If $A \subseteq B$ and $p \in S(B)$, the sequence $(a_i : i \in I)$ is a *Morley sequence in p over A* if it is B -indiscernible, $a_i \models p$ and $\text{tp}(a_i/B, a_j : j < i)$ does not divide over A for all $i \in I$. If $A = B$, we simply call it a *Morley sequence in p* .

Fact 5.7. [5, Corollary 3.2.5] or [4, Proposition 16.12] *If $\text{tp}(b/cd)$ does not divide over d and $\text{tp}(a/cbd)$ does not divide over bd , then $\text{tp}(ab/cd)$ does not divide over d .*

For the rest of the section we fix a hyperdefinable set X over \emptyset . We call a type $p(x)$ an *X -type* if it implies $x \in X$.

The following theorem generalizes [5, Theorem 2.4.7] to the local hyperdefinable context. Note that in the classical development the forking properties for hyperimaginaries are deduced from the corresponding properties for representatives. Here we cannot do this, as the ambient theory may well not be simple. So we have to work with hyperimaginaries in X throughout.

Theorem 5.8. *The following are equivalent:*

- (1) *Symmetry holds on X : For all $a, b, c \in X$, $\text{tp}(a/bc)$ does not divide over b if and only if $\text{tp}(c/ab)$ does not divide over b .*
- (2) *Transitivity holds on X : If $a, b, c, d \in X$, then $\text{tp}(a/bcd)$ does not divide over b if and only if $\text{tp}(a/bc)$ does not divide over b and $\text{tp}(a/bcd)$ does not divide over bc .*
- (3) *Local character holds on X : There is κ such that for all countable $a \in X$ and $A \subset X$ there is $A_0 \subseteq A$ with $|A_0| \leq \kappa$ such that $\text{tp}(a/A)$ does not divide over A_0 . In fact, we can take $\kappa = 2^{|T|}$.*
- (4) *$D(\cdot, \Phi, \Psi) < \infty$ for any partial X -type $\Phi(x, y)$ and inconsistency witness Ψ for Φ .*
- (5) *For any $A \subseteq B \subset X$, a partial X -type $\pi(x, B)$ does not divide over A if and only if there is a Morley sequence I in $\text{tp}(B/A)$ such that $\{\pi(x, B') : B' \in I\}$ is consistent.*

If any of these conditions is satisfied, then for all $A \subseteq B \subset X$ and $a \in X$ the type $\text{tp}(a/B)$ does not divide over A if and only if

$$D(\text{tp}(a/B), \Phi, \Psi) = D(\text{tp}(a/A), \Phi, \Psi)$$

for all (Φ, Ψ) . Moreover, Extension holds on X : For any partial X -type $\pi(x)$ over B , if π does not divide over A then it has a completion which does not divide over A .

Proof: (1) \Rightarrow (2) Clearly, if $\text{tp}(a/bcd)$ does not divide over b , it does not divide over bc and $\text{tp}(a/bc)$ does not divide over b . Conversely, suppose that $\text{tp}(a/bcd)$ does not divide over bc and $\text{tp}(a/bc)$ does not divide over b . By symmetry, $\text{tp}(d/abc)$ does not divide over bc and $\text{tp}(c/ab)$ does not divide over b . By Fact 5.7 $\text{tp}(cd/ab)$ does not divide over b , so again by symmetry $\text{tp}(a/bcd)$ does not divide over b .

(2) \Rightarrow (4) Suppose there is a partial X -type Φ and an inconsistency witness Ψ for Φ such that $D(x = x, \Phi, \Psi) = \infty$. Put $I = \{\pm 1, \pm(1 + \frac{1}{n}) : n > 0\}$ and choose a sequence $(b_i, a_i^j : i \in I, j < \omega)$ as given by Lemma 5.5. Let $A^- = \{b_i a_i^0 : i < -1\}$ and $A^+ = \{b_i a_i^0 : i > 1\}$. Then $\text{tp}(b_1/A^-A^+)$ does not divide over A^- and $\text{tp}(b_1/A^-A^+a_{-1})$ does not divide over A^-A^+ , since the former is finitely satisfiable in A^- and the latter in A^+ . However, $(a_{-1}^j : j < \omega)$ witnesses that $\Phi(x, a_{-1})$, and hence $\text{tp}(b_1/A^-A^+a_{-1})$, divides over A^- , contradicting transitivity.

(4) \Rightarrow (3) Assume (4). First, we note that for $A \subseteq B \subset X$, if

$$D(\text{tp}(a/B), \Phi, \Psi) = D(\text{tp}(a/A), \Phi, \Psi)$$

for all (Φ, Ψ) , then $\text{tp}(a/B)$ does not divide over A . This is obvious, as if some A -indiscernible sequence $(B_i : i < \omega)$ in $\text{tp}(B/A)$ witnesses dividing, we can take $\Phi(x, y) = \text{tp}(a, B)$ and $\Psi = \text{tp}(B_1, B_2, \dots, B_n)$ for $n < \omega$ sufficiently large. Then Ψ is an n -inconsistency witness (clearly, we may restrict to countable B), and

$$D(\text{tp}(a/B), \Phi, \Psi) < D(\text{tp}(a/A), \Phi, \Psi).$$

Given $\text{tp}(a/A)$ it is hence enough to take $A_0 \subseteq A$ big enough such that

$$D(\text{tp}(a/A), \Phi, \Psi) = D(\text{tp}(a/A_0), \Phi, \Psi)$$

for all (Φ, Ψ) . There are only $2^{|T|}$ such pairs, so we need at most that many parameters.

(3) \Rightarrow (4) Suppose $D(x = x, \Phi, \Psi) = \infty$. Then for any cardinal κ we can find an indiscernible sequence $(b_i, a_i^j : i \leq \kappa^+, j < \omega)$ as in Lemma 5.5. Since $\Phi(x, a_i^0)$ divides over $\{a_j^0 : j < i\}$ for all $i \leq \kappa^+$, the type $\text{tp}(b_{\kappa^+}/a_i^0 : i < \kappa^+)$ divides over any subset of its domain of cardinality $\leq \kappa$.

(4) \Rightarrow (5). Assume (4). Given $a_E \in X$ and $A \subseteq B = b_E \subset X$, for any pair (Φ, Ψ) and any formula $\varphi(y, b)$ we can adjoin either

$$\exists yz [xEy \wedge zEb \wedge \varphi(y, z)] \quad \text{or} \quad \exists yz [xEy \wedge zEb \wedge \neg \varphi(y, z)]$$

and preserve $D(\cdot, \Phi, \Psi)$ -rank. By compactness we can thus complete $\text{tp}(a_E/A)$ to an E -type p over B of the same $D(\cdot, \Phi, \Psi)$ -rank. In particular, no Φ -instance in p divides over A with Ψ as inconsistency witness.

Coding finitely many pairs $(\Phi_i, \Psi_i : i < n)$ in a single one, one obtains an extension p such that no Φ_i -instance Ψ_i -divides for any $i < n$; by compactness we can do this for all pairs (Φ, Ψ) simultaneously and obtain an extension which does not divide over A . Take $B = X^{\mathfrak{M}} \supset A$ for some sufficiently saturated model \mathfrak{M} . Then a sequence $(a_i : i < \omega) \subset B$ such that $a_i \models p \upharpoonright_{(A, a_j : j < i)}$ is a Morley sequence in $\text{tp}(a/A)$.

This shows in particular that if $\pi(x, B)$ does not divide over A , then there is a Morley sequence I in $\text{tp}(B/A)$ such that $\{\pi(x, B') : B' \in I\}$ is consistent.

Conversely, suppose that $\pi(x, B)$ divides over A , as witnessed by an A -indiscernible sequence $(B_i : i < \omega)$ in $\text{tp}(B/A)$ with $\bigcup_{i < \omega} \pi(x, B_i)$ inconsistent. Take any Morley sequence I in $\text{tp}(B/A)$. By [5, Corollary 2.2.8] (which is shown there for real tuples, but transfers easily to hyperimaginaries) we may assume that $B_i \hat{\ } I$ is A -conjugate to I for all $i \in I$ and that $(B_i : i < \omega)$ is indiscernible over AI . If $\bar{\pi}(x) = \bigcup_{B' \in I} \pi(x, B')$ were consistent, then $(B_i : i < \omega)$ would witness that

$$D(\bar{\pi}(x) \wedge \pi(x, B_0), \pi(x, y), \Psi) < D(\bar{\pi}(x), \pi(x, y), \Psi)$$

for some inconsistency witness Ψ . But by A -conjugacy the two ranks must be equal, a contradiction.

(5) \Rightarrow (1) Let us first show *Extension*. If $A \subseteq B \subset X$ and $\pi(x, B)$ is a partial X -type which does not divide over A , let $(B_i : i < \alpha)$ be a very long Morley sequence in $\text{tp}(B/A)$. Consider any realization $a \models \bigwedge_{i < \alpha} \pi(x, B_i)$. Since α is large, there is an infinite subset $J \subset \alpha$ such that $p(aAX) = \text{tp}(aAB_i)$ is constant for $i \in J$. Then $p(xAB)$ is a completion of π which does not divide over A , as witnessed by $(B_i : i \in J)$.

Now given $a, b, c \in X$ such that $\text{tp}(a/bc)$ does not divide over b , let $B = X^{\mathfrak{M}} \ni bc$ for some sufficiently saturated model \mathfrak{M} , and p an extension of $\text{tp}(a/bc)$ to B which does not divide over b . Choose a sequence $(a_i : i < \omega) \subset B$ such that $a_i \models p \upharpoonright_{(bc, a_j : j < i)}$. This is a Morley sequence in $\text{tp}(a/bc)$ over b . Then $(a_i : i < \omega)$ is a Morley sequence in $\text{tp}(a/b)$, and $a_i \models \text{tp}(a/bc)$ for all $i < \omega$. Hence $\text{tp}(c/ba)$ does not divide over b , and symmetry holds.

Finally we show that if (1) – (5) hold and $\text{tp}(a/B)$ does not divide over A for $A \subseteq B \subset X$ and $a \in X$, then $D(\text{tp}(a/A), \Phi, \Psi) \geq n$ implies $D(\text{tp}(a/B), \Phi, \Psi) \geq n$ for all (Φ, Ψ) . For $n = 0$ this is obvious. So suppose $D(\text{tp}(a/A), \Phi, \Psi) \geq n + 1$. Then there is $(d_i : i < \omega)$ indiscernible over A such that $\bar{d} \models \Psi$ for all $\bar{d} \subset (d_i : i < \omega)$ of the right length, and $D(\text{tp}(a/A) \wedge \Phi(x, d_i), \Phi, \Psi) \geq n$ for all $i < \omega$. Let q be a completion

of $\text{tp}(a/A) \wedge \Phi(x, d_0)$ with $D(q, \Phi, \Psi) \geq n$. Clearly, we may assume $a \models q$, and that $\text{tp}(d_0/aB)$ does not divide over aA . As $\text{tp}(a/B)$ does not divide over A , by symmetry and transitivity $\text{tp}(ad_0/B)$ does not divide over A , and $\text{tp}(a/d_0B)$ does not divide over d_0A . By induction hypothesis,

$$D(\text{tp}(a/d_0A), \Phi, \Psi) \geq n \quad \text{implies} \quad D(\text{tp}(a/d_0B), \Phi, \Psi) \geq n.$$

As $\text{tp}(d_0/B)$ does not divide over A and $(d_i : i < \omega)$ is A -indiscernible, we may assume that it remains indiscernible over B . But then it witnesses

$$D(\text{tp}(a/B), \Phi, \Psi) \geq D(\text{tp}(a/d_0B), \Phi, \Psi) + 1 \geq n + 1. \quad \square$$

Definition 5.9. An A -hyperdefinable set X is *simple (over A)* if it satisfies any of the conditions of Theorem 5.8 when we adjoin A to the language. If X is simple over A and $a, b, c \in X$, we shall say that a and c are independent over Ab , written $a \perp_{Ab} c$, if $\text{tp}(a/Abc)$ does not divide over Ab .

Note that we only allow tuples and parameters from $A \cup X$. If X is stably embedded, we can of course allow parameters from anywhere.

Remark 5.10. If X is merely hyperdefinable, it may be simple although no definable or even type-definable imaginary set in the ambient structure is simple.

If X is simple, it is now standard to extend the notions of dividing and independence to hyperimaginaries in X^{heq} . Moreover, we can develop basic simplicity theory (canonical bases, the independence theorem, stratified ranks, generic types, stabilizers, see [4, 5]) within X^{heq} , replacing models \mathfrak{M} by subsets $X^{\text{heq}} \cap \mathfrak{M}^{\text{heq}}$.

Proposition 5.11. *Let X and Y be orthogonal \emptyset -hyperdefinable sets such that X is simple over \emptyset . If $A \subset Y$ is a set of parameters, then X is simple over A , and over $\text{bdd}_Y(A)$. In particular, let Z be a set hyperdefinable over some parameters $A \subset X \cup Y$. If Z is X -internal within $X \cup Y$, then Z is simple over A ; if $Z \subseteq (X \cup Y)^{\text{heq}}$ is almost X -internal, then Z is simple as well.*

Proof: Simplicity over A is obvious from orthogonality; simplicity over $\text{bdd}_Y(A)$ follows. Now if Z is X -internal within $X \cup Y$, then $Z \subset (X \cup \text{bdd}_Y(A))^{\text{heq}}$ by Corollary 4.2, and must be simple as well; if $Z \subseteq (X \cup Y)^{\text{heq}}$ is almost X -internal within $X \cup Y$, it is X -internal within $\text{bdd}_Y(A)$ by Corollary 4.5. \square

6. GROUPS INTERPRETABLE IN ORTHOGONAL SETS

Recall that two hyperdefinable subgroups H_1 and H_2 of some group G are *commensurable* if $H_1 \cap H_2$ has bounded index both in H_1 and in H_2 .

Theorem 6.1. *Suppose X and Y are orthogonal \emptyset -hyperdefinable sets in a structure \mathfrak{M} , and G is an \emptyset -hyperdefinable group in $(X \cup Y)^{heq}$. If X is simple over \emptyset , there is an \emptyset -hyperdefinable normal X -internal subgroup N of G such that the quotient G/N is Y -internal. N is unique up to commensurability.*

Proof: Let us first show uniqueness: If N' is a second such group, then $N/(N \cap N')$ and $N'/(N \cap N')$ are X -internal and Y -internal, and hence bounded by orthogonality of X and Y . Thus N and N' are commensurable.

By Theorem 3.2 every element $g \in G$ is of the form $(g_X, g_Y)_E$ for some $g_X \in X^{heq}$ and $g_Y \in Y^{heq}$, both bounded over g , and some type-definable equivalence relation E with bounded classes, depending on $\text{tp}(g)$. Hence $\text{tp}(g/g_Y)$ is X -internal and $\text{tp}(g/g_X)$ is Y -internal. Now if $h = (h_X, h_Y)_E$ and $gh = ((gh)_X, (gh)_Y)_E$, then

$$(gh)_X \in \text{bdd}_X(g_X, g_Y, h_X, h_Y),$$

whence $(gh)_X \in \text{bdd}_X(g_X, h_X)$ by Corollary 2.8. Similarly $(gh)_Y \in \text{bdd}_Y(g_Y, h_Y)$.

Now X is simple, as is $\text{tp}(g, h/g_Y, h_Y)$ for any $g, h \in G$ by Proposition 5.11. Hence we can consider $g, h \in G$ such that $g \perp_{g_Y, h_Y} h$. Then for any stratified local rank D

$$\begin{aligned} D(gh/(gh)_Y) &\geq D(gh/(gh)_Y, g_Y, h_Y, g) = D(h/g_Y, h_Y, g) \\ (\dagger) \quad &= D(h/g_Y, h_Y) = D(h/h_Y), \end{aligned}$$

where the first equality holds as $(gh)_Y \in \text{bdd}_Y(g_Y, h_Y)$, and the last equality follows from $h \perp_{h_Y} g_Y$ by orthogonality (Example 5.2). Similarly

$$D(gh/(gh)_Y) \geq D(g/g_Y).$$

Now suppose G is a subset of $(X^m \times Y^n)/E$, where m, n are at most countable. Then if $g = (\bar{x}_g, \bar{y}_g)_E \in G$, we have $g \perp_{g_Y} \bar{y}_g$ by X -internality of $\text{tp}(g/g_Y)$ and orthogonality, and $g_Y \in \text{bdd}(\bar{y}_g)$, whence $D(g/g_Y) = D(g/\bar{y}_g)$. By compactness, there is a G -type $p((\bar{x}, \bar{y})_E)$ implying that $D((\bar{x}, \bar{y})_E/\bar{y})$ is maximal for all local stratified ranks. But if $g, h \models p$ with $g \perp_{g_Y, h_Y} h$, then we must have equality in (\dagger) , which

implies that g , h and gh are pairwise independent over $g_Y, h_Y, (gh)_Y$. Put

$$S = \text{stab}(g/g_Y, h_Y, (gh)_Y).$$

By [3, Lemme 1.2 and Remarque 1.3] the element g is generic in the coset Sg , and this coset is hyperdefinable over $\text{bdd}(g_Y, h_Y, (gh)_Y)$. By orthogonality, $g \perp_{g_Y} h_Y, (gh)_Y$. This implies in particular

$$D(S) = D(Sg) = D(g/g_Y, h_Y, (gh)_Y) = D(g/g_Y) = D(p).$$

Moreover, S is X -internal since $\text{tp}(g/g_Y)$ is.

Suppose that S is not commensurable with S^h for some $h \in G$. Then SS^h is still X -internal, with

$$D(SS^h) \geq D(S) = D(p)$$

for every stratified local rank D , and for at least one such rank D_0 we have $D_0(SS^h) > D_0(p)$. Choose $g' \in SS^h$ with $D_0(g'/h) = D_0(SS^h)$. By pre-multiplying with a generic element of S and post-multiplying with a generic element of S^h , the inequality (\dagger) implies that we may assume $D(g'/h) \geq D(p)$ for every stratified local rank D . However, $\text{tp}(g'/h)$ is X -internal, so $g'_Y \in \text{bdd}_Y(g')$ implies $g'_Y \in \text{bdd}_Y(h)$ by Lemma 2.8 and Proposition 4.2. Thus

$$D(g'/g'_Y) \geq D(g'/h) \geq D(p) \quad \text{and} \quad D_0(g'/g'_Y) \geq D_0(g'/h) > D_0(p),$$

contradicting our choice of p . Hence S is commensurable with all its conjugates, and the locally connected component N of S is normal in G [5, Definition 4.5.15 and Corollary 4.5.16]. In fact the same proof, with NZ instead of SS^h , shows that if Z is an X -internal hyperdefinable subset of G , then Z is covered by boundedly many cosets of N . In particular, for any $g' \in G$ the type $\text{tp}(g'/g'_Y)$ is covered by boundedly many cosets of N . But then $g'N \in \text{bdd}(g'_Y)$, and G/N is almost Y -internal, whence Y -internal by Corollary 4.5. \square

Corollary 6.2. *Suppose X and Y are orthogonal type-definable sets over \emptyset in a structure \mathfrak{M} , and G is a type-interpretable group over \emptyset in $(X \cup Y)^{\text{eq}}$. If X is simple over \emptyset , there is a normal X -internal subgroup N of G type-interpretable over \emptyset , such that the quotient G/N is Y -internal. N is unique up to commensurability. If X is definable and supersimple, then we can take N relatively interpretable.*

Proof: The first part is obvious from Theorem 6.1, as a hyperdefinable subgroup of a type-interpretable group is again type-interpretable.

If X is definable and supersimple, N must be contained in a definable X -internal set \bar{X} by [5, Lemma 3.4.17]; note that \bar{X} will also

be supersimple. So N is the intersection of definable supergroups by [5, Theorem 5.5.4], one of which, say N_0 , must be contained in \bar{X} by compactness. Then N_0 is X -internal. As above, N_0 must be commensurable with all its G -conjugates; moreover, commensurability is uniform by compactness (or [5, Lemma 4.2.6]). By [5, Theorem 4.2.4] there is a relatively interpretable normal subgroup \bar{N} commensurable with N_0 . So \bar{N} is X -internal, and G/\bar{N} is Y -internal. \square

Question 6.3. If X and Y are orthogonal type-definable sets (or even definable sets), X is simple and G is a relatively definable group in $(X \cup Y)^{eq}$, can we find a relatively definable normal X -internal subgroup N such that G/N is Y -internal?

Question 6.4. What can we say if neither X nor Y is simple? Is it true that in every hyperdefinable subgroup of $((X \cup Y)^{heq})$ there is a maximal normal hyperdefinable X -internal subgroup N_X , a maximal normal hyperdefinable Y -internal subgroup N_Y , an X -internal hyperdefinable local group G_X , a Y -internal hyperdefinable local group G_Y and a hyperdefinable locally bounded equivalence relation E on $G_X \times G_Y$ such that $G/(N_X N_Y)$ is isogenous, or even isomorphic, to $(G_X \times G_Y)/E$?

REFERENCES

- [1] Itai Ben Yaacov. *Simplicity in compact abstract theories*. J. Math. Logic 3(2):163–191, 2003.
- [2] Alessandro Berarducci and Marcello Mamino. *Groups definable in two orthogonal sorts*. Preprint. arXiv:1304.1380.
- [3] Thomas Blossier, Amador Martín Pizarro and Frank O. Wagner. *A la recherche du tore perdu*. Submitted.
<http://hal.archives-ouvertes.fr/hal-00758982>
- [4] Enrique Casanovas. *Simple Theories and Hyperimaginaries*. Lecture Notes in Logic 39. Cambridge University Press, 2011.
- [5] Frank O. Wagner. *Simple Theories*. Mathematics and Its Applications 503. Kluwer Academic Publishers, 2000.

UNIVERSITÉ DE LYON; CNRS; UNIVERSITÉ LYON 1; INSTITUT CAMILLE JORDAN UMR5208, 43 BD DU 11 NOVEMBRE 1918, 69622 VILLEURBANNE CEDEX, FRANCE

E-mail address: `wagner@math.univ-lyon1.fr`