



HAL
open science

A new look at the Julg-Valette homotopy for groups acting on trees.

Pierre Julg

► **To cite this version:**

Pierre Julg. A new look at the Julg-Valette homotopy for groups acting on trees.. 2013. hal-00876928v1

HAL Id: hal-00876928

<https://hal.science/hal-00876928v1>

Preprint submitted on 25 Oct 2013 (v1), last revised 11 Feb 2014 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A new look at the Julg-Valette homotopy for groups acting on trees.

Pierre Julg

MAPMO (Université d'Orléans et CNRS)

In [JV1] [JV2] we proved the K-theoretic amenability of locally compact groups acting on trees (with amenable stabilizers) by combining two ingredients:

1) We construct a very simple Fredholm module associated to the group action on a tree. It defines a class γ in Kasparov's ring $KK_G(\mathbf{C}, \mathbf{C})$.

2) We show that $\gamma = 1$ by constructing a homotopy using the simple but non trivial fact that the distance kernel on the set of vertices of the tree is of conditionnally negative type. Or equivalently the existence of an affine action of G on the ℓ^2 space of edges of the tree.

Note that the two ingredients are of different nature. The first can be generalized to other situations such as Bruhat-Tits buildings [JV3][KS1], or (hyper)bolic spaces [KS2]. The second is very specific to trees, or generalizations (e.g. CAT(0) cubic complexes cf [B]).

We present here a perhaps more natural proof of the same result. It is inspired by the construction by Pytlik and Szwarc [PS] of a family of uniformly bounded representations of a free group, generalized by Valette [V1] to groups acting on trees. We have had this new version for quite a long time in our private notes. We thank Amaury Freslon and Jacek Brodzki for convincing us that making these notes available could be useful to others.

1. Notations:

Let $X = (X^0, X^1)$ be a tree. There is no orientation on X so that the set X^1 of edges is just a subset of $X^0 \times X^0$ stable by the map $(x, y) \mapsto (y, x)$. By hypothesis, for any x and y in X^0 there is a unique path joining x to y . We denote by $[x, y]$ the set of vertices lying between x and y and $d(x, y)$ the number of edges between x and y . Let G be a group acting on X . In other words G acts on X^0 in such a way that the subset

$X^1 \subset X^0 \times X^0$ is stable. We consider the Hilbert space $\ell^2(X^0)$, $(\delta_x, x \in X^0)$ its canonical Hilbert basis

and π_0 the unitary representation arising from the action of G on the set X^0 defined by $\pi_0(g)\delta_x = \delta_{gx}$. Let $\ell^2(X^1)^-$ be the quotient of $\ell^2(X^1)$ by the subspace generated by the vectors $\delta_{(x,y)} + \delta_{(y,x)}$ for $(x, y) \in X^1$ and π_1 the unitary representation of G on $\ell^2(X^1)^-$ defined by $\pi_1(g)\delta_{(x,y)} = \delta_{(gx,gy)}$

For $x \in X^0$ let $V(x)$ be the set of neighbours of x . Let the cardinality of $V(x)$ be denoted $q_x + 1$. We assume for simplicity that q_x is bounded. We define the following bounded operators S and Q on $\ell^2(X^0)$ by the following formulae:

$$S\delta_x = \sum_{y \in V(x)} \delta_y$$

$$Q\delta_x = q_x \delta_x$$

Note that the operators Q and S only depend on the tree structure on X and therefore commute with the unitaries $\pi_0(g)$.

2. The Pytlik-Szwarc operator.

We now choose an origin $x_0 \in X^0$. Let p_0 the orthogonal projection onto the vector δ_{x_0} . Let us define the operator P on $\ell^2(X^0)$ by

$$\begin{aligned} P\delta_{x_0} &= 0 \\ P\delta_x &= \delta_{x'} \end{aligned}$$

where for $x \neq x_0$ we define x' to be the unique neighbour of x lying between x_0 and x .

Proposition. 1 *The operator P such defined is bounded on $\ell^2(X^0)$ and satisfies:*

$$\begin{aligned} PP^* &= Q + p_0 \\ P + P^* &= S \end{aligned}$$

Proof: For $x \neq x_0$, one has $P^*\delta_x = \sum_{y \in V(x) \setminus \{x'\}} \delta_y$ so that clearly $(P + P^*)\delta_x = S\delta_x$. Similarly $(P + P^*)\delta_{x_0} = P^*\delta_{x_0} = \sum_{y \in V(x_0)} \delta_y = S\delta_{x_0}$.

On the other hand, for $x \neq x_0$ and $y \in V(x) \setminus \{x'\}$ one has $P\delta_y = \delta_x$ so that $PP^*\delta_x = q_x\delta_x$, whereas for any $y \in V(x_0)$, $P\delta_y = \delta_{x_0}$ so that $PP^*\delta_{x_0} = (q_{x_0} + 1)\delta_{x_0}$

Corollary. 1 *Let $T_t = 1 - tP + ((1 - t^2)^{1/2} - 1)p_0$ for any $t \in [0, 1]$ Then the operator T_t satisfies the following formula: $T_t T_t^* = 1 - tS + t^2Q$ and therefore commutes with the unitaries $\pi_0(g)$, $g \in G$.*

Indeed we have $T_t = 1 - tP + \alpha p_0$ where α satisfies $\alpha^2 + 2\alpha + t^2 = 0$. A straightforward calculation (using the obvious fact that $Pp_0 = 0$) yields $T_t T_t^* = 1 - t(P + P^*) + t^2 PP^* + (2\alpha + \alpha^2)p_0$ and the result follows from the proposition.

3. Construction of new representations.

Let us consider the space $D(X^0)$ of finitely supported functions on X^0 as a dense subspace of $\ell^2(X^0)$. Clearly $D(X^0)$ is stable by the operators P , P^* and $\pi_0(g)$ for $g \in G$.

Lemma. 1 *For any complex number z the operator*

$$(1 - zP)^{-1} = \sum_{k=0}^{\infty} z^k P^k$$

is defined on $D(X^0)$ and one has

$$(1 - zP)^{-1}\delta_x = \sum_{y \in [x_0, x]} z^{d(y, x)} \delta_y$$

Proof. Let the elements of $[x_0, x]$ be denoted $x_0, x_1, \dots, x_n = x$. Then $P^k\delta_x = \delta_{x_{n-k}}$ for $k \leq n$ and 0 for $k > n$. Note that $d(x_{n-k}, x) = k$. This makes the statement straightforward.

Theorem. 1 *(Pytlik-Szwarc) Let z be a complex number such that $|z| < 1$. For any $g \in G$ the operator $\rho_z(g) = (1 - zP)^{-1}\pi_0(g)(1 - zP)$ extends to a bounded operator on $\ell^2(X^0)$, thus defining a representation ρ_z of G in $\ell^2(X^0)$. The operator $\rho_z(g) - \pi_0(g)$ is a finite rank operator and the representation ρ_z is uniformly bounded, i.e.*

$$\sup_{g \in G} \|\rho_z(g)\| < \infty$$

Theorem. 2 For any real number such that $0 < t < 1$ the operator $\tilde{\rho}_t(g) = T_t^{-1}\pi_0(g)T_t$ extends to a unitary operator on $\ell^2(X^0)$, thus defining a unitary representation $\tilde{\rho}_t$ of G in $\ell^2(X^0)$. The operator $\tilde{\rho}_t(g) - \pi_0(g)$ is a finite rank operator. The uniformly bounded representation ρ_t is equivalent to the unitary representation $\tilde{\rho}_t$.

Proof: Let us prove that $\rho_z(g) - \pi_0(g)$ has finite rank and a norm bounded independently from g . It is enough to consider the operator

$$\rho_z(g)\pi_0(g)^{-1} - 1 = -z(1 - zP)^{-1}(P - P')$$

where P' is defined just as P but replacing x_0 by gx_0 . It is clear the the above operator is supported on the finite dimensional subspace generated by the δ_x for $x \in [x_0, gx_0]$. That subspace is indeed stable by P and P' which have restrictions of norm 1. Therefore the norm of $(1 - zP)^{-1}(P - P')$ is at most $2 \sum |z|^k = 2(1 - |z|)^{-1}$. This proves theorem 1.

To deduce theorem 2 note that $\tilde{\rho}_t(g) = u_t^{-1}\rho_t(g)u_t$ where $u_t = (1 - p_0) + (1 - t^2)^{1/2}p_0$ is an invertible operator which differs from the identity by a compact operator. It remains to show that $\tilde{\rho}_z(g)$ is unitary. Since it is invertible it is enough to compute $\rho_t(g)^*\rho_t(g) = T_t^*\pi_0(g)^{-1}(T_tT_t^*)^{-1}\pi_0(g)T_t$ which is equal to 1 since $T_tT_t^*$ commutes to $\pi_0(g)$ by the corollary to the proposition above.

Remark. The link with the original Julg-Valette approach is given by an easy computation: the kernel of the (densely defined) operator $(T_tT_t^*)^{-1}$ is

$$\langle T_t^{-1}\delta_x, T_t^{-1}\delta_y \rangle = t^{d(x,y)} = e^{-\lambda d(x,y)}$$

if $t = e^{-\lambda}$.

4. The limit when t tends to 1.

Let us now calculate the limit of $\tilde{\rho}_t(g)$ when $t \rightarrow 1$. Let us recall the definition of the Julg-Valette map $F : \ell^2(X^0) \rightarrow \ell^2(X^1)^-$:

$$F\delta_{x_0} = 0$$

$$F\delta_x = \delta_{(x,x')}$$

We have $Fp_0 = 0$, $F^*F = 1 - p_0$ and $FF^* = 1$.

Lemma. 2 Let $b : \ell^2(X^1)^- \rightarrow \ell^2(X^0)$ be the coboundary operator defined by $b\delta_{(x,y)} = \delta_y - \delta_x$. Then the operators F and P are related by the formulae:

$$1 - P = bF + p_0$$

$$(1 - P)F^* = b$$

Indeed, $(1 - P)\delta_x = \delta_x - \delta_{x'} = bF\delta_x$ if $x \neq x_0$ and $(1 - P)\delta_{x_0} = \delta_{x_0}$. The second formula follows from the first since $(1 - P)F^* = bFF^* = b$.

Remark. Let $c(x, y) = \sum \delta_{(x_i, x_{i+1})}$ if the elements of $[x, y]$ are denoted $x_0 = x, x_1, \dots, x_n = y$. This is the cocycle realizing explicitly the Haagerup property for groups acting properly on trees: $\|c(x, y)\|^2 = d(x, y)$. On has $bc(x, y) = \delta_y - \delta_x$, hence by the second formula above, $c(x, y) = F(1 - P)^{-1}(\delta_y - \delta_x)$.

Corollary. 2 *The operator $(1 - P)^{-1}b$ extends to a bounded operator and one has:*

$$(1 - P)^{-1}b = F^*$$

It follows indeed from the lemma that $(1 - P)F^* = b$.

Proposition. 2 *For any $g \in G$ the unitary operator $\tilde{\rho}_t(g)$ ($0 < t < 1$) converges strongly to*

$$\tilde{\rho}_1(g) = F^*\pi_1(g)F + p_0$$

when $t \rightarrow 1$.

Proof: Let us first prove that $F\tilde{\rho}_t(g)F^*$ strongly converges to $\pi_1(g)$. One has

$$F\tilde{\rho}_t(g)F^* = FT_t^{-1}\pi_0(g)T_tF^* = F(1 - tP)^{-1}\pi_0(g)(1 - tP)F^*$$

which evaluated on δ_e ($e \in X^1$) converges to $F(1 - P)^{-1}\pi_0(g)(1 - P)F^*$ i.e.

$$F(1 - P)^{-1}\pi_0(g)bFF^* = F(1 - P)^{-1}b\pi_1(g) = FF^*\pi_1(g) = \pi_1(g).$$

We deduce that $(1 - p_0)\tilde{\rho}_t(g)(1 - p_0)$ converges strongly to $F^*\pi_1(g)F$.

On the other hand we check that $\tilde{\rho}_t(g)\delta_{x_0} \rightarrow \delta_{x_0}$ when $t \rightarrow 1$. We have indeed $T_t\delta_{x_0} = (1 - t^2)^{1/2}\delta_{x_0}$ so that $(1 - tP)^{-1}\pi_0(g)T_t\delta_{x_0} = (1 - t^2)^{1/2}(1 - tP)^{-1}\delta_{gx_0} = (1 - t^2)^{1/2} \sum t^{d(y, gx_0)}\delta_y$ where the sum is extended to the vertices y of the segment $[x_0, gx_0]$. Finally $T_t^{-1}\pi_0(g)T_t\delta_{x_0} = t^{d(x_0, gx_0)} + (1 - t^2)^{1/2} \sum t^{d(y, gx_0)}\delta_y$ where x_0 is now excluded from the sum. Hence the result.

As a consequence $\tilde{\rho}_t(g)p_0$ converges normally to p_0 , and since $\tilde{\rho}_t$ is unitary, replacing g by g^{-1} we also have that $p_0\tilde{\rho}_t(g)$ converges normally to p_0 .

The proposition clearly follows.

5. Classes in KK_G -theory.

Let us consider the Hilbert space $\ell^2(X^0)$ equipped with the two representations $\tilde{\rho}_t$ and π_0 , which differ by compact operators. The triple $(\tilde{\rho}_t, \pi_0, Id)$ defines an element of the Kasparov group $KK_G(\mathbf{C}, \mathbf{C})$. It is independent of the value on $t \in [0, 1]$ since the operators $\tilde{\rho}_t(g)$ are strongly continuous in t . Now when $t = 0$ we have $\tilde{\rho}_0 = \pi_0$ so that the element is equal to 0. On the other hand when $t = 1$ we have $\tilde{\rho}_1(g) = F^*\pi_1(g)F + p_0$ so that the element is equal to $1 - \gamma$ where γ is defined as in Julg-Valette by the triple (π_0, π_1, F) .

Corollary. 3 *We have the equality $\gamma = 1$ in the Kasparov group $KK_G(\mathbf{C}, \mathbf{C})$.*

Bibliography:

- [B] Jacek Brodzki, *A differential complex for groups acting on CAT(0) cube complex.*
- [JV1] Pierre Julg, Alain Valette, *K-moyennabilité pour les groupes opérant sur les arbres*, C. R. Acad. Sci. Paris Sér. I Math. 296 (1983), no. 23, 977980.
- [JV2] Pierre Julg, Alain Valette, *K-theoretic amenability for $SL_2(\mathbf{Q}_p)$, and the action on the associated tree*, J. Funct. Anal. 58 (1984), no. 2, 194215.

[JV3] Pierre Julg, Alain Valette, *Fredholm modules associated to Bruhat-Tits buildings*. Miniconferences on harmonic analysis and operator algebras (Canberra, 1987), 143155, Proc. Centre Math. Anal. Austral. Nat. Univ., 16, Austral. Nat. Univ., Canberra, 1988.

[KS1] Gennadi Kasparov, Georges Skandalis, *Groups acting on buildings, operator K-theory, and Novikov's conjecture*, K-Theory 4 (1991), no. 4, 303337.

[KS2] Gennadi Kasparov, Georges Skandalis, *Groups acting properly on "bolic" spaces and the Novikov conjecture*. Ann. of Math. (2) 158 (2003), no. 1, 165206.

[PS] Tadeusz Pytlik, Ryszard Szwarc, *An analytic family of uniformly bounded representations of free groups*, Acta Math. 157 (1986), no. 3-4, 287309.

[V1] Alain Valette, *Cocycles d'arbres et représentations uniformément bornées*. C. R. Acad. Sci. Paris Sér. I Math. 310 (1990), no. 10, 703708.

[V2] Alain Valette, *Les représentations uniformément bornées associées un arbre réel*, Bull. Soc. Math. Belg. Sér. A 42 (1990), no. 3, 747760.

[V3] Alain Valette, *Addendum: "Uniformly bounded representations associated with a real tree"*. Bull. Soc. Math. Belg. Sér. A 44 (1992), no. 1, 101102.