



HAL
open science

Central limit theorem for eigenvectors of heavy tailed matrices

Florent Benaych-Georges, Alice Guionnet

► **To cite this version:**

Florent Benaych-Georges, Alice Guionnet. Central limit theorem for eigenvectors of heavy tailed matrices. 2013. hal-00877058v1

HAL Id: hal-00877058

<https://hal.science/hal-00877058v1>

Preprint submitted on 25 Oct 2013 (v1), last revised 30 May 2014 (v4)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

CENTRAL LIMIT THEOREM FOR EIGENVECTORS OF HEAVY TAILED MATRICES

FLORENT BENAYCH-GEORGES, ALICE GUIONNET

ABSTRACT. We consider the eigenvectors of symmetric matrices with independent heavy tailed entries, such as matrices with entries in the domain of attraction of α -stable laws, or adjacency matrices of Erdős-Rényi graphs. We denote by $U = [u_{ij}]$ the eigenvectors matrix (corresponding to increasing eigenvalues) and prove that the bivariate process

$$B_{s,t}^n := \frac{1}{\sqrt{n}} \sum_{\substack{1 \leq i \leq ns \\ 1 \leq j \leq nt}} (|u_{ij}|^2 - \frac{1}{n}) \quad (0 \leq s, t \leq 1),$$

converges in law to a non trivial Gaussian process. An interesting part of this result is the $\frac{1}{\sqrt{n}}$ rescaling, proving that from this point of view, the eigenvectors matrix U behaves more like a permutation matrix (as it was proved in [14] that for U a permutation matrix, $\frac{1}{\sqrt{n}}$ is the right scaling) than like a Haar-distributed orthogonal or unitary matrix (as it was proved in [15] that for U such a matrix, the right scaling is 1).

1. INTRODUCTION

During the last decade, many breakthroughs were achieved in the study of random matrices belonging to the GUE universality-class, that is Hermitian matrices with independent and equidistributed entries (modulo the symmetry constraint) with enough finite moments. The first key result about such matrices is due to Wigner [36] in the fifties who showed that the macroscopic behavior of their eigenvalues is universal and asymptotically described by the semi-circle distribution. However, it took a long time to get more precise information on the local behavior of the eigenvalues, and for instance about the asymptotic distribution of their spacings. Even though local results were conjectured, for instance by Dyson and Mehta [27], it is only in the nineties that rigorous results were derived, such as the convergence of the joint probability distribution of eigenvalues in an interval of size of order N^{-1}

2000 *Mathematics Subject Classification.* 15A52;60F05.

Key words and phrases. Random matrices, heavy tailed random variables, eigenvectors, central limit theorem.

FBG: florent.benaych-georges@parisdescartes.fr, MAP 5, UMR CNRS 8145 - Université Paris Descartes, 45 rue des Saints-Pères 75270 Paris Cedex 6, France.

AG: aguionne@ens-lyon.fr, CNRS & École Normale Supérieure de Lyon, Unité de mathématiques pures et appliquées, 46 allée d'Italie, 69364 Lyon Cedex 07, France and MIT, Mathematics Department, 77 Massachusetts Av, Cambridge MA 02139-4307, USA.

Research supported by Simons Foundation.

or the fluctuations of the largest eigenvalues, see [34]. Yet these results were restricted to Gaussian ensembles for which the joint law of the eigenvalues is known. Recently, these results were shown to be universal, that is to hold also for matrices with independent non Gaussian entries, provided they have enough finite moments [17, 20, 22, 23, 32]. Such a simple question as the convergence of the law of a single spacing was open, even in the GUE case, until recently when it was solved by Tao [33]. Once considering non Gaussian matrices, it is natural to wonder about the behavior of the eigenvectors and whether they are delocalized (that is go to zero in L^∞ norm) as for GUE matrices. This was indeed shown by Erdős, Schlein and Yau [18].

Despite the numerous breakthroughs concerning random matrices belonging to the GUE universality-class, not much is yet known about other matrices. A famous example of such a matrix is given by the adjacency matrix of an Erdős-Rényi graph. Its entries are independent (modulo the symmetry hypothesis) and equal to one with probability $p = p(N)$, zero otherwise. If pN goes to infinity fast enough, the matrix belongs to the GUE universality class [16]. However if pN converges to a finite constant non zero constant, the matrix behaves quite differently, more like a “heavy tailed random matrix”, *i.e.* a matrix filled with independent entries which have no finite second moment. Also in this case, it is known that, once properly normalized, the empirical measure of the eigenvalues converges weakly almost surely but the limit differs from the semi-circle distribution [35, 5, 4, 7, 12, 8]. Moreover, the fluctuations of the empirical measure could be studied [8, 28, 24, 25]. It turns out that it fluctuates much more than in the case of matrices from the GUE universality-class, as fluctuations are square root of the dimension bigger. However, there is no result about the local fluctuations of the eigenvalues except in the case of matrices with entries in the domain of attraction of an α -stable law in which case it was shown [2, 31] that the largest eigenvalues are much bigger than the others, converge to a Poisson distribution and have localized eigenvectors. About localization and delocalization of the eigenvectors, some models are conjectured [13, 30] to exhibit a phase transition; eigenvalues in a compact would have more delocalized eigenvectors than outside this compact. Unfortunately, very little could be proved so far in this direction. Only the case where the entries are α -stable random variables could be tackled [12]; it was shown that for $\alpha > 1$ the eigenvectors are delocalized whereas for $\alpha < 1$ and large eigenvalues, a weak form of localization holds.

In this article, we study another type of properties of the eigenvectors of a random matrix. Namely we consider the bivariate process

$$G_{s,t}^n := \sum_{\substack{1 \leq i \leq ns \\ 1 \leq j \leq nt}} (|u_{ij}|^2 - \frac{1}{n}) \quad (0 \leq s \leq 1, \quad 0 \leq t \leq 1),$$

where $U = [u_{ij}]$ is an orthogonal matrix whose columns are the eigenvectors of an Hermitian random matrix A . In the case where A is a GUE matrix [15], and then a more general matrix in the GUE universality-class [6], it was shown that this process converges in law towards a bivariate Brownian bridge. Here, we investigate the same process in the case where A is a heavy tailed random matrix and show that it fluctuates much more, namely it

is $\frac{1}{\sqrt{n}}G^n$ which converges in law. The limit is a Gaussian process whose covariance depends on the model and is not trivial (note that when the u_{ij} 's are the entries of a uniformly distributed random permutation, *i.e.* of a somehow sparse matrix, the process $\frac{1}{\sqrt{n}}G^n$ also converges in law, towards the bivariate Brownian bridge [14]).

More precisely, we consider a real symmetric random $n \times n$ matrix A that can be either a Wigner matrix with exploding moments (which includes the adjacency matrix for Erdős-Rényi graphs) or a matrix with i.i.d. entries in the domain of attraction of a stable law (or more generally a matrix satisfying the hypotheses detailed in Hypothesis 2.1). We then introduce an orthogonal matrix $U = [u_{ij}]$ whose columns are the eigenvectors of A so that we have $A = U \text{diag}(\lambda_1, \dots, \lambda_n)U^*$. We then define the bivariate processes

$$B_{s,t}^n := \frac{1}{\sqrt{n}} \sum_{\substack{1 \leq i \leq ns \\ 1 \leq j \leq nt}} (|u_{ij}|^2 - \frac{1}{n}) \quad (0 \leq s \leq 1, \quad 0 \leq t \leq 1)$$

and

$$C_{s,\lambda}^n := \frac{1}{\sqrt{n}} \sum_{\substack{1 \leq i \leq ns \\ 1 \leq j \leq n; \lambda_j \leq \lambda}} (|u_{ij}|^2 - \frac{1}{n}) \quad (0 \leq s \leq 1, \quad \lambda \in \mathbb{R})$$

and prove, in Theorem 2.4, that both of these processes (with a little technical restriction on the domain of B) converge in law to (non trivial) Gaussian processes linked by the relation

$$B_{s, F_{\mu_\Phi}(\lambda)} = C_{s,\lambda},$$

where $F_{\mu_\Phi}(\lambda) = \mu_\Phi((-\infty, \lambda])$ denotes the cumulative distribution function of the limit spectral law μ_Φ of A , *i.e.*

$$(1) \quad F_{\mu_\Phi}(\lambda) = \lim_{n \rightarrow \infty} F_n(\lambda), \quad \text{with} \quad F_n(\lambda) := \frac{1}{n} |\{i; \lambda_i \leq \lambda\}|.$$

The idea of the proof is the following one. We first notice that for any $s \in [0, 1]$, the function $\lambda \mapsto C_{s,\lambda}^n$ is the cumulative distribution function of the random signed measure $\nu_{s,n}$ on \mathbb{R} defined by

$$(2) \quad \nu_{s,n} := \frac{1}{\sqrt{n}} \sum_{1 \leq i \leq ns} \sum_{j=1}^n \left(|u_{ij}|^2 - \frac{1}{n} \right) \delta_{\lambda_j}$$

(*i.e.* that for any $\lambda \in \mathbb{R}$, $C_{s,\lambda}^n = \nu_{s,n}((-\infty, \lambda])$). Then, we introduce the Cauchy transform $X^n(s, z) := \int_{\lambda \in \mathbb{R}} \frac{d\nu_{s,n}(\lambda)}{z - \lambda}$ of $\nu_{s,n}$ and prove (Proposition 2.8) that the process $(X^n(s, z))_{s,z}$ converges in law to a limit Gaussian process $(H_{s,z})$. This convergence is proved thanks to the classical CLT for martingales (Theorem 6.3 of the Appendix) together with the Schur complement formula and fixed points characterizations like the ones of the papers [5, 4, 8]. Then to deduce the convergence in law of the process $(C_{s,\lambda}^n)_{s,\lambda}$, we use the idea that the cumulative distribution function of a signed measure is entirely determined by its Cauchy transform. In fact, as the measures $\nu_{s,n}$ of (2) are random, things are slightly more

complicated, and we need to prove a tightness lemma for the process $(C_{s,\lambda}^n)_{s,\lambda}$ (specifically Lemma 6.1 of the Appendix, first applied to the process $(B_{s,t}^n)$ and then transferred to $(C_{s,\lambda}^n)$ by Formula (3) below). This lemma reduces the problem to the proof of the unicity of the possible limits for $(C_{s,\lambda}^n)$. Then, we use the formula

$$\int_{\lambda \in \mathbb{R}} \frac{C_{s,\lambda}^n}{(z - \lambda)^2} d\lambda = -X^n(s, z)$$

and Lemma 6.2 of the appendix to be able to claim that $(C_{s,\lambda}^n)_{s,\lambda}$ has a unique limit point. The result proved for $(C_{s,\lambda}^n)$ can then be transferred to $(B_{s,t}^n)$ thanks to Formula (3) below, where $F_n(\lambda)$, defined at (1) above, converges to the deterministic limit $F_{\mu_{\mathfrak{q}}}(\lambda)$:

$$(3) \quad C_{s,\lambda}^n = B_{s,F_n(\lambda)}^n.$$

Organization of the paper. The main results are stated in Section 2. In Section 3, we give a proof of Theorem 2.4, based on Proposition 2.8, which is proved in Section 4. Proposition 2.6 is proved in Section 5. At last, some technical results are proved or recalled in the Appendix.

Notations. For u, v depending implicitly on n , we write $u \ll v$ when $u/v \rightarrow 0$ as $n \rightarrow \infty$. For x a random variable, $\text{Var}(x)$ denotes the *variance* of x , i.e. $\mathbb{E}[|x|^2] - |\mathbb{E}x|^2$. Power functions are defined on $\mathbb{C} \setminus \mathbb{R}_-$ via the standard determination of the argument on this set taking values in $(-\pi, \pi)$. The set \mathbb{C}^+ (resp. \mathbb{C}^-) denotes the open upper (resp. lower) half plane and for any $z \in \mathbb{C}$, $\text{sgn}_z := \text{sign}(\Im z)$.

2. MAIN RESULTS

Although technical, the model introduced in Hypothesis 2.1 below has the advantage to be general enough to contain several models of interest.

Hypothesis 2.1. Let, for each $n \geq 1$, $A_n = [a_{ij}]$ be an $n \times n$ real symmetric random matrix whose sub-diagonal entries are some i.i.d. copies of a random variable a (depending implicitly on n) such that:

- The random variable a can be decomposed into $a = b + c$ such that as $n \rightarrow \infty$,

$$(4) \quad \mathbb{P}(c \neq 0) \ll n^{-1}$$

$$(5) \quad \text{Var}(b) \ll n^{-1/2}$$

Moreover, if the b_i 's are independent copies of b ,

$$(6) \quad \lim_{K \rightarrow \infty} \lim_{n \rightarrow \infty} \mathbb{P} \left(\sum_{i=1}^n (b_i - \mathbb{E}(b_i))^2 \geq K \right) = 0.$$

- For any $\varepsilon > 0$ independent of N , the random variable a can be decomposed into $a = b_\varepsilon + c_\varepsilon$ such that

$$(7) \quad \limsup_{n \rightarrow \infty} n \mathbb{P}(c_\varepsilon \neq 0) \leq \varepsilon$$

for all $k \geq 1$, $n \mathbb{E}[(b_\varepsilon - \mathbb{E} b_\varepsilon)^{2k}]$ has a finite limit $C_{\varepsilon,k}$ as $n \rightarrow \infty$.

- For ϕ_n the function defined on the closure $\overline{\mathbb{C}^-}$ of $\mathbb{C}^- := \{\lambda \in \mathbb{C}; \Im \lambda < 0\}$ by

$$(8) \quad \phi_n(\lambda) := \mathbb{E} [\exp(-i\lambda a^2)],$$

we have the convergence, uniform on compact subsets of $\overline{\mathbb{C}^-}$,

$$(9) \quad n(\phi_n(\lambda) - 1) \longrightarrow \Phi(\lambda),$$

for a certain function Φ defined on $\overline{\mathbb{C}^-}$.

- The function Φ of (9) admits the decomposition

$$(10) \quad \Phi(z) = \int_0^\infty g(y) e^{i\frac{y}{z}} dy$$

where $g(y)$ is a function such that for some constants $K, \gamma > -1, \kappa \geq 0$, we have

$$(11) \quad |g(y)| \leq K 1_{y \leq 1} y^\gamma + K 1_{y \geq 1} y^\kappa, \quad \forall y > 0.$$

- The function Φ of (9) also either has the form

$$(12) \quad \Phi(x) = -\sigma(ix)^{\alpha/2}$$

or admits the (other) decomposition, for x, y non zero:

$$(13) \quad \Phi(x+y) = \iint_{(\mathbb{R}^+)^2} e^{i\frac{x}{v} + i\frac{y}{v'}} d\tau(v, v') + \int_{\mathbb{R}^+} e^{i\frac{x}{v}} d\mu(v) + \int_{\mathbb{R}^+} e^{i\frac{y}{v'}} d\mu(v')$$

for some complex measures τ, μ on respectively $(\mathbb{R}^+)^2$ and \mathbb{R}^+ such that for all $b > 0$, $\int e^{-bv} d|\mu|(v)$ is finite and for some constants $K > 0$, $-1 < \gamma \leq 0$ and $\kappa \geq 0$, and

$$(14) \quad \frac{d|\tau|(v, v')}{dv dv'} \leq K (v^\gamma 1_{v \in]0, 1]} + v^\kappa 1_{v \in]1, \infty[}) (v'^\gamma 1_{v' \in]0, 1]} + v'^\kappa 1_{v' \in]1, \infty[}).$$

Remark 2.1. When Φ satisfies (12) (e.g. for Lévy matrices), (13) holds as well. Indeed, for all $x, y \in \mathbb{C}^+$ (with a constant C_α that can change at every line),

$$(15) \quad \begin{aligned} \Phi(x^{-1} + y^{-1}) &= C_\alpha \left(\frac{1}{x} + \frac{1}{y}\right)^{\alpha/2} = C_\alpha \frac{1}{x^{\alpha/2}} \frac{1}{y^{\alpha/2}} (x+y)^{\alpha/2} \\ &= C_\alpha \int_0^\infty dw \int_0^\infty dw' \int_0^\infty dv w^{\alpha/2-1} w'^{\alpha/2-1} v^{-\alpha/2-1} e^{iwx+iw'y} (e^{iv(x+y)} - 1) \end{aligned}$$

(where we used the formula $z^{\alpha/2} = C_\alpha \int_{t=0}^{+\infty} \frac{e^{itz} - 1}{t^{\alpha/2+1}} dt$ for any $z \in \mathbb{C}^+$ and $\alpha \in (0, 2)$, which can be proved with the residues formula) so that (13) holds with $\mu = 0$ and $\tau(v, v')$ with density with respect to Lebesgue measure given by

$$(16) \quad C_\alpha \int_0^{+\infty} u^{-\alpha/2-1} \{(v-u)^{\alpha/2-1} (v'-u)^{\alpha/2-1} 1_{0 \leq u \leq v \wedge v'} - v^{\alpha/2-1} v'^{\alpha/2-1}\} du.$$

Unfortunately τ does not satisfy (14) as its density blows up at $v = v'$: we shall treat both case separately.

Examples of random matrices satisfying Hypothesis 2.1 are defined as follows.

Definition 2.2 (Models of symmetric heavy tailed matrices). Let $A = (a_{i,j})_{i,j=1,\dots,n}$ be a random symmetric matrix with i.i.d. sub-diagonal entries.

1. We say that A is a **Lévy matrix** of parameter α in $]0, 2[$ when $A = X/a_n$ where the entries x_{ij} of X have absolute values in the domain of attraction of α -stable distribution, more precisely

$$(17) \quad \mathbb{P}(|x_{ij}| \geq u) = \frac{L(u)}{u^\alpha}$$

with a slowly varying function L , and

$$a_n = \inf\{u : P(|x_{ij}| \geq u) \leq \frac{1}{n}\}$$

($a_n = \tilde{L}(n)n^{1/\alpha}$, with $\tilde{L}(\cdot)$ a slowly varying function).

2. We say that A is a **Wigner matrix with exploding moments** with parameter $(C_k)_{k \geq 1}$ whenever the entries of A are centered, and for any $k \geq 1$

$$(18) \quad n \mathbb{E}[(a_{ij})^{2k}] \xrightarrow[n \rightarrow \infty]{} C_k,$$

with for a constant $C > 0$. We assume that there exists a unique measure m on \mathbb{R}^+ such that for all $k \geq 0$,

$$(19) \quad C_{k+1} = \int x^k dm(x).$$

The following proposition has been proved at Lemmas 1.3, 1.8 and 1.11 of [8].

Proposition 2.3. *Both Lévy matrices and Wigner matrices with exploding moments satisfy Hypothesis 2.1:*

- For Lévy matrices, the function Φ of (9) is given by formula

$$(20) \quad \Phi(\lambda) = -\sigma(i\lambda)^{\alpha/2}$$

for some constant $\sigma \geq 0$, the function g of (10) is $g(y) = C_\alpha y^{\frac{\alpha}{2}-1}$, with $C_\alpha = -\sigma i^{\alpha/2}$.

- For Wigner matrices with exploding moments, the function Φ of (9) is given by

$$(21) \quad \Phi(\lambda) = \int \underbrace{\frac{e^{-i\lambda x} - 1}{x}}_{:= -i\lambda \text{ for } x=0} dm(x),$$

for m the measure of (19), the function g of (10) is

$$(22) \quad g(y) = - \int_{\mathbb{R}^+} \underbrace{\frac{J_1(2\sqrt{xy})}{\sqrt{xy}}}_{:= 1 \text{ for } xy=0} dm(x),$$

for J_1 the Bessel function of the first kind defined by $J_1(s) = \frac{s}{2} \sum_{k \geq 0} \frac{(-s^2/4)^k}{k!(k+1)!}$, and the measures τ and μ of (13) are absolutely continuous with densities

$$(23) \quad \frac{d\tau(v, v')}{dv dv'} := \int \frac{J_1(2\sqrt{vx})J_1(2\sqrt{v'x})}{\sqrt{vv'}} dm(x) \quad \text{and} \quad \frac{d\mu(v)}{dv} := - \int \frac{J_1(2\sqrt{vx})}{\sqrt{v}} dm(x).$$

One can easily see that our results also apply to complex Hermitian matrices: in this case, one only needs to require Hypothesis 2.1 to be satisfied by the absolute value of non diagonal entries and to have a_{11} going to zero as $N \rightarrow \infty$.

A Lévy matrix whose entries are truncated in an appropriate way is a Wigner matrix with exploding moments [5, 26, 35]. The recentered version¹ of the adjacency matrix of an Erdős-Rényi graph, *i.e.* of a matrix A such that

$$(24) \quad A_{ij} = 1 \text{ with probability } p/n \text{ and } 0 \text{ with probability } 1 - p/n,$$

is also an exploding moments Wigner matrix, with $\Phi(\lambda) = p(e^{-i\lambda} - 1)$ (the measure m is $p\delta_1$). In this case the fluctuations were already studied in [28].

It has been proved in [8] (see also [35, 28]) that under Hypothesis 2.1, the empirical spectral law

$$(25) \quad \mu_n := \frac{1}{n} \sum_{j=1}^n \delta_{\lambda_j}$$

converges weakly in probability to a deterministic probability measure μ_Φ that depends only on Φ , *i.e.* that for any continuous bounded function $f : \mathbb{R} \rightarrow \mathbb{C}$, we have the almost sure convergence

$$(26) \quad \frac{1}{n} \text{Tr } f(A) = \frac{1}{n} \sum_{j=1}^n f(\lambda_j) \xrightarrow[n \rightarrow \infty]{} \int f(x) d\mu_\Phi(x).$$

We introduce $F_{\mu_\Phi}(\lambda) := \mu_\Phi((-\infty, \lambda])$, cumulative distribution function of μ_Φ and define the set $E_\Phi \subset [0, 1]$ by

$$(27) \quad E_\Phi := \{0\} \cup F_{\mu_\Phi}(\mathbb{R}) \cup \{1\}.$$

In the case of Lévy matrices, it has been proved in [4, Theorem 1.3] that μ_Φ has no atoms (because it is absolutely continuous), so that $E_\Phi = [0, 1]$.

We introduce the eigenvalues $\lambda_1 \leq \dots \leq \lambda_n$ of A and an orthogonal matrix $U = [u_{ij}]$ such that $A = U \text{diag}(\lambda_1, \dots, \lambda_n) U^*$. We assume U defined in such a way that the rows

¹The recentering has in fact asymptotically no effect on the spectral measure A as it is a rank one perturbation.

of the matrix $[|u_{ij}|]$ are exchangeable (this is possible² because A is invariant, in law, by conjugation by any permutation matrix). Then define the bivariate processes

$$B_{s,t}^n := \frac{1}{\sqrt{n}} \sum_{\substack{1 \leq i \leq ns \\ 1 \leq j \leq nt}} (|u_{ij}|^2 - \frac{1}{n}) \quad (0 \leq s \leq 1, \quad 0 \leq t \leq 1)$$

and

$$C_{s,\lambda}^n := \frac{1}{\sqrt{n}} \sum_{\substack{1 \leq i \leq ns \\ 1 \leq j \leq n; \lambda_j \leq \lambda}} (|u_{ij}|^2 - \frac{1}{n}) \quad (0 \leq s \leq 1, \quad \lambda \in \mathbb{R}).$$

The following theorem is the main result of this article. We endow $D([0,1]^2)$ and $D([0,1] \times \mathbb{R})$ with the Skorokhod topology and $D([0,1] \times E_\Phi)$ with the topology induced by the Skorokhod topology on $D([0,1]^2)$ by the projection map from $D([0,1]^2)$ onto $D([0,1] \times E_\Phi)$ (see Section 4.1 of [6] for the corresponding definitions).

Theorem 2.4. *As $n \rightarrow \infty$, the joint distribution of the processes*

$$(B_{s,t}^n)_{(s,t) \in [0,1] \times E_\Phi} \quad \text{and} \quad (C_{s,\lambda}^n)_{(s,\lambda) \in [0,1] \times \mathbb{R}}$$

converges weakly to the joint distribution of some centered Gaussian processes

$$(B_{s,t})_{(s,t) \in [0,1] \times E_\Phi} \quad \text{and} \quad (C_{s,\lambda})_{(s,\lambda) \in [0,1] \times \mathbb{R}}$$

vanishing on the boundaries of their domains and satisfying the relation

$$(28) \quad B_{s, F_{\mu_\Phi}(\lambda)} = C_{s,\lambda}$$

for all $s \in [0,1]$, $\lambda \in \mathbb{R}$. Moreover, the process the process $(B_{s,t})$ is continuous on $[0,1] \times E_\Phi$.

Remark 2.5. Note that the limit of $B_{s,t}^n$ is only given here when $t \in E_\Phi$, *i.e.* when t is not in the ‘‘holes’’ of $F_{\mu_\Phi}(\mathbb{R})$. But as these holes result from the existence of some atoms in the limit spectral distribution of A , the variations of $B_{s,t}^n$ when t varies in one of these holes may especially depend on the way we choose the columns of A for eigenvalues with multiplicity larger than one. By the results of [15], in the case where the atoms of μ_Φ result in atoms (with asymptotically same weight) of μ_n , the choice we made here should lead to a limit process $(B_{s,t})_{(s,t) \in [0,1]^2}$ which would interpolate $(B_{s,t})_{(s,t) \in [0,1] \times E_\Phi}$ with some Brownian bridges in these ‘‘holes’’, namely for when $t \in [0,1] \setminus E_\Phi$.

The following proposition insures that the $\frac{1}{\sqrt{n}}$ scaling in the definitions of $B_{s,t}^n$ and $C_{s,\lambda}^n$ is the right one.

Proposition 2.6. *If the function $\Phi(z)$ of (9) is not identically equal to z , then for any fixed $s \in (0,1)$, the covariance of the process $(B_{s,t})_{t \in E_\Phi}$ (hence also that of $(C_{s,\lambda})_{\lambda \in \mathbb{R}}$) is not identically null.*

²Such a matrix U can be defined, for example, by choosing some orthogonal bases of all eigenspaces of A with uniform distributions, independently with each other and independently of A (given its eigenspaces of course).

Remark 2.7. One could wonder if the covariance might vanish uniformly on some compact in the t variable, hence giving some support to the belief that the eigenvectors could behave more alike the eigenvectors of GUE for “small” eigenvalues (in the latter case the covariance should vanish). Unfortunately, it does not seem that the covariance should be so closely related with the localization/delocalization properties of the eigenvectors. Indeed, let us consider Lévy matrices with $\alpha \in (1, 2)$. Their eigenvectors are delocalized [12], so that one could expect the covariance of the process $(B_{s,t})_{t \in E_\Phi}$ to vanish. This is in contradiction with the fact that such matrices enter our model, hence have eigenvectors satisfying Theorem 2.4 and Proposition 2.6.

To prove Theorem 2.4, a key step will be to prove the following proposition, which also allows to make the variance of the limiting processes in Theorem 2.4 more explicit.

Let us define, for $z \in \mathbb{C} \setminus \mathbb{R}$ and $s \in [0, 1]$,

$$(29) \quad X^n(s, z) := \frac{1}{\sqrt{n}} \left(\text{Tr} \left(P_s \frac{1}{z - A} \right) - s_n \text{Tr} \frac{1}{z - A} \right),$$

where P_s denotes the diagonal matrix with diagonal entries $1_{i \leq ns}$ ($1 \leq i \leq n$) and

$$s_n := \frac{1}{n} \text{Tr} P_s = \frac{\lfloor ns \rfloor}{n}.$$

Proposition 2.8. *The distribution of the random process*

$$(X^n(s, z))_{s \in [0, 1], z \in \mathbb{C} \setminus \mathbb{R}}$$

converges weakly in the sense of finite marginals towards the distribution of a centered Gaussian process

$$(30) \quad (H_{s,z})_{s \in [0, 1], z \in \mathbb{C} \setminus \mathbb{R}}$$

with a covariance given by (51).

As it will appear from the proofs that the process $(C_{s,\lambda})$ of Theorem 2.4 and the process $(H_{s,z})$ from the previous proposition are linked by the formula

$$(31) \quad \int_{\lambda \in \mathbb{R}} \frac{C_{s,\lambda}}{(z - \lambda)^2} d\lambda = -H_{s,z} \quad (s \in [0, 1], z \in \mathbb{C} \setminus \mathbb{R}),$$

the covariance of $C_{s,\lambda}$ (hence of $B_{s,t}$ by (28)) can be deduced from that of the process $H_{s,z}$ as follows (the proof of this proposition is a direct application of (31) and of Formula (66) of the Appendix).

Proposition 2.9. *For any $s, s' \in [0, 1]$ and any $\lambda, \lambda' \in \mathbb{R}$ which are not atoms of μ_Φ , we have*

$$(32) \quad \mathbb{E}[C_{s,\lambda} C_{s',\lambda'}] = \frac{1}{\pi^2} \lim_{\eta \downarrow 0} \int_{-\infty}^{\lambda} \int_{-\infty}^{\lambda'} \mathbb{E}[\Im(H_{s,E+i\eta}) \Im(H_{s',E'+i\eta})] dE dE'.$$

When λ or λ' is an atom of μ_Φ , the covariance can be obtained using (32) and the right continuity of $C_{s,\lambda}$ in λ .

3. PROOF OF THEOREM 2.4

We introduce the cumulative distribution function

$$(33) \quad F_n(\lambda) := \frac{1}{n} |\{j; \lambda_j \leq \lambda\}|$$

of the empirical spectral law μ_n defined at (25). We shall use the following formula several times: for all $s \in [0, 1]$ and $\lambda \in \mathbb{R}$,

$$(34) \quad C_{s,\lambda}^n = B_{s,F_n(\lambda)}^n.$$

We know, by Lemma 6.1 of the appendix, that the sequence $(\text{distribution}(B^n))_{n \geq 1}$ is tight and has all its accumulation points supported by the set of continuous functions on $[0, 1]^2$. As F_n converges to F_{μ_Φ} for the Skorokhod topology, it follows that the sequences

$$\tilde{B}^n := (B_{s,t}^n)_{(s,t) \in [0,1] \times E_\Phi} \quad \text{and} \quad (C_{s,\lambda}^n = B_{s,F_n(\lambda)}^n)_{(s,\lambda) \in [0,1] \times \mathbb{R}}$$

are tight in their respective spaces. To prove the theorem, it suffices to prove that the sequence $(\text{distribution}(\tilde{B}^n, C^n))_{n \geq 1}$ has only one accumulation point (which is Gaussian centered, vanishing on the boundaries, supported by continuous functions as far as the first component is concerned and satisfying (28)). So let $((\tilde{B}_{s,t})_{(s,t) \in [0,1] \times E_\Phi}, (C_{s,\lambda})_{(s,\lambda) \in [0,1] \times \mathbb{R}})$ be a pair of random processes having for distribution such an accumulation point. By (34), we have

$$\tilde{B}_{s,F_{\mu_\Phi}(\lambda)} = C_{s,\lambda}$$

for all $s \in [0, 1]$, $\lambda \in \mathbb{R}$. Hence it suffices to prove that the distribution of C is totally prescribed and Gaussian centered.

First, let us note that one can suppose that along the corresponding subsequence, the distribution of $((B_{s,t}^n)_{(s,t) \in [0,1]^2}, (C_{s,\lambda}^n)_{(s,\lambda) \in [0,1] \times \mathbb{R}})$ converges weakly to the distribution of a pair (B, C) of processes such that B is continuous and vanishing on the boundary of $[0, 1]^2$ (not that the difference with what was supposed above is that now, t varies in $[0, 1]$ and not only in E_Φ). Again, by (34), we have

$$(35) \quad B_{s,F_{\mu_\Phi}(\lambda)} = C_{s,\lambda}$$

for all $s \in [0, 1]$, $\lambda \in \mathbb{R}$. Hence the process C is continuous in s and continuous in λ at any λ which is not an atom of the (non random) probability measure μ_Φ . Hence it follows from Lemma 6.2 of the appendix that it suffices to prove that the distribution of the process

$$\left(X(s, z) := \int_{\lambda \in \mathbb{R}} \frac{C_{s,\lambda}}{(z - \lambda)^2} d\lambda \right)_{s \in [0,1], z \in \mathbb{C} \setminus \mathbb{R}}$$

is totally prescribed (and Gaussian centered). This distribution is the limit distribution, along our subsequence, of the process

$$(36) \quad \left(\int_{\lambda \in \mathbb{R}} \frac{C_{s,\lambda}^n}{(z - \lambda)^2} d\lambda \right)_{s \in [0,1], z \in \mathbb{C} \setminus \mathbb{R}}.$$

But by Lemma 3.1 below, the process of (36) is simply (the opposite of) the process $(X^n(s, z))_{s, z}$, defined above at (29). As Proposition 2.8 states that (regardless to the subsequence considered) the distribution of the process $(X^n(s, z))_{s, z}$ converges weakly to a Gaussian centered limit, this concludes the proof of Theorem 2.4.

Lemma 3.1. *For any $s \in [0, 1]$ and any $z \in \mathbb{C} \setminus \mathbb{R}$, we have*

$$(37) \quad \int_{\lambda \in \mathbb{R}} \frac{C_{s, \lambda}^n}{(z - \lambda)^2} d\lambda = -X^n(s, z).$$

Proof. Let us introduce, for $s \in [0, 1]$, the random signed measure $\nu_{s, n}$ on \mathbb{R} defined by

$$\nu_{s, n} := \frac{1}{\sqrt{n}} \sum_{1 \leq i \leq ns} \sum_{j=1}^n \left(|u_{ij}|^2 - \frac{1}{n} \right) \delta_{\lambda_j}.$$

Then for any $\lambda \in \mathbb{R}$, $C_{s, \lambda}^n = \nu_{s, n}((-\infty, \lambda])$. Moreover, by Fubini, we know that for any finite signed measure m on \mathbb{R} ,

$$(38) \quad \int_{\lambda \in \mathbb{R}} \frac{m((-\infty, \lambda])}{(z - \lambda)^2} d\lambda = - \int_{\lambda \in \mathbb{R}} \frac{dm(\lambda)}{z - \lambda}.$$

Hence

$$\int_{\lambda \in \mathbb{R}} \frac{C_{s, \lambda}^n}{(z - \lambda)^2} d\lambda = - \int_{\lambda \in \mathbb{R}} \frac{d\nu_{s, n}(\lambda)}{z - \lambda}.$$

On the other hand, we have

$$\begin{aligned} X^n(s, z) &= \frac{1}{\sqrt{n}} \left(\sum_{1 \leq i \leq ns} \left(\frac{1}{z - A} \right)_{ii} - \frac{1}{n} \sum_{1 \leq i \leq ns} \sum_{i=1}^n \frac{1}{z - \lambda_j} \right) \\ &= \frac{1}{\sqrt{n}} \left(\sum_{1 \leq i \leq ns} \sum_{i=1}^n |u_{ij}|^2 \frac{1}{z - \lambda_j} - \frac{1}{n} \sum_{1 \leq i \leq ns} \sum_{i=1}^n \frac{1}{z - \lambda_j} \right) \\ &= \frac{1}{\sqrt{n}} \sum_{1 \leq i \leq ns} \sum_{i=1}^n \left(|u_{ij}|^2 - \frac{1}{n} \right) \frac{1}{z - \lambda_j} \\ &= \int_{\lambda \in \mathbb{R}} \frac{d\nu_{s, n}(\lambda)}{z - \lambda}. \end{aligned}$$

This concludes the proof. \square

4. PROOF OF PROPOSITION 2.8

To prove Proposition 2.8, one needs to prove that the distribution of any linear combination of the $X^n(s, z)$'s ($s \in [0, 1]$, $z \in \mathbb{R}$) converges weakly. For $s = 0$ or 1 , $\nu_{s, n}$ is null, as $X^n(s, z)$, hence we can focus on $s \in (0, 1)$. Any such linear combination can be written

$$M^n := \sum_{i=1}^p \alpha_i X^n(s_i, z_i),$$

for some α_i 's in \mathbb{C} , some s_i 's in $[0, 1]$ and some complex non real numbers z_i .

We want to prove that M^n converges in law to a certain complex centered Gaussian variable. We are going to use the CLT for martingales stated at Theorem 6.3 of the appendix. Indeed, for \mathcal{F}_k^n the σ -algebra generated by the first k rows (or columns) of the symmetric matrix A , the sequence $(M_k^n := \mathbb{E}[M^n | \mathcal{F}_k^n])_{k=0, \dots, n}$ is a centered martingale (to see that it is centered, just use the fact that as A is invariant, in law, by conjugation by any permutation matrix, for all z , the expectation of $(\frac{1}{z-A})_{jj}$ does not depend on j).

Then, denoting $\mathbb{E}[\cdot | \mathcal{F}_k^n]$ by \mathbb{E}_k , and defining

$$Y_k := (\mathbb{E}_k - \mathbb{E}_{k-1})(M^n)$$

(which depends implicitly on n), we need to prove that for any $\varepsilon > 0$,

$$(39) \quad L^n(\varepsilon) := \sum_{k=1}^n \mathbb{E}(|Y_k|^2 1_{|Y_k| \geq \varepsilon}) \xrightarrow{n \rightarrow \infty} 0,$$

and that the sequences

$$\sum_{k=1}^n \mathbb{E}_{k-1}(|Y_k|^2) \quad \text{and} \quad \sum_{k=1}^n \mathbb{E}_{k-1}(Y_k^2)$$

converge in probability towards some deterministic limits. As $\overline{X^n(s, z)} = X^n(s, \bar{z})$, it is in fact enough to fix $s, s' \in (0, 1)$ and $z, z' \in \mathbb{C} \setminus \mathbb{R}$ and to prove that for

$$(40) \quad Y_k := (\mathbb{E}_k - \mathbb{E}_{k-1})(X^n(s, z)) \quad \text{and} \quad Y'_k := (\mathbb{E}_k - \mathbb{E}_{k-1})(X^n(s', z')),$$

we have (39) for any $\varepsilon > 0$ and that

$$\sum_{k=1}^n \mathbb{E}_{k-1}(Y_k Y'_k)$$

converges in probability towards a deterministic constant. We introduce the notation

$$G := \frac{1}{z - A} \quad \text{and} \quad G' := \frac{1}{z' - A}.$$

Let $A^{(k)}$ be the symmetric matrix with size $n - 1$ obtained by removing the k -th row and the k -th column of A and set $G^{(k)} := \frac{1}{z - A^{(k)}}$. Note that $\mathbb{E}_k G^{(k)} = \mathbb{E}_{k-1} G^{(k)}$, so that Y_k , which is equal to $\frac{1}{\sqrt{n}}(\mathbb{E}_k - \mathbb{E}_{k-1})(\text{Tr}(P_s G) - s_n \text{Tr} G)$, can be rewritten

$$Y_k = \frac{1}{\sqrt{n}}(\mathbb{E}_k - \mathbb{E}_{k-1})((\text{Tr}(P_s G) - \text{Tr}(P_s^{(k)} G^{(k)})) - s_n(\text{Tr} G - \text{Tr} G^{(k)}))$$

Then, (39) is obvious by Formula (68) of the appendix (indeed, $L^n(\varepsilon)$ is null for n large enough). Let us now apply Formula (67) of the appendix. We get

$$(41) \quad Y_k = \frac{1}{\sqrt{n}}(\mathbb{E}_k - \mathbb{E}_{k-1}) \left(\frac{1_{k \leq ns} - s_n + \mathbf{a}_k^* G^{(k)} (P_s^{(k)} - s_n) G^{(k)} \mathbf{a}_k}{z - a_{kk} - \mathbf{a}_k^* G^{(k)} \mathbf{a}_k} \right).$$

Then by Lemma 7.7 of [8], one can neglect the non diagonal terms in the expansions of the quadratic forms in (41), so that one can replace Y_k by $\frac{1}{\sqrt{n}}(\mathbb{E}_k - \mathbb{E}_{k-1})(f_k)$, with

$$(42) \quad f_k := f_k(z, s) = \frac{1_{k \leq ns} - s_n + \sum_j \mathbf{a}_k(j)^2 \{G^{(k)}(P_s^{(k)} - s_n)G^{(k)}\}_{jj}}{z - \sum_j \mathbf{a}_k(j)^2 G_{jj}^{(k)}}.$$

It follows that

$$(43) \quad \sum_{k=1}^n \mathbb{E}_{k-1}(Y_k Y_k') = \frac{1}{n} \sum_{k=1}^n \mathbb{E}_{k-1}[f_k f_k'] - \mathbb{E}_{k-1}[f_k] \mathbb{E}_{k-1}[f_k'] + o(1),$$

where f_k' is defined as f_k in (42), replacing the function s by s' and z by z' .

Let us denote by $\mathbb{E}_{\mathbf{a}_k}$ the expectation with respect to the randomness of the k -th column of A (*i.e.* the conditional expectation with respect to the σ -algebra generated by the a_{ij} 's such that $k \notin \{i, j\}$). Note that $\mathbb{E}_{k-1} = \mathbb{E}_{\mathbf{a}_k} \circ \mathbb{E}_k = \mathbb{E}_k \circ \mathbb{E}_{\mathbf{a}_k}$, hence

$$(44) \quad \mathbb{E}_{k-1}[f_k f_k'] - \mathbb{E}_{k-1}[f_k] \mathbb{E}_{k-1}[f_k'] = \mathbb{E}_{\mathbf{a}_k}[\mathbb{E}_k f_k \mathbb{E}_k f_k'] - \mathbb{E}_k \mathbb{E}_{\mathbf{a}_k} f_k \times \mathbb{E}_k \mathbb{E}_{\mathbf{a}_k} f_k'.$$

For each $s \in (0, 1)$ let us define \mathbb{C}_s^2 to be the set of pairs (z, \tilde{z}) of complex numbers such that

$$(\Im z > 0 \text{ and } -\frac{\Im z}{1-s} < \Im \tilde{z} < \frac{\Im z}{s}) \quad \text{or} \quad (\Im z < 0 \text{ and } \frac{\Im z}{s} < \Im \tilde{z} < -\frac{\Im z}{1-s}).$$

Note that \mathbb{C}_s^2 is the set of pairs (z, \tilde{z}) of complex numbers such that $\Im z \neq 0$ and both $\Im(z + (1-s)\tilde{z})$ and $\Im(z - s\tilde{z})$ have the same sign as $\Im z$.

Lemma 4.1. *For any fixed $z \in \mathbb{C} \setminus \mathbb{R}$ and any fixed $s \in (0, 1)$, as $n, k \rightarrow \infty$ in such a way that $k/n \rightarrow u \in [0, 1]$, we have the convergence in probability*

$$\lim_{N \rightarrow \infty} \mathbb{E}_{\mathbf{a}_k}[f_k(z, s)] = L_u(z, s) := - \int_0^{+\infty} \frac{1}{t} \partial_{\tilde{z}, \tilde{z}=0} e^{i \operatorname{sgn}_z t(z + \tilde{z}(1_{u \leq s-s}))} e^{\rho_{z, \tilde{z}, s}(t)} dt,$$

where for $s \in (0, 1)$ fixed, $(z, \tilde{z}, t) \mapsto \rho_{z, \tilde{z}, s}(t)$ is the unique function defined on $\mathbb{C}_s^2 \times \mathbb{R}_+$, analytic in its two first variables and continuous in its second one, taking values into $\{z \in \mathbb{C}; \Re z \leq 0\}$, solution of

$$\rho_{z, \tilde{z}, s}(t) = t \int_0^\infty g(ty) (s e^{iy \operatorname{sgn}_z \tilde{z}} + (1-s)) e^{iy \operatorname{sgn}_z (z - s\tilde{z})} e^{\rho_{z, \tilde{z}, s}(y)} dy$$

where g is the function introduced at (10).

Proof. We use the fact that for $z \in \mathbb{C} \setminus \mathbb{R}$,

$$(45) \quad \frac{1}{z} = -i \operatorname{sgn}_z \times \int_0^{+\infty} e^{\operatorname{sgn}_z itz} dt,$$

where sgn_z has been defined above by $\text{sgn}_z = \text{sgn}(\Im z)$. Hence by (42),

$$f_k = -i \text{sgn}_z \int_0^{+\infty} \{1_{k \leq ns} - s_n + \sum_j (G^{(k)}(P_s^{(k)} - s_n) G^{(k)})_{jj} \mathbf{a}_k(j)^2\} e^{i \text{sgn}_z t(z - \sum_j (G^{(k)})_{jj} \mathbf{a}_k(j)^2)} dt.$$

Let us define $G^{(k)}(\cdot, \cdot)$ on $\mathbb{C}_{s_n}^2$ by

$$G^{(k)}(z, \tilde{z}) := \frac{1}{z + \tilde{z}(P_s^{(k)} - s_n) - A^{(k)}}$$

(note that $G^{(k)}(\cdot, \cdot)$ is well defined the remark following the definition of \mathbb{C}_s^2). Then for any fixed $z \in \mathbb{C} \setminus \mathbb{R}$,

$$G_k(z)(P_s^{(k)} - s_n)G_k(z) = -\partial_{\tilde{z}, \tilde{z}=0} G^{(k)}(z, \tilde{z}).$$

Hence

$$\begin{aligned} & \{1_{k \leq ns} - s_n + \sum_j (G^{(k)}(P_s^{(k)} - s_n) G^{(k)})_{jj} \mathbf{a}_k(j)^2\} e^{i \text{sgn}_z t(z - \sum_j (G^{(k)})_{jj} \mathbf{a}_k(j)^2)} \\ &= \frac{1}{it \text{sgn}_z} \partial_{\tilde{z}, \tilde{z}=0} e^{i \text{sgn}_z t\{z + \tilde{z}(1_{k \leq ns} - s_n) - \sum_j (G^{(k)}(z, \tilde{z})_{jj} \mathbf{a}_k(j)^2\}} \end{aligned}$$

and

$$(46) \quad f_k = - \int_0^{+\infty} \frac{1}{t} \partial_{\tilde{z}, \tilde{z}=0} e^{i \text{sgn}_z t\{z + \tilde{z}(1_{k \leq ns} - s_n) - \sum_j (G^{(k)}(z, \tilde{z})_{jj} \mathbf{a}_k(j)^2\}} dt.$$

Thus for ϕ_n as defined as in (18) by $\phi_n(\lambda) = \mathbb{E} e^{-i\lambda a_{11}^2}$, we have (permuting \mathbb{E} and $\int_0^{+\infty}$ because $z \in \mathbb{C} \setminus \mathbb{R}$ is fixed and for each j , $-G^{(k)}(z, \tilde{z})_{jj}$ has imaginary part with the same sign as z for \tilde{z} small enough),

$$\mathbb{E}_{\mathbf{a}_k}(f_k) = - \int_0^{+\infty} \frac{1}{t} \partial_{\tilde{z}, \tilde{z}=0} e^{i \text{sgn}_z t(z + \tilde{z}(1_{k \leq ns} - s_n))} \prod_j \phi_n(\text{sgn}_z t G^{(k)}(z, \tilde{z})_{jj}) dt$$

Now, by (18), we have $n(\phi_n - 1) \rightarrow \Phi$ as $n \rightarrow \infty$. As $\Re(i \text{sgn}_z z) < 0$, the integrals are well dominated at infinity and the integral

$$\int_0^{+\infty} \frac{1}{t} \partial_{\tilde{z}, \tilde{z}=0} e^{i \text{sgn}_z t(z + \tilde{z}(1_{k \leq ns} - s_n))} e^{\frac{1}{n} \sum_j \Phi(\text{sgn}_z t G^{(k)}(z, \tilde{z})_{jj})} dt$$

is well converging at the origin as the derivative in \tilde{z} is of order t since Φ is analytic on \mathbb{C}^- where $G^{(k)}(z, \tilde{z})_{jj}$ almost surely takes its values, while being uniformly bounded, hence

$$\mathbb{E}_{\mathbf{a}_k}(f_k) = o(1) - \int_0^{+\infty} \frac{1}{t} \partial_{\tilde{z}, \tilde{z}=0} e^{i \text{sgn}_z t(z + \tilde{z}(1_{k \leq ns} - s_n))} e^{\frac{1}{n} \sum_j \Phi(\text{sgn}_z t G^{(k)}(z, \tilde{z})_{jj})} dt.$$

We therefore basically need to compute the asymptotics of

$$\rho_{z, \tilde{z}, s}^n(t) := \frac{1}{n} \sum_j \Phi(\text{sgn}_z t G^{(k)}(z, \tilde{z})_{jj}).$$

Note that by definition of Φ , for any $\lambda \in \overline{\mathbb{C}^-}$, $\Re \Phi(\lambda) \leq 0$. Thus $\rho_{z, \tilde{z}, s}^n(t)$ is analytic in $z \in \mathbb{C} \setminus \mathbb{R}$, and uniformly bounded on compact subsets of $\mathbb{C} \setminus \mathbb{R}$ and takes values in

$\{z \in \mathbb{C}; \Re z \leq 0\}$. By Montel's theorem, all limit points of this function for uniform convergence on compact subsets will satisfy the same property. Now, notice that by Schur complement formula and the usual removal of the non diagonal terms (Lemma 7.7 of [8] again), that for $n \gg 1$,

$$G^{(k)}(z, \tilde{z})_{jj} \simeq \frac{1}{z + \tilde{z}(1_{k \leq ns} - s_n) - \sum_{\ell} a_{\ell\ell}^2 G^{(k,j)}(z, \tilde{z})_{\ell\ell}}$$

where $G^{(k,j)}$ is the resolvent where two rows and columns have been suppressed. We can now proceed as usual to write that by invariance of the law of A by conjugation by permutation matrices, for all j ,

$$\mathbb{E}[\Phi(\operatorname{sgn}_z tG^{(k)}(z, \tilde{z})_{jj})] = \begin{cases} \mathbb{E}[\Phi(\operatorname{sgn}_z tG^{(k)}(z, \tilde{z})_{11})] & \text{if } j \leq ns, \\ \mathbb{E}[\Phi(\operatorname{sgn}_z tG^{(k)}(z, \tilde{z})_{nn})] & \text{if } j > ns, \end{cases}$$

so that by concentration arguments, see [8, Appendix], $\rho_{z, \tilde{z}, s}^n(t)$ self-averages so that for $n \gg 1$,

$$\begin{aligned} \rho_{z, \tilde{z}, s}^n(t) &\simeq \mathbb{E}\left[\frac{1}{n} \sum_j \Phi(\operatorname{sgn}_z tG^{(k)}(z, \tilde{z})_{jj})\right] \\ &= s \mathbb{E}[\Phi(\operatorname{sgn}_z tG^{(k)}(z, \tilde{z})_{11})] + (1-s) \mathbb{E}[\Phi(\operatorname{sgn}_z tG^{(k)}(z, \tilde{z})_{nn})]. \end{aligned}$$

On the other side, with the function g introduced in the hypothesis at (10), we have

$$\begin{aligned} \mathbb{E}[\Phi(\operatorname{sgn}_z tG^{(k)}(z, \tilde{z})_{11})] &= t \int_0^\infty g(ty) e^{iy \operatorname{sgn}_z (z + \tilde{z}(1-s_n))} \prod_j \phi_n(y \operatorname{sgn}_z G^{(k,1)}(z, \tilde{z})_{jj}) dy \\ &\sim t \int_0^\infty g(ty) e^{iy \operatorname{sgn}_z (z + \tilde{z}(1-s_n))} e^{\rho_{z, \tilde{z}, s}^n(y)} dy \\ \mathbb{E}[\Phi(\operatorname{sgn}_z tG^{(k)}(z, \tilde{z})_{nn})] &= t \int_0^\infty g(ty) e^{iy \operatorname{sgn}_z (z - s_n \tilde{z})} \prod_j \phi_n(\operatorname{sgn}_z y G^{(k,1)}(z, \tilde{z})_{jj}) dy \\ &\sim t \int_0^\infty g(ty) e^{iy \operatorname{sgn}_z (z - s_n \tilde{z})} e^{\rho_{z, \tilde{z}, s}^n(y)} dy \end{aligned}$$

so that we deduce that the limit points $\rho_{z, \tilde{z}, s}(t)$ of $\rho_{z, \tilde{z}, s}^n(t)$ satisfy

$$\rho_{z, \tilde{z}, s}(t) = t \int_0^\infty g(ty) (s e^{iy \operatorname{sgn}_z \tilde{z}} + (1-s)) e^{iy \operatorname{sgn}_z (z - s \tilde{z})} e^{\rho_{z, \tilde{z}, s}(y)} dy.$$

Let us now prove that for each fixed $s \in (0, 1)$, there exists a unique function satisfying this equation and the conditions stated in the lemma. So let us suppose that we have two solutions $\rho_{z, \tilde{z}, s}(t)$ and $\tilde{\rho}_{z, \tilde{z}, s}(t)$. Then

$$\Delta_{z, \tilde{z}}(t) := \rho_{z, \tilde{z}, s}(t) - \tilde{\rho}_{z, \tilde{z}, s}(t)$$

satisfies

$$\Delta_{z,\tilde{z}}(t) = t \int_0^\infty g(ty) (s e^{iy \operatorname{sgn}_z \tilde{z}} + (1-s)) e^{iy \operatorname{sgn}_z (z-s\tilde{z})} (e^{\rho_{z,\tilde{z},s}(y)} - e^{\tilde{\rho}_{z,\tilde{z},s}(y)}) dy,$$

hence for

$$\delta(z, \tilde{z}) := \min\{\operatorname{sgn}_z \Im(z + (1-s)\tilde{z}), \operatorname{sgn}_z \Im(z - s\tilde{z})\} > 0,$$

we have

$$|\Delta_{z,\tilde{z}}(t)| \leq t \int_0^\infty |g(ty)| e^{-\delta(z,\tilde{z})y} |\Delta_{z,\tilde{z}}(y)| dy$$

Thus by the hypothesis made on g at (11),

$$|\Delta_{z,\tilde{z}}(t)| \leq \underbrace{K t^{\gamma+1} \int_0^\infty y^\gamma e^{-\delta(z,\tilde{z})y} |\Delta_{z,\tilde{z}}(y)| dy}_{:= I_1(z,\tilde{z})} + \underbrace{K t^{\kappa+1} \int_0^\infty y^\kappa e^{-\delta(z,\tilde{z})y} |\Delta_{z,\tilde{z}}(y)| dy}_{:= I_2(z,\tilde{z})}$$

It follows that the numbers $I_1(z, \tilde{z})$ and $I_2(z, \tilde{z})$ defined above satisfy

$$\begin{aligned} I_1(z, \tilde{z}) &\leq K \left(I_1(z, \tilde{z}) \int_0^\infty y^{2\gamma+1} e^{-\delta(z,\tilde{z})y} dy + I_2(z, \tilde{z}) \int_0^\infty y^{\gamma+\kappa+1} e^{-\delta(z,\tilde{z})y} dy \right), \\ I_2(z, \tilde{z}) &\leq K \left(I_1(z, \tilde{z}) \int_0^\infty y^{\gamma+\kappa+1} e^{-\delta(z,\tilde{z})y} dy + I_2(z, \tilde{z}) \int_0^\infty y^{2\kappa+1} e^{-\delta(z,\tilde{z})y} dy \right). \end{aligned}$$

For $\delta(z, \tilde{z})$ large enough, the integrals above are all strictly less than $\frac{1}{4K}$, so that $I_1(z, \tilde{z}) = I_2(z, \tilde{z}) = 0$. It follows that for $\Im z$ large enough and $\Im \tilde{z}$ small enough, both solutions coincide. By analytic continuation, unicity follows. \square

Getting back to (43) and (44), we shall now, as in [8], analyze

$$L_k^n(s, z; s', z') := \mathbb{E}_{k-1}[f_k f'_k].$$

Let us first define the measure $\tilde{\tau} := \tau + \delta_0 \otimes \mu + \mu \otimes \delta_0$ on $(\mathbb{R}^+)^2$ for τ and μ the measures introduced at (13) or at Remark 2.1. We always have, for $x, y \in \mathbb{C}^+$,

$$(47) \quad \Phi(x^{-1} + y^{-1}) = \iint_{(\mathbb{R}_+)^2} e^{i(xv+yv')} d\tilde{\tau}(v, v')$$

Lemma 4.2. *Let us fix $s_1, s_2 \in (0, 1)$. As $k, n \rightarrow \infty$ in such a way that k/n goes to $u \in [0, 1]$, $L_k^n(s_1, z; s_2, z')$ tends to*

$$L_u(s_1, z; s_2, z') :=$$

$$\iint_{\mathbb{R}_+^2} \partial_{\tilde{z}, \tilde{z}=0} \partial_{\tilde{z}', \tilde{z}'=0} e^{i \operatorname{sgn}_z t(z+\tilde{z}(1_{u \leq s_1}-s_1)) + i \operatorname{sgn}_{z'} t'(z'+\tilde{z}'(1_{u \leq s_2}-s_2)) + \rho_u(s_1, t, z, \tilde{z}; s_2, t', z', \tilde{z}')} \frac{dt dt'}{tt'}$$

where the function

$$(t, z, \tilde{z}, t', z', \tilde{z}') \in \mathbb{R}_+ \times \mathbb{C}_{s_1}^2 \times \mathbb{R}_+ \times \mathbb{C}_{s_2}^2 \mapsto \rho_u(s_1, t, z, \tilde{z}; s_2, t', z', \tilde{z}')$$

is characterized as follows : $\rho_u(s_1, t, z, \tilde{z}; s_2, t', z', \tilde{z}') = \rho_u(s_2, t', z', \tilde{z}'; s_1, t, z, \tilde{z})$ and if, for example, $s_1 \leq s_2$, then for $\gamma_1 = s_1, \gamma_2 = s_2 - s_1, \gamma_3 = 1 - s_2$,

$$\rho_u(s_1, t, z, \tilde{z}; s_2, t', z', \tilde{z}') = \sum_{\beta=1}^3 \gamma_\beta \rho_u^\beta(s_1, t, z, \tilde{z}; s_2, t', z', \tilde{z}')$$

where the functions

$$(t, z, \tilde{z}, t', z', \tilde{z}') \in \mathbb{R}_+ \times \mathbb{C}_{s_1}^2 \times \mathbb{R}_+ \times \mathbb{C}_{s_2}^2 \longmapsto \rho_u^\beta(s_1, t, z, \tilde{z}; s_2, t', z', \tilde{z}') \quad (\beta = 1, 2, 3)$$

are the unique analytic in $z, \tilde{z}, z', \tilde{z}'$ and continuous in t and in t' functions such that

$$\begin{aligned} \rho_u^\beta(s, t_1, z_1, \tilde{z}_1; s_2, t_2, z_2, \tilde{z}_2) &= \iint e^{\sum_{r=1,2} \text{sgn}_{z_r} \frac{iv_r}{t_r} (z_r + \tilde{z}_r (1_{\beta \leq r} - s_r)) + u} \mathbf{1}_{u > s_\beta} \rho_u(z_1, \tilde{z}_1, s_1, \frac{v_1}{t_1}; z_2, \tilde{z}_2, s_2, \frac{v_2}{t_2}) \\ &\quad e^{(1-u) \mathbf{1}_{u > s_\beta} \rho_{z_1, \tilde{z}_1, s_1}(v_1/t_1) + (1-u) \mathbf{1}_{u > s_\beta} \rho_{z_2, \tilde{z}_2, s_2}(v_2/t_2)} d\tilde{\tau}(v_1, v_2). \end{aligned}$$

Proof. Of course, $L_k^n(s_1, z; s_2, z') = L_k^n(s_2, z'; s_1, z)$. Let us suppose for example that $s_1 \leq s_2$. For short, we write $s := s_1$ and $s' := s_2$. Note that

$$L_k^n(s, z; s', z') := \mathbb{E}_k[\mathbb{E}_{\mathbf{a}_k}[f_k \tilde{f}_k]]$$

where f_k'' is defined as f_k' replacing the matrix A by the matrix

$$(48) \quad A' = [a'_{ij}]_{1 \leq i, j \leq N}$$

defined by the the fact that the a'_{ij} 's such that $i > k$ and $j > k$ are i.i.d. copies of a_{11} (modulo the fact that A' is symmetric), independent of A , for all other pairs (i, j) , $a'_{ij} = a_{ij}$. We define, for $(z', \tilde{z}) \in \mathbb{C}_{s_n}^2$,

$$G'^{(k)}(z', \tilde{z}) := \frac{1}{z' + \tilde{z}(P_{s'}^{(k)} - s'_n) - A'^{(k)}}.$$

First recall the following formula for f_k established at Equation (46):

$$f_k = - \int_0^{+\infty} \frac{1}{t} \partial_{\tilde{z}, \tilde{z}=0} e^{i \text{sgn}_z t \{z + \tilde{z}(1_{k \leq n} s - s_n) - \sum_j G^{(k)}(z, \tilde{z})_{jj} \mathbf{a}_k(j)^2\}} dt.$$

In the same way, we find

$$f_k'' = - \int_0^{+\infty} \frac{1}{t'} \partial_{\tilde{z}, \tilde{z}=0} e^{i \text{sgn}_{z'} t' \{z' + \tilde{z}(1_{k \leq n} s' - s'_n) - \sum_j G'^{(k)}(z', \tilde{z})_{jj} \mathbf{a}_k(j)^2\}} dt.$$

Hence we have

$$\begin{aligned}
L_k^n(s, z; s', z') &= \iint_{\mathbb{R}_+^2} \partial_{\tilde{z}, \tilde{z}=0} \partial_{z', \tilde{z}'=0} e^{i \operatorname{sgn}_z t \{z + \tilde{z}(1_{k \leq ns - s_n})\} + i \operatorname{sgn}_{z'} t' \{z' + \tilde{z}'(1_{k \leq ns' - s'_n})\}} \times \\
&\quad \prod_j \phi_n(\operatorname{sgn}_z t G^{(k)}(z, \tilde{z})_{jj} + \operatorname{sgn}_{z'} t' G'^{(k)}(z', \tilde{z}')_{jj}) \frac{dt dt'}{tt'} \\
&\simeq \iint_{\mathbb{R}_+^2} \partial_{\tilde{z}, \tilde{z}=0} \partial_{z', \tilde{z}'=0} e^{i \operatorname{sgn}_z t \{z + \tilde{z}(1_{k \leq ns - s_n})\} + i \operatorname{sgn}_{z'} t' \{z' + \tilde{z}'(1_{k \leq ns' - s'_n})\}} \times \\
&\quad \exp(\rho_k^n(s, t, z, \tilde{z}; s', t', z', \tilde{z}')) \frac{dt dt'}{tt'}
\end{aligned}$$

with

$$\rho_k^n(s, t, z, \tilde{z}; s', t', z', \tilde{z}') := \frac{1}{n} \sum_j \Phi(\operatorname{sgn}_z t G^{(k)}(z, \tilde{z})_{jj} + \operatorname{sgn}_{z'} t' G'^{(k)}(z', \tilde{z}')_{jj}).$$

We can derive the asymptotics of this term as before thanks to (47). We have to separate the j 's according to whether $j \leq k$ or $j > k$ and $k \geq sn$ or $k \geq s'n$. Assume $j \leq k$ so that $a'_{j\ell} = a_{j\ell}$ for all ℓ . Then

$$\begin{aligned}
&\Phi(\operatorname{sgn}_z t G^{(k)}(z, \tilde{z})_{jj} + \operatorname{sgn}_{z'} t' G'^{(k)}(z', \tilde{z}')_{jj}) \\
&\simeq \iint_{\mathbb{R}_+^2} e^{\operatorname{sgn}_z \frac{iv}{t} \{z + \tilde{z}(1_{j \leq ns - s}) - \sum_\ell a_{\ell\ell}^2 G^{(k,j)}(z, \tilde{z})_{\ell\ell}\}} e^{\operatorname{sgn}_{z'} \frac{iv'}{t'} \{z' + \tilde{z}'(1_{j \leq ns' - s'}) - \sum_\ell a_{\ell\ell}^2 G'^{(k,j)}(z', \tilde{z}')_{\ell\ell}\}} d\tilde{\tau}(v, v')
\end{aligned}$$

thus using the concentration around the expectation, we get

$$\begin{aligned}
&\Phi(\operatorname{sgn}_z t G^{(k)}(z, \tilde{z})_{jj} + \operatorname{sgn}_{z'} t' G'^{(k)}(z', \tilde{z}')_{jj}) \\
&\simeq \iint_{\mathbb{R}_+^2} e^{\operatorname{sgn}_z \frac{iv}{t} \{z + \tilde{z}(1_{j \leq ns - s})\} + \operatorname{sgn}_{z'} \frac{iv'}{t'} \{z' + \tilde{z}'(1_{j \leq ns' - s'})\}} e^{\rho_k^n(s, \frac{v}{t}, z, \tilde{z}, s', \frac{v'}{t'}, z', \tilde{z}')} d\tilde{\tau}(v, v')
\end{aligned}$$

Assume now that $j > k$ so that $a_{\ell\ell}$ and $a'_{\ell\ell}$ are independent if $\ell \geq k$. Then

$$\begin{aligned}
&\Phi(\operatorname{sgn}_z t G^{(k)}(z, \tilde{z})_{jj} + \operatorname{sgn}_{z'} t' G'^{(k)}(z', \tilde{z}')_{jj}) \\
&\simeq \iint_{\mathbb{R}_+^2} e^{\operatorname{sgn}_z \frac{iv}{t} \{z + \tilde{z}(1_{j \leq ns - s}) - \sum_\ell a_{\ell\ell}^2 G^{(k,j)}(z, \tilde{z})_{\ell\ell}\}} e^{\operatorname{sgn}_{z'} \frac{iv'}{t'} \{z' + \tilde{z}'(1_{j \leq ns' - s'}) - \sum_\ell a'_{\ell\ell}{}^2 G'^{(k,j)}(z', \tilde{z}')_{\ell\ell}\}} d\tilde{\tau}(v, v')
\end{aligned}$$

and using again the concentration around the expectation, we get

$$\begin{aligned}
&\Phi(\operatorname{sgn}_z t G^{(k)}(z, \tilde{z})_{jj} + \operatorname{sgn}_{z'} t' G'^{(k)}(z', \tilde{z}')_{jj}) \\
&\simeq \iint_{\mathbb{R}_+^2} e^{\operatorname{sgn}_z \frac{iv}{t} \{z + \tilde{z}(1_{j \leq ns - s})\} + \operatorname{sgn}_{z'} \frac{iv'}{t'} \{z' + \tilde{z}'(1_{j \leq ns' - s'})\}} \times \\
&\quad e^{\frac{k}{n} \rho_k^n(s, \frac{v}{t}, z, \tilde{z}, s', \frac{v'}{t'}, z', \tilde{z}') + (1 - \frac{k}{n}) \{\rho_{z, \tilde{z}, s}(\frac{v}{t}) + \rho_{z', \tilde{z}', s'}(\frac{v'}{t'})\}} d\tilde{\tau}(v, v')
\end{aligned}$$

From these equations, we see that, under our hypothesis that $s_1 \leq s_2$, we have 3 possible regimes, each of them giving rise to a limit point and an equation. Hence the whole system is governed by 3 equations. The proof of the fact that under analyticity hypotheses, the limit points are uniquely described by the resulting equations goes along the same lines as the proofs in Section 5.2 of [8], sketched as follows. First, we have to consider separately the case where Φ satisfies (12) and the case where Φ satisfies (13). In the case where Φ satisfies (12), the proof is very similar to the proof of the corresponding case in Section 5.2 of [8] and to the detailed proof of the uniqueness for Lemma 4.1 of the present paper, using (14) instead of (11). The case where Φ satisfies (13) is a little more delicate. As in Lemma 5.1 of [8], one first needs to notice that considered as functions of t, t', t'' , the limit points satisfy an Hölder bound, using essentially the facts that for any $2\kappa \in (0, \alpha/2)$

$$(49) \quad \limsup_{N \geq 1} \mathbb{E} \left[\left(\sum |a_{1i}|^2 \right)^{2\kappa} \right] < \infty,$$

and that for any $\beta \in (\alpha/1, 1]$, there exists a constant $c = c(\alpha, \beta)$ such that for any x, y in \mathbb{C}^- ,

$$(50) \quad |x^{\frac{\alpha}{2}} - y^{\frac{\alpha}{2}}| \leq c|x - y|^\beta (|x| \wedge |y|)^{\frac{\alpha}{2} - \beta}.$$

Then one has to interpret the equation relating the limit points associated to the 3 possible regimes as a fixed point equation for a strictly contracting function in a space of Hölder functions: the key argument, to prove that the function is contracting, is to use the estimates given in Lemma 5.7 of [12]. \square

This concludes the proof of Proposition 2.8 and it follows from this that the covariance of the process $H_{s,z}$ is given by

$$(51) \quad C(s, z; s', z') := \mathbb{E}[H_{s,z}H_{s',z'}] = \int_0^1 du(L_u(s, z; s', z') - L_u(z, s)L_u(z', s'))$$

with the functions L defined in Lemmas 4.1 and 4.2.

5. PROOF OF PROPOSITION 2.6

Let us now prove that the limit covariance of $(C_{s,\lambda})$ is not identically zero (hence this is also the case for $(B_{s,t})$ by (28)). Using Lemma 6.1 and (34), one easily sees that $(C_{s,\lambda}^n)$ is uniformly bounded in L^4 . It follows that

$$(52) \quad \text{Var}(C_{s,\lambda}) = \mathbb{E}[(C_{s,\lambda})^2] = \lim_{n \rightarrow \infty} \mathbb{E}[(C_{s,\lambda}^n)^2].$$

Thus we shall prove that the limit of $\mathbb{E}[(C_{s,\lambda}^n)^2]$ is not identically zero.

First note that for $x_1, \dots, x_n \in \mathbb{C}$ such that $x_1 + \dots + x_n = 0$, for any $0 \leq \ell \leq n$, we have

$$(53) \quad \sum_{i=1}^{\ell} x_i = \sum_{i=1}^n \alpha_i x_i \quad \text{for } \alpha_i := \begin{cases} 1 - \frac{\ell}{n} & \text{if } i \leq \ell, \\ -\frac{\ell}{n} & \text{if } i > \ell, \end{cases}$$

and that $\alpha_1 + \dots + \alpha_n = 0$. Note also that for (X_1, \dots, X_n) an exchangeable random vector and $\alpha_1, \dots, \alpha_n \in \mathbb{C}$ such that $\alpha_1 + \dots + \alpha_n = 0$, we have

$$(54) \quad \mathbb{E} \sum_{i, i'} \alpha_i \alpha_{i'} X_i X_{i'} = \sum_i \alpha_i^2 \mathbb{E}[X_1(X_1 - X_2)].$$

It follows easily from (53) and (54) that if we also suppose that almost surely, the X_i 's sum up to zero, then for any $0 \leq \ell \leq n$, we have

$$(55) \quad \mathbb{E} \sum_{\substack{1 \leq i \leq \ell \\ 1 \leq i' \leq \ell}} X_i X_{i'} = n \left(\frac{\ell}{n} - \frac{\ell^2}{n^2} \right) \mathbb{E}[X_1(X_1 - X_2)].$$

Let us now fix $s \in (0, 1)$ and $\lambda \in \mathbb{R}$ and apply (55) with $X_i = \sum_{j; \lambda_j \leq \lambda} (|u_{ij}|^2 - n^{-1})$ and $\ell = \lfloor ns \rfloor$. For $\tilde{s} := \lfloor ns \rfloor / n$, we get

$$(56) \quad \text{Var}(C_{s, \lambda}^n) = \mathbb{E}[(C_{s, \lambda}^n)^2] = (\tilde{s} - \tilde{s}^2) \mathbb{E}[X_1(X_1 - X_2)]$$

Note also that as each $|u_{ij}|^2$ has expectation n^{-1} ,

$$\mathbb{E}[X_1(X_1 - X_2)] = \mathbb{E} \left[\sum_{\substack{j; \lambda_j \leq \lambda \\ j'; \lambda_{j'} \leq \lambda}} |u_{1j}|^2 |u_{1j'}|^2 - |u_{1j}|^2 |u_{2j'}|^2 \right]$$

Moreover, by exchangeability of the rows of U (which is true even conditionally to the λ_j 's) and the fact that its columns have norm one, we easily see that for any j, j' ,

$$n(n-1) \mathbb{E}[1_{\lambda_j, \lambda_{j'} \leq \lambda} |u_{1j}|^2 |u_{2j'}|^2] + n \mathbb{E}[1_{\lambda_j, \lambda_{j'} \leq \lambda} |u_{1j}|^2 |u_{1j'}|^2] = 1,$$

so that

$$\mathbb{E}[X_1(X_1 - X_2)] = O\left(\frac{1}{n}\right) + \left(1 - \frac{1}{n}\right) \mathbb{E} \left[\sum_{j, j'; \lambda_j, \lambda_{j'} \leq \lambda} (|u_{1j}|^2 |u_{1j'}|^2 - n^{-2}) \right].$$

By (56), we deduce that

$$\mathbb{E}[(C_{s, \lambda}^n)^2] = O\left(\frac{1}{n}\right) + \left(1 - \frac{1}{n}\right) (\tilde{s} - \tilde{s}^2) \mathbb{E} \left[\sum_{j, j'; \lambda_j, \lambda_{j'} \leq \lambda} (|u_{1j}|^2 |u_{1j'}|^2 - n^{-2}) \right],$$

so that

$$(57) \quad \mathbb{E}[(C_{s, \lambda}^n)^2] = O\left(\frac{1}{n}\right) + (\tilde{s} - \tilde{s}^2) \mathbb{E} \left[\left(\sum_{j; \lambda_j \leq \lambda} (|u_{1j}|^2 - n^{-1}) \right)^2 \right]$$

Note that

$$\sum_{j; \lambda_j \leq \lambda} |u_{1j}|^2 = \mu_{n, \mathbf{e}_1}((-\infty, \lambda])$$

where $\mu_{n,\mathbf{e}_1} := \sum_{j=1}^n |u_{1j}|^2 \delta_{\lambda_j}$ is the empirical spectral law of A according to the first vector \mathbf{e}_1 of the canonical basis, also defined by the fact that for any test function f ,

$$\int f(x) d\mu_{n,\mathbf{e}_1}(x) = (f(A))_{11}.$$

Thus for μ_n defined by (25), we have

$$\sum_{j: \lambda_j \leq \lambda} (|u_{1j}|^2 - n^{-1}) = (\mu_{n,\mathbf{e}_1} - \mu_n)((-\infty, \lambda])$$

Hence by (57),

$$(58) \quad \mathbb{E}[(C_{s,\lambda}^n)^2] = O\left(\frac{1}{n}\right) + (\tilde{s} - \tilde{s}^2) \mathbb{E}[\{(\mu_{n,\mathbf{e}_1} - \mu_n)((-\infty, \lambda])\}^2].$$

Let us now suppose that for a certain $s \in (0, 1)$, we have $\text{Var}(C_{s,\lambda}) = 0$ for all $\lambda \in \mathbb{R}$. To conclude the proof, we shall now exhibit a contradiction. By (52) and (58), we know that for all λ , $\mathbb{E}[\{(\mu_{n,\mathbf{e}_1} - \mu_n)((-\infty, \lambda])\}^2] \rightarrow 0$ as $n \rightarrow \infty$. Thus by dominated convergence, as $n \rightarrow \infty$,

$$\int_{\lambda \in \mathbb{R}} \frac{\|(\mu_{n,\mathbf{e}_1} - \mu_n)((-\infty, \lambda])\|_{L^2}}{|z - \lambda|^2} d\lambda \rightarrow 0$$

and by the triangular inequality, we deduce that as $n \rightarrow \infty$,

$$\int_{\lambda \in \mathbb{R}} \frac{(\mu_{n,\mathbf{e}_1} - \mu_n)((-\infty, \lambda])}{(z - \lambda)^2} d\lambda \xrightarrow{L^2} 0.$$

But by (38), for any $z \in \mathbb{C} \setminus \mathbb{R}$, with the notation $G(z) := (z - A)^{-1}$,

$$\begin{aligned} \int_{\lambda \in \mathbb{R}} \frac{(\mu_{n,\mathbf{e}_1} - \mu_n)((-\infty, \lambda])}{(z - \lambda)^2} d\lambda &= - \int_{\lambda \in \mathbb{R}} \frac{d(\mu_{n,\mathbf{e}_1} - \mu_n)(\lambda)}{z - \lambda} \\ &= \frac{1}{n} \text{Tr } G(z) - G(z)_{11}. \end{aligned}$$

We deduce that as $n \rightarrow \infty$,

$$(59) \quad \frac{1}{n} \text{Tr } G(z) - G(z)_{11} \xrightarrow{L^2} 0.$$

Note that by exchangeability of the rows of A , the LHT of (59) is centered, hence converges in probability to zero. By (26), we deduce that $G(z)_{11}$ converges in probability to the Stieltjes transform $G_{\mu_\Phi}(z)$ of the limit empirical spectral law μ_Φ of A . By the Schur complement formula (see [1, Lem. 2.4.6]) and the asymptotic vanishing of non diagonal terms in the quadratic form (Lemma 7.7 of [8]), we deduce that, for $A^{(1)}$ the matrix obtained after having removed the first row and the first column to A and $G^{(1)}(z) := (z - A^{(1)})^{-1}$,

$$(60) \quad z - \sum_{j=2}^n |a_{1j}|^2 G^{(1)}(z)_{jj}$$

converges in probability to $1/G_{\mu_\Phi}(z)$.

But one can easily deduce from our hypotheses that the random variable of (60) converges in distribution to a random variable which is non constant (because $\Phi(z)$ is not identically equal to z).

6. APPENDIX

6.1. A tightness lemma for bivariate processes. Let us endow the space $D([0, 1]^2)$ with the Skorokhod topology (see [6] for the definitions).

Let $M = [m_{ij}]_{1 \leq i, j \leq n}$ be a random bistochastic matrix depending implicitly on n . We define the random process of $D([0, 1]^2)$

$$S^n(s, t) := \frac{1}{\sqrt{n}} \sum_{\substack{1 \leq i \leq ns \\ 1 \leq j \leq nt}} \left(m_{ij} - \frac{1}{n} \right).$$

Lemma 6.1. *Let us suppose that M is, in law, invariant under left multiplication by any permutation matrix. Then the process S^n is C -tight in $D([0, 1]^2)$, i.e. the sequence $(\text{distribution}(S^n))_{n \geq 1}$ is tight and has all its accumulation points supported by the set of continuous functions on $[0, 1]^2$. Moreover, the process S^n is uniformly bounded in L^4 .*

Proof. Let us prove that for all $0 \leq s < s' \leq 1$, $0 \leq t < t' \leq 1$,

$$(61) \quad \mathbb{E}[(\Delta_{s, s', t, t'} S^n)^4] \leq \frac{7}{n} + 6(s' - s)^2(t' - t)^2(1 - (s' - s))^2,$$

where $\Delta_{s, s', t, t'} S^n$ denotes the increment of S^n on $[s, s'] \times [t, t']$, i.e.

$$(62) \quad \Delta_{s, s', t, t'} S^n := \frac{1}{\sqrt{n}} \sum_{\substack{ns < i \leq ns' \\ nt \leq j \leq nt'}} \left(m_{ij} - \frac{1}{n} \right).$$

As S^n vanishes on the boundary on $[0, 1]^2$, according to [9, Th. 3], (61) will imply the lemma.

To prove (61), we fix $0 \leq s < s' \leq 1$, $0 \leq t < t' \leq 1$. Let us now introduce some notation (where the dependence on n will be dropped for readability). We define the sets

$$I := \{i \in \{1, \dots, n\}; ns < i \leq ns'\} \text{ and } J := \{j \in \{1, \dots, n\}; nt < j \leq nt'\},$$

the numbers $(\alpha_i)_{1 \leq i \leq n}$ defined by

$$\alpha_i := \begin{cases} -\frac{1}{\sqrt{n}} \frac{|I|}{n} & \text{if } i \notin I \\ \frac{1}{\sqrt{n}} \left(1 - \frac{|I|}{n} \right) & \text{if } i \in I \end{cases}$$

and the exchangeable random vector (implicitly depending on n) (X_1, \dots, X_n) defined by

$$X_i = \sum_{j \in J} m_{ij}.$$

Note that

$$\Delta_{s,s',t,t'} S^n = \frac{1}{\sqrt{n}} \left(\left(\sum_{i \in I} X_i \right) - \frac{|I||J|}{n} \right)$$

and that as columns of M sum up to one, $|J| = \sum_{j \in J} \sum_{i=1}^n m_{ij} = \sum_{i=1}^n X_i$, hence

$$\Delta_{s,s',t,t'} S^n = \frac{1}{\sqrt{n}} \left(\sum_{i \in I} X_i - \frac{|I|}{n} \sum_{i=1}^n X_i \right) = \sum_{i=1}^n \alpha_i X_i.$$

Thus by exchangeability of the X_i 's, we have

$$\begin{aligned} \mathbb{E}[(\Delta_{s,s',t,t'} S^n)^4] &= \mathbb{E}(X_1^4) \text{sum}_4(\alpha) + 4 \mathbb{E}(X_1^3 X_2) \text{sum}_{3,1}(\alpha) + 3 \mathbb{E}(X_1^2 X_2^2) \text{sum}_{2,2}(\alpha) \\ &\quad + 6 \mathbb{E}(X_1^2 X_2 X_3) \text{sum}_{2,1,1}(\alpha) + \mathbb{E}(X_1 X_2 X_3 X_4) \text{sum}_{1,1,1,1}(\alpha), \end{aligned}$$

with

$$\begin{aligned} \text{sum}_4(\alpha) &:= \sum_{i=1}^n \alpha_i^4, \quad \text{sum}_{3,1}(\alpha) := \sum_{i \neq j} \alpha_i^3 \alpha_j, \quad \text{sum}_{2,2}(\alpha) := \sum_{i \neq j} \alpha_i^2 \alpha_j^2, \\ \text{sum}_{2,1,1}(\alpha) &:= \sum_{\substack{i,j,k \\ \text{pairwise } \neq}} \alpha_i^2 \alpha_j \alpha_k, \quad \text{sum}_{1,1,1,1}(\alpha) := \sum_{\substack{i,j,k,\ell \\ \text{pairwise } \neq}} \alpha_i \alpha_j \alpha_k \alpha_\ell. \end{aligned}$$

As the α_i 's sum up to zero, we have

$$\begin{aligned} \text{sum}_{3,1}(\alpha) &= \sum_i (\alpha_i^3 \sum_{j \neq i} \alpha_j) = -\text{sum}_4(\alpha), \\ \text{sum}_{2,1,1}(\alpha) &= \sum_i (\alpha_i^2 \sum_{j \neq i} (\alpha_j \sum_{k \notin \{i,j\}} \alpha_k)) = \sum_i (\alpha_i^2 \sum_{j \neq i} (\alpha_j (-\alpha_i - \alpha_j))) \\ &= -\text{sum}_{3,1}(\alpha) - \text{sum}_{2,2}(\alpha) = \text{sum}_4(\alpha) - \text{sum}_{2,2}(\alpha) \\ \text{sum}_{1,1,1,1}(\alpha) &= -3 \text{sum}_{2,1,1}(\alpha) = 3 \text{sum}_{2,2}(\alpha) - 3 \text{sum}_4(\alpha) \end{aligned}$$

Thus

$$\begin{aligned} \mathbb{E}[(\Delta_{s,s',t,t'} S^n)^4] &= \text{sum}_4(\alpha) (\mathbb{E}(X_1^4) - 4 \mathbb{E}(X_1^3 X_2) + 6 \mathbb{E}(X_1^2 X_2 X_3) - 3 \mathbb{E}(X_1 X_2 X_3 X_4)) \\ &\quad + \text{sum}_{2,2}(\alpha) (3 \mathbb{E}(X_1^2 X_2^2) - 6 \mathbb{E}(X_1^2 X_2 X_3) + 3 \mathbb{E}(X_1 X_2 X_3 X_4)), \end{aligned}$$

Now, as for all i , $|\alpha_i| \leq \frac{1}{\sqrt{n}}$, we have $\text{sum}_4(\alpha) \leq \frac{1}{n}$, and as for all i , $0 \leq X_i \leq 1$ (because the rows of s sum up to one), we have

$$\mathbb{E}[(\Delta_{s,s',t,t'} S^n)^4] \leq \frac{7}{n} + 3 \text{sum}_{2,2}(\alpha) (\mathbb{E}(X_1^2 X_2^2) + \mathbb{E}(X_1 X_2 X_3 X_4)).$$

To conclude the proof of (61), we shall prove that

$$(63) \quad \text{sum}_{2,2}(\alpha) \leq (s' - s)^2$$

and

$$(64) \quad \mathbb{E}(X_1^2 X_2^2) + \mathbb{E}(X_1 X_2 X_3 X_4) \leq 2(t' - t)^2.$$

Let us first check (63). We have

$$\text{sum}_{2,2}(\alpha) \leq \left(\sum_i \alpha_i^2 \right)^2 = \left\{ (n - |I|) \frac{|I|^2}{n^3} + |I| \frac{1}{n} (1 - |I|/n)^2 \right\}^2 = \left\{ \frac{|I|}{n} \left(1 - \frac{|I|}{n} \right) \right\}^2,$$

which gives (63). Let us now check (64). As $0 \leq X_i \leq 1$, it suffices to prove that

$$(65) \quad \mathbb{E}(X_1 X_2) \leq (t' - t)^2.$$

We have

$$\mathbb{E}(X_1 X_2) = \sum_{j, j' \in J} \mathbb{E}(m_{1j} m_{2j'}),$$

so it suffices to prove that uniformly on $j, j' \in \{1, \dots, n\}$, $\mathbb{E}(m_{1j} m_{2j'}) \leq \frac{1}{n(n-1)}$. We get this as follows: using the exchangeability of the rows of M and the fact that its rows sum up to one, we have, for any $j, j' \in \{1, \dots, n\}$,

$$1 = \mathbb{E} \left(\left(\sum_i m_{ij} \right) \left(\sum_{i'} m_{i'j'} \right) \right) = n(n-1) \mathbb{E}(m_{1j} m_{2j'}) + n \mathbb{E}(m_{1j} m_{1j'}).$$

This concludes the proof. \square

6.2. Injectivity of the Cauchy transform for certain classes of functions.

Lemma 6.2. *Let f be a real valued bounded càdlàg function on \mathbb{R} vanishing at infinity with at most countably many discontinuity points. Then f is entirely determined by the function*

$$K_f(z) := \int \frac{f(\lambda)}{(z - \lambda)^2} d\lambda \quad (z \in \mathbb{C} \setminus \mathbb{R}).$$

More precisely, for any $\lambda \in \mathbb{R}$, we have

$$(66) \quad f(\lambda) = \lim_{\tilde{\lambda} \downarrow \lambda} \lim_{\eta \downarrow 0} \frac{1}{\pi} \int_{-\infty}^{\tilde{\lambda}} \Im K_f(E + i\eta) dE.$$

f is cont. at $\tilde{\lambda}$

Proof. Let us introduce the Cauchy transform of f , defined, on $\mathbb{C} \setminus \mathbb{R}$, by $H_f(z) := \int \frac{f(\lambda)}{z - \lambda} d\lambda$.

It is well known that at any $\tilde{\lambda}$ where f is continuous, we have

$$f(\tilde{\lambda}) = \lim_{\eta \downarrow 0} -\frac{1}{\pi} \Im H_f(\tilde{\lambda} + i\eta).$$

Then, the result follows because for all $\tilde{\lambda} \in \mathbb{R}$, $\eta > 0$,

$$-H_f(\tilde{\lambda} + i\eta) = \int_{-\infty}^{\tilde{\lambda}} K_f(E + i\eta) dE.$$

□

6.3. CLT for martingales. Let $(\mathcal{F}_k)_{k \geq 0}$ be a filtration such that $\mathcal{F}_0 = \{\emptyset, \Omega\}$ and let $(M_k)_{k \geq 0}$ be a square-integrable complex-valued martingale starting at zero with respect to this filtration. For $k \geq 1$, we define the random variables

$$Y_k := M_k - M_{k-1} \quad v_k := \mathbb{E}[|Y_k|^2 | \mathcal{F}_{k-1}] \quad \tau_k := \mathbb{E}[Y_k^2 | \mathcal{F}_{k-1}]$$

and we also define

$$v := \sum_{k \geq 1} v_k \quad \tau := \sum_{k \geq 1} \tau_k \quad L(\varepsilon) := \sum_{k \geq 1} \mathbb{E}[|Y_k|^2 1_{|Y_k| \geq \varepsilon}].$$

Let now everything depend on a parameter n , so that $\mathcal{F}_k = \mathcal{F}_k^n$, $M_k = M_k^n$, $Y_k = Y_k^n$, $v = v^n$, $\tau = \tau^n$, $L(\varepsilon) = L^n(\varepsilon)$, \dots

Then we have the following theorem. It is proved in the real case at [10, Th. 35.12]. The complex case can be deduced noticing that for $z \in \mathbb{C}$, $\Re(z)^2$, $\Im(z)^2$ and $\Re(z)\Im(z)$ are linear combinations of z^2 , \bar{z}^2 , $|z|^2$.

Theorem 6.3. *Suppose that for some constants $v \geq 0$, $\tau \in \mathbb{C}$, we have the convergence in probability*

$$v^n \xrightarrow[n \rightarrow \infty]{} v \quad \tau^n \xrightarrow[n \rightarrow \infty]{} \tau$$

and that for each $\varepsilon > 0$,

$$L^n(\varepsilon) \xrightarrow[n \rightarrow \infty]{} 0.$$

Then we have the convergence in distribution

$$M_n^n \xrightarrow[n \rightarrow \infty]{} Z,$$

where Z is a centered complex Gaussian variable such that $\mathbb{E}(|Z|^2) = v$ and $\mathbb{E}(Z^2) = \tau$.

6.4. Some linear algebra lemmas. Let $\|\cdot\|_\infty$ denote the operator norm of matrices associated with the canonical Hermitian norm.

Lemma 6.4. *Let $A = [a_{ij}]$ be an $n \times n$ Hermitian matrix, $z \in \mathbb{C} \setminus \mathbb{R}$, $G := (z - A)^{-1}$, P be a diagonal matrix. For $1 \leq k \leq n$ we denote by $A^{(k)}$, $P^{(k)}$ be the matrices with size $n - 1$ obtained by removing the k -th row and the k -th column of A and P and set $G^{(k)} := (z - A^{(k)})^{-1}$. Then*

$$(67) \quad \mathrm{Tr}(PG) - \mathrm{Tr}(P^{(k)}G^{(k)}) = \frac{P_{kk} + \mathbf{a}_k^* G^{(k)} P^{(k)} G^{(k)} \mathbf{a}_k}{z - a_{kk} - \mathbf{a}_k^* G^{(k)} \mathbf{a}_k},$$

with \mathbf{a}_k the k -th column of A where the diagonal entry has been removed. Moreover,

$$(68) \quad |\mathrm{Tr}(PG) - \mathrm{Tr}(P^{(k)}G^{(k)})| \leq \frac{5\|P\|_\infty}{|\Im z|}.$$

Proof. • Let us first prove (67). By linearity, one can suppose that P has only one nonzero diagonal entry, say the i th one, equal to one. Using the well known formula

$$((z - A)^{-1})_{ii} - 1_{i \neq k} ((z - A^{(k)})^{-1})_{ii} = \frac{G_{ki}G_{ik}}{G_{kk}},$$

we have

$$\begin{aligned} \mathrm{Tr}(PG) - \mathrm{Tr}(P^{(k)}G^{(k)}) &= ((z - A)^{-1})_{ii} - 1_{i \neq k} ((z - A^{(k)})^{-1})_{ii} \\ &= \frac{G_{ki}G_{ik}}{G_{kk}} \\ &= \frac{((z - A)^{-1}P(z - A)^{-1})_{kk}}{((z - A)^{-1})_{kk}} \\ &= \frac{\partial_t|_{t=0}((z - A - tP)^{-1})_{kk}}{((z - A)^{-1})_{kk}} \end{aligned}$$

Let \log denote the determination of the log on $\mathbb{C} \setminus \mathbb{R}^-$ vanishing at one. Then

$$\begin{aligned} \mathrm{Tr}(PG) - \mathrm{Tr}(P^{(k)}G^{(k)}) &= \partial_t|_{t=0} \log\{((z - A - tP)^{-1})_{kk}\} \\ &= \partial_t|_{t=0} \log \frac{1}{z - a_{kk} - t1_{k=i} - \mathbf{a}_k^*(z - X^{(k)} - tP^{(k)})^{-1}\mathbf{a}_k} \\ &= -\partial_t|_{t=0} \log(z - a_{kk} - t1_{k=i} - \mathbf{a}_k^*(z - X^{(k)} - tP^{(k)})^{-1}\mathbf{a}_k) \\ &= \frac{-\partial_t|_{t=0}(z - a_{kk} - t1_{k=i} - \mathbf{a}_k^*(z - X^{(k)} - tP^{(k)})^{-1}\mathbf{a}_k)}{(z - a_{kk} - t1_{k=i} - \mathbf{a}_k^*(z - X^{(k)} - tP^{(k)})^{-1}\mathbf{a}_k)|_{t=0}} \\ &= \frac{P_{kk} + \mathbf{a}_k^*G^{(k)}PG^{(k)}\mathbf{a}_k}{z - a_{kk} - \mathbf{a}_k^*G^{(k)}\mathbf{a}_k} \end{aligned}$$

• Let us now prove (68) (the proof does not use (67)). One can suppose that $k = 1$. Let us introduce

$$\tilde{A} := \begin{bmatrix} a_{11} & 0 & \cdots & 0 \\ 0 & & & \\ \vdots & & A^{(1)} & \\ 0 & & & \end{bmatrix}$$

and define \tilde{G} and $\tilde{G}^{(1)}$ as G and $G^{(1)}$ with \tilde{A} instead of A . We have

$$|\mathrm{Tr}(PG) - \mathrm{Tr}(P^{(1)}G^{(1)})| \leq |\mathrm{Tr}(P(G - \tilde{G}))| + |\mathrm{Tr}(P\tilde{G}) - \mathrm{Tr}(P^{(1)}\tilde{G}^{(1)})| + |\mathrm{Tr}(P^{(1)}(G^{(1)} - \tilde{G}^{(1)}))|.$$

Let us treat the terms of the RHT separately.

The third term is null because $\tilde{A}^{(1)} = A^{(1)}$. We have

$$|\operatorname{Tr}(P(G - \tilde{G}))| \leq \|P(G - \tilde{G})\|_\infty \operatorname{rank}(G - \tilde{G})$$

which is $\leq \frac{4\|P\|_\infty}{|\Im z|}$ by the resolvent formula. At last, as P is diagonal and the matrix $z - \tilde{A}$ can be inverted by blocs, we have

$$|\operatorname{Tr}(P\tilde{G}) - \operatorname{Tr}(P^{(1)}\tilde{G}^{(1)})| = |P_{11}\tilde{G}_{11}| \leq \frac{\|P\|_\infty}{|\Im z|}.$$

□

REFERENCES

- [1] G. Anderson, A. Guionnet, O. Zeitouni *An Introduction to Random Matrices*. Cambridge studies in advanced mathematics, **118** (2009).
- [2] A. Auffinger, G. Ben Arous, S. Péché *Poisson convergence for the largest eigenvalues of heavy tailed random matrices*, Ann. Inst. Henri Poincaré Probab. Stat., **45**, 2009, 589–610
- [3] Z. D. Bai, J.W. Silverstein *Spectral analysis of large dimensional random matrices*, Second Edition, Springer, New York, 2009.
- [4] S. Belinschi, A. Dembo, A. Guionnet *Spectral measure of heavy tailed band and covariance random-matrices*, Comm. Math. Phys., **289**, 2009, 1023–1055.
- [5] G. Ben Arous and A. Guionnet *The spectrum of heavy tailed random matrices*. Comm. Math. Phys. **278** (2008), no. 3, 715–751.
- [6] F. Benaych-Georges *Eigenvectors of Wigner matrices: universality of global fluctuations* Random Matrices Theory Appl. Vol. 1 (2012), no. 4, 23 pp.
- [7] F. Benaych-Georges, T. Cabanal-Duvillard *Marchenko-Pastur Theorem and Bercovici-Pata bijections for heavy-tailed or localized vectors* ALEA, Lat. Am. J. Probab. Math. Stat. Vol. 9 (2012), no. 2, 685–715.
- [8] F. Benaych-Georges, A. Guionnet, C. Male *Central limit theorems for linear statistics of heavy tailed random matrices*. arXiv.
- [9] P.J. Bickel, M.J. Wichura *Convergence criteria for multiparameter stochastic processes and some applications*, Ann. Math. Statist., 42(5):1656–1670, 1971.
- [10] P. Billingsley *Probability and measure*, Wiley, third edition.
- [11] C. Bordenave, P. Caputo, D. Chafaï *Spectrum of non-Hermitian heavy tailed random matrices*, Comm. Math. Phys., **307**, 2011, 513–560.
- [12] C. Bordenave, A. Guionnet *Localization and delocalization of eigenvectors for heavy-tailed random matrices*, to appear in Prob. Th. rel. Fields.
- [13] J.-P. Bouchaud, P. Cizeau, *Theory of Lévy matrices* Phys. Rev. E 50 (1994).
- [14] G. Chapuy *Random permutations and their discrepancy process*, in 2007 Conference on Analysis of Algorithms, AofA 07, Discrete Math. Theor. Comput. Sci. Proc. AH (2007), pp. 415–426.
- [15] C. Donati-Martin, A. Rouault *Truncations of Haar unitary matrices, traces and bivariate Brownian bridge*, Random Matrices: Theory and Application (RMTA) Vol. 01, No. 01.
- [16] L. Erdős, A. Knowles, H.T. Yau, J. Yin, *Spectral statistics of Erdős-Rényi Graphs II: Eigenvalue spacing and the extreme eigenvalues*, Comm. Math. Phys., **314**, 2012, 587–640.
- [17] L. Erdős, J. Ramírez, B. Schlein, H.T. Yau *Universality of sine-kernel for Wigner matrices with a small Gaussian perturbation*, Electron. J. Probab., **15**, 2010, no. 18, 526–603.
- [18] L. Erdős, B. Schlein, H.T. Yau *Semicircle law on short scales and delocalization of eigenvectors for Wigner random matrices*, Ann. Probab., **37**, 2009, 815–852.
- [19] L. Erdős, H.T. Yau, J. Yin *Rigidity of eigenvalues of generalized Wigner matrices*, Adv. Math., **229**, 2012, 3, 1435–1515.

- [20] L. Erdős, H.T. Yau, J. Yin *Universality for generalized Wigner matrices with Bernoulli distribution*, J. Comb., **2**, 2011, 15–81
- [21] L. Erdős, B. Schlein, H.T. Yau *Universality of random matrices and local relaxation flow*, Invent. Math., **185**, 2011, 75–119
- [22] L. Erdős, J. Ramirez, B. Schlein, T. Tao, V. Vu, H.T. Yau *Bulk universality for Wigner Hermitian matrices with subexponential decay*, Math. Res. Lett., **17**, **2010**, 667–674
- [23] L. Erdős, J. Ramirez, S. Péché B. Schlein, H.T. Yau *Bulk universality for Wigner matrices*, Comm. Pure Appl. Math., **63**, 2010, 895–925
- [24] A. M. Khorunzhy, B. A. Khoruzhenko, L.A. Pastur *Asymptotic properties of large random matrices with independent entries*, J. Math. Phys. **37** (1996) 5033–5060.
- [25] O. Khorunzhy, M. Shcherbina, V. Vengerovsky *Eigenvalue distribution of large weighted random graphs*, J. Math. Phys. **45** (2004), no. 4, 1648–1672.
- [26] C. Male *The limiting distributions of large heavy Wigner and arbitrary random matrices*, arXiv:1111.4662v3 preprint.
- [27] M. Mehta, *Random matrices*, Pure and Applied Mathematics (Amsterdam), **142**, Third, Elsevier/Academic Press, Amsterdam, 2004.
- [28] M. Shcherbina, B. Tirozzi *Central limit theorem for fluctuations of linear eigenvalue statistics of large random graphs*, J. Math. Phys., **51**, 2010, 023523, 20
- [29] J. W. Silverstein *Weak convergence of random functions defined by the eigenvectors of sample covariance matrices*. Ann. Probab. **18** (1990), no. 3, 1174–1194.
- [30] F. Slanina *Localization of eigenvectors in random graphs*, Eur. Phys. B, 2012, 85:361.
- [31] A. Soshnikov *Poisson statistics for the largest eigenvalue of Wigner random matrices with heavy tails*. Electron. Comm. Probab. **9** (2004) 82–91.
- [32] T. Tao, V. Vu *Random matrices: universality of local eigenvalue statistics*, Acta Math., **206** (2011), 127–204.
- [33] T. Tao *The asymptotic distribution of a single eigenvalue gap of a Wigner matrix*, Probab. Theory Related Fields, **157**, 2013, 81–106.
- [34] C. Tracy, H. Widom *Level-spacing distributions and the Airy kernel*. Comm. Math. Phys. **159** (1994), no. 1, 151–174.
- [35] I. Zakharevich *A generalization of Wigner’s law*, Comm. Math. Phys., **268**, 2006, 2, 403–414.
- [36] E.P. Wigner *On the distribution of the roots of certain symmetric matrices*, Ann. Math., **67**, 1958, 325–327.