



Structure and yield of all-sized and even-sized conifer-dominated stands on fertile sites

E Lähde, O Laiho, Y Norokorpi, T Saksa

► To cite this version:

E Lähde, O Laiho, Y Norokorpi, T Saksa. Structure and yield of all-sized and even-sized conifer-dominated stands on fertile sites. *Annales des sciences forestières*, 1994, 51 (2), pp.97-109. <hal-00882935>

HAL Id: hal-00882935

<https://hal.science/hal-00882935v1>

Submitted on 11 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

Original article

Structure and yield of all-sized and even-sized conifer-dominated stands on fertile sites

E Lähde, O Laiho, Y Norokorpi, T Saksa

The Finnish Forest Research Institute, Box 18, FIN-01301 Vantaa, Finland

(Received 4 June 1993; accepted 22 September 1993)

Summary — The material studied consisted of 807 sample plots located in SW Finland. The data were inventoried (in 1951–1953) using a systematic temporary circular plot line survey. Each plot represented a particular stand. An all-sized stand structure (*ie* a stem distribution resembling an inverted letter J) amounted to 62% of sample plots whereas 25% were even-sized (resembling a normal distribution) and 13% were irregularly uneven-sized. The number of stems per ha in all-sized stands was nearly twice that of even-sized stands. The mean annual increment increased linearly with an increase in volume. Consequently, there was no difference in increment between all-sized and even-sized stands because the volume of the growing stock in the latter group was greater than in the former. The mean annual increment in stands with equal average volumes was, however, greater in all-sized mixed stands than in even-sized conifer stands. The relative growth in all-sized mixed stands was approximately 25% higher than in corresponding even-sized stands.

stem distribution / stand structure / yield / all-sized stand / even-sized stand / fertile site

Résumé — **Structure et rendement de peuplements irréguliers et réguliers à majorité de conifères sur terrains fertiles.** Le matériel d'étude est constitué de 807 placettes d'échantillonnage réparties dans le sud-ouest de la Finlande (fig 1). Les données étaient inventoriées (1951-1953) au moyen de placettes circulaires temporaires systématiques alignées. Chaque placette représentait un peuplement particulier. Des placettes d'échantillonnage (62%) avaient une structure de peuplement irrégulière (c'est-à-dire avec une distribution des diamètres des tiges ressemblant à la lettre J inversée), 25% une structure régulière (ressemblant à une distribution normale des tiges) et 13% une structure irrégulière aux diamètres inégaux. Le nombre de tiges par hectare dans les structures irrégulières atteignait presque le double de celui des structures régulières (fig 2). L'accroissement annuel moyen augmentait de façon linéaire avec l'augmentation en volume (fig 3). Par conséquent, il n'y avait pas de différence d'accroissement entre les peuplements irréguliers et les peuplements réguliers car le volume du matériel sur pied croissant du dernier groupe était plus grand que celui du premier groupe (tableau I). L'accroissement annuel moyen de peuplements à volume moyen égal était toutefois plus grand dans des peuplements mélangés irréguliers que dans des peuplements de conifères réguliers (tableau II). La croissance relative des peuplements mélangés irréguliers était environ 25% plus élevée que celle des peuplements correspondants réguliers.

distribution des tiges / structure de peuplement / rendement / peuplement irrégulier / peuplement régulier / terrain fertile

INTRODUCTION

Even-sized (even-aged) forestry and all-sized (uneven-aged) forestry are the 2 main principles applied in the tending of forests. All-sizedness (all-agedness) is the term applied to stands whose stem distribution more or less resembles an inverted letter J, i.e. the number of trees in the diameter classes diminishes as the diameter increases (de Liocourt, 1898; Baker, 1934; Meyer, 1952; Alexander and Edminster, 1978; Curtis, 1978; Gibbs, 1978; Daniel *et al*, 1979). Oliver and Larson (1990) refer to stands whose stem distribution resembles an inverted letter J by the name multicohort stands. In this study such stands are referred to as all-sized stands. Even-sizedness (even-agedness) is loosely defined as referring to stands whose stem distribution roughly resembles a normal distribution (Curtis, 1978; Gibbs, 1978; Gingrich, 1978; Daniel *et al*, 1979). This approach is applied in this study as well. A more precise criterion involves the application of a certain range of diameter classes (Smith, 1962; Daniel *et al*, 1979). This range may be, for instance, 12 or 20 cm (Lähde *et al*, 1991, 1992).

Diameter at breast height is the generally used means of depicting age differences. However, tree age and tree size do not always correlate very well. In fact, tree growth has been observed to be more dependent on size than age (Cajander, 1934; Sarvas, 1944; Vuokila, 1970; Indermühle, 1978). On being released from the dominance of larger trees, the trees forming the understorey generally attain the same size as those that have always been free to grow (Cajander, 1934; Näslund, 1944; Hawley, 1946; Hatcher, 1967; Schütz, 1969; Indermühle, 1978; Nilsen and Haveræen, 1983; Klensmeden, 1984). When diameter is used as the criterion for structure it is more appropriate to use the terms all-sized and even-sized instead of all- or uneven-aged and even-aged. The termino-

logy used in this study complies with this view.

Both practicing foresters as well as researchers have participated in the age-old debate on which of the 2 principles is better (Hassenkamp, 1955; Børset, 1963; Carbonnier, 1978; Mikola, 1984; Viitala, 1986). The prevailing opinion has varied from country to country – even in cycles of a few decades (Mustian, 1978).

In Fennoscandia, the raising of even-sized forest stands became the standard practice around the middle of this century. During the following decades the opposition to silviculture aimed at all-sized stand structure was extremely strong in the Nordic countries, especially in Sweden and Finland. Thereafter, however, forest use has become diversified. The role of forests in recreation, protection of the environment and the landscape as well as in other aspects of multiple-use has received increasing attention. Growing interest has been directed to the ways in which forests are treated (Smith, 1972, 1975; Gould, 1975; Vrabec, 1977; Doolittle, 1978; Gibbs, 1978; Lundqvist, 1984). For instance, in the United States the general trend has led to the development of silvicultural regimes aimed at raising structurally all-sized forests (Gingrich, 1967; Leak *et al*, 1969; Gibbs, 1978; Hann and Bare, 1979). In the case of the Nordic countries this stage has only just begun (Lähde *et al*, 1985; Hagner, 1992a, 1992b; Haveræen, 1992; Lähde, 1992; Larsen, 1992).

Although the dispute over the main policy to be followed in the raising of forests has been an on-going issue for decades, there are only few long-term silvicultural experiments comparing the differences between the opposed policies. Separate studies are, however, available on the 2 options. The number of studies conducted on the raising of even-sized stands clearly exceeds that of all-sized silviculture despite the fact that natural development results in highly diverse

forest structure (Ilvessalo, 1920a, 1920b; Sirén, 1955; Kammerlander, 1978; Larsen, 1980; Heinzelman, 1981; Solomon *et al*, 1986; Norokorpi, 1992). This diversity is often accompanied by all-sizedness (Ussva, 1932; Pobedinski, 1988; Lähde *et al*, 1991, 1992).

With the exception of some studies (*eg* Böhmer, 1957; Mitscherlich, 1963; Kern, 1966; Hasse and Ek, 1981; Pretzsch, 1985; Solomon *et al*, 1986; Haight and Gets, 1987; Lundqvist, 1989), the comparison of the differences between these 2 silvicultural policies has remained at the level of rough estimations only. Several simulation models have also been developed for different growing stocks in the case of all-sized silviculture (Eyre and Zillgitt, 1953; Trimble, 1961, 1970; Hart, 1964; Marquis *et al*, 1969; Mayer, 1969; Frank and Björkom, 1973; Adams and Ek, 1974; Hladik, 1975; Leak and Graber, 1976). Inventories of timber resources represent a hitherto little appreciated means of obtaining comparative data on forest yield and structure.

The material chosen for this study consists of part of the data collected in the course of the 3rd national forest inventory carried out in Finland. In the study all-sized and even-sized stands of conifers and of mixed species are compared in terms of occurrence, structure and yield on fertile sites. In accordance with the concept generally accepted in the Nordic countries, the hypotheses applied in the study are: (1) that mixed broad-leaved-coniferous stands grow better than conifer stands; and (2) that even-sized stands grow better than all-sized stands.

MATERIALS AND METHODS

The 3rd national forest inventory (1951–1953) in Finland was conducted as a systematic temporary circular plot line survey (Ilvessalo, 1951). Until then the forests of Finland had generally

been treated with various forms of light selection felling and thinning from above. Dimension felling has also been widely used (Ilvessalo, 1956).

Southwestern Finland (fig 1) was chosen for this study because of its uniform climatic conditions. The material was collected from fertile mineral soil sites (*Myrtillus* site type or more fertile, see Cajander, 1949).

The mean dominant height (100 thickest trees/ha) was also measured on most of the plots. Because the age of dominant trees was not exactly determined, the site (height) index of the plot stands could not be estimated. However, dominant height measured may give a good base for comparisons of site quality. According to Indermühle (1978) biological age is not characteristic in uneven-aged forests because growing in suppression causes an overestimation of age. In general, the site index is difficult to estimate in uneven-aged forests (Andreassen, 1992).

The stands were in thinning, preparatory or regeneration cutting stages, with a volume of at least 40 m³/ha. The growing stock was generally dominated by Norway spruce (*Picea abies* L Karst), with admixtures of Scots pine (*Pinus sylvestris* L) and broad leaved species (*Betula pendula* Roth 20%, *B. pubescens* Ehrh 50% and others, mainly *Alnus incana*, 30%). The silvicultural state had to be good or satisfactory or the stands had to have been untreated for many years (Ilvessalo, 1951).

Each sample plot represented a particular stand, *ie* it was located entirely within one stand (Ilvessalo, 1951). Thus, the structure on any plot could not be admixture of different stands. The size of the plots was 0.1 ha (1 000 m²) and all trees with dbh (diameter at breast height) over 10 cm were measured. Small trees (dbh 2–10 cm) were tallied from within a concentric circle with an area of 0.01 ha (100 m²).

In this study the trees were divided according to dbh into 4 cm diameter classes as follows: 1 = 2–6; 2 = 6–10; 3 = 10–14; 4 = 14–18; 5 = 18–22; 6 = 22–26; 7 = 26–30; 8 = 30–34; and 9 = > 34 cm. Broad leaved trees of vegetative origin were not tallied as they were not assumed to be capable of developing into actual trees. The sample plots were individually classified according to the structure as follows (applying the classifications used by Smith (1962) and Daniel *et al* (1979)):

J: All-sized. Stem distribution resembling an inverted letter J; trees present in at least the 4 smallest




Fig 1. Study area in southwestern Finland.

diameter classes, with the mode in the first or second class (502 sample plots);

E: Even-sized. Stem distribution resembling a normal distribution; mode in neither of the 2 smallest diameter classes nor at either end of the distribution (197 sample plots);

O: Others (irregularly uneven-sized) (108 sample plots, only some main results are given).

Classifications of another kind have also been used in describing the stand structure. Leemans (1991) and Szwagrzyk (1992), for instance, used

the age, height, dbh, and exact tree location as characters.

A total of 807 sample plots were studied (table 1). The structural groups were divided into 2 sub-groups on the basis of the stem number of broad-leaved species. The division is generally made according to volume or basal area. In stands that were all-sized in structure the number of small trees is, however, important for stand development. The species groups were as follows:

Table 1. Parameters ($\bar{x} \pm se$) of the stands studied. Results of 2-way analysis of variance and Tukey's test.

<i>Stand structure</i>	<i>Tree species group</i>	<i>Sample plots</i>	<i>Relative growth (%)</i>	<i>Volume (m³/ha)</i>	<i>Mean increment (m³/ha/a)</i>
All-sized	Conifer	160	3.6 (0.1 ^a)	155 (5 ^a)	5.2 (0.1 ^a)
	Mixed	342	4.3 (1.1 ^b)	137 (3 ^b)	5.5 (0.1 ^a)
Even-sized	Conifer	119	3.1 (0.1 ^c)	187 (6 ^c)	5.4 (0.2 ^a)
	Mixed	78	3.4 (0.1 ^{ac})	177 (7 ^c)	5.7 (0.2 ^a)
<i>F</i> -value, structure			37.98 ***	54.28 ***	1.87
<i>F</i> -value, tree species			15.94 ***	8.31 **	4.07
Interaction			4.29	0.62	0.15

Values with different small letters differ statistically ($p < 0.05$) from each other.

A: Conifer stands. No more than 120 broad-leaves trees per ha (average = 78).

B. Mixed (broad leaved–coniferous) stands. More than 120 broad leaved trees per ha (average = 698; basal area 30%).

Comparison of the yield between different stand groups is presented as a mean annual increment (excluding bark) for the total material and for the same average volume class, and as a relative growth (%). The effect of stand structure and tree species composition on different stand parameters was analysed with 2-way analysis of variance. The differences between different group means were tested with Tukey's test. The dependence between mean annual increment and volume in stands with different structure and tree species composition was analysed with regression analysis.

RESULTS

Structure, species composition and stem number

The stem number in mixed stands was higher than in conifer stands (fig 2). The proportion of broad-leaved species in the mixed stands averaged 34–41%. The broad-leaved species particularly increased the proportion of small trees. Broad-leaved trees

with diameters over 30 cm were observed only on some sample plots. The stem number in all-sized stands was 2 329 stems/ha; this was more than twice as much as the 1 079 stems recorded in even-sized stands and almost twice as much as in irregularly uneven-sized stands where the figure was 1 173.

In all-sized stands, on average, the stem distribution of both Norway spruce and broad-leaved species resembled an inverted letter J just as the overall stem distribution did. Correspondingly, the stem distribution of the various tree species in even-sized stands resembled a normal distribution in the same way as the overall stem distribution did (fig 2). The proportion of Scots pine was greater in these (15%) than it was in all-sized stands (8%). The difference in the overall number of stems between the structural groups was a consequence of the number of spruce and broad-leaved species.

Yield

The relative growth (mean of 5 previous years, %) in mixed stands was higher than




Fig 2. Average stem distribution in stands of different structure and tree species composition. Observe the scales showing stem number in the subfigures. Legends: dbh – classes: 1 = 2–6; 2 = 6–10; ..., 9 = > 34 cm. Total stem numbers/ha are given in the subfigures.

in conifer stands (table I). However, only the largest difference (19%), in all-sized stands, was statistically significant ($p < 0.01$). The relative growth in both all-sized conifer and mixed stands was significantly ($p < 0.01$) higher than in the corresponding even-sized stands.

The volume of the growing stock in mixed stands was less than that in conifer stands. This was also the case when comparing all-sized stands with even-sized stands. The differences in mean annual increment averages between these groups were not statistically significant (table I). A




Fig 3. Dependence between mean annual increment and volume in all-sized (J) and even-sized (E) stands. A = conifer and B = mixed stands.

positive linear dependence applied between mean annual increment and volume (fig 3). When the volume in all-sized mixed stands, for instance, rose from 100 m³/ha to 300 m³/ha, the mean annual increment rose by ≈ 3 m³/ha.

The dominant height (\bar{x} , se, m) in all-sized and even-sized stands is presented in the following setting:

All-sized	Conifer	19.1 ± 0.2
	Mixed	18.3 ± 0.2
Even-sized	Conifer	20.0 ± 0.3
	Mixed	20.1 ± 0.3

The dominant height in all-sized stands differed significantly ($p < 0.05$) from that in even-sized stands. The difference between

conifer and mixed stands was significant only in the all-sized group.

The mean annual increment in all-sized mixed stands with equal average volume ($152 \pm 15 \text{ m}^3/\text{ha}$) was a quarter higher ($1.2 \text{ m}^3/\text{ha}$, under bark) than that of even-sized conifer stands (table II). The difference in growth for these structure groups was statistically significant ($p < 0.05$). The over-bark (average bark 16%; Ilvessalo, 1956) mean annual increment was $7.0 \text{ m}^3/\text{ha}$ in this all-sized mixed stand group.

The dominant height (\bar{x} , se, m) of these groups was as follows:

All-sized	Pure conifer	18.6 ± 0.4
	Mixed	19.2 ± 0.3
Even-sized	Pure conifer	19.3 ± 0.5
	Mixed	20.8 ± 0.3

Table II. Comparison of mean annual increment (\bar{x} , se, m^3/ha) in stands of different structure and tree species composition and with equal average volumes (mean $152 \pm 15 \text{ m}^3/\text{ha}$). Results of 2-way analyses of variance and Tukey's test (log transformation).

Stand structure	Tree species groups	Sample plots	Mean annual increment (excluding bark)
All-sized	Conifer	25	5.3 (0.2 ^{ab})
	Mixed	63	5.9 (0.2 ^b)
Even-sized	Conifer	21	4.7 (0.4 ^a)
	Mixed	20	5.2 (0.3 ^{ab})
<i>F</i> -value, structure			4.16*
<i>F</i> -value, tree species			2.91
Interaction			0.02

Values with different small letters differ statistically ($p < 0.05$) from each other.

The dominant height in even-sized mixed stands differed significantly ($p < 0.05$) from all-sized stands.

DISCUSSION

The first hypothesis set for this study (i.e. that mixed (broad leaved-coniferous) stands were better than conifer stands) was confirmed fairly well in all-sized stands. According to Frivold (1982) the yield in mixed stands of birch and spruce in Norway was better than in pure spruce stands. However, the proportion of birch should be clearly decreased at the age of 40 yr on fertile sites and at the age of 70 yr on barren sites. In Central Europe mixed stands of birch and spruce grew better than pure spruce stands (Otto, 1986).

Previous studies (Phares, 1978) have also shown that species composition plays a role in the yield and development of structurally different stands. The trees in mixed stands are more closely spaced than in conifer stands (Frivold, 1982; Mielikäinen, 1985). Mixed stands also use better the growing space available in the soil in that different tree species, especially spruce and birches, have different rooting depths (Laitakari, 1927, 1934; Sirén, 1955).

The relative growth in stands of diverse structure under corresponding conditions to this study has generally varied within the range of 2–4% (Barth, 1929; Näslund, 1944; Böhmer, 1957; Nilsen and Haverlaen, 1983; Lundqvist, 1989). Barth (1929) reported a growth figure 1.7 m^3 greater in a Norway spruce-dominated stand of diverse structure than the average for the forest region in question. According to Böhmer (1957), the growth of Norway spruce in an irregularly uneven-aged stand was equal to the average growth of an even-aged stand.

Indermühle (1978) found the yield in a spruce-dominated all-sized stand in sou-

thern Germany to be surprisingly high. Ekhart *et al* (1961) and Mayer (1969) stated in Austria that an all-sized stand grows better than an even-sized one. Smith and De-Bald (1978) concluded from several materials in North America that the yield in all-sized forests is slightly higher than that in even-sized ones. Hasse and Ek (1981) have observed all-sized stands in broad leaved forests of North-America to produce more commercial timber than even-sized stands do although their total yield hardly differs. In general, stand structure had little influence on the yield of forest (Burger, 1942; Smith, 1962; Mitscherlich, 1963; Kern, 1966; Gingrich, 1967; Hladik, 1975; Lundqvist, 1989). According to some studies the volume increment in even-sized stands is higher than in all-sized ones (Walker, 1956; Trimble and Manthy, 1966; Trimble and McClung, 1966; McCauley and Trimble, 1972, 1975).

In a simulation-based study, Pukkala and Kolström (1988) estimated the yield of an all-sized (uneven-aged) stand of Norway spruce in southern Finland to be 5 m³/ha/a. They compared the simulated growth estimate to the yield tables compiled by Koivisto (1954) for repeatedly treated even-sized stands. However, Koivisto's material also included all-sized stands since the majority of forest stands of that time were all-sized in structure (Lähde *et al*, 1992). Later Kolström (1992) estimated that growth in all-sized stands is about the same as growth in even-sized stands.

When sample plots carrying equivalent average volumes were compared in this study, it was observed that the increment of all-sized mixed stands was greater than that of even-sized conifer stands. The relative growth was greater in all-sized stands than in even-sized stands independently of volume. The difference in dominant height was the inverse. Thus, the hypothesis of even-sized stands possessing greater growth than all-sized stands was not confirmed. The result was quite the opposite.

ACKNOWLEDGMENTS

We thank M Hagner, JP Schütz, K Andreassen, LH Frivold and 2 anonymous reviewers for manuscript review.

REFERENCES

- Adams DM, Ek AR (1974) Optimizing the management of uneven-aged forest stands. *Can J For Res* 4, 274-287
- Alexander RR, Edminster CB (1978) Regulation and control under uneven-aged management. *In: Uneven-aged silviculture management in the United States*, US Dep Agric, For Serv Timber management research, Washington, DC Gen Tech Rep WO-24, 217-230
- Andreassen K (1992) Volume production and problem with the stability of the stand structure in uneven-aged and multi-layered spruce (*Picea abies* Karst) forest in Norway. *In: Silvicultural alternatives. Proc Internordic Workshop June 22-25 1992* (M Hagner, ed). Swedish University of Agricultural Sciences. Department of Silviculture, Reports, No 35, 23-28
- Baker FS (1934) *Theory and Practice of Silviculture*. McGraw-Hill Book Company. New York and London, 502 p
- Barth A (1929) Skjermforyngelsen i produksjons – økonomisk belysning. *Acta For Fenn* 34, 1-33
- Böhmer JG (1957) *Bledningsskog II. Tidsskrift for skogbruk* 65, 203-247
- Børset O (1963) Skogskjøtsel og skogpleie gjennom 50 år. Norges landbrukshøgskole, Institutt for skogskjøtsel. 8 p
- Burger H (1942) Holz, Blattmenge und Zuwachs. VI. Ein Plenterwald mittlerer Standortsgüte. Der bernische Staatswald Toppwald im Emmental. *Mitt Schweiz Anstalt forst Versuchswesen* 22, 377-445
- Cajander AK (1949) Forest types and their significance. *Acta For Fenn* 56, 1-71
- Cajander EK (1934) Kuusen taimistojen vapautamisen jälkeisestä pituuskasvusta. *Commun Inst For Fenn* 19, 1-53
- Carbonnier C (1978) Skogarnas vård och förtyngning. *In: Skogshögskolan 150 år. Sveriges*

- Lantbruksuniversitet. Allmänna skrifter nr 2. Uppsala, 85-126
- Curtis RO (1978) Growth and yield in uneven-aged stands. In: *Uneven-aged silviculture and management in the United States*, US Dep Agric, For Serv. Timber management research, Washington, DC Gen Tech Rep WO-24, 186-201
- Daniel TW, Helms JA, Baker FS (1979) *Principles of Silviculture*. 2nd ed, McGraw-Hill Book Company. 500 p
- Doolittle WT (1978) Constraints on forest management in the eastern hardwood region. In: *Uneven-aged silviculture and management in the United States*, US Dep Agric, For Serv Timber management research, Washington, DC Gen Tech Rep WO-24, 63-66
- Ekhart G, Frauendorfer R, Nather J (1961) Die Wälder der Gemeinde Julbach, unter besonderer Berücksichtigung der stufig aufgebauten Mischwälder. *Mitt Forstl Bundes-Vers Anst Mariabrunn* 58, 1-93
- Eyre FH, Zillgitt WM (1953) Partial cuttings in northern hardwoods of the Lake States. *US Dep Agric, Tech Bull* 1076, 124 p
- Frank RM, Björkbom JC (1973) *A Silvicultural Guide for Spruce-Fir in the Northeast*. US Dep Agric, For Serv Gen Tech Rep NE6, 23 p
- Frivold LH (1982) Bestandsstruktur og produksjon i blandingskog av bjørk (*Betula verrucosa* Ehrh, *B pubescens* Ehrh) og gran (*Picea abies* (L) Karst) i Sydøst-Norge. Summary: Stand structure and yield of mixed stands of birch (*Betula verrucosa* Ehrh, *B pubescens* Ehrh) and spruce (*Picea abies* (L) Karst) in South East Norway. *Meld Nor Landbrukshøgsk* 61, 18, 108 p
- Gibbs CB (1978) Uneven-aged silviculture and management? Even-aged silviculture and management? Definitions and differences. In: *Uneven-aged silviculture and management in the United States*. US Dep Agric, For Serv Timber management research. Washington, DC Gen Tech Rep WO-24, 18-24
- Gingrich SF (1967) Measuring and evaluating stocking and stand density in upland hardwood forests in the Central States. *For Sci* 13, 38-53
- Gingrich SF (1978) Growth and yield. In: *Uneven-aged silviculture and management in the United States*. US Dep Agric, For Serv. Timber management research. Washington, DC Gen Tech Rep WO-24, 115-124
- Gould EM (1975) Aesthetics and land use. Proc Logging Residue Conf, Morgantown West Virg. June 4-6, 1975
- Hagner M (1992a) Basic ideas behind the new trends in Swedish forestry. In: *Silvicultural alternatives. Proc Internordic Workshop June 22-25 1992* (M Hagner, ed). Swedish University of Agricultural Sciences, Department of Silviculture, Reports No 35, 10-12
- Hagner M (1992b) Biologiskt och ekonomiskt resultat i fältförsök med plockhuggning kombinerad med plantering. Abstract: Biological and economical results from experiments with selective felling combined with enrichment planting. The Swedish University of Agricultural Sciences. Department of Silviculture. Working paper 63, 1-52
- Haight RG, Getz WM (1987) Fixed and equilibrium endpoint problems in uneven-aged stand management. *For Sci* 33, 908-931
- Hann DW, Bare BB (1979) *Uneven-aged forest management: State of the art (or science?)* US Dep Agric, For Serv Gen Tech Res INT-50, 18 p
- Hart AC (1964) *The Penobscot management-intensity demonstration plots*. US Dep Agric, For Serv Res Pas NE-25, 24 p
- Hasse WD, Ek AR (1981) A simulated comparison of yield for even- versus uneven-aged management of northern hardwood stands. *J Env Manag* 12, 236-246
- Hassenkamp W (1955) Das Scheitern der Plenterwirtschaft in Skandinavien. *Forstarchiv* 26, 241-251
- Hatcher RJ (1967) Balsam fir advance growth after cutting in Quebec. *For Chron* 40, 86-92
- Haveraaren O (1992) The background of the new trend in silviculture in Norway. In: *Silvicultural alternatives. Proc Internordic Workshop June 22-25 1992* (M Hagner, ed). Swedish University of Agricultural Sciences. Department of Silviculture. Reports, No 35, 13-14
- Hawley RC (1946) *The Practice of Silviculture*. 5th Ed. John Wiley and Sons, Inc New York, 354 p
- Heinselman ML (1981) Fire and succession in the conifer forests of northern North America. In: *Forest succession: Concepts and application* (DC West, HH Shugart, DB Botkin, eds). Springer-Verlag New York, 374-405
- Hladik M (1975) Die Holzproduktion im gemischten Tannen- und Buchenwald bei einer Kahl-

- schlag- und Plenterwirtschaft. Lesnickej fakulty VSLD vo Zvolene, *Zborník vedeckých Prac* 7-2, 55-72
- Ilvessalo Y (1920a) Tutkimuksia metsätyyppien taksatoorisesta merkityksestä nojautuen etupäässä kotimaiseen kasvutaulujen laatimistyöhön. Referat: Untersuchungen über die taxatorische Bedeutung der Waldtypen, hauptsächlich auf den Arbeiten für die Aufstellung der neuen Ertragstafeln Finnlands fassend. *Acta For Fenn* 15, 1-157
- Ilvessalo Y (1920b) Kasvu- ja tuototaulukot Suomen eteläpuoliskon mänty-, kuusi- ja koivumetsille. Referat: Ertragstafeln für die Kiefern, Fichten- und Birkenbestände in der Südhälfte von Finland. *Acta For Fenn* 15, 1-94
- Ilvessalo Y (1951) III valtakunnan metsien arviointi. Suunnitelma ja maastotyön ohjeet. Summary: Third national forest survey of Finland. Plan and instructions for field work. *Commun Inst For Fenn* 39, 1-67
- Ilvessalo Y (1956) Suomen metsät vuosista 1921–24 vuosiin 1951–53. Kameen valtakunnan metsien inventointiin perustuva tutkimus. Summary: The forests of Finland from 1921–1924 to 1951–1953. A survey based on three national forest inventories. *Commun Inst For Fenn* 47, 1-227
- Indermühle MP (1978) Struktur-, Alters-, und Wachstumsuntersuchungen in einem Fichten-Plenterwald des subalpinen Stufe. Beiheft zu den Zeitschriften des Schweizerischen Forstvereins 60, 98 p
- Kammerlander H (1978) Aufbau, Verjüngung und Verbißgefährdung der Plenterwälder im Raum Kufstein/Tirol. *Schweiz Zeit Forstwesen* 129, 711-726
- Kern KG (1966) Wachstum und Umweltfaktoren im Schlag- und Plenterwald. Vergleichende ertragskundlich-ökologische Untersuchungen in zwei Ta-Fi-(Bu)-Plenterwaldbeständen und zwei benachbarten Fi-Schlagwäldern des Hochschwarzwaldes. Bayerischer Landwirtschaftsverlag, München Basel Wien, 232 p
- Klensmeden U (1984) Stamvis blädning. Några studier på två försöksytor i Dalarna. Examensarbete 1984-1986. Sveriges lantbruksuniversitet. Institutionen för skogsskötsel. Umeå, 38 p
- Koivisto P (1954) Kasvu- ja tuototaulukoita. Summary: Growth and yield tables. *Commun Inst For Fenn* 5, 1-49
- Kolström T (1992) Dynamics of uneven-aged stands of Norway spruce: A model approach. Metsäntutkimuslaitoksen tiedonantoja 411. The Finnish Forest Research Institute, Research Papers 411, 29 p + appendixes
- Lähde E (1992) The background ideas to the new trends in silviculture. In: *Silvicultural alternatives. Proc Internordic Workshop June 22–25 1992* (M Hagner, ed). Swedish University of Agricultural Sciences. Department of Silviculture Reports, No 35, 17-19
- Lähde E, Laiho O, Norokorpi Y, Saksa T (1991) The structure of advanced virgin forest in Finland. *Scand J For Res* 6 (4), 527-537
- Lähde E, Laiho O, Norokorpi Y, Saksa T (1992) Stand structure of thinning and mature conifer-dominated forests in boreal zone. In: *Silvicultural alternatives. Proc Internordic Workshop June 22–25 1992* (M Hagner, ed) Swedish University of Agricultural Sciences. Department of Silviculture, Reports, No 35, 58-65
- Lähde E, Norokorpi Y, Oikarinen M (1985) Mikkelin ekoläänin metsien vaihtoehtoiset käsittelymallit. Metsäntutkimuslaitoksen tiedonantoja 180. *The Finnish Forest Research Institute. Research Papers* 180, 67 p
- Laitakari E (1927) Männyn juuristo. Morfologinen tutkimus. Summary: The root system of pine (*Pinus silvestris*). A morphological investigation. *Acta For Fenn* 33 (1), 1-380
- Laitakari E (1934) Koivun juuristo. Summary: The root system of birch (*Betula verrucosa* and *odorata*). *Acta For Fenn* 41(2), 1-216
- Larsen JA (1980) *The boreal ecosystem*. Academic Press, New York, 500 p
- Larsen JB (1992) New trends and on-going research in alternative silvicultural methods in Denmark. In: *Silvicultural alternatives. Proc Internordic Workshop June 22–25 1992* (M Hagner, ed). Swedish University of Agricultural Sciences. Department of Silviculture, Reports, No 35, 15-16
- Leak WB, Graber RE (1976) Seedling input, death, and growth in uneven-aged northern hardwoods. *Can J For Res* 6 (3), 368-374
- Leak WB, Solomon DS, Filip SM (1969) A silvicultural guide for northern hardwoods in the Northeast. US Dep Agric, For Serv Res Pap NE-143, 34 p
- Leemans R (1991) Canopy gaps and establishment patterns of spruce (*Picea abies* (L) Karst)

- in two old-growth coniferous forests in central Sweden. *Vegetatio* 93, 157-165
- de Liocourt F (1988) De l'aménagement des sapinières. *Bull Soc For Franche-Comté Belfort* 6, 396-405
- Lundqvist L (1984) Blädning och etappvis slutavverkning – litteraturstudie. *Sver Skogsvård-sförb Tidskr* 6, 27-40
- Lundqvist L (1989) Blädning i granskog. Strukturförändringen, volymtillväxt, inväxning och föryngring på försöksytor skötta med stamvis blädning. Summary: Use of the selection system in Norway spruce forests – changes in the stand structure, volume increment, ingrowth and regeneration on experimental plots managed with single-tree selection. Dissertation. Sveriges landbruksuniversitet. Institutionen för skogsskötsel. Umeå, 99 p
- Marquis DA, Solomon DS, Björkbom JC (1969) A silvicultural guide for paper birch in the Northeast. *US Dep Agric, For Serv Res Pap NE-130*, 47 p
- Mayer H (1969) Long term rationalization of forest enterprises by way of silviculture. *Allg Forst-zeitschr* 23, 40-45
- McCauley OD, Trimble GR Jr (1972) Forestry returns evaluated for uneven-aged management in two Appalachian woodlots. *USDA For Serv Res Pap NE-244*, 12 p + illus
- McCauley OD, Trimble GR Jr (1975) Site quality in Appalachian hardwoods: the biological and economic response under selection silviculture. *USDA For Serv Res Pap NE-312*, 22 p + illus
- Meyer HA (1952) Structure, growth and drain in balanced uneven-aged forests. *J For* 50, 85-92
- Mielikäinen K (1985) Koivusekoituksen vaikutus kuusikon rakenteeseen ja kehitykseen. Summary: Effect of an admixture of birch on the structure and development of Norway spruce stands. *Commun Inst For Fenn* 133, 79 p
- Mikola P (1984) Harsintametsätalous. Summary: Selection system. *Silva Fennica* 18, 293-301
- Mitscherlich G (1963) Untersuchungen in Schlag- und Plenterwäldern. *Allg Forst Jagdz* 134 (1), 1-12
- Mustian AP (1978) History and philosophy of silviculture management systems in use today. In: Uneven-aged silviculture and management in the United States. *US Dep Agric, For Serv Timber Management Research Gen Tech Rep WO-24*, 1-17
- Näslund M (1944) Den gamla norrländska granskogens reaktionsförmåga efter genomhug-gning. Referat: Die Reaktionsfähigkeit des alten norrländischen Fichtenwaldes nach Dur-chauung. *Medd Statens skogsförsöksanstalt* 33, 1-194
- Nilsen P, Haveraaen O (1983) Årringbredder hos gjenstående traer etter hogst i eldre granskog. Rapport fra Norsk Institutt for skogforsk-ning 9/83, 1-16
- Norokorpi Y (1992) Natural structure and deve-lopment of forests as a basis for alternative silvicultural methods in northern Finland. In: *Silvicultural alternatives. Proc Internordic Workshop June 22-25 1992* (M Hagner, ed). Swedish University of Agricultural Sciences. Department of Silviculture. Reports No 35, 78-85
- Oliver CD, Larson BC (1990) *Forest stand dyna-mics*. McGraw-Hill, New York, 467 p
- Otto HJ (1986) Standortliche Voraussetzungen, Ziele und Waldbautechnik in Fichten-Buchen-Mischbeständen des Harges. *Allg Forst Jagdztg* 157, 10, 199-196, 11, 214-222
- Phares RE (1978) Research gaps and research needs for the eastern forest regions. In: *Un-even-aged silviculture and management in the United States*. US Dep Agric, For Serv Timber Management Research. Washington, DC Gen Tech Rep WO-24, 152-157
- Pobedinski AV (1988) Comparative evaluation of even-aged and uneven-aged stands (in Russia). *Lesnoe Khozyaistvo* 2, 40-43
- Pretzsch H (1985) Die Fichten-Tannen-Buchen-Plenterwaldversuche in den ostbayerischen Forstämtern Freyung und Bodenmais. *Forst-archiv* 56, 3-9
- Pukkala T, Kolström T (1988) Simulation of the development of Norway spruce stands using a transition matrix. *For Ecol Manage* 25, 255-267
- Sarvas R (1944) Tukkipuun harsintojen vaikutus Etelä-Suomen yksityismetsiin. Referat: Ein-wirkung der Sägestamplenterungen auf die Privatwälder Südfinnlands. *Commun Inst For Fenn* 33, 1-268
- Schütz JP (1969) Étude des phénomènes de la croissance en hauteur et en diamètre du sapin (*Abies alba* Mill) et de l'épicéa (*Picea abies* Karst) dans deux peuplements jardinés et une forêt vierge. Supplément aux organes de la Société forestière suisses, 44, 1-114

- Sirén G (1955) The development of spruce forest on raw humus sites in northern Finland and its ecology. *Acta For Fenn* 62 (4), 1-408
- Smith DM (1962) *The practice of silviculture*. John Wiley and Sons. New York, 578 p
- Smith DM (1972) The continuing evolution of silvicultural practice. *J For* 70 (2), 89-92
- Smith DM (1975) Dangers in over-simplification of forestry. *Maine For Rev* 8, 3-4
- Smith HC, DeBald PS (1978) Economics of even-aged and uneven-aged silviculture and management in eastern hardwoods. In: *Uneven-aged silviculture and management in the United States*. US Dep Agric, For Serv Timber management research. Washington, DC Gen Tech Rep WO-24, 125-141
- Solomon DS, Hosmer RA, Hayslett HT Jr (1986) A two-stage matrix model for predicting growth of forest stands in the northeast. *Can J For Res* 16 (3), 521-528
- Szwagrzyk J (1992) Small scale spatial patterns of trees in a mixed *Pinus sylvestris* – *Fagus sylvatica* forest. *For Ecol Manage* 51, 301-315
- Trimble GR Jr (1961) Managing mountain hardwoods – a ten-year appraisal. US Dep Agric, For Serv NEFES Pap 143, 25 p
- Trimble GR Jr (1968) Research findings on even-aged management in hardwoods in the Appalachian area of the Central States. Monongahela NF Timber Management Conf Proc, 15 p
- Trimble GR Jr (1970) 20 years of intensive uneven-aged management. Effect on growth, yield, and species composition in two hardwood stands in West Virginia. US Dep Agric, For Serv Res Pap NE-154, 12 p
- Trimble GR Jr, Manthy RS (1966) Implications of even-aged management on timber. In: Trends from selection cutting to even-aged management. Soc Am For, Allegheny Sect Proc 1965, 62-75
- Trimble GR Jr, McClung L (1966) Clearcutting profitable. *WV Conserv* 30, 2-4
- Uppskattning av Sveriges skogstillgångar verkställd åren 1923-1929 (1932) Redogörelse avgiven av riksskogstaxeringsnämnden. Del I. *Statens Offentliga Utredningar* 26, 1-254
- Viitala R (1986) Skandinavien eri-ikäisrakenteista metsistä. In: Eriakenteisen metsän hoidosta (R Viitala, ed). Metsäntutkimuslaitoksen tiedonantoja 219. *The Finnish Forest Research Institute. Department of Silviculture. Research Papers* 219, 49-77
- Vrablec JJ (1977) Public lands – from concept to practice. US Dep Agric, For Serv Gen Tech Rep NE-29, 77-81
- Vuokila Y (1970) Harsintaperiaate kasvatushakkuissa. Summary: Selection from above in intermediate cuttings. *Acta For Fenn* 110, 1-45
- Walker N (1956) Growing stock volumes in unmanaged and managed forests. *J For* 54, 378-383