



# Dating natural gaps in the holm oak forest (*Quercus ilex* L) in Fango MAB Reserve (Corsica) by reading rings of maquis components

C Panaïotis, R Loisel, G Paradis

## ► To cite this version:

C Panaïotis, R Loisel, G Paradis. Dating natural gaps in the holm oak forest (*Quercus ilex* L) in Fango MAB Reserve (Corsica) by reading rings of maquis components. *Annales des sciences forestières*, 1995, 52 (5), pp.477-487. <hal-00883015>

**HAL Id: hal-00883015**

**<https://hal.science/hal-00883015v1>**

Submitted on 11 May 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

Original article

# Dating natural gaps in the holm oak forest (*Quercus ilex* L) in Fango MAB Reserve (Corsica) by reading rings of maquis components

C Panaïotis <sup>1</sup>, R Loisel <sup>2</sup>, G Paradis <sup>1</sup>

<sup>1</sup> Université de Corse, Cevaren, botanique et écologie, BP 52, 20250 Corte;

<sup>2</sup> Faculté des sciences et techniques, laboratoire de botanique et d'écologie méditerranéenne,  
case 461, 13397 Marseille cedex 13, France

(Received 15 July 1994; accepted 2 December 1994)

**Summary** — This work is related to the study of natural gaps in a *Quercus ilex* Mediterranean forest in Corsica, France. The aim was to find a way to precisely date the *volis* to obtain a chronological sample which corresponded to the time of vegetation opening due to the fall of a holm oak and up to complete vegetation closure. The best marker appeared to be *Phillyrea latifolia*, a typical maquis tree found in association with the holm oak, which has a high capacity to produce stem sprouts after the breakage. In order to date the *chablis*, rings from the stem sprouts must be counted and a careful observation of the other species must be made to confirm results.

**Corsica / *Quercus ilex* / maquis / natural gap / dendrochronology**

**Résumé** — Datation des trouées naturelles en forêt de chênes verts (*Quercus ilex* L) dans la réserve MAB du Fango (Corse) par la lecture des cernes des essences du maquis. Ce travail a trait à l'étude des trouées naturelles dans une forêt méditerranéenne à *Quercus ilex* en Corse. Il s'agit de trouver un moyen de dater précisément les *volis* afin d'obtenir un échantillonnage chronologique, correspondant à l'ouverture de la végétation, par chute d'un vieux chêne vert, jusqu'à la fermeture complète du milieu. Le meilleur marqueur de ces événements se révèle être *Phillyrea latifolia*, arbre du maquis en association avec le chêne vert, qui présente une excellente capacité à rejeter de tige après cassure. Il suffit de compter les cernes de ces rejets pour dater les *chablis*, tout en prenant en compte les autres espèces du maquis pour confirmer les résultats.

**Corse / *Quercus ilex* / maquis / trouée naturelle / dendrochronologie**

## INTRODUCTION

Gaps from natural treefalls have already been studied: (i) in tropical forests (Oldeman, 1974; Florence, 1981; Huc and Rosalina, 1981; Rollet, 1983; Riéra and Alexandre, 1988; Hartshorn, 1989); and (ii) in temperate forests (Falinski, 1977; Walter, 1979a, 1979b; Faille *et al*, 1984a, 1984b; Collins and Pickett, 1987; Koop and Hilgen, 1987; Lemée *et al*, 1991). However, no work has yet been performed on Mediterranean forests, in particular holm oak forests (*Quercus ilex* L.).

In the Mediterranean basin, the frequency and intensity of disturbances over the centuries (fires, firewood cutting, coal mining, grazing, *etc*) have made it impossible to study large modern forest areas, which have evolved naturally over a long period of time. Barbéro (1990) is the only author who noted the existence of *chablis* in southern France and regeneration in these natural openings.

*Chablis* is defined as (Oldeman, 1990) "the uprooting of a tree, the uprooted tree, the inaccessible heap of broken or surviving vegetation and the branches, the opening (gap) in the forest canopy." The author distinguishes it from the *volis* which is "the breaking of a tree trunk (most often by storm), the broken and fallen upper part of the tree, the mass of vegetation and branches and the gap».

This study focused on natural gaps. It is rarely possible to date gaps directly.

Faille *et al* (1984a, b) and Koop and Hilgen (1987) used a natural known disturbance (hurricane of 1967) as a base and dated their *chablis* before or after this event by comparing trunk decomposition on the ground.

In tropical forests, settling by opportunistic species creates lines of seeds along the trunks on the ground. Brokaw (1982) sug-

gested that new tree age class then appears. Dating the event seems to cause problems, but no explanation was provided on the method used. Riéra (1986) stated that the estimation of the *chablis* age is quite difficult. Rollet (1983) considered 4 age classes for gaps: "very recent, recent, old and very old", without explaining his criteria for identifying these classes.

A synchronistic analysis of natural gaps at different ages (from vegetation opening to complete vegetation closure) was undertaken. The purpose of this work was to discover 1 or more markers which made it possible to precisely date the event, to study natural regeneration in holm oak forests in the meso-Mediterranean strata.

## REGION AND STUDY AREA

This study was performed in the Fango Valley (Haute-Corse), an area classified as a man and biosphere reserve since 1977 (Viale and Frontier, 1979) due to the ancient nature of the holm oak forests stands.

The valley extends from the Paglia Orba (altitude 2 525 m) to the sea (approximately 10 km). The climate is subhumid Mediterranean with an average annual rainfall of 720 mm (forest ranger lodge, altitude 192 m) and an average annual temperature of 14.6°C. On the shady side of the valley, the state forest covers an area of 4 318 ha (fig 1).

The forest lies essentially on rhyolites (Vellutini, 1973). The studied forest has not been exploited since 1827 (ONF, 1992), in the Perticato district. *Volis* are principally located in the lower part of the vale, on a 15 ha area with an altitude ranging from 300 to 450 m. The old holm oak forest is at least 150 years old and it grows on brown acid soil (Roche and Roux, 1976).




Fig 1. Localisation of the state forest of the Fango.

## MATERIALS AND METHODS

The forest structure consists of a mosaic of large holm oaks (cover from 50 to 75%; dbh from 0.40 to 1 m; height from 15 to 20 m) and clumps (total density: 460 t/ha;  $g = 28 \text{ m}^2/\text{ha}$ ) with a high maquis below (height to 7 m) (M'hamedi, 1994; Panaiotis, 1994).

The age of these *Quercus ilex* causes *volis*, which are often single. Ten *volis* were selected, located in areas with relatively similar ecological conditions. Their dating was first determined by the decomposition state of the wood. We wanted to obtain the whole range of possible ages: from the gap of the 1st year to the period of complete vegetation closure, when trunks on the ground are entirely decomposed.

After localization of these differently aged gaps, we noticed the very high capacity of some maquis species to produce sprouts from a previous breakage. This "gap filling" can occur in different ways (figs 2, 3, 4).


In the forest, samples are cut at the base of these sprouts (figs 2, 3, 4: s is the collected stem section). The samples were cut again in the laboratory to obtain small 1 to 1.5 cm thick rounds. They were pumiced with very thin sandpaper to allow ring reading with a focusing glass.

## RESULTS AND DISCUSSION

The number of rings in each sample makes it possible to estimate their age (table I) using classic dendrochronology techniques (Martin, 1974).

Before observing sprouting, *Fraxinus ornus* L was the intended age marker in these gaps. This hypothesis was based on the fact that this tree acts as an opportunistic species, and would thus take advantage of these natural clearings to root in large quantities due its anemophilous nature.


Nevertheless, the surface of these clearings was not large enough (120 m<sup>2</sup> at the largest) to permit real settling to occur. Furthermore, it was often present in the undergrowth (see table III, where plot 4C corresponds to the control forest plot without canopy opening). It characterizes a phytosociological subassociation with the holm oak: the *Quercetum ilicis gallo provinciale*.




**Fig 2.** Recent *volis* (5 years). Holm oak fall has induced the breakage of *Phillyrea latifolia* stems. Lightning favors the emergence of sprouts at the break. Cuts are taken at the base of these sprouts. s is the section of the stems collected. The number of rings in this section yields *volis* age.

**Table I.** Age (years), diameter (cm) and height (cm) of the different stems harvested in the *volis*.

	Age	Diam	Height		Age	Diam	Height
Volis I				Volis V			
<i>Fraxinus ornus</i> 1	7	0.8	117	<i>Fraxinus ornus</i> 1	40	1.4	520
<i>Fraxinus ornus</i> 2	13	1.2	178	<i>Fraxinus ornus</i> 2	14	1.2	120
<i>Fraxinus ornus</i> 3	16	1.2	173	<i>Phillyrea latifolia</i> 1	16	3.1	450
<i>Phillyrea latifolia</i> 1	7	1.5	280	<i>Phillyrea latifolia</i> 2	16	3.7	410
<i>Phillyrea latifolia</i> 2	8	1.3	90	Volis VII			
<i>Phillyrea latifolia</i> 3	7	0.8	80	<i>Fraxinus ornus</i>	17	1.6	290
<i>Phillyrea latifolia</i> 4	8	2.2	90	<i>Phillyrea latifolia</i> 1	18	4.2	400
Volis II				<i>Phillyrea latifolia</i> 2	18	3.7	350
<i>Fraxinus ornus</i> 1	6	0.7	125	<i>Phillyrea latifolia</i> 3	18	3.5	450
<i>Fraxinus ornus</i> 2	5	0.8	70	<i>Quercus ilex</i>	17	1.8	250
<i>Phillyrea latifolia</i> 1	6	1.1	205	<i>Arbutus unedo</i>	18	2.8	310
<i>Phillyrea latifolia</i> 2	6	1.2	160	Volis VIII			
<i>Arbutus unedo</i>	6	1.6	70	<i>Fraxinus ornus</i> 1	13	1.1	230
Volis III				<i>Fraxinus ornus</i> 2	13	1.5	260
<i>Fraxinus ornus</i>	26	4.4	500	<i>Phillyrea latifolia</i> 1	6	1.4	230
<i>Phillyrea latifolia</i> 1	14	2.5	350	<i>Phillyrea latifolia</i> 2	18	3.1	440
<i>Phillyrea latifolia</i> 2	14	2.2	360	<i>Phillyrea latifolia</i> 3	6	1.6	210
<i>Viburnum tinus</i>	14	1.4	210	Volis IX			
<i>Quercus ilex</i>	14	1.1	120	<i>Phillyrea latifolia</i> 1	4	1.2	180
Volis IV				<i>Phillyrea latifolia</i> 2	2	0.4	45
<i>Fraxinus ornus</i> 1	12	1.3	140	<i>Phillyrea latifolia</i> 3	2	0.6	90
<i>Fraxinus ornus</i> 2	14	0.8	138	Volis X			
<i>Phillyrea latifolia</i> 1	12	2.8	400	<i>Phillyrea latifolia</i> 1	19	2.8	470
<i>Phillyrea latifolia</i> 2	12	2.2	330	<i>Phillyrea latifolia</i> 2	19	3.3	310
<i>Phillyrea latifolia</i> 3	11	3.1	390	<i>Phillyrea latifolia</i> 3	19	4.4	520

**Fig 3.** Middle-aged *volis* (12 years). Holm oak fall has induced the breakage of *Phillyrea latifolia* stems. Lightning favors the emergence of sprouts at the break. Cuts are taken at the base of these sprouts. *s* is the section of the stems collected. The number of rings in this section yields *volis* age.



**Fig 4.** Old *volis* (16 years). Holm oak fall has induced the breakage of *Phillyrea latifolia* stems. Lightning favors the emergence of sprouts at the break. Cuts are taken at the base of these sprouts. *s* is the section of the stems collected. The number of rings in this section yields *volis* age.

*ornetosum* (Allier and Lacoste, 1980; Gamisans, 1991).

In spite of its deciduous nature, the ash tree presents some false and double rings. It can wait for a long time for a favorable moment for upward growth (an opening in the canopy). It sometimes infiltrates through the maquis to reach the sunlight (see table I, *volis* V, a 40-year-old shoot with a diameter of 1.4 cm). Consequently, the rings are very close together and hard to distinguish. Nevertheless, one can observe a clear increase in the 1st rings corresponding to a sudden outburst of sunlight (Lémée, 1985) (table I: *volis* III, the sample of *Fraxinus ornus* shows widely spaced rings at 14 years old). The dating of the gaps cannot only be

performed by reading *Fraxinus ornus* rings; this tree could be used as an indirect temporal marker.

Dating based on sprouts from *Phillyrea latifolia* L stems yielded results which were consistent with the state of decomposition of the trees lying on the ground. Rings for *Phillyrea latifolia* stems are well-defined. The new sprout takes advantage of the whole shrub root system and of the intense lighting available through the gap. The diameter increase is important enough to localize the sequence corresponding to 1-year rings: this species presents double rings due to several annual growth periods.

Estimation of *volis* age was essentially based on *Phillyrea latifolia*, which seems to

**Table II.** Estimation of *volis* ages (in years).

<i>Volis</i>	VI	IX	II	I	IV	III	V	VII	VIII	X
Age	0	2	5	7	12	14	16	17	18	19

respond best to shoot breakage caused by holm oak fall (table II).

*Volis VI* corresponds to a holm oak fall in the spring 1993. The delay before *Phillyrea latifolia* emits a stem probably depends on the season during which the *volis* occurs. It can be estimated to within 1 year at the longest, when growth the following spring corresponds to the "naught point" of rings. The 1st visible ring corresponds to the 1st year of sprout growth. Initial *P. latifolia* sprout rings are better defined than seedling rings (of *Quercus ilex*, for example) which are hard to date at 1 year. At *volis IX*, sample 1 in table I is 4 years old. The 2 other samples (2 and 3) are 2 years old. Thus, the opening is 2 years old, as further shown by the closing rate and the decomposition rate of the trunk compared to others. Sample 1 comes from a preexistent bough.

*Volis V* roughly indicates the closing period of these gaps. Sixteen years seem to be necessary for gap cicatrization in *volis* under 100 m<sup>2</sup>, with a closing rate of 95% (the closing rate corresponds to the vertical crown projection of the forest strata [A1;A2] and of the higher shrub strata [a1]) (see table III).

*Volis VII* presents almost no evidence of decayed holm oak wood on the ground. Twenty years seem to be necessary for trunk decomposition in the biotic and abiotic conditions of the vale.

The other maquis species (with the exception of *Phillyrea latifolia*) occasionally yield complementary information to help estimate more precisely *volis* age.

– *Erica arborea* L always breaks in the stem. The sprouts are emitted from the stump. These sprouts are not good markers because they do not necessarily occur at the same time as opening. They can appear after the main stem droops due to insufficient light.

– *Arbutus unedo* L sprouts well from the stem but does not occur as frequently as *Phillyrea latifolia* (see table III). The probability for this species to be damaged by the oak fall is therefore low. Nevertheless, it is a good potential temporal marker (dating to be compared with the one provided by *Phillyrea*).

– *Viburnum tinus* L is rare in the undergrowth (table III). It is not a good marker: its stems bend down but do not break due to small diameter. Only 1 sample was collected in the 10 *volis*.

– *Quercus ilex* is often present in shrub from near the *volis*. Cutting it would certainly yield interesting information (increase in ring growth), but this would eliminate the only study element (sometimes 2 or 3 individuals).

Tree penetration with the increment borer is often very difficult and ring reading is increasingly difficult due to tannins (Zhang, 1987).

The seedlings, sometimes numerous, are anterior to the tree fall. The acorns present at the time of the event (seed bank), or posterior to the fall (mast coming from closely seed trees), probably benefit from the light to germinate or grow faster (currently under study). Consequently, they cannot provide precise information for *volis* dating.

## CONCLUSION

*Phillyrea latifolia* is the best marker for natural gaps occurring after the fall of large holm oaks in the Fango forest. This species has numerous good features:

– It is the maquis species which skirts the holm oak in dense forests the longest, due to its forest behavior and its capacity to grow higher than *Arbutus unedo* and *Erica arborea*.



Table III. Phytosociological relevés of 10 *volis* (October 93) (relevés 1 to 10) and of the forest control plot (May 93) (relevé 11).

Relevés (table) Relevés ( <i>volis</i> )	1 VI	2 IX	3 II	4 I	5 IV	6 III	7 V	8 VII	9 VII	10 X	11 4C
Tree cover (+10 m)	—	—	20	20	20	—	—	5	10	50	50
Tree cover (10–5 m)	15	—	5	—	20	60	20	75	10	40	—
Shrub cover (5–1.5 m)	10	30	20	10	30	30	80	10	40	10	60
Shrub cover (1.5–0.3 m)	10	40	5	10	20	20	5	10	10	5	5
Herb cover (<0.3 m)	5	20	5	5	5	5	1	10	10	5	5
Moss cover	10	25	5	5	30	10	20	40	40	60	10
Litter cover	70	30	50	50	40	50	50	40	80	70	60
Rocky	*	*	—	—	*	—	—	—	—	**	*
Stony	**	*	**	**	**	**	**	*	**	**	**
Shingly	*	—	*	***	*	*	*	—	**	—	*
Area (m <sup>2</sup> )	120	110	170	90	100	160	95	95	130	88	225
Slope (°)	35	20	30	35	35	35	35	35	35	30	40
Exposure	nw	nw	wnw	w	nw	w	nw	nw	nw	nw	wnw
Altitude (m)	345	310	315	410	355	320	315	345	395	380	315
Volis height (m)	0.8	0.2	1.5	1.5	0.2	1.5	2	0	1.3	0	—
Diameter of the trunk basis (m)	0.82	0.28	0.89	0.70	0.70	0.76	0.60	0.73	0.89	0.57	—
Closing rate (%)	10	20	25	10	50	70	95	95	75	100	—
Age of the <i>volis</i> (in years)	0	2	5	7	12	14	16	17	18	19	—
Max height (in m)	12	6	15	10	10	10	10	15	10	15	15
Mean shrub height (m)	0.5	5	3	1.2	3	5	5	5	3.5	8	5
Number of species	13	15	11	11	14	11	8	15	13	11	14
<b>Trees</b>											
<i>Quercus ilex</i>	—	—	2.1	2.1	2.1	—	—	1.1	2.2	3.3	3.3
<i>Quercus ilex</i>	2.3	—	1.1	—	1.1	1.2	2.1	1.1	2.1	—	—
<i>Fraxinus ornus</i>	—	—	—	—	+	—	—	—	—	—	—
<i>Phillyrea latifolia</i>	—	—	—	—	2.2	3.2	2.1	2.2	—	1.3	—
<i>Arbutus unedo</i>	—	—	1.1	—	1.1	2.2	—	2.3	—	2.1	—
<i>Erica arborea</i>	—	—	+	—	—	—	—	2.3	—	2.2	—
<b>Shrubs</b>											
<i>Arbutus unedo</i>	—	—	2.2	—	2.2	+	3.2	—	2.2	—	2.2
<i>Arbutus unedo</i>	—	1.1	1.1	+	1.1	—	—	—	1.2	—	1.1
<i>Erica arborea</i>	—	1.2	—	—	1.1	+	2.2	+	1.1	—	2.3
<i>Erica arborea</i>	+	+	+	2.1	+	—	+	+	+	—	1.1

<i>Phillyrea latifolia</i>	a1	1.1	2.3	2.3	1.1	1.1	1.2	2.3	2.2	1.1	2.3	2.2	2.1
<i>Phillyrea latifolia</i>	a2	1.1	2.2	1.1	2.1	2.1	2.1	2.3	—	1.1	1.2	1.1	+
<i>Quercus ilex</i>	a1	+	—	—	2.1	—	+	1.1	—	+	+	+	—
<i>Quercus ilex</i>	a2	+	+	—	—	—	1.2	1.1	1.1	1.1	+	1.3	—
<i>Smilax aspera</i>	a1	—	1.2	+	—	—	2.3	2.4	2.4	+	1.1	+	1.3
<i>Smilax aspera</i>	a2	1.1	1.3	+	—	—	2.2	2.2	—	+	1.1	—	+
<i>Rubus ulmifolius</i>	a1	—	1.3	1.2	—	—	—	—	—	2.2	—	—	1.2
<i>Rubus ulmifolius</i>	a2	—	3.3	1.1	+	—	1.1	—	—	1.1	—	—	+
<i>Fraxinus ornus</i>	a1	+	—	—	—	—	—	—	+	+	—	—	—
<i>Fraxinus ornus</i>	a2	+	1.1	+	1.1	1.1	1.1	1.1	1.1	—	+	—	1.1
<i>Viburnum tinus</i>	a2	+	—	—	—	+	+	+	—	—	—	—	—
<i>Rubia peregrina</i>	a2	—	—	—	—	—	—	—	—	—	—	—	—
<i>Hedera helix</i>	a1	—	—	—	—	—	—	—	—	+	—	—	—
<i>Myrtus communis</i>	a1	—	—	—	—	—	—	—	1.1	—	—	—	—
<b>Herbs</b>													
<i>Quercus ilex</i>		1.1	+	1.1	1.1	1.1	1.1	—	+	+	1.1	1.1	1.1
<i>Fraxinus ornus</i>		+	+	1.1	1.1	1.1	1.1	1.1	+	1.1	1.1	+	+
<i>Phillyrea latifolia</i>		+	+	+	+	+	+	—	+	+	+	+	1.1
<i>Arbutus unedo</i>		+	+	+	+	+	—	—	+	+	+	+	+
<i>Erica arborea</i>		+	+	—	+	+	1.1	+	+	+	+	+	+
<i>Viburnum tinus</i>		+	1.1	+	—	—	—	—	—	—	—	—	—
<i>Juniperus oxycedrus</i>		—	+	—	—	—	—	—	—	—	—	—	—
<i>Rubia peregrina</i>		—	1.2	+	+	+	+	+	+	1.2	+	—	1.1
<i>Smilax aspera</i>		+	1.2	+	—	—	+	1.1	+	1.1	+	—	+
<i>Rubus ulmifolius</i>		—	+	+	—	—	+	+	—	+	+	—	—
<i>Hedera helix</i>		—	—	—	—	—	—	—	+	+	—	—	—
<i>Galium rotundifolium</i>		+	—	—	—	—	+	1.2	1.2	1.2	1.2	+	1.2
<i>Cyclamen repandum</i>		—	—	—	+	—	—	—	—	—	—	—	—
<i>Cyclamen hederifolium</i>		—	—	—	—	—	—	—	—	—	—	—	—
<i>Brachypodium retusum</i>		—	1.2	—	1.1	—	—	—	—	—	—	—	—
<i>Ruscus aculeatus</i>		+	—	—	—	—	—	—	—	—	—	—	—
<i>Hieracium gr. pilosella</i>		—	—	—	—	—	—	—	—	+	—	—	—
<i>Selaginella denticulata</i>		—	—	+	—	—	—	—	—	—	—	—	—
<i>Polypodium cambricum</i>		+	+	—	—	—	+	—	—	+	+	+	+
<i>Asplenium adiantum-nigrum</i>		+	+	—	+	+	—	+	—	+	+	+	+
<i>Asplenium trichomanes</i>		+	—	—	+	—	—	—	—	—	—	1.2	—
Moss		2.3	2.3	2.3	2.2	2.3	2.3	3.3	1.3	2.3	3.4	4.4	2.3

Abund-dominance: + rare; 1 < 5%; 2, 5 to 25%; 3, 25 to 50%; 4, 50 to 75%; 5, > 75%. Sociability: 1 individually; 2 in group; 3 in troop; 4 small colonies; 5 in crops. Stones: — none; \* < 10%; \*\* 10 to 25%; \*\*\* 25 to 50%; \*\*\*\* > 50%.

- Its nearly systematic presence gives it more of a chance to be damaged.
- Its exceptional stem sprouting capacity yields the “naught point” of the gap.
- The smooth aspect of the sprout bark compared to the primary trunk makes them easy to recognize even 20 years after the event.
- Finally, its rings are very well-defined.

Nevertheless, more sampling and data from other species (*Arbutus unedo*, *Fraxinus ornus*, *Quercus ilex*) would be necessary to confirm these results.

The precise dating of the *volis* in the Mediterranean forest is possible due to the excellent sprouting capacity of its species, especially *Phillyrea latifolia*. The *volis* can be accurately dated to within approximately 1 year, which is in accordance with results obtained in other forests.

The study of natural gaps in the Fango MAB Reserve may make it possible for us to better understand the dynamics of holm oak forests, which have been disturbed over the centuries.

## ACKNOWLEDGMENTS

This work is part of a global research program on the study of the potential regeneration of aging holm oak stands (*Quercus ilex* L) in Corsica (Fango Man & Biosphere Reserve) supported by the Ministry of Environment (SRETIE-EGPN).

Thanks are due to the Regional Natural Park of Corsica for their participation in this project. We are grateful to I Barros who improved the English.

## REFERENCES

- Allier C, Lacoste A (1980) Maquis et groupements végétaux de la série du chêne vert dans le bassin du Fango (Corse). *Ecologia Mediterranea* 5, 59-82
- Barbéro M (1990) Méditerranée : bioclimatologie, sclérophylle, sylvigénèse. *Ecologia Mediterranea* XVI, 1-12
- Brokaw NVL (1982) The definition of tree fall gap and its effects on measures of forest dynamics. *Biotropica* 14, 158-160
- Collins BS, Pickett STA (1987) Influence of canopy opening on the environment and herb layer in a northern hardwoods forest. *Vegetatio* 70, 3-10
- Faillie A, Lémée G, Pontailier JY (1984a) Dynamique des clairières d'une forêt inexploitée (réserves biologiques de la forêt de Fontainebleau). I. Origine et état actuel des ouvertures. *Acta Oecologica, Oecol Gener* 5, 35-51
- Faillie A, Lémée G, Pontailier JY (1984b) Dynamique des clairières d'une forêt inexploitée (réserves biologiques de la forêt de Fontainebleau). II. Fermeture des clairières actuelles. *Acta Oecologica, Oecol Gener* 5, 181-199
- Falinski JB (1977) Research on vegetation and plant population dynamics conducted by the Białowieża Geobotanical Station of the Warsaw University in the Białowieża primeval forest 1952-1977. *Phytocoenosis* 6, 148 p
- Florence J (1981) Chablis et sylvigénèse dans une forêt dense humide sempervirente du Gabon. Thèse, univ Strasbourg, 261 p
- Gamisans J (1991) *La végétation de la Corse. Compléments aux prodrome de la flore de la Corse* (D Jeanmonod, HM Burdet, eds), Conservatoire et Jardin botanique de Genève, 391 p
- Gamisans J, Jeanmonod D (1993) *Catalogue des plantes vasculaires de la Corse. Compléments aux prodrome de la flore de la Corse* (D Jeanmonod, HM Burdet, eds), Conservatoire et Jardin Botanique de Genève, 258 p
- Hartshorn GS (1989) Gap-phase dynamics and tropical tree species richness. In: *Tropical forests, botanical dynamics, speciation and diversity* (LB Holm-Nielsen, IC Nielsen, H Baslev, eds), Academic Press, New York, 380 p
- Huc, Rosalina (1981) Chablis and primary forest dynamics in Sumatra. *Biotrop TFR* 1-2-2, Bogor, Indonesia, 13 p
- Koop H, Hilgen P (1987) Forest dynamics and regeneration mosaic shifts in unexploited beech (*Fagus sylvatica*) stands at Fontainebleau (France). *For Ecol Manage* 20, 135-150
- Lémée G (1985) Rôle des arbres intolérants à l'ombrage dans la dynamique d'une hêtraie naturelle (forêt de Fontainebleau). *Acta Oecologica, Oecol Plant* 6, 3-20
- Lémée G, Faillie A, Pontailier JY (1991) Dynamique linéaire et cyclique d'une forêt inexploitée : cas des réserves biologiques de la forêt de Fontainebleau. *Coll Phytosocio (Berlin)* XX, 273-282
- Martin P (de) (1974) *Analyse des cernes. Dendrochronologie et dendroclimatologie*. Masson, Paris, France, 78 p

- M'hamed M (1994) Contribution à l'étude des potentialités de régénération du chêne vert (*Quercus ilex* L) dans les trouées naturelles – réserve MAB du Fango (Haute-Corse). DESS, univ Corse, Corsica, France, 70 p
- Oldeman RAA (1974) *L'architecture de la forêt guyanaise*, mém Orstom n° 73, Orstom, Paris, France, 214 p
- Oldeman RAA (1990) *Forests: elements of sylvology*. Springer-Verlag, Berlin, 624 p
- ONF (1992) Historique de la forêt domaniale du Fango (de 1827 à nos jours). Doc interne, Bastia, 26 p, cartes
- Panaïotis C (1994) Diversité structurale des formations forestières à chêne vert (*Quercus ilex* L) et des maquis de la forêt domaniale du Fango (réserve de la biosphère). *Trav Sci Parc Nat Rég Rés Nat Corse* 48, 1-68
- Riéra B (1986) À propos des chablis en forêt guyanaise. Piste Saint-Elie. *Mém du Mus Nat Hist Nat Série A*, Paris, France, 132, 109-114
- Riéra B, Alexandre DY (1988) Surface des chablis et temps de renouvellement en forêt dense tropicale. *Acta Oecologica, Oecol Gener* 9, 211-220
- Roche D, Roux C (1976) *Les sols d'une séquence bioclimatique méditerranéenne montagnarde en Corse cristalline*. Mém DEA, CEPE-CNRS Montpellier, France, 112 p
- Rollet B (1983) La régénération naturelle dans les trouées : un processus général de la dynamique des forêts tropicales humides. *Bois et Forêts des Tropiques* 201, 3-34; 202, 19-24
- Vellutini PJ (1973) Étude géologique de la vallée du Fango (Corse). Les rhyolites ignimbritiques et leur substrat paléozoïque. Thèse doct d'État, Aix-en-Provence, France, 214 p, carte, annexes
- Viale D, Frontier S (1979) Synthèse et conclusions. In: *Projet de création d'une réserve de la biosphère dans la vallée du Fango (Haute Corse), Étude préalable*, APEEM, 4<sup>e</sup> partie, 21 p
- Walter JM (1979a) Étude des structures spatiales en forêt alluviale rhénane. I. Problèmes structuraux et données expérimentales. *Oecologia Plantarum* 14, 345-359
- Walter JM (1979b) Étude des structures spatiales en forêt alluviale rhénane. V. L'architecture forestière observée. *Oecologia Plantarum* 14, 401-410
- Zhang SH (1987) Contribution à l'étude de la croissance en diamètre du chêne vert (*Quercus ilex* L) en relation avec le climat. Thèse de doct, USTL de Montpellier, France, 183 p