



HAL
open science

Differential effects of N, P and K on photosynthesis and partitioning of N in *Pinus pinaster* needles

Charles R. Warren, John F. Mcgrath, Mark A. Adams

► **To cite this version:**

Charles R. Warren, John F. Mcgrath, Mark A. Adams. Differential effects of N, P and K on photosynthesis and partitioning of N in *Pinus pinaster* needles. *Annals of Forest Science*, 2005, 62 (1), pp.1-8. hal-00883856

HAL Id: hal-00883856

<https://hal.science/hal-00883856>

Submitted on 11 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Differential effects of N, P and K on photosynthesis and partitioning of N in *Pinus pinaster* needles

Charles R. WARREN^{a,b*}, John F. MCGRATH^c, Mark A. ADAMS^{a,b}

^a Department of Botany, University of Western Australia, Crawley WA 6009, Australia

^b Forest Science Centre, The University of Melbourne/Natural Resources and Environment, Water St, Creswick VIC 3363, Australia

^c Department of Conservation and Land Management, 17 Dick Perry Avenue, Kensington WA 6152, Australia

(Received 10 June 2003; accepted 3 October 2003)

Abstract – We investigated the response of one-year-old *Pinus pinaster* Ait. growing on a sandy, nutrient-poor soil to the factorial addition of N, P and K. We tested two hypotheses: (1) fertiliser application increases rates of light-saturated photosynthesis (A_{\max}) and leaf-level water-use efficiency (WUE, as indicated by carbon isotope composition), and (2) greater concentrations of N and Rubisco explain fertiliser-promoted increases in A_{\max} and WUE. Two years after fertiliser application, the height of *P. pinaster* was significantly greater in plots to which P was added (127 ± 7 cm; mean \pm SE of three replicate plots) compared to those without added P (70 ± 10 cm), but was unaffected by addition of N or K. The fraction of N present as Rubisco varied between 4 and 18% and was unrelated to N concentrations, but positively related to P and K concentrations. A_{\max} and $\delta^{13}\text{C}$ did not vary significantly among treatments and were unrelated to concentrations of Rubisco, N, P or K. We conclude that fertiliser application two years previously has little effect on current A_{\max} and $\delta^{13}\text{C}$ of *P. pinaster*, and thus the previously reported effects of fertiliser on gas exchange are likely ephemeral.

nitrogen / phosphorus / potassium / photosynthesis / rubisco

Résumé – Effets différentiels de N, P et K sur la photosynthèse et le fractionnement de N dans les aiguilles de *Pinus pinaster*. La réponse de plant de *Pinus pinaster* Ait., poussant sur un sol sableux et pauvre, a été étudiée en fonction de l'addition de N, P et K. Deux hypothèses ont été testées : (1) la fertilisation accroît le taux de saturation lumineuse de la photosynthèse (A_{\max}) et le niveau d'efficacité de l'eau (WUE, évalué par la composition isotopique du carbone), (2) une plus grande concentration de N et de rubisco explique que la fertilisation accroît A_{\max} et WUE. Deux années après l'application de la fertilisation, la hauteur de *P. pinaster* était significativement plus grande dans les placeaux où P avait été ajouté (127 ± 7 cm, moyenne \pm SE de 3 répétitions), comparativement à ceux sans addition de P (70 ± 10 cm) mais n'était pas affectée par l'addition de N et K. La fraction de N présente sous la forme de rubisco variait entre 4 et 18 % et n'était pas expliquée par les concentrations en N, mais était positivement expliqué par les concentrations en P et K. A_{\max} et $\delta^{13}\text{C}$ ne variaient pas significativement avec le traitement et n'était pas reliés aux concentrations en rubisco, N, P ou K. On conclue que l'application d'un fertilisant deux ans auparavant a peu d'effet sur les valeurs actuelles de A_{\max} et $\delta^{13}\text{C}$ de *P. pinaster* et alors les effets rapportés auparavant des fertilisants sur les échanges gazeux sont probablement éphémères.

azote / phosphore / potassium / photosynthèse / rubisco

Abbreviations: A_{\max} , maximum rate of light-saturated photosynthesis at ambient CO_2 ; CE, capillary electrophoresis; C_i , intercellular CO_2 concentration; Chl, chlorophyll; g_s , stomatal conductance; LAI, leaf area index; PPFD, photosynthetic photon flux density; Rubisco, ribulose-1,5-bisphosphate carboxylase/oxygenase; WUE, water-use efficiency; $\delta^{13}\text{C}$, stable carbon isotope composition.

1. INTRODUCTION

Application of fertiliser often increases the growth of *Pinus pinaster* Ait. and other conifers [24, 38]. The mass and area of foliage increases following fertiliser application, and this partially explains the growth response [9, 46]. Fertiliser application also increases rates of photosynthesis in some cases [9, 47, 48], but not all [10, 51]. Water-use efficiency (WUE) can be increased by fertiliser addition [18, 21, 47] and on water-limited sites may be a further explanation of the growth response.

However, there are too few studies to unequivocally identify either increased rates of photosynthesis and/or greater WUE as primary causes of the growth response to fertiliser application for field-grown conifers.

Studies on plant nutrition often focus on nitrogen (N) since it is the nutrient required in the largest quantity and is generally the nutrient most likely to limit carbon gain [14]. Among species there is often a strong positive correlation between maximum rates of photosynthesis (A_{\max}) and N [16, 19, 56]. This relationship owes much to the large proportion of N (up to 75%)

* Corresponding author: crwarren@unimelb.edu.au

present in the chloroplasts, much of it in thylakoid membranes and soluble proteins of the Calvin cycle, particularly the enzyme Rubisco (ribulose-1,5-bisphosphate carboxylase/oxygenase; EC 4.1.1.39) [16]. Within conifers, N and photosynthesis are positively reported for some species [35, 39]. More generally, however, relationships between N and photosynthesis are less consistent for conifers than for non-conifers, and may be weak or even negative [33, 34, 41, 43, 45, 47, 52, 53].

Strong relationships between phosphorus (P) and A_{\max} have been observed for conifers, for example in *Pinus radiata* [15, 47], *P. pinaster* [6, 30] and *Pinus strobus* [43]. A partial explanation may lie in a positive relationship between the concentration of P and amount of Rubisco, as observed in *P. pinaster* [55] and in herbaceous species [11, 23, 26, 42]. In *Picea sitchensis*, Rubisco activity per unit leaf mass was increased by P addition [13]. Similarly, positive relationships between maximum rates of carboxylation (V_{cmax}) and P were reported for *P. pinaster* [30] and *Pinus taeda* [28].

Concentrations of other nutrients, such as potassium (K), may also be related to A_{\max} [4, 5, 22]. Growth is expected to improve in response to added K on some, especially sandy, soils owing to the high mobility of K in both soil and plant and generally large plant requirements. K is the most abundant univalent cation in plant cells and plays a significant part in regulating stomatal function [32]. Hence, one means by which K deficiency reduces photosynthesis is by decreasing stomatal conductance [40, 50]. More generally, transpiration increases (and WUE decreases) if K is in poor supply [7, 29]. In these cases, changes in stomatal conductance are not responsible for the decline in photosynthesis; instead the large K requirement for protein synthesis [27] is probably responsible for strong correlations between concentrations of K and proteins such as Rubisco [20].

It is possible that the responses of conifer growth to N, P and K fertiliser share a mechanistic basis in their effects on photosynthesis and Rubisco concentration. Most studies relating N, P and K nutrition to photosynthesis and Rubisco have focussed on a handful of well-studied herbaceous species. Of the few studies on conifers, none have included direct quantification of Rubisco. Previously we reported relationships among addition of P fertiliser, growth, and photosynthesis of *P. pinaster* [55]. Here we extend those observations using a factorial N, P, K fertiliser trial at the same site and with one-year-old *P. pinaster*. We measured the concentration of Rubisco and other major N fractions in seeking to confirm the putative common basis for relationships between N, P, K, A_{\max} and WUE (as estimated by stable carbon isotope composition [17]).

2. MATERIALS AND METHODS

2.1. Field site

The field study was conducted in south-west Western Australia, approximately 20 km north of Bullsbrook (latitude 31° 67' S, longitude 116° 02' E, 40 m above sea level) at a site we have described previously [55]. The region has a Mediterranean climate with cool, wet winters and hot, dry summers, in July (mid-winter), the mean daily temperature is 8.7 °C and the maximum is 17.6 °C. In January (mid-summer), the mean daily minimum temperature is 17.0 °C and the maximum is 33.1 °C. Mean annual rainfall at Bullsbrook is 692 mm, with 84% of the total annual rainfall between May and October. Potential annual

evaporation [8, 36, 44] is around 1700 mm. The soil at the site is a P-deficient yellow siliceous sand to a depth of at least 3 m. Prior to being planted with *P. pinaster*, the site was occupied by *Banksia* woodland typical of the many seasonally dry and nutrient-poor sites in this area.

In 1995 the site was cleared and planted with seedlings of *P. pinaster* at 1500 stems ha⁻¹. In August 1996, a fertiliser trial was established in a randomised complete block design with eight factorial combinations of N, P and K replicated in three blocks. Plots were around 0.06 ha. Fertiliser was applied by hand to the soil surface. Elemental rates of fertiliser application were: N 275 kg ha⁻¹ as urea, P at 140 kg ha⁻¹ as double superphosphate, and K at 248 kg ha⁻¹ as muriate of potash. All plots received Cu (2 kg ha⁻¹), Zn (4 kg ha⁻¹), Mn (2 kg ha⁻¹).

2.2. Height and photosynthesis measurements

In winter 1998, the height of all trees was measured to the nearest 5 cm. The maximum rate of light-saturated photosynthesis (A_{\max}) was measured on five trees per plot several weeks after the first substantial (> 50 mm) winter rainfall on a warm (c. 20 °C maximum) cloudless day. This period was chosen because photosynthesis was unlikely to be limited by either soil or atmospheric water deficits, and foliage nutrient concentrations are more stable than during the spring growth flush. Measurements were made with an open, infra-red gas analyser (LCA-4, Analytical Development Co, Hoddesdon, England) at 340–370 μmol mol⁻¹ CO₂, 20–25 °C leaf temperature and >1500 μmol m⁻² s⁻¹ PPFD. Two one-year-old fascicles (four needles) were laid across the leaf chamber. Photosynthesis, transpiration and C_i were allowed to stabilise before measurements were taken, this generally required about five minutes. Immediately following measurement of photosynthesis, foliage was dissected out of the leaf chamber for measurement of area and mass. Additional samples were collected from the five trees per plot used for photosynthesis measurement. One sample was promptly frozen and stored at -20 °C, another sample was dried and ground to a fine powder in a ball mill.

2.3. N and stable carbon isotope composition ($\delta^{13}\text{C}$)

Dried samples were analysed for N and $\delta^{13}\text{C}$ by ANCA-MS. Samples (5 to 6 mg dry mass) were combusted to CO₂ and N₂ in the presence of O₂ (Roboprep-CN, Europa Scientific, Crewe, UK) before passing into a mass spectrometer (Tracermass, Europa Scientific). $\delta^{13}\text{C}$ (in ‰ units) was calculated with respect to the PDB standard: $\{=[^{13}\text{C}/^{12}\text{C}_{\text{sample}}]/(^{13}\text{C}/^{12}\text{C}_{\text{standard}}) - 1\} \times 1000$.

2.4. P and K

Dried foliage was digested in sulphuric acid/hydrogen peroxide as described previously [2]. P was analysed colorimetrically by the ascorbic-reduced vanado-molybdophosphoric acid method [37]. K was determined by flame photometry.

2.5. Chlorophylls

Chlorophylls were extracted from frozen samples with *n*, *n*-dimethylformamide and analysed colourimetrically using the extinction coefficients of Wellburn [58].

2.6. Rubisco

Proteins were extracted from frozen samples, as described by Warren et al. [56]. Needles were ground to a fine powder in a slurry of acid-washed sand, insoluble polyvinylpyrrolidone (15% w/w) and extraction buffer (10 mL g⁻¹ FW) (80 mM Tris-HCl, 0.1 M β-mercaptoethanol, 2% (w/v) SDS, and 15% (v/v) glycerol). The extract was centrifuged for five minutes in a microfuge and the supernatant used for protein analysis. Proteins were denatured by heating at 95–100 °C for 10 min in a water bath. Benzoic acid was used as an internal reference. Capillary electrophoresis (CE) was performed with a Bio-Rad




Figure 1. Relationships between fertiliser treatment and (a) height of trees at age 3, (b) needle N concentration, (c) needle P concentration, (d) needle N:P ratio, (e) needle K concentration. Fertiliser was applied to one-year-old trees and their height was measured and foliage collected two years later. Data are means of three replicate plots. Error bars are ± 1 SE and Fisher's LSD (0.05) for fertiliser treatment is indicated.

3000 system (Bio-Rad, Hercules, CA, USA) controlled by a computer equipped with CE-3000 software (Bio-Rad). The separation of proteins was performed in CE-SDS protein run buffer (Bio-Rad) in an uncoated fused-silica capillary (50 μm i.d. \times 24 cm long, 19.4 cm effective length). Electrophoresis was conducted at 20 $^{\circ}\text{C}$ and a constant voltage of 15 kV. The detector was set at 220 nm and sample was injected electrophoretically at 10 kV for 5 s. The capillary was rinsed sequentially between successive electrophoretic runs with 0.1 M NaOH (90 s), 0.1 M HCl (60 s), CE-SDS protein run buffer (120 s). Dry weight fractions were measured on a separate subsample of needles and the concentration of Rubisco was calculated on a dry weight basis.

2.7. Statistical analysis

Characteristics were compared between nutrient treatments by three-way fully factorial ANOVA. Where treatment means are reported, they are the mean of three replicate plots ± 1 SE.

3. RESULTS

3.1. Growth

Two years after fertiliser was applied, the height of *P. pinaster* varied between 55 and 137 cm and was significantly greater ($P = 0.0003$) in plots receiving P (127 ± 7 cm; mean \pm SE of three replicate plots) than in those without added P (70 ± 10 cm) (Fig. 1a). Addition of N and/or K fertiliser did not affect height ($P > 0.05$).

3.2. Needle concentrations of N, P and K

Needle concentrations of N, P and K were significantly greater in plots to which that element was added compared to those not receiving fertiliser ($P < 0.05$) (Fig. 1). Addition of P fertiliser decreased concentrations of N ($P = 0.0005$), but increased concentrations of K ($P = 0.02$). Ratios of N:P were




Figure 2. The relationship between the percentage of N present as Rubisco and (a) needle N concentration, (b) needle P concentration, (c) needle K concentration. Fertiliser was applied to one-year-old trees and foliage was collected two years later. Each point is a single measurement.

significantly affected by addition of P fertiliser ($P = 0.0009$) (Fig. 1). In plots without added P, the N:P ratio was 19 or greater, whereas it was 14 or less in plots to which P was added. On the other hand, addition of neither N nor K affected N:P ratios ($P > 0.05$).

3.3. Needle concentrations of Rubisco and chlorophyll

The fraction of total N present as Rubisco varied between 4 and 18% and was unrelated to needle concentrations of N, but weakly and positively related to P ($P = 0.015$, $R^2 = 0.25$) and K ($P = 0.05$, $R^2 = 0.16$, Fig. 2). Similarly, concentrations of




Figure 3. The relationship between the ratio of Rubisco to chlorophyll and (a) needle N concentration, (b) needle P concentration, (c) needle K concentration. Fertiliser was applied to one-year-old trees and foliage was collected two years later. Each point is a single measurement.

Rubisco per unit chlorophyll varied between 4 and 14 mmol mol^{-1} and were unrelated to needle concentrations of N or K, and weakly related to P ($P = 0.04$, $R^2 = 0.18$, Fig. 3).

3.4. $\delta^{13}\text{C}$ and A_{max}

The $\delta^{13}\text{C}$ signature of *P. pinaster* needles did not vary between treatments ($P > 0.05$), and was unrelated to needle concentrations of N, P, K or Rubisco ($P > 0.05$, Fig. 4). There was no significant difference in specific leaf area (SLA, leaf area per unit dry mass) among treatments ($P > 0.05$; $\text{SLA} = 2.94 \pm 0.03 \text{ m}^2 \text{ kg}^{-1}$, mean \pm SE), and thus trends in A_{max} were the




Figure 4. Relationship between stable carbon isotope composition ($\delta^{13}\text{C}$) and (a) needle N concentration, (b) needle P concentration, (c) needle K concentration, and (d) Rubisco concentration. Fertiliser was applied to one-year-old trees and foliage was collected two years later. Each point is a single measurement.

same on mass and area bases. Data are presented solely on a mass basis for ease of comparison with nutrient concentration data. A_{max} varied between 33 and 74 $\text{nmol g}^{-1} \text{s}^{-1}$, and did not vary significantly among treatments ($P > 0.05$). A_{max} was unrelated to needle concentrations of N, P, K or Rubisco (Fig. 5).

4. DISCUSSION

Height growth of *P. pinaster* was severely limited by the supply of P, but not by N or K (Fig. 1a). That growth was limited by P is supported by the very high N:P ratio of 19 or greater in plots without added P [1, 25]. The absence of a significant N limitation finds further support in our observation that N:P ratios were unaffected by addition of N. Addition of P fertiliser increased height by almost 50%, whereas A_{max} was unaffected – an argument against any lasting role of A_{max} in the growth response. Concentrations of N, P, and K were maintained within a relatively small range, and there was no correlation between A_{max} and concentrations of N, P, or K (Fig. 5). These findings suggest that “fine-tuning” of growth with nutrient supply ensured that all needles produced were photosynthetically competent. A similar lack of photosynthetic response to nutrient addition has been reported in *Pseudotsuga menziesii* [10], *Pinus*

strobus [43], *Pinus taeda* [51] and *Pinus pinaster* [55]. Together these findings cast doubt on whether photosynthesis plays a general and/or lasting role in the response of growth to fertiliser addition. The absence of differences in WUE (as indicated by $\delta^{13}\text{C}$) is almost certainly a function of the similarity of A_{max} among treatments. We cannot exclude the possibility that small yet significant increases in WUE and/or A_{max} were not detected owing to the inherently high variability of A_{max} and WUE. Nevertheless, modelling studies with *P. radiata* suggest that increased rates of photosynthesis due to fertiliser explain only 10% of the increase of net photosynthesis within the canopy [31]. Increases in leaf area and changes in biomass partitioning probably explain the bulk of the growth response [9, 46].

Two years had elapsed since fertiliser was applied in the present study, and it may well be the case that A_{max} and WUE were greater immediately following fertiliser application, but in the two subsequent years declined to control levels. Our results are further evidence that fertiliser-promoted increases in A_{max} and WUE are often transient, lasting for only the first season after fertiliser application [57]. Increases in A_{max} and WUE beyond a year or so are unlikely unless increased growth of needles and other tissues are matched by continuing fertiliser inputs [12].




Figure 5. Relationship between the maximum rate of photosynthesis (A_{\max}) and (a) needle N concentration, (b) needle P concentration, (c) needle K concentration, and (d) Rubisco concentration. Fertiliser was applied to one-year-old trees and A_{\max} was measured two years later. Each point is a single measurement.

Contrary to our hypothesis, there was no evidence that differences in either relative or absolute concentrations of Rubisco could explain differences in A_{\max} and growth. Strong relationships among Rubisco, A_{\max} and growth are expected where N is a limiting element, whereas no relationship is expected in studies such as this where there is little evidence for a significant N limitation (Fig. 1a; see also N:P ratios, Fig. 1d). Trends in the allocation of N to Rubisco perhaps reflect the dual role of the enzyme in storage and photosynthesis [49]. For example, Rubisco concentration varied three-fold but this did not translate into differences in A_{\max} (Fig. 5), suggesting that Rubisco specific activity decreased with increasing Rubisco concentration. Positive relationships between K or P and allocation of N to Rubisco (Figs. 2 and 3) are consistent with recent suggestions that Rubisco accumulates in parallel with storage of P and K owing to the relatively fixed stoichiometry of these elements in plant matter and consequent “futility” of storing one without the other [55]. Alternatively one might argue that this positive relationship is simply a consequence of the high K (and to a lesser extent P) requirement of protein synthesis [20, 27]. Irrespective of its cause, we note that increased allocation of N to Rubisco was associated with low N:P and N:K ratios, which we might contrast with the increased allocation of N to amino acids commonly observed when N is in excess relative to elements such as K or P [3, 54].

5. CONCLUSIONS

Our results with *P. pinaster*, and those from other conifers, cast doubt on whether the growth response to fertiliser is a function of greater A_{\max} and/or WUE. By co-ordinating growth with nutrient supply, *P. pinaster* rarely produces needles that are nutrient deficient (with respect to photosynthesis). Allocation of N to Rubisco is plastic and modified by the supply of P and K, in partial support of our hypothesis. However, despite differences in allocation of N to Rubisco, there were no differences in A_{\max} . These findings argue against limitation of photosynthesis by Rubisco, and provide tentative support for the hypothesised role of Rubisco as a store of N.

Acknowledgements: The authors thank Ian Dumbrell and Keith Mungham from the Western Australian Department of Conservation and Land Management for establishing and maintaining the fertiliser trial. Charles Warren was supported by a Department of CALM/UWA postgraduate scholarship. The Australian Research Council is acknowledged for financial support.

REFERENCES

- [1] Adams M.B., Allen H.L., Nutrient proportions in foliage of semi-mature loblolly pine, *Plant Soil* 86 (1985) 27–34.

- [2] Adams M.A., Attiwill P.M., Nutrient cycling and nitrogen mineralization in eucalypt forests of south-eastern Australia. I. Nutrient cycling and nitrogen turnover, *Plant Soil* 92 (1986) 319–339.
- [3] Adams M.A., Attiwill P.M., Wang L.M., Effects of phosphorus supply on growth and nitrogen fractions in xylem sap and foliage of *Eucalyptus regnans* (F. Muell.), *E. nitens* (Maiden) and *E. globulus* (Labill.) seedlings: implications for herbivory, *Trees* 9 (1995) 324–331.
- [4] Barnes J.D., Pfirmann T., Steiner K., Lutz C., Busch U., Kuchenhoff H., Payer H.D., Effects of elevated CO₂, O₃ and potassium deficiency on Norway spruce [*Picea abies* (L.) Karst] seasonal changes in photosynthesis and non-structural carbohydrate content, *Plant Cell Environ.* 18 (1995) 1345–1357.
- [5] Bednarz C.W., Oosterhuis D.M., Physiological changes associated with K deficiency in cotton, *J. Plant Nutr.* 22 (1999) 303–313.
- [6] Ben Brahim M., Loustau D., Gaudillère J.P., Saur E., Effects of phosphate deficiency on photosynthesis and accumulation of starch and soluble sugars in one-year-old seedlings of maritime pine (*Pinus pinaster*), *Ann. Sci. For.* 53 (1996) 801–810.
- [7] Bradbury I.K., Malcolm D.C., The effect of phosphorus and potassium on transpiration, leaf diffusive resistance and water-use efficiency in Sitka spruce (*Picea sitchensis*) seedlings, *J. Appl. Ecol.* 14 (1977) 631–641.
- [8] Bristow K.L., Campbell G.S., On the relationship between incoming solar radiation and daily maximum and minimum temperature, *Agric. For. Meteorol.* 31 (1984) 159–166.
- [9] Brix H., Effects of thinning and nitrogen fertilization on growth of Douglas-fir: relative contribution of foliage quantity and efficiency, *Can. J. For. Res.* 13 (1983) 167–175.
- [10] Brix H., Ebell L.F., Effects of nitrogen fertilization on growth, leaf area and photosynthesis rate in Douglas-fir, *For. Sci.* 15 (1969) 189–196.
- [11] Brooks A., Effects of phosphorus nutrition on ribulose-1,5-bisphosphate carboxylase activation, photosynthetic quantum yield and amounts of some Calvin cycle metabolites in spinach leaves, *Aust. J. Plant Physiol.* 13 (1986) 221–237.
- [12] Carlyle J.C., Nutrient management in a *Pinus radiata* plantation after thinning: the effect of nitrogen fertilizer on soil nitrogen fluxes and tree growth, *Can. J. For. Res.* 25 (1995) 1673–1683.
- [13] Chandler J.W., Dale J.E., Photosynthesis and nutrient supply in needles of Sitka spruce [*Picea sitchensis* (Bong.) Carr.], *New Phytol.* 125 (1993) 101–111.
- [14] Chapin F.S. III., Bloom A.J., Field C.B., Waring R.H., Plant responses to multiple environmental factors, *BioSciences* 37 (1987) 49–57.
- [15] Conroy J.P., Milham P.J., Reed M.L., Barlow E.W.R., Increases in phosphorus requirements for CO₂-enriched pine species, *Plant Physiol.* 92 (1990) 977–982.
- [16] Evans J.R., Photosynthesis and nitrogen relationships in leaves of C₃ plants, *Oecologia* 78 (1989) 9–19.
- [17] Farquhar G.D., Ehleringer J.R., Hubick K.T., Carbon isotope discrimination and photosynthesis, *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 40 (1989) 503–537.
- [18] Field C.B., Merino J., Mooney H.A., Compromises between water-use efficiency and nitrogen-use efficiency in five species of Californian evergreens, *Oecologia* 60 (1983) 384–389.
- [19] Field C., Mooney H.A., The photosynthesis-nitrogen relationship in wild plants, in: Givnish T.J. (Ed.), *On the economy of plant form and function*, Cambridge University Press, Cambridge, UK, 1986, pp. 25–55.
- [20] Flaig H., Mohr H., Assimilation of nitrate and ammonium by the Scots pine (*Pinus sylvestris*) seedling under conditions of high nitrogen supply, *Physiol. Plant.* 84 (1992) 568–576.
- [21] Guehl J.-M., Fort C., Ferhi A., Differential response of leaf conductance, carbon isotope discrimination and water-use efficiency to nitrogen deficiency in maritime pine and pedunculate oak plants, *New Phytol.* 131 (1995) 149–157.
- [22] Houtman F., Godbold D.L., Shasheng W., Hüttermann A., Gas exchange in *Populus maximowiczii* in relation to Potassium and phosphorus nutrition, *J. Plant Physiol.* 135 (1990) 675–679.
- [23] Jacob J., Lawlor D.W., Dependence of photosynthesis of sunflower and maize leaves on phosphate supply, ribulose-1,5-bisphosphate carboxylase/oxygenase activity, and ribulose-1,5-bisphosphate pool size, *Plant Physiol.* 98 (1992) 801–807.
- [24] Keay J., Turton A.G., Campbell N.A., Some effects of nitrogen and phosphorus fertilization of *Pinus pinaster* in Western Australia, *For. Sci.* 14 (1968) 408–417.
- [25] Koerselman W., Meulman A.F.M., The vegetation N:P ratio: a new tool to detect the nature of nutrient limitation, *J. Appl. Ecol.* 33 (1996) 1441–1450.
- [26] Lauer M.J., Pallardy S.G., Belvins D.G., Randall D.D., Whole leaf carbon exchange characteristics of phosphate deficient soybeans (*Glycine max* L.), *Plant Physiol.* 91 (1989) 848–854.
- [27] Leigh R.A., Wyn Jones R.G., A hypothesis relating critical potassium concentrations for growth to the distribution and functions of this ion in the plant cell, *New Phytol.* 97 (1984) 1–13.
- [28] Lewis J.D., Griffin K.L., Thomas R.B., Strain B.R., Phosphorus supply affects the photosynthetic capacity of loblolly pine grown in elevated carbon dioxide, *Tree Physiol.* 14 (1994) 1229–1244.
- [29] Lösch R., Jensen C.R., Andersen M.N., Diurnal courses and factorial dependencies of leaf conductance and transpiration of differently potassium fertilized and watered field grown barley plants, *Plant Soil* 140 (1992) 205–224.
- [30] Loustau D., Ben Brahim M., Gaudillère J.P., Dreyer E., Photosynthetic responses to phosphorus nutrition in two-year-old maritime pine seedlings, *Tree Physiol.* 19 (1999) 707–715.
- [31] McMurtrie R.E., Rook D.A., Kelliher F.M., Modeling the yield of *Pinus radiata* on a site limited by water and nitrogen, *For. Ecol. Manage.* 30 (1990) 381–413.
- [32] MacRobbie E.A.C., Signal transduction and ion channels in guard cells, *Phil. Trans. Roy. Soc. Lond. B.* 353 (1998) 1475–1488.
- [33] Maier C.A., Teskey R.O., Internal and external control of net photosynthesis and stomatal conductance of mature eastern white pine (*Pinus strobus*), *Can. J. For. Res.* 22 (1992) 1387–1394.
- [34] Medlyn B.E., Loustau D., Delzon S., Temperature response of a biochemically based model of photosynthesis. I. Seasonal changes in mature maritime pine (*Pinus pinaster* Ait.), *Plant Cell Environ.* 25 (2002) 1155–1165.
- [35] Mitchell A.K., Hinkley T.M., Effects of foliar nitrogen concentration on photosynthesis and water use efficiency in Douglas-fir, *Tree Physiol.* 12 (1993) 403–410.
- [36] Monteith J.L., Evaporation and environment, *Symp. Soc. Exp. Biol.* 19 (1965) 205–234.
- [37] Murphy J., Riley H.P., A modified single solution method for the determination of phosphate in natural waters, *Anal. Chim. Acta* 27 (1962) 31–36.
- [38] Nambiar E.K.S., Management of forests under nutrient and water stress, *Water Air Soil Pollut.* 54 (1990) 209–230.
- [39] Oleksyn J., Tjoelker M.G., Lorenc-Plucinska G., Konwinska A., Zytowski R., Karolewski P., Reich P.B., Needle CO₂ exchange, structure and defence traits in relation to needle age in *Pinus heldreichii* Christ – a relict of tertiary flora, *Trees* 12 (1997) 82–89.
- [40] Peaslee D.E., Moss D.N., Stomatal conductivities in K-deficient leaves of maize (*Zea mays* L.), *Crop Sci.* 8 (1968) 427–430.
- [41] Porté A., Loustau D., Variability of the photosynthetic characteristics of mature needles within the crown of a 25-year-old *Pinus pinaster*, *Tree Physiol.* 18 (1998) 223–232.
- [42] Rao M., Terry N., Leaf phosphate status, photosynthesis and carbon partitioning in sugar beet. I. Changes in growth, gas exchange, and Calvin cycle enzymes, *Plant Physiol.* 90 (1989) 814–819.
- [43] Reich P.B., Schoettle A.W., Role of phosphorus and nitrogen in photosynthetic and whole plant carbon gain and nutrient use efficiency in eastern white pine, *Oecologia* 77 (1988) 25–33.
- [44] Running S.W., Ramakrishna R.N., Hungerford R.D., Extrapolation of synoptic meteorological data in mountainous terrain and its use for simulating forest evapotranspiration and photosynthesis, *Can. J. For. Res.* 17 (1987) 472–483.

- [45] Schoettle A.W., Smith W.K., Interrelationships among light, photosynthesis and nitrogen in the crown of mature *Pinus contorta* ssp. *latifolia*, *Tree Physiol.* 19 (1999) 13–22.
- [46] Sheriff D.W., Responses of carbon gain and growth of *Pinus radiata* stands to thinning and fertilizing, *Tree Physiol.* 16 (1996) 527–536.
- [47] Sheriff D.W., Nambiar E.K.S., Fife D.N., Relationships between nutrient status, carbon assimilation and water use efficiency in *Pinus radiata* needles, *Tree Physiol.* 2 (1986) 73–88.
- [48] Smolander H., Oker-Blom P., The effect of nitrogen content on the photosynthesis of Scots pine needles and shoots, *Ann. Sci. For.* 46 (1989) 473s–475s.
- [49] Stitt M., Schulze E.-D., Does Rubisco control the rate of photosynthesis and plant growth? An exercise in molecular ecophysiology, *Plant Cell Environ.* 17 (1994) 465–487.
- [50] Terry N., Ulrich A., Effects of phosphorus deficiency on the photosynthesis and respiration of leaves of sugar beet, *Plant Physiol.* 51 (1973) 43–47.
- [51] Teskey R.O., Gholz H.L., Cropper W.P., Influence of climate and fertilisation on net photosynthesis of mature slash pine, *Tree Physiol.* 14 (1994) 1215–1227.
- [52] Teskey R.O., Whitehead D., Linder S., Photosynthesis and carbon gain by pines, *Ecol. Bull.* 43 (1994) 35–49.
- [53] Vapaavuori E.M., Vuorinen A.H., Aphalo P.J., Smolander H., Relationships between net photosynthesis and nitrogen in Scots pine: seasonal variation in seedlings and shoots, *Plant Soil* 168–169 (1995) 263–270.
- [54] Warren C.R., Adams M.A., Capillary electrophoresis for the determination of major amino acids and sugars in foliage: application to the nitrogen nutrition of sclerophyllous species, *J. Exp. Bot.* 51 (2000) 1147–1157.
- [55] Warren C.R., Adams M.A., P affects growth and partitioning of N to Rubisco in *Pinus pinaster*, *Tree Physiol.* 22 (2002) 11–19.
- [56] Warren C.R., Adams M.A., Chen Z., Is photosynthesis related to concentrations of nitrogen and Rubisco in leaves of Australian native plants? *Aust. J. Plant Physiol.* 27 (2000) 407–416.
- [57] Warren C.R., McGrath J.F., Adams M.A., Water availability and carbon isotope discrimination in conifers, *Oecologia* 127 (2001) 476–486.
- [58] Wellburn A.R., The spectral determination of chlorophylls *a* and *b*, as well as total carotenoids, using various solvents with spectrophotometers of different resolution, *J. Plant. Physiol.* 144 (1994) 307–313.