



**HAL**  
open science

## Divergence among species and populations of Mediterranean pines in biomass allocation of seedlings grown under two watering regimes

María Regina Chambel, José Climent, Ricardo Alía

### ► To cite this version:

María Regina Chambel, José Climent, Ricardo Alía. Divergence among species and populations of Mediterranean pines in biomass allocation of seedlings grown under two watering regimes. *Annals of Forest Science*, 2007, 64 (1), pp.87-97. hal-00884061

**HAL Id: hal-00884061**

**<https://hal.science/hal-00884061>**

Submitted on 11 May 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Divergence among species and populations of Mediterranean pines in biomass allocation of seedlings grown under two watering regimes

María Regina CHAMBEL, José CLIMENT, Ricardo ALÍA\*

Instituto Nacional de Investigación y Tecnología Agraria y Alimentaria (INIA), Apto 8111, 28080 Madrid, Spain

(Received 27 February 2006; accepted 4 April 2006)

**Abstract** – Seedlings of four populations each of *Pinus pinaster*, *P. halepensis*, *P. canariensis* and *P. pinea* were grown in controlled conditions to evaluate both inter- and intra-specific differences in response to watering. We submitted half of the plants to a moderate water stress and after 22 weeks, we recorded height, stem diameter and root, stem and leaves dry weight. Patterns and amounts of phenotypic changes, including changes in biomass allocation, were analysed. We found a scant response in *P. canariensis*, *P. pinaster* and *P. halepensis* presented high population divergence for phenotypic changes, and *P. pinea* showed marked allocational shifts and no population divergence. The phenotypic changes observed within species are interpreted as a plastic response. The variation encountered within *P. halepensis* and *P. pinaster* may be indicative of specialisation to either resource-rich or resource-poor habitats, being populations from favourable sites more plastic. *P. pinea* exhibited a very uniform plastic response, indicating generalist behaviour.

phenotypic changes / early testing / pine / drought stress / ontogeny

**Résumé** – Divergences parmi les espèces et populations de pins méditerranéens pour l'allocation de biomasse chez des semis poussant sous deux régimes d'alimentation hydrique. Des semis de quatre populations de *Pinus pinaster*, de *P. halepensis*, de *P. canariensis*, et de *P. pinea* ont été élevés en conditions contrôlées pour évaluer au niveau inter- et intra-spécifique les différences de réponse au régime d'alimentation hydrique. Nous avons soumis la moitié des plants à un stress hydrique modéré et après 22 semaines nous avons mesuré leur hauteur, le diamètre de la tige et des racines, le poids sec de la tige et des feuilles. Les modèles et l'importance des changements phénotypiques, incluant les variations d'allocation de biomasse ont été analysés. Nous avons trouvé une faible réponse pour *P. canariensis*; *P. pinaster* et *P. halepensis* ont présenté une importante divergence des populations au plan des changements phénotypiques, et *P. pinea* a montré une modification sensible au plan de l'allocation de biomasse sans divergence de population. Les changements phénotypiques observés chez les espèces ont été interprétés comme une réponse en terme de plasticité. Les variations rencontrées chez *P. halepensis* et *P. pinaster* peuvent être l'indice d'une spécialisation pour des habitats riches ou pauvres en terme de ressources. *P. pinea* a présenté une plasticité uniforme de réponse, révélant un comportement généraliste.

changements phénotypiques / test précoce / stress hydrique / ontogénie

## 1. INTRODUCTION

How do plants modify their phenotypes according to environment? This question has been the focus of interest in science in general and in forest science in particular during the last two hundred years [23]. In most cases, those phenotypic changes due to genotype by environment interactions were considered as a source of error in most breeding and genetic evaluation programs, and several techniques have been developed to deal with this topic [20,38,47]. However, recently new perspectives were open to the analyses of those changes, when seen from a more general point of view, and taking into consideration different evolutionary implications. In this framework, considerable research efforts are being dedicated to the study of phenotypic plasticity, i.e. the ability of a genotype (in a broad sense: species, population, family or clone, see [37] for a general discussion on the topic) to alter its morphology and physiology in response to changes in the environmental

conditions [36,40]. These changes are not inherently adaptive; in particular those related to resource limitation might represent inevitable responses of the organisms [10,28,51]. In fact, individuals faced with low resource levels during growth inevitably grow less. Nevertheless, phenotypic responses to different environments may also include specific developmental and functional adjustments that increase fitness in those environments [7,13,27,44,49]. According to the optimal partitioning theory, plants respond to stressful environmental conditions by shifting carbon allocation to the organs collecting the most limiting resource, a form of plasticity conducive to growth maximization [5,11,42]. However adjustments in biomass allocation also occur as a natural consequence of growth and development (ontogenetic drift sensu Evans [12]), reflecting a shift in plant priorities along an ontogenetic trajectory [50]. In many cases, developmental stage and environment alter the functional relationship between traits [33]. As a consequence, conclusions regarding morphological adjustments in response to a given stress treatment may differ

\* Corresponding author: alia@inia.es

**Table I.** Identification, location and relevant ecological features of the populations used in the present study.

Species	Population (code)	Latitude	Longitude	Altitude (m)	Annual rainfall (mm)	Summer rainfall (mm) <sup>1</sup>	Growth conditions <sup>2</sup>
<i>Pinus pinaster</i>	Arenas de S. Pedro (PR-AR )	40° 12' N	05° 06' W	750	1 190	105	F
	Cómpeta (PR-CP)	36° 51' N	03° 55' W	900	700	24	U
	Leiria (PR-LE)	39° 45' N	08° 55' W	60	910	50	F
	Coca (PR-CC)	41° 14' N	04° 31' W	810	470	67	U
<i>Pinus halepensis</i>	Villa de Ves (PH-VV)	39° 10' N	01° 14' W	850	490	90	F
	North Euboia (PH-NE)	38° 59' N	23° 30' E	40	674	67	F
	Alcantud (PH-AL)	40° 34' N	02° 19' W	950	660	94	F
	Ses Salines (PH-SS)	39° 17' N	03° 02' E	10	300	27	U
<i>Pinus canariensis</i>	Vilaflor (PC-VI)	28° 11' N	16° 38' W	2100	450	1	U
	Barlovento (PC-BA)	28° 47' N	17° 51' W	1900	950	12	F
	Punta Gorda (PC-PG)	28° 47' N	17° 58' W	800	550	3	F
	Tirajana (PC-TI)	27° 53' N	15° 36' W	950	300	0	U
<i>Pinus pinea</i>	Tordesillas (PA-TO)	41° 30' N	04° 57' W	680	470	75	U
	Tarazona de la Mancha (PA-TM)	39° 17' N	01° 55' W	700	400	56	U
	Cartaya (PA-CA)	37° 22' N	07° 11' W	82	510	18	F
	Palafrugell (PA-PL)	41° 57' N	03° 06' W	100	660	94	F

<sup>1</sup> Summer months include June, July and August. <sup>2</sup> F = Favourable growth conditions, U = Unfavourable growth conditions, based on ecological data from [12].

dramatically if ontogenetic changes in phenotypic expression are also taken into consideration [9, 19, 29, 39, 46].

Still, little is known about the trade-offs between phenotypic changes and ecotypic differentiation in long-lived organisms that must face a changing environment. Mediterranean pines sensu Klaus [22] (*Pinus pinaster*, *P. halepensis*, *P. brutia*, *P. pinea*, *P. canariensis*, *P. roxburghii* and *P. heldreichii*) constitute an interesting group of species to address these questions. They form a well-defined phylogenetic group [26], exhibit marked differences in life-history traits [43], have different evolutionary histories, and presently occupy different ecological niches [4], with marked differences in water availability [14].

The objectives of this research were to check for differences among closely related species and populations within species in the degree and nature of morphological changes in response to watering regimes at early developmental stages. Our work hypothesis is that phenotypic change is a trait on itself, different from the trait under evaluation in each environment, and thus subject of genetic control at different organization levels [7, 35, 37].

To attain the proposed objectives, we used four Mediterranean pines with different degrees of drought tolerance, each represented by four populations. We examined growth and morphology of seedlings of these four species grown under two water regimes and assessed the effect of water availability on growth rates and on the allometric relationships between biomass compartments independently from the effect of ontogenetic drift. Furthermore, to avoid confounding the effect

of watering treatments with ontogenetic shifts in biomass allocation, besides using plant size as a proxy to ontogeny in allometric analysis, we also used a categorical morphological scale to characterise the ontogenetic stage of each seedling. Based on our results, we discuss the use of short-term experiments under controlled conditions to evaluate phenotypic changes and the relationship among our observations and phenotypic plasticity facing drought.


## 2. MATERIAL AND METHODS

### 2.1. Plant material

We used four Mediterranean pine species (*Pinus pinaster*, *P. halepensis*, *P. canariensis* and *P. pinea*), each represented by four populations. The seeds were collected in natural populations, selected to cover a wide range of environments, including material from populations with both favourable and unfavourable environmental or genetic growth conditions (Tab. I). So as to draw conclusions at the population level and ensure repeatability of the experiment, we used an equilibrated mix of seeds collected from random samples of 25 to 30 open-pollinated individuals per population, separated by a minimum of 100 m to reduce consanguinity.

### 2.2. Experimental design

The study was conducted in a growth chamber with controlled temperature and photoperiod. After seed germination, we



**Figure 1.** Ontogenetic scores. (0) Cotyledonary stage, (1) emergence of the epicotyl rosette, (2) epicotyl elongation, (3) formation of axillary buds, (4) elongation of axillary long shoots, (5) formation of secondary axillary long shoots, (6) occurrence of dwarf shoots, (7) formation of a terminal bud.

transplanted thirty-six seedlings per population into 250 cm<sup>3</sup> individual plastic containers, filled with peat and vermiculite (4:1, v/v) and placed them inside the growth chamber. Half of the seedlings (18 per population) were randomly assigned to each of the water-stress treatments. A split-split plot design (population within species within water stress treatment), with two replicates of nine seedlings each, was used to control the effect of competition between neighbours and to compensate for the light intensity gradient across the chamber. Plants were maintained in the growth chamber for twenty-two weeks, following a protocol that included nine weeks of long photoperiod and high temperature, followed by a progressive decrease of both photoperiod and temperature to induce bud rest. Similar protocols have proven to significantly accelerate the maturation rate in maritime pine seedlings, leading to higher correlations with mature behaviour [24, 30]. Plants were watered to field capacity approximately every two days, except during the water stress treatment, as detailed below. The water stress treatment started in the ninth week from transplant and lasted for six weeks, coinciding with the period of high temperature. During this period, the water supply was withdrawn from half of the plants until the water content of each individual container reached 30% of field capacity (determined by weight). This watering level was maintained approximately constant until the end of the stress period. The remaining plants were watered as described previously.

Seedling height was measured before and after the water stress period, and height growth during the interval (HGD) was computed both for stressed and non-stressed plants. After twenty-two weeks in the growth chamber, the plants were harvested. Diameter at root collar (D) and seedling height (H) of every plant were measured. The seedlings were then partitioned into roots, stems, and leaves for biomass assessment [32]. All plant parts were oven-dried for 48 h at 80°C and then weighted. Dry weights of leaves (LDW), stems (SDW) and roots (RDW) were obtained and total dry weight (TDW) was computed from these values.

Shoot ontogeny was followed throughout the growth period, using a categorical, seven-level scale (Fig. 1), inspired by the works of Lester [25] and Williams [52] and based on the heteroblasty of shoot

development [21]. Seedlings were assigned to the values 0 for the cotyledonary stage, (1) for emergence of the epicotyl rosette, (2) for epicotyl elongation, (3) for formation of axillary buds, (4) for elongation of axillary long shoots, (5) for formation of secondary axillary long shoots, (6) for occurrence of dwarf shoots (either on the main shoot or on lateral branches) and (7) for formation of a terminal bud. Higher scores reflected a more developed ontogenetic stage, allowing the comparison of different species, even when transition from level to level may not be continuous for all plants.

### 2.3. Statistical analysis

Growth (H, D, HGD) and biomass (TDW, RDW, SDW, LDW) variables were analysed with the general linear model approach to analysis of variance, with type III sum of squares, using SAS software. The model terms were fitted according to the hierarchical design of the experiment, considering populations as nested within species. In addition to this analysis, an ANOVA was carried out for each species separately to evaluate different trends at the population level. A significant effect of the water stress treatment in this analysis indicates the existence of phenotypic changes in response to drought for the trait considered and a significant genotype by environment interaction indicates the existence of differences among population or species for those changes [36]. Whenever the treatment factor was significant, the difference between mean phenotype of each species or population in the two environments considered was evaluated with a *t*-test.

Besides plotting standard reaction norms, we represented graphically the position of each population in the space defined by its mean phenotype under the water stress treatment (on the *x*-axis) and under the non-stress treatment (on the *y*-axis) following Pigliucci and Schlichting [31]. This way, each population is represented by a single point and, if the two axes are in the same scale, the main diagonal represents the line of null phenotypic change, that corresponds to a flat reaction norm and the tangent (slope) of the angle  $\alpha$ , formed between the line connecting each point to the origin and the *x*-axis

can be interpreted as an index quantifying phenotypic change. The main advantage of this index, when compared to the most common methods based on the difference between mean phenotypic values in each environment (e.g. [34]), represented in this biplot by the orthogonal distance to the main diagonal, is that the slope is reflecting the change in relative terms, more significant from a biological point of view. Besides, this index also reflects the direction of the response (slope higher or lower than one), which has obvious biological relevance [48]. We will further refer to this index as angular phenotypic change index (APCI).

For the study of biomass allocation, we performed an allometric analysis through the regression of the natural logarithms of each biomass component (LDW, SDW and RDW) and the sum of the other two components [29, 32]. Changes in allocational patterns were assessed by comparison of the slopes and intercepts corresponding to different watering levels [37]. When, for a given species or population within species, a strong linear relation between biomass compartments existed and the two lines of regression corresponding to the two water treatments overlapped, the slope of those lines will differ only if the water stress treatment caused significant changes in the relative growth rates of leaves, shoots and roots. Ontogenetic scores were analysed using logistic regression based on maximum likelihood estimations (procedure CATMOD of SAS).

### 3. RESULTS

#### 3.1. Inter-specific variation

Species accounted for the highest proportion of the variability encountered in the analysis of most traits, as expected (Tab. II). Nevertheless, at this level, treatment effect was significant or highly significant for all the variables analysed. Species x treatment interaction was not significant either for height at the end of the experiment (H) or for height growth during the drought period (HGD), but it was significant for diameter (D,  $p < 0.05$ ) and most of the biomass-related traits (Tab. II). Considering the overall species effect (including the four populations together), the same ranking was found in both treatments for height and diameter growth, total biomass and leaf and stem biomass, with *Pinus pinea* attaining the highest values, followed by *P. canariensis*, *P. pinaster* and *P. halepensis*.

Height growth was significantly reduced during the drought period in the stressed plants of all four species ( $p < 0.001$ ). At the end of the experiment, the seedlings of *Pinus canariensis* and *Pinus pinaster* showed no differences for biomass-related traits (TDW, RDW, SDW and LDW) while the other three species showed significant or highly significant reductions in both traits due to the imposed drought.

The allometric analysis revealed that the water stress induced changes in the proportions of biomass allocated to each plant compartment that were independent of plant size, i.e. the existence of changes in allocation patterns (differences on the slope, interception or both) in response to drought, in all four species. In general, these changes affected mainly roots and leaves. Different allometric trajectories for stems were found only in *Pinus halepensis* and *P. canariensis*, while the four

**Table II.** Proportion of the variance due to treatment, species and treatment by species interaction in the inter-specific analysis and significance of the corresponding F tests.

	Treatment	Species	Treatment × Species
RDW	0.281**	0.100***	0.056**
SDW	0.206***	0.614***	0.005
LDW	0.154***	0.523***	0.027***
TDW	0.208***	0.403***	0.024**
H	0.208***	0.723***	0.010
D	0.162***	0.765***	0.011**
HGD	0.637***	0.115***	0.004

RDW: root dry weight; SDW: stem dry weight; LDW: leaf dry weight; TDW: total dry weight; H: height; D: diameter; HGD: height growth during drought. Significance levels \*\*  $p < 0.05$ , \*\*\*  $p < 0.001$ .

species displayed allocational changes for leaves and all except *P. canariensis* for roots. Therefore, *P. halepensis* exhibited the highest degree of change in biomass allocation, followed by *P. pinea*, *P. pinaster* and finally by *P. canariensis*.

We found sharp differences among species for seedling maturation, as evaluated by the ontogenetic scores. *Pinus pinaster* attained the highest mean score, with a high proportion of seedlings bearing axillary dwarf shoots, followed by *P. halepensis* and *P. pinea* (not significantly different) and by *P. canariensis*, with a very low score. After 22 weeks, no plant attained the highest score on the scale, corresponding to the formation of a true terminal bud covered with cataphylls. Nevertheless, some *P. pinaster* seedlings showed a terminal rosette of short primary needles, closely resembling a terminal bud. Although statistically the ontogenetic scores of *P. halepensis* and *P. pinea* were not significantly different, individual plants of both groups were in fact very different. A small proportion of the *P. halepensis* seedlings formed dwarf shoots (level 6), while most were in level 5 and some remained in level 4. On the contrary, the relatively high score attained in *P. pinea* was exclusively due to the abundant secondary branching (level 5). The seedlings of this species formed the most uniform group with regard to ontogeny. In general, however, ontogenetic scores proved to be relatively stable within each species; only for *P. halepensis* we found significant differences among populations and water stress treatments ( $p < 0.001$  and  $p < 0.05$ , respectively). The relationship between ontogenetic score and plant dry weight was different for each species and generally weak, especially in *P. pinea* ( $r^2 = 0.02$ ).

#### 3.2. Intra-specific variation

Water stress treatment accounted for the highest proportion of the variability in *Pinus pinaster* in all traits except leaf biomass, while differences among populations were highly significant for all biomass components and for total height ( $p < 0.001$ ). Population x treatment interaction was highly significant for all biomass related variables ( $p < 0.001$ ), but

**Table III.** Proportion of the variance due to treatment, population and treatment by population interaction in the intra-specific analysis and significance of the corresponding F tests.

	<i>P. pinaster</i>			<i>P. halepensis</i>			<i>P. canariensis</i>			<i>P. pinea</i>		
	T	P	T × P	T	P	T × P	T	P	T × P	T	P	T × P
RDW	0.44***	0.20***	0.31***	0.74***	0.12***	0.11***	0.00	0.04	0.28	0.04**	0.01	0.00
SDW	0.51***	0.24***	0.21***	0.67***	0.16***	0.11***	0.58**	0.16	0.14	0.13***	0.02	0.01
LDW	0.25**	0.31***	0.39***	0.71***	0.13***	0.13***	0.00	0.19	0.32	0.19***	0.02	0.01
TDW	0.35***	0.26***	0.34***	0.74***	0.12***	0.12***	0.01	0.13	0.31	0.13***	0.01	0.00
H	0.69***	0.24***	0.02	0.51***	0.42***	0.03	0.11	0.24**	0.01	0.83***	0.08	0.06
D	0.46**	0.16	0.28**	0.86***	0.02	0.05**	0.45	0.30	0.09	0.16***	0.03	0.00
HGD	0.91***	0.03	0.03	0.65***	0.15***	0.00	0.62***	0.03	0.02	0.62***	0.04***	0.01

T: treatment; P: population; T × P: treatment × population; RDW: root dry weight; SDW: stem dry weight; LDW: leaf dry weight; TDW: total dry weight; H: height; D: diameter; HGD: height growth during drought. Significance levels \*\*  $p < 0.05$ , \*\*\*  $p < 0.001$ .

not significant for height growth (Tab. III). This species displayed striking differences among populations in response to the drought stress treatment. Population PR-LE showed a remarkable change in total biomass and dry weight components (Tab. IV, Fig. 2), presenting the highest values of the angular phenotypic change index (APCI) found in this study (ranging from 2.71 for RDW to 1.31 for H, Fig. 2). On the contrary, populations PR-AR and PR-CP proved to be markedly stable for all traits under study (Tab. IV, Fig. 2). Population PR-CC exhibited significant changes only for height growth (reduced H in the stress treatment). In most cases, the response to drought was not reflected in different allometric relations between the biomass components. This species displayed significant differences in biomass allocational patterns only for stems in population PR-CP (different slopes) and for leaves in population PR-AR (different intercepts).

In *Pinus halepensis*, the water stress treatment accounted for the highest proportion of variability in all variables considered; population × treatment interaction followed trends similar to those described for *P. pinaster* (Tab. IV). Populations PH-VV and PH-NE exhibited significant changes for all the growth and biomass related variables, with population PH-VV reaching higher values of APCI in all cases (Tab. IV, Fig. 2). Population PH-AL exhibited significant reductions of diameter growth, TDW and RDW (APCI of 1.30 and 1.35, respectively), while population PH-SS showed no significant phenotypic changes for any of the traits considered. Contrasting with these results, population PH-AL showed the highest degree of change in biomass allocation, shifting allometric trajectories of all biomass compartments as a consequence of drought. Populations PH-NE and PH-SS exhibited significant changes in biomass allocation to stems while population PH-VV presented changes for leaves.

In the Canary Island pine, neither the water stress treatment nor the population accounted for significant proportions of the variance, with a few exceptions (SDW and HGD for treatment and H for population). No population × treatment interaction was found in this species. When considering the four populations separately, this species still displayed the lowest levels of phenotypic change. APCI values were in general amongst the

lowest observed in the present study (Fig. 2), with significant differences between treatments only for RDW in population PC-PG. Nevertheless, populations PC-VI and PC-BA revealed shifts on the allometric trajectories for both leaves and roots, indicating that in this species the changes in biomass allocation patterns prevailed over the changes in growth variables.

The proportion of variability due to the water stress treatment in *Pinus pinea* was generally lower than that found in *P. pinaster* and *P. halepensis*, while the effect of population and that of population × treatment interaction were not significant in any case (Tab. IV). The populations of this species exhibited the highest levels of change in biomass allocation patterns, with the allometric curves fitted for roots and leaves overlapping completely and showing significantly different trajectories between drought treatments in all four populations (Fig. 3). Therefore, indicating that also in this species the allocational shifts overcome growth differences. Worthy of note is the fact that the direction of the response was identical in all cases; allocation to roots increased at the expenses of the above ground biomass components as a consequence of drought.

#### 4. DISCUSSION

The present paper is focused on the morphological response of seedlings from close related species (and populations within species) to two contrasting watering regimes. The observed responses raise some questions worthy of a close look: the significance of those phenotypic changes from the perspective of the phenotypic plasticity of populations and the relationship among the observed changes and species differences regarding life history and ecology.

Until present, little information was available on the morphological changes induced by water stress during the initial developmental stages of Mediterranean pines and its variability within and between species. The direct comparison of the responses among species or populations within species poses some experimental difficulties related to the artificial design of the common stress treatments combined with the different tolerance of the species or populations under study. However, this is the only possible way to isolate genetic effects from

**Table IV.** Means ( $\pm$  standard errors) for biomass components and growth variables.

			RDW (mg)	SDW (mg)	LDW (mg)	TDW (mg)	H (mm)	D (mm)	HGD (mm)	
<i>P. pinaster</i>	PR-AR	No-stress	479 $\pm$ 84	188 $\pm$ 31	573 $\pm$ 126	1 240 $\pm$ 226	59.5 $\pm$ 6.6	2.3 $\pm$ 0.22	31.9 $\pm$ 4.4	
		Stress	490 $\pm$ 49	193 $\pm$ 19	816 $\pm$ 78	1 499 $\pm$ 141	52.2 $\pm$ 1.7	2.4 $\pm$ 0.13	15.9 $\pm$ 1.7	
	PR-CP	No-stress	471 $\pm$ 101	180 $\pm$ 39	704 $\pm$ 121	1 355 $\pm$ 251	49.1 $\pm$ 6.0	2.3 $\pm$ 0.15	25.9 $\pm$ 4.6	
		Stress	429 $\pm$ 30	162 $\pm$ 13	661 $\pm$ 64	1 251 $\pm$ 99	39.3 $\pm$ 2.6	2.2 $\pm$ 0.09	19.2 $\pm$ 2.1	
	PR-LE	No-stress	1058 $\pm$ 96	346 $\pm$ 31	1 500 $\pm$ 151	2 904 $\pm$ 275	66.6 $\pm$ 4.0	2.9 $\pm$ 0.10	36.0 $\pm$ 2.0	
		Stress	391 $\pm$ 52	169 $\pm$ 19	667 $\pm$ 72	1 226 $\pm$ 136	52.7 $\pm$ 4.4	2.1 $\pm$ 0.13	19.1 $\pm$ 1.4	
	PR-CC	No-stress	508 $\pm$ 134	194 $\pm$ 41	714 $\pm$ 207	1 416 $\pm$ 377	62.0 $\pm$ 3.9	2.3 $\pm$ 0.23	32.2 $\pm$ 2.1	
		Stress	438 $\pm$ 76	132 $\pm$ 18	611 $\pm$ 81	1 181 $\pm$ 167	47.3 $\pm$ 5.0	2.1 $\pm$ 0.12	21.1 $\pm$ 2.8	
	<i>P. halepensis</i>	PH-VV	No-stress	704 $\pm$ 51	174 $\pm$ 15	876 $\pm$ 66	1 754 $\pm$ 128	50.5 $\pm$ 2.1	2.3 $\pm$ 0.08	27.0 $\pm$ 1.4
			Stress	356 $\pm$ 36	99 $\pm$ 8	487 $\pm$ 36	942 $\pm$ 76	39.7 $\pm$ 1.5	1.7 $\pm$ 0.05	18.3 $\pm$ 2.0
PH-NE		No-stress	760 $\pm$ 71	166 $\pm$ 19	787 $\pm$ 61	1 714 $\pm$ 147	60.2 $\pm$ 3.6	2.2 $\pm$ 0.09	32.8 $\pm$ 2.2	
		Stress	540 $\pm$ 47	120 $\pm$ 9	574 $\pm$ 35	1 234 $\pm$ 79	49.8 $\pm$ 2.8	1.8 $\pm$ 0.07	23.4 $\pm$ 1.9	
PH-AL		No-stress	695 $\pm$ 44	154 $\pm$ 15	796 $\pm$ 58	1 644 $\pm$ 107	42.3 $\pm$ 1.8	2.2 $\pm$ 0.09	21.8 $\pm$ 1.2	
		Stress	516 $\pm$ 52	114 $\pm$ 9	630 $\pm$ 43	1 261 $\pm$ 100	38.1 $\pm$ 1.7	1.9 $\pm$ 0.08	14.1 $\pm$ 1.8	
PH-SS		No-stress	482 $\pm$ 51	98 $\pm$ 14	528 $\pm$ 57	1 108 $\pm$ 118	43.8 $\pm$ 2.6	2.0 $\pm$ 0.08	27.2 $\pm$ 1.9	
		Stress	482 $\pm$ 37	103 $\pm$ 8	535 $\pm$ 44	1 120 $\pm$ 85	39.9 $\pm$ 2.3	1.9 $\pm$ 0.07	17.8 $\pm$ 1.9	
<i>P. canariensis</i>		PC-VI	No-stress	634 $\pm$ 62	204 $\pm$ 21	882 $\pm$ 64	1 720 $\pm$ 141	70.7 $\pm$ 5.4	2.9 $\pm$ 0.10	37.7 $\pm$ 3.0
			Stress	562 $\pm$ 55	184 $\pm$ 14	954 $\pm$ 55	1 700 $\pm$ 112	70.2 $\pm$ 2.9	2.8 $\pm$ 0.10	25.6 $\pm$ 3.2
	PC-BA	No-stress	687 $\pm$ 60	253 $\pm$ 24	1 101 $\pm$ 95	2 040 $\pm$ 164	68.2 $\pm$ 3.8	2.9 $\pm$ 0.10	34.7 $\pm$ 2.7	
		Stress	605 $\pm$ 80	191 $\pm$ 16	918 $\pm$ 66	1 715 $\pm$ 145	62.7 $\pm$ 4.3	2.8 $\pm$ 0.10	28.2 $\pm$ 2.9	
	PC-PG	No-stress	553 $\pm$ 42	193 $\pm$ 14	825 $\pm$ 61	1 571 $\pm$ 111	65.9 $\pm$ 4.8	2.9 $\pm$ 0.07	33.2 $\pm$ 2.6	
		Stress	694 $\pm$ 44	195 $\pm$ 13	984 $\pm$ 70	1 873 $\pm$ 111	63.6 $\pm$ 3.5	2.8 $\pm$ 0.08	25.5 $\pm$ 2.6	
	PC-TI	No-stress	603 $\pm$ 48	227 $\pm$ 17	893 $\pm$ 59	1 724 $\pm$ 116	77.4 $\pm$ 4.0	2.8 $\pm$ 0.08	41.8 $\pm$ 2.9	
		Stress	642 $\pm$ 55	198 $\pm$ 12	856 $\pm$ 53	1 696 $\pm$ 112	73.3 $\pm$ 3.5	2.6 $\pm$ 0.08	27.3 $\pm$ 4.1	
	<i>P. pinea</i>	PA-TO	No-stress	783 $\pm$ 49	334 $\pm$ 28	1 690 $\pm$ 172	2 808 $\pm$ 237	73.7 $\pm$ 2.3	3.1 $\pm$ 0.10	32.9 $\pm$ 1.2
			Stress	708 $\pm$ 63	291 $\pm$ 17	1 218 $\pm$ 67	2 217 $\pm$ 139	66.8 $\pm$ 1.7	2.9 $\pm$ 0.06	26.5 $\pm$ 1.6
PA-TM		No-stress	742 $\pm$ 70	299 $\pm$ 29	1 367 $\pm$ 131	2 409 $\pm$ 226	76.2 $\pm$ 2.9	3.1 $\pm$ 0.17	33.6 $\pm$ 1.7	
		Stress	693 $\pm$ 76	270 $\pm$ 18	1 198 $\pm$ 82	2 160 $\pm$ 169	72.7 $\pm$ 1.9	2.9 $\pm$ 0.06	25.9 $\pm$ 1.3	
PA-CA		No-stress	719 $\pm$ 46	317 $\pm$ 23	1 416 $\pm$ 143	2 452 $\pm$ 201	80.2 $\pm$ 3.2	3.1 $\pm$ 0.12	37.6 $\pm$ 2.2	
		Stress	597 $\pm$ 58	248 $\pm$ 13	1 171 $\pm$ 66	2 016 $\pm$ 128	67.1 $\pm$ 1.9	2.9 $\pm$ 0.06	27.4 $\pm$ 1.8	
PA-PL		No-stress	727 $\pm$ 48	293 $\pm$ 28	1 365 $\pm$ 103	2 385 $\pm$ 171	73.3 $\pm$ 2.6	3.0 $\pm$ 0.10	32.8 $\pm$ 2.0	
		Stress	691 $\pm$ 89	265 $\pm$ 24	1 077 $\pm$ 89	2 033 $\pm$ 195	66.1 $\pm$ 2.3	2.7 $\pm$ 0.08	21.4 $\pm$ 1.5	

RDW: root dry weight (mg); SDW: stem dry weight (mg); LDW: leaf dry weight (mg); TDW: total dry weight (mg); H: height (mm); D: diameter (mm); HGD: height growth during drought (mm).


treatment effects. Similar approaches are commonly used in common garden provenance studies in and in most plasticity studies [6, 16, 45].

The levels of phenotypic change detected in this study with analysis of variance (GLM), angular phenotypic change index (APCI) and allometric analysis were not always coincident. In fact, these analyses evaluate different types of response, which can even be considered as different traits [37]. Both analysis of variance and APCI compare the mean response per species or populations in each environment at a given age. The allometric analysis detects shifts in biomass allocation priorities, independent of plant size, i.e. changes in the allometric trajec-

ories of a given species or population, induced in this case, by the water stress.

Regarding the relationship between the observed changes and the phenotypic plasticity of the material under study, if we consider that plasticity must be evaluated in the same material and at similar ontogenic stage, we can conclude that, for comparison at the species level, this is not the case in our study given that sharp differences in ontogeny were found among species at the end of the experiment. However, when comparisons are made at the intraspecific level, both the similarity of ontogenetic scores and the overlapping of plant biomass ranges support the idea that the observed phenotypic changes are in fact plastic responses to an imposed drought stress [32].





**Figure 3.** Allometric trajectories for roots (left column) and leaves (right column) of the four populations of *Pinus pinea* studied. Dots and continuous lines represent stressed plants, triangles and dotted lines represent non-stressed plants.



and the reviewers for their helpful critical comments. This study was supported by the project INIA-DGB CC03-048, and a grant from the AECl to the first author.

## REFERENCES

- [1] Alía R., Gil L., Pardos J.A., Performance of 43 *Pinus pinaster* Ait. Provenances on 5 locations in central Spain, *Silvae Genet.* 44 (1995) 75–81.
- [2] Alía R., Moro J., Denis J.B., Performance of *Pinus pinaster* provenances in Spain: interpretation of the genotype by environment interaction, *Can. J. For. Res.* 27 (1997) 1548–1559.
- [3] Balaguer L., Martínez-Ferri E., Valladares F., Pérez-Corona M.E., Baquedano F.J., Castillo F.J., Manrique E., Population divergence in the plasticity of the response of *Quercus coccifera* to the light environment, *Funct. Ecol.* 15 (2001) 124–135.
- [4] Barbéro M., Loisel R., Quézel P., Richardson D.M., Romane F., Pines of the Mediterranean basin, in: Richardson D.M. (Ed), *Ecology and biogeography of Pinus*, Cambridge University Press, 1998 pp. 153–170.
- [5] Bloom A.J., Chapin F.S., Mooney H.A., Resource limitation in plants – an economic analogy, *Annu. Rev. Ecol. Syst.* 16 (1985) 363–392.
- [6] Bouvet J.-M., Vigneron P., Saya A., Phenotypic plasticity of growth trajectory and ontogenic allometry in response to density for eucalyptus hybrid clones and families, *Ann. Bot.* 96 (2005) 811–821.
- [7] Bradshaw A.D., Evolutionary significance of phenotypic plasticity in plants, *Adv. Genet.* 13 (1965) 115–155.
- [8] Climent J., Gil L., Pérez E., Pardos J.A., Efecto de la procedencia en la supervivencia de plántulas de *Pinus canariensis* Sm. en medio árido., *Invest. Agr. Sist. Recur. For.* 11 (2002) 171–180.
- [9] Coleman J.S., McConnaughay K.D.M., Ackerly D.D., Interpreting phenotypic variation in plants, *Trends Ecol. Evol.* 9 (1994) 187–191.
- [10] Curt T., Coll L., Prévosto B., Baladier P., Kunstler G., Plasticity in growth, biomass allocation and root morphology in beech seedlings as induced by irradiance and herbaceous competition, *Ann. For. Sci.* 62 (2005) 51–60.
- [11] Chapin F.S., Bloom A.J., Field C.B., Waring R.H., Plant responses to multiple environmental factors, *BioScience* 37 (1987) 49–57.
- [12] Evans G.C., *The quantitative analysis of plant growth*, University of California Press, Berkeley, California, USA, 1972.
- [13] Galloway L.F., Maternal effects provide phenotypic adaptation to local environmental conditions, *New Phytol.* 166 (2005) 93–100.
- [14] Gandullo J., Sánchez Palomares O., *Estaciones ecológicas de los pinares españoles*, ICONA, Ministerio de Agricultura, Pesca y Alimentación, Madrid, 1994.
- [15] Gómez A., Aguiriano E., Alía R., Bueno M.A., Análisis de los recursos genéticos de *Pinus pinea* L. en España mediante microsatélites del cloroplasto, *Invest. Agrar. Sist. Recur. For.* 11 (2002) 145–154.
- [16] Griffith T.M., Sultan S.E., Shade tolerance plasticity in response to neutral vs. green shade cues in *Polygonum* species of contrasting ecological breadth, *New Phytol.* 166 (2005) 141–148.
- [17] Harfouche A., Baradat P., Durel C.E., Variabilité intraspécifique chez le pin maritime (*Pinus pinaster* Ait.) dans le sud-est de la France. I. Variabilité des populations autochtones et des populations de l'ensemble de l'aire de l'espèce, *Ann. Sci. For.* 52 (1995) 307–328.
- [18] Hopkins E.R., Butcher T.B., Provenance comparisons of *Pinus pinaster* Ait. in western Australia, *CALMScience* 1 (1993) 55–105.
- [19] Huber H., Lukács S., Watson M.A., Spatial structure of stoloniferous herbs: an interplay between structural blue-print, ontogeny and phenotypic plasticity, *Plant Ecol.* 141 (1999) 107–115.
- [20] Johnsen O., Fossdal C.G., Nagy N., Molmann J., Daehlen O.G., Skroppa T., Climatic adaptation in *Picea abies* progenies is affected by the temperature during zygotic embryogenesis and seed maturation, *Plant. Cell Environ.*
- [21] Jones C.S., An essay on juvenility, phase change and heteroblasty in seed plants, *Int. J. Plant Sci.* 160 (1999) S105–S111.
- [22] Klaus W., Mediterranean pines and their history, *Plant Syst. Evol.* 162 (1989) 133–163.
- [23] Langle O., Two hundred years genecology, *Taxon* 20 (1971) 653–722.
- [24] Lascoux D.M., Kremer A., Dormling I., Growth and phenology of 1-year-old maritime pine (*Pinus pinaster*) seedlings under continuous light: implications for early selection, *Can. J. For. Res.* 23 (1993) 1325–1336.
- [25] Lester D.T., Developmental patterns of axillary meristematic activity in seedlings of *Pinus*, *Bot. Gaz.* 129 (1968) 206–210.
- [26] López G.G., Kamiya K., Harada K., Phylogenetic relationships of Diploxylon pines (Subgenus *Pinus*) based on plastid sequence data, *International. J. Plant Sci.* 163 (2002) 737–747.
- [27] Merilä J., Laurila A., Lindgren B., Variation in the degree and costs of adaptive phenotypic plasticity among *Rana temporaria* populations, *J. Evol. Biol.* 17 (2004) 1132–1140.
- [28] Meyers L.A., Bull J.J., Fighting change with change: Adaptive variation in an uncertain world, *Trends Ecol. Evol.* 17 (2002) 551–557.
- [29] Müller I., Schmid B., Weiner J., The effect of nutrient availability on biomass allocation patterns in 27 species of herbaceous plants, *Perspect. Plant Ecol. Evol. Syst.* 3 (2000) 115–127.
- [30] Nguyen A., Dormling I., Kremer A., Characterization of *Pinus pinaster* seedling growth in different photoperiods and thermoperiods in a phytotron as a basis for early selection, *Scand. J. For. Res.* 10 (1995) 129–139.
- [31] Pigliucci M., Schlichting C.D., Reaction norms of *Arabidopsis*. IV. Relationships between plasticity and fitness, *Heredity* 76 (1996) 427–436.
- [32] Poorter H., Nagel O., The role of biomass allocation in the growth response of plants to different levels of light, CO<sub>2</sub>, nutrients and water: a quantitative review, *Aust. J. Plant Physiol.* 27 (2000) 595–607.
- [33] Preston K.A., Ackerly D.D., The evolution of allometry in modular organisms, in: Pigliucci M., Preston K.A. (Eds.), *Phenotypic Integration: studying the ecology and evolution of complex phenotypes*, Oxford University Press, 2003.
- [34] Scheiner S.M., Genetics and evolution of phenotypic plasticity, *Annu. Rev. Ecol. Syst.* 24 (1993) 35–68.
- [35] Scheiner S.M., Lyman R.F., The genetics of phenotypic plasticity. II. Response to selection, *J. Evol. Biol.* 4 (1991) 23–50.
- [36] Schlichting C.D., The evolution of phenotypic plasticity in plants, *Annu. Rev. Ecol. Syst.* 17 (1986) 667–693.
- [37] Schlichting C.D., Pigliucci M., *Phenotypic evolution – A reaction norm perspective*, Sinauer Associates, Sunderland, MA, 1998.
- [38] Shelbourne C.J.A., Genotype-environmentinteraction: its study and its implications in forest tree improvement, in *SABRAO Joint Symposium*, 1972, Tokyo, Government Forest Experiment Station.
- [39] Strand J.A., Weisner S.E.B., Phenotypic plasticity – contrasting species-specific traits induced by identical environmental constraints, *New Phytol.* 163 (2004) 449–451.
- [40] Sultan S.E., Evolutionary implications of phenotypic plasticity in plants, *Evol. Biol.* 21 (1987) 127–178.
- [41] Sultan S.E., Phenotypic plasticity for plant development, function, and life-history, *Trends Plant Sci.* 5 (2000) 537–542.
- [42] Sultan S.E., Phenotypic plasticity in plants: a case study in ecological development, *Evol. Dev.* 5 (2003) 25–33.
- [43] Tapias R., Climent J., Pardos J.A., Gil L., Life histories of Mediterranean pines, *Plant Ecol.* 171 (2004) 53–68.

- [44] Valladares F., Balaguer L., Martinez-Ferri E., Perez-Corona M.E., Manrique E., Plasticity, instability and canalization: is the phenotypic variation in seedlings of sclerophyll oaks consistent with the environmental unpredictability of Mediterranean ecosystems? *New Phytol.* 156 (2002) 457–467.
- [45] Valladares F., Dobarro I., Sanchez-Gomez D., Pearcy R.W., Photoinhibition and drought in Mediterranean woody saplings: scaling effects and interactions in sun and shade phenotypes, *J. Exp. Bot.* 56 (2005) 483–494.
- [46] Valladares F., Wright S.J., Lasso E., Kitajima K. Pearcy R.W., Plastic phenotypic response to light of 16 congeneric shrubs from a Panamanian rainforest, *Ecology* 81 (2000) 1925–1936.
- [47] Van Eeuwijk F.A., Malosetti M., Yin X., Struik P.C., Stam P., Statistical models for genotype by environment data: from conventional ANOVA models to eco-physiological QTL models. *Austr. J. Agric. Res.* (2005) 12.
- [48] Via S., Adaptive phenotypic plasticity – Target or by-product of selection in a variable environment, *Am. Nat.* 142 (1993) 352–365.
- [49] Wang G.G., Bauerle W.L., Effects of light intensity on the growth and energy balance of photosystem II electron transport in *Quercus alba* seedlings, *Ann. For. Sci.* 63 (2006) 111–118.
- [50] Weiner J., Allocation, plasticity and allometry in plants, *Perspect. in Plant Ecol. Evol. Syst.* 6 (2004) 207–215.
- [51] Wells C., Pigliucci M., Adaptive phenotypic plasticity: the case of heterophylly in aquatic plants, *Perspect. Plant Ecol. Evol. Syst.* 3/1 (2000) 1–18.
- [52] Williams C.G., The influence of shoot ontogeny on juvenile-mature correlations in loblolly pine, *For. Sci.* 33 (1987) 411–422.