

Tillage management effects on pesticide fate in soils. A review

Lionel Alletto, Yves Coquet, Pierre Benoit, Djilali Heddadj, Enrique Barriuso

► To cite this version:

Lionel Alletto, Yves Coquet, Pierre Benoit, Djilali Heddadj, Enrique Barriuso. Tillage management effects on pesticide fate in soils. A review. Agronomy for Sustainable Development, 2010, 30 (2), 10.1051/agro/2009018 . hal-00886521

HAL Id: hal-00886521 https://hal.science/hal-00886521

Submitted on 11 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés. Agron. Sustain. Dev. 30 (2010) 367–400 © INRA, EDP Sciences, 2009 DOI: 10.1051/agro/2009018 Available online at: www.agronomy-journal.org



Review article

Tillage management effects on pesticide fate in soils. A review

Lionel ALLETTO^{1*}, Yves COQUET², Pierre BENOIT², Djilali HEDDADJ³, Enrique BARRIUSO²

¹ Université de Toulouse - École d'ingénieurs de Purpan, Agronomy Department, 75 voie du TOEC, BP 57 611, 31076 Toulouse Cedex 3, France
² UMR 1091 INRA/AgroParisTech Environment and Arable Crops, Institut National de la Recherche Agronomique/Institut National des Sciences et Industries du Vivant et de l'Environment, 78850 Thiverval-Grignon, France

³ Chambres d'agriculture de Bretagne, Recherche appliquée - Pôle agronomie, avenue du Général Borgnis Desbordes, BP 398, 56009 Vannes Cedex, France

(Accepted 4 May 2009)

Abstract – Reducing tillage intensity through the implementation of conservation practices is a way to reach a more sustainable agriculture. Reducing tillage is indeed an efficient way to control soil erosion and to decrease production costs. Nonetheless, the environmental impact of reduced tillage is not well known because conservation techniques may induce strong changes in soil physicochemical properties and biological activity. Knowledge on the fate of applied pesticides under conservation practices is particularly important from this point of view. We review here the advances in the understanding, quantification and prediction of the effects of tillage on pesticide fate in soils. We found the following major points: (1) for most dissipation processes such as retention, degradation and transfer, results of pesticide behaviour studies in soils are highly variable and sometimes contradictory. This variability is partially explained by the multiplicity of processes and contributive factors, by the variety of their interactions, and by their complex temporal and spatial dynamics. In addition, the lack of a thorough description of tillage systems and sampling strategy in most reports hampers any comprehensive interpretation of this variability. (2) Implementation of conservation tillage induces an increase in organic matter content at the soil surface and its gradual decrease with depth. This, in turn, leads to an increase in pesticide retention in the topsoil layer. (3) Increasing retention of pesticides in the topsoil layer under conservation tillage decreases the availability of the pesticides for biological degradation. This competition between retention and degradation leads to a higher persistence of pesticides in soils, though this persistence can be partially compensated for by a more intensive microbial activity under conservation tillage. (4) Despite strong changes in soil physical properties under conservation tillage, pesticide transfer is more influenced by initial soil conditions and climatic conditions than by tillage. Conservation tillage systems such as no-tillage improve macropore connectivity, which in turn increases pesticide leaching. We conclude that more knowledge is needed to fully understand the temporal and spatial dynamics of pesticide in soil, especially preferential flows, in order to improve the assessment of pesticide risks, and their relation to tillage management.

conventional tillage / conservation tillage / herbicide / retention / degradation / transport / soil carbon

1 2 2.1 2.2 2.3 2.4 2.5 2.6 3 3.1 3.2 3.3

Contents

^{*} Corresponding author: lionel.alletto@purpan.fr

L. Alletto et al.

4	Transfer of pesticides	
	4.1 Volatilisation	382
	4.2 Leaching	383
	4.3 Runoff	387
	4.4 Summary and recommendations	393
5	Conclusion	394

1. INTRODUCTION

Tillage can be defined as any mechanical operation on the soil and crop residues that aims at providing a suitable seedbed where crop seeds are sown. Soil preparation has always been an important component of traditional agriculture. Since 6000 BC, a wide variety of farming tools have been developed, from the simple digging stick to the paddle-shaped spade that could be pulled by humans or animals (Lal et al., 2007). In the 18th century, the mouldboard plough, a curved board that cuts and rotates the soil at a significant working depth, of more than 20 cm, was introduced in England. This tool was particularly efficient in incorporating organic residues, fertilisers and lime, in controlling weeds and helping the growing process. In the early part of the 19th century, the "initial" model of the mouldboard plough received several improvements to reach its "standardised" version by 1870, similar to current models (Hanson, 2006; Lal et al., 2007).

In the United States, between 1910 and 1930, the use of the mouldboard plough widely expanded, especially in the Great Plains of the Middle West. In Europe, its use started to be significant after 1945. These changes in tillage practices were accompanied by significant progresses in seed selection and by the development of the use of chemicals for plant protection, allowing a significant increase in crop yields. For example, wheat yields in France increased from 0.8-1 tons per hectare in 1945 to more than 8 tons in 1985. Because of its success, the mouldboard plough became a centrepiece of traditional agriculture.

In parallel with this growth of agricultural production, the widespread use of the mouldboard plough led to severe problems of soil and environmental degradation. One of the most well-known environmental problems was the drastic increase in wind erosion known as the "Dust Bowl" in 1930 in the US Great Plains (Lal et al., 2007; Masutti, 2004). Faced with this major problem, depriving some areas of any production capacity, the American reaction was swift. In 1935, the Soil Conservation Service - now called the Natural Resources Conservation Service - was created within the United States Department of Agriculture. Many important research programmes and communication efforts aimed at promoting the adoption of soil conservation techniques, grouped under the term "conservation tillage". These techniques are defined as any tillage and planting system that leaves at least 30% of the soil surface covered by crop residue after planting to reduce soil erosion by water, or at least 1.1 tons of crop residue/ha to reduce soil erosion by wind. A broad spectrum of farming methods can thus be classified as conservation techniques: from tillage techniques with a high working depth, such as sub-soiling, to reduced tillage of 0-15 cm or direct drilling techniques without any preliminary soil tillage, called "nosurfaces sown without ploughing (%)



Figure 1. Evolution of cultivated surfaces without ploughing in France between 1994 and 2006. Whatever the crop, the trend is an increase of surfaces cultivated without ploughing and winter crops are more often sown without ploughing than spring crops (Agreste, 2004, 2008).

tillage". This last type of tillage technique remained less developed until 1940 with the discovery of hormonal herbicides such as 2,4-dichlorophenoxy acetic acid (2,4D) that allowed farmers to control weeds without ploughing. Fairly quickly, these North American techniques thrived in the South American countries, particularly in Brazil where water erosion was significant (Bernoux et al., 2006), and in Argentina and Chile. Tillage practices and their codes used in this review are reported in Table I.

In Europe, because soil erosion was not as great as in the US, the interest of farmers in conservation tillage was limited. In the years 1970–1980, some attempts to develop these techniques were faced with decreases in crop yields, while, at the same time, farmers were encouraged to produce more to earn more. At the beginning of the 1990s, with the Rio conference (1992) which laid down the foundations for a sustainable agriculture, the concerns moved towards considering the vulnerability of water, soil and landscape resources. Moreover, the emergence of agricultural product surpluses led to lower prices. The control by farmers of their income more than ever implied lower production costs and a further increase in productivity, especially as the size of farms expanded. Under this economic context, soil conservation tillage, deemed to be faster and less costly in time and energy than conventional tillage, finally started to become attractive.

Progressively, surfaces sown in France without ploughing increased for all crops (Fig. 1). Spring crops are, however, more frequently sown after ploughing than winter crops. Moreover, the development of conservation techniques varies according to the regions. For example, in the south-west part of France – Aquitaine and Midi-Pyrénées regions –, more than

Table I. Tillage practice types used in this review article.

Code	Tillage practice	% soil surface covered by crop residues [†]	Agricultural tool
CT	Conventional tillage	< 30% (0–10%)	Mouldboard plough
			Disk plough
			Spading machine
CnT	Conservation tillage		
MT _{Dk}	Mulch tillage	> 30% (30–50%)	Disk harrow
MT_{Ch}	Mulch tillage	> 30% (30–50%)	Chisel plough, harrow
SS	Sub-soiling	> 30% (30-60%)	Sub-soiler, deep ripper, paratill
RT	Ridge tillage	> 30% (40-60%)	Ridger
ST	Strip tillage	> 30% (40-60%)	Strip-till
NT	No-tillage (or direct drilling)	> 50% (depending on crop residues)	Seeding drill (with disc openers)

[†] (CTIC, 2006).

Number of herbicide treatments



Figure 2. Mean number of herbicide treatments applied in 2006 for several crops sown after ploughing (conventional tillage) or without ploughing (conservation tillage). Note the systematic increase of herbicide treatments with the implementation of conservation tillage (Agreste, 2004, 2008).

75% of the wheat surfaces were not ploughed in 2006 against 50% in the whole country (Agreste, 2008).

One of the main roles of tillage is to provide an efficient control of weeds. Indeed, tillage influences weed populations by the combined effects of mechanical destruction of weed seedlings and by changing the vertical distribution of weed seeds in soil (Peigné et al., 2007). It also changes the soil climatic conditions which control weed dormancy, germination and growth. Reducing tillage intensity generally tends to increase the concentration of weeds in the topsoil (Moonen and Barberi, 2004; Torresen et al., 2003; Vasileiadis et al., 2007) and is very often associated with an increase in herbicide use (Fig. 2).

Although many studies have been conducted, mainly on the North American continent, the environmental fate of pesticides under conservation tillage presents many contradictions and remains, finally, poorly understood (Aubertot et al., 2005; Réal et al., 2007). Publication of articles concerning the effects of tillage practices on pesticide fate in soils, water and air started slowly during the 1970s and showed a rapid increase at the end of the 1980s followed by a peak rate of publication at the end of the 1990s (Fig. 3a). One-third of the published articles were found in one journal, and the first three journals were from the US, illustrating the great interest of US scientists in this subject (Fig. 3b).





Figure 3. Articles published on the effects of tillage practices on pesticide fate in soils, water and air. (*a*) Time series of articles found per year; and (*b*) major source journals found in (a). The plot includes the top 8 journals representing 115 articles out of the 144 references cited in the review.

Reducing tillage intensity leads to significant and complex changes in soil physical, chemical and biological properties, most often interrelated with each other, thus affecting the fate of the applied pesticides. A first review, conducted by Locke and Bryson (1997), dealt with herbicide interactions in soils under plant residue-managed farming systems. Following their work, we sought to assess the current state of knowledge of the effects of tillage on the processes involved in the fate of pesticides in soils cultivated with arable crops. Figure 4 summarises



Figure 4. Relationships between the processes conditioning the fate of pesticides in soils, water and air and the soil factors modified by tillage operations.

the main changes in soil properties due to tillage practices and their effects on the mechanisms involved in pesticide fate. As shown in this figure, understanding the effects of tillage on pesticide fate in soils implies a comprehensive evaluation of all the interactions between the different dissipation processes and of all the different soil factors affected by tillage operations. Some of these interactions are now well known, but most of them are still poorly understood. In this review, for each of these processes, we paid particular attention to the understanding of the mechanisms involved, their relative importance and the origin of the contradictory results found. The lists of the molecules, mainly herbicides, mentioned in this review are shown in Table II.

2. INTERCEPTION – RETENTION

Pesticide interception depends on several factors such as the type of application (pre- or post-emergence), the presence of a

crop and, if any, plant growth stage and species, the presence of a mulch, etc. Because interception of pesticides by plant foliage and its effect on the environment have not been widely studied (Reddy and Locke, 1996), the subject developed in this part focuses mainly on the interception and retention by the mulch and by the soil. In some cases, due to the difficulty of formally separating interception and retention, these processes are discussed simultaneously.

2.1. Mulch effect

The accumulation of organic residues on soil surface (mulch) in conservation tillage generally leads to an increase in the interception of applied pesticides, more particularly for apolar pesticides or pesticides with a low polarity (Locke, 1992; Novak et al., 1996; Reddy and Locke, 1998; Reddy et al., 1997a, b; Zablotowicz et al., 2000). This interception

Table II. Chemical and common names for pesticides mentioned in this review.

Common name	Nature	Chemical name
Acetochlor	Herbicide	2'-ethyl-6'-methyl-N-(ethoxymethyl)-2-chloroacetylanilide
Acifluorfen	Herbicide	5-[2-chloro-4-(trifluoromethyl)phenoxy]-2-nitrobenzoic acid
Alachlor	Herbicide	2-chloro-N-(2, 6-diethylphenyl)-N-(methoxymethyl)acetamide
Atrazine	Herbicide	(6-chloro-N-ethyl)-N'-(1-methylethyl)-1,3,5-triazine-2,4-diamine
Bentazone	Herbicide	3-isopropyl-1H-2,1,3-benzonthiadiazain-(4)3H-one 2,2-dioxide
Carbofuran	Insecticide	2,3-dihydro-2,2-dimethyl-7-benzofuranyl methylcarbamate
Chlorimuron	Herbicide	Ethyl-2-[[[[4-chloro-6-methoxy-2-pyrimidinyl)amino]carbonyl]amino]sulfonyl]benzoic acid
Clomazone	Herbicide	2-[(2-chlorophenyl)methyl]-4,4-dimethyl-3-isoxazolidinone
Clopyralid	Herbicide	3,6-dichloropicolinic acid
Chlorpyrifos	Insecticide	O, O-diethyl-O-(3,5,6-trichloro-2-pyridyl) phosphorothioate
Cyanazine	Herbicide	2-4-chloro-6-)ethylamino) 1,3,5-triazin-2-yl]amino-2-methylpropanenitrile
Diazinon	Insecticide	O, O-dimethyl O-2-isopropyl-6-methylpyrimidin4-yl phosphorothioate
Dicamba	Herbicide	3,6-dichloro-2-methoxybenzoic acid
Diclofop	Herbicide	(±)-2-(4-(2,4-dichlorophenoxy)phenoxy)propanoic acid
Diclosulam	Herbicide	N-(2,6-dichlorophenyl)-5-ethoxy-7-fluoro-(1,2,4)triazolo(1,5-c)pyrimidine-2-sulfonamide
Diketonitrile	Herbicide	2-cyclopropyl-3-(2-mesyl-4-trifluoromethylphenyl)-3-oxopropanenitrile
Dimethipin	Defoliant	2,3-dihydro-5,6-dimethyl-1,4-dithiin 1,1,4,4-tetraoxide
Fluometuron	Herbicide	N,N-dimethyl-N'-[3-(trifluoromethyl)-phenyl]-urea
Fonofos	Insecticide	<i>O</i> -ethyl <i>S</i> -phenyl ethylphosphonodithioate
Glyphosate	Herbicide	N-(phosphonomethyl)glycine
Imazapic	Herbicide	(RS)-2-(4,5-dihydro-4-isopropyl-4-methyl-5-oxoimidazol-2-yl)-5-methylnicotinic acid
Imazapyr	Herbicide	2-(4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1 <i>H</i> -imidazol-2-yl)-3-pyridine carboxylic acid
Imazaquin	Herbicide	2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-3-quinolinecarboxylic acid
Imazethapyr	Herbicide	(±)-2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1 <i>H</i> -imidiazol-2-yl]-5-ethyl-3-
		pyridinecarboxylic acid
Isoproturon	Herbicide	3-(4-isopropylphenyl)-1,1-dimethylurea
Isoxaflutole	Herbicide	5-cyclopropyl-1,2-isoxazol-4-yl alpha alpha alpha -trifluoro-2-mesyl-p-tolyl ketone
Linuron	Herbicide	3-(3,4-dichlorophenyl)-1-methoxy-1-methylurea
MCPA	Herbicide	4-chloro-2-methyphenoxy acetic acid
Metolachlor	Herbicide	2-chloro-N-(2-ethyl-6-methylphenyl)-N-(2-methoxy-1-methylethyl)acetamide
Metamitron	Herbicide	4-4-amino-3-methyl-6-phenyl-1,2,4-triazin-5(4H)-one
Metribuzin	Herbicide	4-amino-6-(1,1-dimethylethyl)-3-(methylthio)-1,2, 4-triazine-5(4H)-one
Norflurazon	Herbicide	4-chloro-5(methylamino)-2-(3-(trifluoromethyl)phenyl)-3(2H)-pyridazinone
Oxyfluorfen	Herbicide	2-chloro-1-(3-ethoxy-4-nitrophenoxy)-4-(trifluoro-methyl) benzene
Pendimethalin	Herbicide	N-(1-ethylpropyl)-3,4-dimethyl-2,6-dinitrobenzenamine
Propachlor	Herbicide	2-chloro-N-isopropylacetanilide
Simazine	Herbicide	2-chloro-4,6-bis[ethylamino]-s-triazine
Sulfentrazone	Herbicide	N-[2,4-dichloro-5-[4-(difluoromethyl)-4,5-dihydro-3-methyl-5-oxo-1H-1,2,4-triazol-1-
		yl]phenyl]methanesulfonamide
Terbufos	Insecticide	S-(((1,1-dimethylethyl)thio)methyl) O, O-diethyl phosphorodithioate
Terbuthylazine	Herbicide	N-2-tert-butyl-6-chloro-N-4-ethyl-1,3,5-triazine-2,4-diamine
Thidiazuron	Defoliant	N-phenyl-N-1,2,3-thidiazol-5-yl-urea
Tribufos	Defoliant	S,S,S-tributyl phosphorotrithioate
2,4 D	Herbicide	(2,4-dichlorophenoxy)acetic acid
2,4-Dichlorophenol	Herbicide precursor	2,4-Dichlorophenol

depends on the amount and type of crop residues and, for example, with a proportion $\ge 30\%$ of the soil surface covered, it was found to range from 40 to 70% of the applied dose for several preemergence herbicides (Banks and Robinson, 1982; Ghadiri et al., 1984; Isensee and Sadeghi, 1994; Sadeghi and Isensee, 1997). In terms of mass of residues, for a quantity > 4.5 t ha⁻¹, the interception of acetochlor, alachlor and meto-

lachlor was higher than 90% of the applied doses (Banks and Robinson, 1986). Retention processes are closely associated with the interception of pesticides. Indeed, crop residues can have sorption capacities 10 to 60 times higher than soil (Boyd et al., 1990; Reddy et al., 1995b) and can significantly modify the (bio)availability and the migration of pesticides in soil. As a result, a loss of efficacy of some pesticides has been observed (Erbach and Lovely, 1975; Mills et al., 1989; Shelton et al., 1998) which can lead to an increase in the applied doses (Shelton et al., 1998; Worsham, 1991). By penetrating inside crop residues, most of the molecules are physically entrapped within cell wall structures, such as cellulose microfibrils embedded in a lignin-hemicellulose matrix (Dao, 1991) and, most often, lose their activity.

The nature and decomposition degree of crop residues both influence interception and retention of pesticides, but in contrasted ways depending on studies. For example, hairy vetch (Vicia villosa Roth) residues had a higher retention for chlorimuron than rye (Secale cereale L.) residues (Reddy et al., 1995) and a higher retention for fluometuron than wheat (Triticum aestivum L.) residues (Gaston et al., 2001) (Tab. III). This greater sorption by vetch than rye or wheat residues may be related to differences in physical state, that offered a greater surface area for herbicide sorption in the case of the vetch, and in composition of the residues that contained less cellulose and more amino acids (Gaston et al., 2001). Sigua et al. (1993) indicated that interception of atrazine was enhanced with fresh maize residues, due to a combination of a greater hydrophobicity and a higher sorption capacity of the fresh compared with the aged maize residues. In contrast, with metribuzin (Dao, 1991), chlorimuron (Reddy et al., 1995b) and cyanazine (Reddy et al., 1997a), interception was higher with aged residues. In these cases, the increase in sorption due to aging was attributed to both changes in the physical state and in the chemical composition of the crop residues. On one hand, physical alterations of the residues increase their external surface area for herbicide sorption compared with fresh residues. On the other hand, the chemical evolution of the residues during degradation leads to an increase in the lignin/cellulose ratio, resulting in an increase in the sorption of chlorimuron and metribuzin (Dao, 1991; Reddy et al., 1995b).

Interception is not only influenced by the amount of mulch. It is also conditioned by the formulation of the molecules. Compared with the commercial formulation, the use of microencapsulated or granular forms of alachlor led to a decrease in its interception and, finally, the herbicide reached the soil surface more rapidly (Johnson et al., 1989; Sadeghi et al., 1998).

Once intercepted by the mulch, the molecule can be washed off from the crop residues to reach the soil surface. This washoff depends on the timing between pesticide treatment and the first rainfall and its intensity. On maize residues, Martin et al. (1978) reported a range of 30 to 60% wash-off by the first 5 mm of water for alachlor, atrazine, cyanazine and propachlor, which was equivalent to the wash-off by the next 30 mm of water. At the end of their experiment, most of the applied chemicals had been washed off from the mulch. Timing between treatment and first rainfall also strongly influences the quantity of pesticides that could be washed from the mulch. It was found to be the highest after heavy rainfall occurring within two weeks following the treatment (Sadeghi and Isensee, 1996; 1997). Depending on rainfall quantity and intensity, between 70 and 96% of intercepted atrazine were released between the 1st and 3rd weeks after treatment (Ghadiri et al., 1984; Isensee and Sadeghi, 1994). However, this washoff directly depends on the retention capacities of the pesticide on crop residues (Gaston et al., 2001). For example, metolachlor, once intercepted, seems to form stronger and less reversible bonds with crop residues than acetochlor or alachlor (Banks and Robinson, 1986). In some cases, a gradual release of the pesticide from the residues by wash-off may lead to an increase in its efficacy (Dao, 1991), but could also increase crop injuries (Barnes et al., 1989; Loux et al., 1989; Mills and Witt, 1991).

2.2. Organic carbon content effect

One of the main changes related to the implementation of conservation tillage compared with conventional tillage is the redistribution of organic carbon in the soil (Balesdent et al., 1990; Tebrügge and During, 1999). Generally, organic carbon content increases in surface soil due to the presence and decomposition of the mulch and gradually decreases with depth (Lal et al., 1994; Pinheiro et al., 2004; Six et al., 1999). For most pesticides, organic carbon content and adsorption are positively correlated, resulting in a higher adsorption in surface soil under conservation tillage than under conventional tillage (Tab. III). For molecules with low sorption capacity, such as bentazon (Gaston et al., 1996) or diclosulam (Lavorenti et al., 2003), the effect of tillage may not be significant. Very few studies aimed at analysing the effects of tillage on pesticide sorption kinetics. For acifluorfen (Gaston and Locke, 2000), chlorimuron (Reddy et al., 1995a) and cyanazine (Reddy et al., 1997b), no effect of tillage system was found on sorption kinetics and most of the sorption occurred during the first hour of contact. For some molecules, such as alachlor (Locke, 1992) and sulfentrazone (Reddy and Locke, 1998), sorption was faster under conservation tillage and was positively correlated with higher soil organic matter contents.

Tillage management also modifies pesticide desorption. For sulfentrazone and alachlor, whatever the tillage system, desorption was found to be hysteretic, but a greater proportion of the molecule, representing approximately 35% of the applied sulfentrazone, remained sorbed under no-tillage (against 20% under conventional tillage) (Locke, 1992; Reddy and Locke, 1998). For cyanazine (Reddy et al., 1997b), desorption was also more reversible under conventional tillage but, when ryegrass residues were added to the soil samples, desorption was similar under conventional and conservation tillage. As indicated by Locke (1992), desorption tends to decline with increasing contact time between soil and pesticide due to diffusion processes within the soil.

However, considering only the quantitative increase in organic carbon content in surface soil under conservation tillage is not sufficient to explain the increase in pesticide sorption found in some studies (Ding et al., 2002b; Novak et al., 1996; Zablotowicz et al., 2000). Although the analytical procedures used to extract soil organic matter can induce artefacts by modifying its molecular structure (Ding et al., 2002b; Salloum et al., 2001), it appears that the nature of soil organic matter

Pesticide	Tillage ¹	Depth.		Soil properties		KD	$K_{ m F}$	K _{OC}	Reference
			Clay	Organic carbon (OC)	рН	_			
		cm		${\rm g}~{\rm kg}^{-1}$		$L kg^{-1}$	$L^{n_{\rm F}} {\rm mg}^{(1-n_{\rm F})} {\rm kg}^{-1}$	$L kg^{-1} OC$	
Acetochlor	NT CT	0-10	-	16.6 13.3	-	2.7 1.7	-	166 126	(Ferri et al., 2002)
	NT	0-10		10.2	5.6*		0.8		
	NT	10-20	-	5.6	5.3*	-	0.5	-	
	NT	20-30	-	4.4	5.7*	-	0.6	-	
Acifluorfen	CT	0-10	-	8.7	5.8*	-	0.8	-	(Gaston and Locke 2000)
Actinuorien	CT	10-20	-	6.4	5.8*	-	0.5	-	(Gaston and Locke, 2000)
	CT	20-30	-	4.9	5.8*	-	0.5	-	
	NT	0-5	290	16.7	5.5*	5.4	5.6	323	
Alachlor	CT	0-5	260	10.2	5.5*	3.5	3.6	342	(Locke, 1992)
	NT	0-5	-	72.5	4.7*	-	6.0	-	
	NT	5-10	-	29.0	5.0*	-	3.6	-	
	NT	10-15	-	29.5	6.2*	-	4.0	-	
	NT	15-20	-	32.5	6.4*	-	3.7	-	
	CT	0-5	-	30.0	5.5*	-	3.6	-	
Alachlor	CT	5-10	-	26.0	6.0^{*}	-	4.0	-	(Clay et al., 1991)
	CT	10-15	-	29.0	5.2*	-	3.9	-	
	CT	15-20	-	28.0	5.7*	-	4.0	-	
	$NT + B^2$		-	8.4	-	0.9	-	110	
	$NT + V^2$		-	8.6	-	1.1	-	130	
	NT		-	6.7	-	1.0	-	143	
Alachlora	CT + B	0.15	-	4.7	-	0.5	-	99	$(\mathbf{X}_{\mathbf{u}\mathbf{v}} \text{ at al} 1007)$
Alaciiloite	CT + V	0-15	-	5.7	-	0.5	-	95	(Aue et al., 1997)
	CT		-	6.3	-	0.6	-	91	
	NT	0-2.5	-	19.1	5.2	7.8	-	-	
	NT	2.5-5	-	15.1	5.0	7.1	-	-	
	NT	5-10	-	10.4	5.8	7.0	-	-	
	NT	10-15	-	10.4	6.2	7.3	-	-	
	NT	15-20	-	10.4	6.3	7.0	-	-	
	NT	20-25	-	8.1	6.5	6.2	-	-	
	NT	25-30	-	7.0	6.7	6.1	-	-	
	NT	30-35	-	6.4	6.8	6.1	-	-	
	NT	35-40	-	5.8	6.9	5.8	-	-	
	CT	0-2.5	-	12.8	5.6	7.4	-	-	
	CT	2.5-5	-	12.8	5.2	1.5	-	-	
Atrazine	CT	3-10 10 15	-	11.0	5.7	7.0 6.7	-	-	(Ghadiri et al., 1984)
11000000	CT	15 20	-	10.4	6.3	6.5	-	-	
	CT	20-25	_	8 1	6.6	6.5	-	-	
	CT	25-30	_	7.0	6.7	6.0	-	_	
	CT	30-35	-	5.2	6.9	5.7	-	-	
	CT	35-40	-	4.7	7.0	5.5	-	-	
	NT	0.15		10.6		17			
Atrazine	CT	0-15	-	9.8	-	1.7	-	-	(Novak et al., 1996)
	NT	0-10	-	10.2	5.6*			-	
	NT	10-20	-	5.6	5.3*			-	
	NT	20-30	-	4.4	5.7*	0.03	0.04	-	
Bentazon	CT	0-10	-	8.7	5.8*	0.05	0.04	-	(Gaston et al., 1996)
Semazon	CT	10-20	-	6.4	5.8*			-	())))
	CT	20-30	-	4.9	5.8*			-	

Table III. Sorption properties of pesticides under different tillage practices.

Table III. Continued.

Pesticide	de Tillage ¹ Depth.			Soil properties		KD	$K_{\rm F}$	K _{OC}	Reference	
	U	1	Clay	Organic	pН		•	00		
				carbon (OC)						
		cm		$g kg^{-1}$		$L kg^{-1}$	$L^{n_{\rm F}} {\rm mg}^{(1-n_{\rm F})} {\rm kg}^{-1}$	$L kg^{-1} OC$		
	NT	0-7.5	280	30.6	6.6*	-	1.0	31		
	CT		250	17.8	6.4*	-	0.6	31		
Chlorimuron	NT		330	46.0	6.2*	-	6.5	142	(Reddy et al. 1995a)	
Chiofinitaton	CT		260	19.2	4.5*	-	2.1	108	(Reduy et al., 1995a)	
	NT		-	15.3	5.4*	-	2.1	141		
	СТ		-	13.1	5.7*	-	1.6	125		
	NT $(9)^3$	0-5	145	23.7	5.1	1.0	-	21		
Clopyralid	CT		145	22.7	5.8	0.3	-	2	(Shang and Arshad, 1998)	
	NT (8)		532	47.0	4.9	1.6	-	51		
	CT (10)	0.5	520	44.5	5.4	2.1	-	32		
Cyanazine	NT (10) CT	0-5	230	21.4	5.2	3.5	-	165	(Reddy et al., 1997b)	
	VT (0)	0.5	145	22.7	5.5	2.2	-	140		
	NI (9) CT	0-3	145	23.7	5.1	U.2 Undetected	-	15 Undetected		
Dicamba			522	47.0	J.0		-	22	(Shang and Arshad, 1998)	
	CT		520	47.0	4.9 5.4	1.4	-	23		
	NT	0-10	355	19.9	5.0*	1.0	1.8	98		
Diclosulam	СТ	0-10	341	12.8	4 4*	2.0	1.0	159	(Lavorenti et al., 2003)	
	$NT + Adv^2$	0-3	-	13.9	5.2	-	2.0	-		
	NT + Adv	3-6	-	8.0	5.5	_	1.0	_		
	NT+V	0-3	_	15.6	5.5	-	1.8	-		
	NT+V	3-6	-	7.3	5.2	-	1.2	-		
	NT+B	0-3	-	11.2	6.1	-	1.8	-		
	NT+B	3-6	-	4.7	5.5	-	1.3	-		
Fluometuron	CT+ Adv	0-3	-	7.4	5.7	-	0.8	-	(Gaston et al., 2001)	
	CT+ Adv	3-6	-	7.0	5.2	-	1.2	-		
	CT+V	0-3	-	7.4	5.5	-	0.9	-		
	CT+V	3-6	-	6.6	5.0	-	1.1	-		
	CT+B	0-3	-	7.8	5.8	-	1.0	-		
	CT+B	3-6	-	5.8	5.1	-	0.8	-		
	NT	0-4	150	20.0	5.1	2.2	-	-		
	NT	4-8	150	10.0	5.5	1.7	-	-		
	NT	8-15	160	9.0	6.5	1.4	-	-		
	CT	0-4	130	13.0	5.5	1.7	-	-		
	CT	4-8	140	13.0	5.7	2.1	-	-		
Fluometuron	CT	8-15	150	11.0	6.3	1.4	-	-	(Brown et al., 1994)	
	NI+V	0-4	150	25.0	4./	3.0	-	-		
	NI+V	4-8	150	11.0	4.9	1.0	-	-		
	NI + V	8-15	150	9.0	5.7	1.3	-	-		
	CT + V	18	140	15.0	5.0	2.1	-	-		
	CT + V	4-0 8 15	150	11.0	5.6	1.6	-	-		
	NT (11)	0-2	150	30.1	5.0	1.0	7.2	-		
	NT (11)	2-5	_	11.8	_	_	2.0	_		
	NT	5-10	_	59	-	_	1.4	_		
	NT	10-25	_	33	-	_	1.4	_		
Fluometuron	СТ	0-2	-	20.3	_	-	1.9	-	(Zablotowicz et al., 2000)	
	CT	2-5	-	11.2	-	-	2.2	-		
	СТ	5-10	-	6.9	-	-	1.0	-		
	СТ	10-25	-	4.4	-	-	1.2	-		
El	NT	0-8	130	24.5	5.1*	2.0	-	80	(0-h	
Fluometuron	СТ	0-8	100	14.2	5.2*	1.3	-	92	(Suba and Essington, 1999)	

Table III. Continued.

Pesticide	Tillage ¹	Depth.		Soil properties		KD	$K_{\rm F}$	K _{OC}	Reference
			Clay	Organic carbon	pН	-			
				(OC)					
		cm		g kg ⁻¹		L kg ⁻¹	$L^{n_{\rm F}} {\rm mg}^{(1-n_{\rm F})} {\rm kg}^{-1}$	L kg ⁻¹ OC	
	NT	0-2	-	13.9	6.7	2.4	-	-	
	NT D 2	2-10	-	6.1	5.9	1.6	-	-	
	NT+Rg ²	0-2	-	19.8	6.0	5.0	-	-	
	NT+Rg	2-10	-	6.9	5.7	1.4	-	-	
	CT	0-2	-	8.0	6.5	1.7	-	-	
	CT	2-10	-	6.0	5.8	1.6	-	-	
	CT+Rg	0-2	-	9.5	6.4	2.1	-	-	
	CI+Kg	2-10	-	/.3	5.7	2.0	-	-	
		0-2	-	12.2	0.5	0.6	-	-	
		2-10	-	0./	0.7	0.4	-	-	
	KI+S'	0-2	-	12.0	0.0	0.7	-	-	
	KI+5 CT	2-10	-	7.1	6.0	0.3	-	-	
	CT	2 10	-	0.4	6.4	0.4	-	-	
Fluometuron	CT	0.2	-	6.4 6.8	6.1	0.4	-	-	(Locke et al., 2005)
	CT+S	2 10	-	0.8	6.6	0.4	-	-	
	RT	0-2	-	10.3	5.5	0.5	-	-	
	RT	2-10		9.5	5.5	0.5	_	_	
	RT+S	0-2	_	8.8	5.5	0.5	_	_	
	RT+S	2-10	-	8.5	5.9	0.5	_	_	
	CT	0-2	-	9.0	5.6	0.4	_	-	
	CT	2-10	-	8.6	5.4	0.4	_	-	
	CT+S	0-2	-	8.7	5.7	0.4	-	-	
	CT+S	2-10	-	8.6	5.5	0.4	-	-	
	NT (9)		145	23.7	5.1	2.0	-	63	
	CT		145	22.7	5.8	0.7	_	40	
MCPA	NT (8)	0-5	532	47.0	J.0 1 Q	73		168	(Shang and Arshad, 1998)
	CT		520	44.5	4.) 5.4	10.4	_	182	
	NT	0.5	520	15.4	5.1	10.1	2.5	220	
	NT	10.15	-	3.5	-	-	5.5	230	
Metolachlor	CT	0-5	-	8.5	-	-	2.0	240	(Ding et al., 2002b)
	СТ	10-15		5.4	-		1.4	240	
	NT	0.9	120	24.5	5.1*	12.1	1.1	406	
Norflurazon		0-8	100	24.3	5.1 5.2*	12.1	-	490	(Suba and Essington, 1999)
		0-8	100	14.2	5.2	0.5	-	430	
	NT	0-2	-	13.9	6.7	2.2	-	-	
	NT NT D	2-10	-	6.1	5.9	1.6	-	-	
	NI+Kg	0-2	-	19.8	6.0	3.6	-	-	
	NI+Kg	2-10	-	0.9	5.7	1.0	-	-	
	CT	0-2	-	8.0	0.5 5 9	1.0	-	-	
Norflurazon		2-10	-	0.0	5.0	1.8	-	-	(Locke et al., 2005)
	CT+Rg	2 10	-	9.5	57	2.5	-	-	
	NT	2-10	-	7.5	5.1	2.1	-	- 20	
	N I CT		280	30.6	6.6°	0.9	1.2	30	
			250	17.8	0.4 [*]	0.8	1.0	47	
Sulfentrazon		0-7.5	260	40.0	0.2	3.2	5.4 2.6	/1	(Reddy and Locke 1998)
Sulfelituzon	NT	0 7.5	200	19.2	4.3 5.0*	2.9	2.0	155	(Reduy and Ebeke, 1990)
	CT		230 210	21. 4 16.0	5.2 5.3*	2.3 1.5	5.5 1.8	90 106	
	01		210	10.0	5.5	1.5	1.0	100	
	NT (9)	0-3	188	27.0	6.6*	4.2	-	155	
	NT	3-10	177	16.7	6.6°	2.2	-	132	
2.4 Dichlorophonel	N I CT	10-25	168	10.3	6.4	1.3	-	122	(Düring et al. 2002)
	CT	0-5	152	11.5	0.1	1./	-	150	(During et al., 2002)
	CT	3-10	155	11.0	0.1	1.0	-	140	
	U	10-25	143	11.5	0.2*	1.4	-	126	

¹ Codes of tillage practices are reported in Table I. ² Adv: weed residues; B: wheat residues; Rg: ray-grass residues; S: rye residues; V: vetch residues.

³ Numbers in parenthesis indicate the age in years of the tillage system.

* pH CaCl₂.

and, hence, its reactivity, are also affected by tillage (Ding et al., 2002a; Sleutel et al., 2007; Tatzber et al., 2008) and by the nature of crop residues (Ding et al., 2006). In soil organic matter collected under a no-tillage system, Preston et al. (1994) found a higher number of reactive functional groups compared with that collected under a conventional tillage system. Ding et al. (2002a) found that humic acids and humin from conservation tillage contained more aliphatic carbon and less aromatic carbon than under conventional tillage. A larger amount of amino acids were also linked to these humic acids under no-tillage, whereas they were more linked to fulvic acids under conventional tillage (Szajdak et al., 2003). Moreover, reactive/recalcitrant peak ratios indicated that humic acids in the topsoil (0-5 cm-depth) were more biologically active under conservation tillage than under conventional tillage. With depth, the soil organic matter composition changes. Humification processes are more advanced in deeper soil layers and lead to an increase in aromaticity with the development of polycondensed rings (Ding et al., 2002a). The reactivity of organic matter fractions may also affect pesticide desorption. As an example, desorption of metolachlor from conventional tillage humic acids was less significant than from conservation tillage humic acids (Ding et al., 2002b). The hysteresis index (which is the ratio of Freundlich exponents for desorption and sorption) was lower for humic acids under conventional tillage, indicating that sorbed metolachlor molecules were more difficult to desorb. This may be caused by higher aromatic carbon contents in conventional tillage humic acids (Ding et al., 2002b; Xing, 2001).

The nature of crop residues also influences the composition of soil organic matter. For example, rye residues tend to form more aromatic and less aliphatic humic acids than a mix of vetch/rye residues (Ding et al., 2006), which could potentially affect pesticide behaviour and efficacy in soil (Ding et al., 2002a; Nanny and Maza, 2001). However, to our current knowledge, little work has been published on the effects of these changes in soil organic matter composition due to tillage on pesticide sorption. Ding et al. (2002b) highlighted a stronger sorption of metolachlor by humin than by humic acids, and humin content was found to be higher under notillage systems (Stearman et al., 1989). For humin, which is a highly condensed form of soil organic matter, several studies showed that sorption was more variable and that sorption isotherms were less linear (Xing and Pignatello, 1997; Yuan and Xing, 1999). In contrast, Stearman et al. (1989) observed a higher sorption of metribuzin and oxyfluorfen on fulvic and humic acids, which are abundant in conventional tillage, than on humin, which is abundant in conservation tillage. Dissolved organic carbon content is most often higher under conservation tillage and is mainly composed of small molecules, with a size similar to that of fulvic acids (Suba and Essington, 1999). In some studies, sorption of pesticides on dissolved organic carbon was found to be an efficient way to increase the mobility of some pesticides in soils, e.g. atrazine and 2,4D, leading to a significant transport of these chemicals through the soil profile (Chin et al., 1990; Gao et al., 1998; Lafrance et al., 1994; Li et al., 2005). In another study, sorption of fluometuron and norflurazon on the dissolved organic carbon formed

under conservation tillage did not lead to a higher leaching of these molecules (Suba and Essington, 1999).

2.3. Soil pH effect

Soil pH may be differently modified by tillage techniques: conservation tillage leads to an increase in pH (Reddy and Locke, 1998), no change (Comia et al., 1994; Rasmussen, 1991) or, more often, to a decrease in pH, especially in surface soil due to the accumulation of organic matter and fertilisers (Arshad et al., 1999; Doran, 1980; Levanon et al., 1994). For many pesticides, sorption is strongly influenced by soil pH and tends to increase when soil pH decreases (Barriuso and Calvet, 1992; Barriuso et al., 1992; Grey et al., 1997). In sandy soils, it has been found that, for similar organic carbon contents between tillage treatments, acidification in conservation techniques could significantly increase the sorption of several molecules according to their pKa (Shang and Arshad, 1998). For s-triazines (weak bases), the decrease in pH causes their protonation, promoting sorption on organic matter (Senesi and Testini, 1982) and decreasing their herbicide activity.

2.4. Other effects

Tillage techniques modify other soil properties (Fig. 4) whose effects on retention mechanisms are still poorly understood. Due to the presence of a mulch, water content in conservation techniques is generally higher than in conventional techniques (Drury et al., 1999; Shelton et al., 1998). This increase in soil water content led, in some cases, to a decrease in pesticide retention (Dao and Lavy, 1991; Walker, 1971). In other cases, it led to an increase in adsorption that could be explained by a decrease in the hydrophobicity of the organic compounds and an access to sorption sites facilitated by the presence of water (Berglof et al., 2000; Ochsner et al., 2006). Soil temperature is also modified by the mulch under conservation tillage. The mulch intercepts light energy, thus reducing temperature at the soil surface in conservation techniques, from 1 to 5 °C (Bragagnolo and Mielniczuk, 1990; Grant et al., 1990; Gupta et al., 1988). However, to our current knowledge, there is no published data about the consequences of soil water content and temperature modifications by tillage management on pesticide retention.

2.5. Sorption of metabolites

Little information is available on tillage effects on metabolite retention. According to their molecular composition, metabolites can have lower sorption capacities, e.g. cyanazine metabolites (Reddy et al., 1997a), or higher sorption capacities, e.g. atrazine and bentazon metabolites (Clay and Koskinen, 1990; Gaston et al., 1996) than the parent compound. For cyanazine metabolites, as observed for the parent compound, sorption was higher under conservation tillage than under conventional tillage.

2.6. Summary and recommendations

Regarding pesticide retention processes, most studies highlighted the role of the mulch in conservation tillage systems which intercepted the molecules. In a general way, tillage systems act on pesticide retention mainly through their effect on the distribution of organic matter in soil. Recent works on the characterisation of organic matter fractions and their reactivity towards pesticides show significant influences of the type of tillage and nature of crop residues. Using recent analytical technologies, these studies on the characterisation of organic compounds found in soils or crop residues are a significant advance in the qualitative evaluation of the effects of agricultural practices on pesticide retention. To progress in this direction, further research should not be focused on tillage management only, but rather on the whole cropping systems. As a matter of fact, other management options such as crop rotation, cover crops and manure application play important roles in the dynamics of soil organic matter. Regarding conservation tillage systems, research efforts should be made to clarify the indirect effects of the mulch via modifications of pH, water and temperature dynamics in the underlying soil.

3. DEGRADATION

Understanding pesticide degradation in soils is a key step for assessing their persistence and their risks of transfer in the environment. Degradation studies can be carried out in the field, thus integrating a set of non-controlled phenomena such as fluctuations in temperature, soil water content or radiation (Tab. IV), or in the laboratory, where soil samples are kept under controlled conditions (Tab. V). In all cases, degradation is conditioned by a combination of factors, i.e. microflora, organic matter, water content, temperature and pH, which are directly influenced by tillage (Fig. 4). Biotic degradation of pesticides is most often seen as quantitatively more important than abiotic degradation. In conservation tillage, the total microbial biomass is generally larger than in conventional tillage (Biederbeck et al., 1997; Dalal et al., 1991; Doran, 1980), but it remains strongly dependent on soil conditions. Fungi populations, which were found to be efficient in pesticide degradation (Barr and Aust, 1994), are generally larger in conservation tillage and their biomass increases as degree of tillage is reduced (Drijber et al., 2000; Kabir, 2005). These biological differences due to tillage have effects, sometimes contradictory, on the degradation of pesticides in the mulch and in the soil.

3.1. Pesticide degradation in the mulch

The interception by the mulch in conservation techniques can modify the persistence of the applied pesticides. These effects are sometimes negligible (Banks and Robinson, 1982), but more often they significantly affect the fate of pesticides in soil. Crop residues may have higher microbial activity and degradation capacity than the soil, thus reducing the concentrations of pesticide that reach the soil surface (Locke et al., 2005; Zablotowicz et al., 1998). Moreover, depending on the nature of the pesticide, interception by the mulch may generate photodegradation, thus reducing the persistence of the molecules (Selim et al., 2003). In other works, crop residues increased the residence time of pesticides because of the competition between retention and degradation processes, and a gradual release of the molecules by desorption was observed (Mazzoncini et al., 1998). By acting as a physical protector of the soil surface, crop residues may also limit the air flow between the soil and the atmosphere and within the soil, thus slowing down the activity of degrading microorganisms (Sorenson et al., 1991).

Depending on the nature of crop residues, the degradation of molecules can be affected by the presence of a mulch, but in contrasted ways. For example, in no-tillage, vetch residues accelerated the degradation of metolachlor by from 1.5 to 3 times, but had no effect on the degradation of atrazine (Teasdale et al., 2003). On the contrary, according to laboratory studies, vetch residues seemed to slow down fluometuron degradation compared with soil samples without vetch residues (Brown et al., 1994; Zablotowicz et al., 1998) or other types of residues such as wheat residues (Gaston et al., 2001), rye residues (Zablotowicz et al., 1998), or ray grass residues (Locke et al., 1995). This slowdown could be due to the abundance of nitrogen in legumes (Fabaceae). This nitrogen would be preferentially used by the microorganisms to the detriment of the N sources contained in pesticides. In the same way, Gan et al. (1996) observed a lower degradation rate of atrazine after an addition of nitrogen. However, experiments on undisturbed soil columns with simulated rainfall, that allowed nitrogen leaching and thus reduced the amount bioavailable in the soil column, relativised these results and no negative effect of vetch residues on fluometuron degradation was found (Gaston et al., 2003). This experiment provided an explanation of the difference between the results obtained in laboratory incubations where leaching of nitrogen is not possible (Gaston et al., 2001) and those obtained in the field where N can leach out through the soil (Brown et al., 1996).

3.2. Pesticide degradation in soil

Tillage practices were found to modify significantly pesticide degradation in soil, but in contrasted ways according to studies. Degradation of several pesticides was indeed found to be slower (Brown et al., 1994; Gaston and Locke, 2000; Otto et al., 1997; Ulbrich et al., 2005), equivalent (Gaynor et al., 1998; Locke et al., 1996, 2005; Monks and Banks, 1993; Reddy et al., 1995b; Renner et al., 1998), or faster (Gaston et al., 2001, 2003, Lavorenti et al., 2003; Levanon et al., 1994; Sadeghi and Isensee, 1997) under conservation tillage than under conventional tillage. Several reasons could explain these differences. To explain a lower degradation under conservation tillage, the most commonly mentioned phenomenon is the competition between retention and degradation. In conservation tillage, pesticide adsorption to the soil solid phase is

Pesticide	Tillage ¹	Depth		Soil properties		DT_{50}^{2}	Reference
			Clay	Organic carbon	pН		
		cm	-	g kg ⁻¹	-	d	
	NT	0-30	-	-	-	< 1	
Alachlor	СТ	0-30	-	-	-	< 1	(Weed et al., 1998)
	NT	0-110	-	-		8-41	
Alachlor	CT	0 110				8 /1	
encapsulated	NT	0.110	-	-	-	4 20	(Gish et al., 1994)
alachlor	IN I CTT	0-110	-	-	-	4-20	
	CI	0-110	-	-	-	4-20	
Atrazine	NT	0-10	-	-	-	50	(Ghadiri et al., 1984)
1 Infulline	CT	0 10	-	-	-	42	(Chadhir et all, 1901)
Atrozina	NT	0-50	-	-	-	71	(Cich et al. 1001)
Attazine	CT	0-70	-	-	-	73	(OISII et al., 1991)
	NT	0-110	-	-	-	36	
Atrazine	СТ	0-110	-	-	-	36	
encapsulated	NT	0-110	_	-	_	110	(Gish et al., 1994)
atrazine	CT	0-110	_	_	_	110	
	NT	0 10				110	
	INI NT D 3	0-10				45, 50, 50	
Atrazine	NI+Kg ⁵	0-10	390	14.5	5.4	40, 33, 54	(Gaynor et al., 2000)
	CI	0-10				46, 34, 54	
	CT+Rg	0-10				35, 59, 24	
	NT		-	-	-	43, 56, 43, 35	
A	RT top	0.10	-	-	-	33, 75, 53, 36	$(0, \dots, 0, 1, 1, 1, 1, 1, 1, 0, 0, 0)$
Atrazine	RT furrow	0-10	-	-	-	31, 47, 53, 35	(Gaynor et al., 1998)
	СТ		-	-	_	33, 62, 58, 47	
	NT		17-26	11.0	6-6.5	35, 02, 30, 17	(Isensee and Sadeghi
Atrazine	CT	0-50	16 24	7.0	1 9 5	27 21 20 18	
			55 1((7.0	4.0-3	37, 21, 29, 10	1994)
Atrazine	N I	0-150	55-166	0.1-9.7	5.3-0.1	149	(Heatwole et al., 1997)
	CI		59-179	1.2-4.4	5.5-6.7	215	
	NT (7) ⁵		-	-	-	12	
Atrazine	CT	0-50	-	-	-	22	(Sadeghi and Isensee 1996)
Auazine	$CT \rightarrow NT$	0-50	-	-	-	-, 21, 32	(Sadegin and Isensee, 1996)
	NT		-	-	-	-, 17, 23	
	$(7) \rightarrow CT$						
	RT		230	23.2	6.0	52, 91	
Clomazon	СТ	0-10				58,83	(Curran et al., 1992)
	NT		260	10.4	7.8	32, 14, 22	
	CT	-	200	10.4	7.8	<i>J2</i> , 14, <i>22</i>	
Chlorimuron		-	200	10.4	7.8	44, 15, 18	(Baughman et al., 1996)
	NI	-	220	8.7	0.3	17,82	-
	CI	-	220	8.7	6.3	14, 22	
Cvanazine	NT	-0-50	-	-	-	13	(Gish et al. 1991)
Cyanazine	CT	0-70	-	-	-	13	
Elmontotunon	NT, NT+Rg	0.2	-	-	-	7 15	(Leghe et al. 2005)
Fluometuron	CT, CT+Rg	0-2	-	-	-	7-13	(Locke et al., 2005)
	NT		150	8.7	5.2	30. 23	
	$NT+V^2$		150	10.4	48	38 19	
Fluometuron	CT	0-8	130	76	5.6	24, 26	(Brown et al., 1996)
	CT V		140	7.0 9.7	5.0	24, 20	
	CI+V	0.10	140	0.7	3.0	30, 23	
	NI (3)	0-10	780	20.3	4.7	66	
Imazapic	СГ	0-10				45	(Ulbrich et al., 2005)
mazapro	NT	0-10	280	27.2	5 8	35	(01011011 01 011, 2000)
	CT	0-10	280	21.5	5.0	32	
	NT (3)	0-10	790	20.2	4 7	53	
Ŧ	CT	0-10	/80	20.3	4./	50	
Imazapyr	NT	0-10				43	(Ulbrich et al., 2005)
	СТ	0-10	280	27.3	5.8	45	
	RT RT	0.10				39 144	
Imazaquin	CT	0-10	230	23.2	6.0	<i>Jj</i> , 1 44 <i>A</i> 1 117	(Curran et al., 1992)
	U					41, 11/	

Table IV. Field studies of pesticide degradation under different tillage practices.

Pesticide	Tillage ¹	Depth		Soil properties		DT_{50}^{2}	Reference
			Clay	Organic carbon	pН	_	
		cm		$g kg^{-1}$		d	
Imazaquin	NT CT	0-20	-	16.9, 19.8	5.9, 6.4	27, 33 53, 22	(Mills and Witt, 1991)
Imazethapyr	RT CT	0-10	230	23.2	6.0	82, 53 122, 56	(Curran et al., 1992)
Imazethapyr	NT CT	0-20	-	16.9, 19.8	5.9, 6.4	36, 40 40, 12	(Mills and Witt, 1991)
Isoproturon	NT RT CT	0-30 0-30 0-30	170 170 170	6.1 5.8 4.5	- -	12 8 15	(Otto et al., 1997)
Metolachlor	NT RT CT	0-30 0-30 0-30	170 170 170	6.1 5.8 4.5	- -	9 26 29	(Otto et al., 1997)
Metolachlor	NT CT	0-150	55-166 59-179	0.1-9.7 1.2-4.4	5.3-6.1 5.5-6.7	45 34	(Heatwole et al., 1997)
Metolachlor	NT RT top RT furrow CT	0-10		- - -	- - -	40, 65, 37, 30 23, 87, 41, 35 28, 42, 40, 28 32, 68, 43, 40	(Gaynor et al., 1998)
Metolachlor	NT NT+Rg CT CT+Rg	0-10 0-10 0-10 0-10	390	14.5	5.4	46, 42, 72 42, 40, 69 42, 45, 79 44, 44, 97	(Gaynor et al., 2000)
Metribuzin	NT NT+Rg CT CT+Rg	0-10 0-10 0-10 0-10	390	14.5	5.4	24, 27, 34 24, 26, 36 23, 29, 32 24, 29, 37	(Gaynor et al., 2000)
Metribuzin	NT CT NT CT	0-5 0-5 0-5 0-5	- - -	13.3 18.0	5.7 5.1	12, 11 5, 17 15, 15 13, 11	(Sorenson et al., 1991)
Terbuthylazine	NT RT CT	0-30 0-30 0-30	170 170 170	6.1 5.8 4.5		25 32 22	(Otto et al., 1997)

¹ Codes of tillage practices are reported in Table I.

² DT₅₀: pesticide half-life.

³ Rg: ray-grass residues; V: vetch residues.

⁴ Commas are used to separate different years of study.

⁵ Numbers in parenthesis indicate the age in years of the tillage system.

generally increased and may lead to a decrease in the availability of the molecules for biological degradation (Zablotowicz et al., 2000). In some cases, microbiological activity in soil can be affected by a lower temperature (Sorenson et al., 1991) or higher soil acidity (Brown et al., 1994) under conservation tillage. Both of these consequences were found to increase sorption of pesticides and thus to reduce their bioavailability. For some authors, repeated fertiliser inputs (Gaynor et al., 1998; Ghadiri et al., 1984), the use of legumes as cover crops (Brown et al., 1994) and the absence of lime could lead to soil acidification and thus mask the effects of tillage practices, particularly for molecules of the s-triazine family whose sorption is highly sensitive to acidity. Other pesticides such as sulfonylureas have their chemical stability reduced by acidification, leading to a faster degradation of these compounds under conservation techniques (Chapman and Cole, 1982). Global soil microbial activity can be correlated with mineralisation of the molecules (Lavorenti et al., 2003), but not systematically (Reddy and Locke, 1998). The increase in soil microbial activity under conservation techniques did not always mean that specific microbial populations involved in the degradation of a molecule were more abundant (Gaston and Locke, 2000). In some cases, crop residues on the soil surface under conservation tillage seemed to disrupt microorganisms' activity

Pesticide	Tillage ¹	Depth	S	oil properti	es	Water content	Temperature	Incubation duration	DT_{50}^{2}	CO_2^2	NER ²	Reference
			Clay	Organic carbon	рН	-						
		cm	٤	g kg ⁻¹		g g ⁻¹	°C	d	d	% applied dose	% applied dose	
	NT	0-10	-	10.2	5.6*				108	5	9	
Acifluorfen	NT	20-30	-	4.4	5.7*	0.35	25	49	165	4	5	(Gaston and
Reindorreit	CT	0-10	-	8.7	5.8*	0.00	25	12	74	6	12	Locke, 2000)
	CT	20-30	-	4.9	5.8*				169	5	3	
Alachlor	NT $(7)^3$	0-5	-	22.0	5.3*	0.35	25	54	6.5	13	54	(Locke et al.,
- Huemor	CT	0-5	-	11.6	5.1*				6.5	7	43	1996)
Alachlor	NT	0-30	-	-	-	-	-	-	≈ 3	-	-	(Weed et al.,
	CT	0-30	-	-	-	-	-	-	≈ 3	-	-	1998)
	NT	0-10	-	10.2	5.6*				50	2	15	
Bentazon	NT	20-30	-	4.4	5.7*	0.30	25	22	87	2	8	(Gaston et al.,
	CT	0-10	-	8.7	5.8*				39	3	20	1996)
	СТ	20-30	-	4.9	5.8*				77	2	9	
Bentazon	NT	0-10	-	-	-	-	-	-	17-234	0.1	15-17	
(soil	NT	10-20	-	-	-	-	-	-	69	0.1	8-12	(Gaston and
columns)	СТ	0-10	-	-	-	-	-	-	12-14			Locke, 1996)
	СТ	10-20	-	-	-	-	-	-	23-35			
	NT (9)		-	15.3	5.4*			48	7	12	63	
	СТ		-	13.1	5.7*				9	17	65	
Bentazon	NT (16)		-	48.0	6.3*			48	9	18	60	
	CT		-	18.9	4.5*				15	14	53	
	NT (18)	0-7.5	-	30.6	6.8*	0.33	25	48	9	14	61	(Wagner et al.,
	CT		-	17.8	6.4*				8	14	62	1996)
	NT (4)		-	20.4	6.3*			48	11	15	59	
	CT		-	17.7	5.0*				11	14	59	
	NT (3)		-	10.2	5.6*			22	50	2	15	
	CT		-	8.7	5.8*				39	3	20	
	NT		280	30.6	6.6*				-	10	22	
	CT		250	17.8	6.4*				-	12	18	
Chlorimuron	NT	0-7.5	330	46.0	6.2*	0.31	-	63	-	11	24	(Reddy et al.,
			260	19.2	4.5* 5.4*				-	14	15	19956)
	N I CT		-	15.5	5.4				-	16	24	
		0.10	-	13.1	5.7	(0	25	110	-	16	24	(I
Diclosulam	NI	0-10	355	19.9	5.0*	60	25	119	6/	14	29	(Lavorenti et al.,
		0-10	341	12.8	4.4	%WHC ³			8/	11	24	2003)
	NI (II)	0-2	-	30.1	-				20	-	24	
	IN I NT	2-3 5 10	-	11.8	-				19	-	23	
	NT	J-10 10.25	-	2.9	-				40 50	-	0	(Zablotowicz
Fluometuron		0.2	-	5.5 20.2	-	0.33	28	25	0	-	9 43	(Zabiolowicz
	CT	0-2	-	20.5	-				9	-	45	et al., 2000)
	CT	2-3 5 10	-	6.0	-				25	-	22	
	CT	J-10 10.25	-	0.9	-				23	-	23	
	NT + A du ⁶	0.2	-	4.4	5.2				94	-	7	
	NT + AdV	0-3	-	7.2	5.2				0	-	-	
	$NT + D^6$	0-3	-	9.1	5.5				44	-	-	(Caston at al
Fluometuron	NI + D'	0-3	-	9.2	0.1 5.7	33 kPa ⁷	25	60	9	-	-	(Gaston et al., 2001)
	CT+Auv CT+V	0-3	-	5.4 4.6	5.7				44 74	-	-	2001)
	CT+P CT+P	0-3	-	4.0 / 0	5.5 5.8				74 23	-	-	
		0.7.5	-	4.9	5.0				2.5	-	-	
	NT + AdV	0-7.5	-	15.9	3.2 5.5				9	-	-	
		0-7.5	-	13.0	3.3 6 1				9	-	-	(Caston et al
Fluometuron	INI+B	0-7.5	-	11.2	0.1 5 7	PS	25	109	9 10	-	-	(Gaston et al., 2002)
	CT+AdV	0-7.5	-	/.4	3.1 5 5				19	-	-	2003)
		0-7.5	-	1.4	5.5 5.0				12	-	-	
	C1+B	0-7.5	-	/.ð	J.8				9	-	-	

 Table V. Laboratory studies of pesticide degradation under different tillage practices.

Pesticide	Tillage ¹	Depth	Soil properties		Water content	Temperature	Incubation duration	DT_{50}^{2}	CO_2^2	NER ²	Reference	
			Clay	Organic carbon	pН							
		cm	g	kg ⁻¹		$g g^{-1}$	°C	d	d	% applied dose	% applied dose	
	NT (11)	0-4	150	20.0	5.1				57	-	-	
	NT	4-8	150	10.0	5.5				79	-	-	
	NT	8-15	160	9.0	6.5		30	84	49	-	-	
	CT	0-4	130	13.0	5.5		50	04	49	-	-	
	CT	4-8	140	13.0	5.7	0.25			55	-	-	
	CT	8-15	150	11.0	6.3				52	-	-	
	NT+V (11)	0-4	150	25.0	4.7	0.25			78	-	-	(Brown et al.,
Fluometuron	NT+V	4-8	150	11.0	4.9				90	-	-	1994)
	NT+V	8-15	150	9.0	5.7				83	-	-	,
	CT+V	0-4	140	15.0	5.0				71	-	-	
	CT+V	4-8	150	15.0	5.0				68	-	-	
	CT+V	8-15	150	11.0	5.6				52	-	-	
T	SS (3)	-	660	16.0	6.4	0.25.0.30	30.25	6	12.16	-	-	(Seifert et al.,
imazaquin	CT	-	000	10.9	0.4	0.25-0.50	50-25	0	12-10	-	-	2001a)
0.16	NT	075	230	21.4	21.4 -	0.20	-	77	-	2	27	(Reddy and
Suifentrazone	CT	0-7.5	210	16.0	-	0.50	-	11	-	2	23	Locke, 1998)

¹ Codes of tillage practices are reported in Table I.

² DT₅₀: pesticide half-life; CO₂: mineralisation; NER: non-extractable residues.

³ Numbers in parenthesis indicate the age in years of the tillage system.

⁴ Hyphens are used to indicate the range of variation of values.

⁵ Water content expressed in % of the water-holding capacity (WHC).

⁶ Adv: weed residues; B: wheat residues; V: vetch residues.

⁷ Water content expressed with the corresponding matric potential (kPa).

* pH CaCl₂.

(Locke and Harper, 1991b; Sorenson et al., 1991), and, compared with conventional tillage, lag phases in the activation of mineralisation may occur (Seifert et al., 2001b). In addition, a greater availability of carbon under conservation tillage compared with conventional tillage can defer the use of pesticides as a source of carbon and thus their degradation in soil (Locke and Harper, 1991b).

Mineralisation is considered as the last step of pesticide degradation, leading to its complete removal from the soil. The mineralisation results are highly contrasted according to pesticides, location sites and incubation conditions (Tab. V) and do not allow any conclusion on an increase or a limitation of mineralisation in conservation tillage. In the same way, non-extractable residue fractions vary widely depending on pesticides, techniques and soils. However, the formation of non-extractable residues is often correlated with the degradation half-life of the molecules (Gaston and Locke, 2000; Lavorenti et al., 2003; Zablotowicz et al., 2000). Locke and Harper (1991b) have also shown that the difference in non-extractable residues of metribuzin between conventional tillage and conservation tillage was mainly due to the coarse fraction of organic matter.

Moreover, although little information is generally mentioned in the literature, the age of the tillage system seems to be a major source of differences in pesticide degradation (Wagner et al., 1996). Sadeghi and Isensee (1996) have evaluated the effect of reversing the tillage of 7-year-old no-tillage and conventional tillage field plots on atrazine degradation. Their results suggested that, after reversing a well-established tillage, time for the new tillage to fully develop its particular effects on atrazine degradation may be shorter for a new conventional tillage than for a new no-tillage. Another major source of degradation variability in field studies is the interannual variability of climatic conditions. It can hide or completely cancel the effects of tillage and it can lead to opposite conclusions depending on the year (Baughman et al., 1996; Mills and Witt, 1991; Sorenson et al., 1991).

In most cases, pesticide degradation forms one or several degradation products. In the same manner as for mother compounds, their degradation can be modified by tillage practices. For example, under a no-tillage system, an accumulation of polar metabolites formed by the degradation of metribuzin was observed, while they were degraded under conventional tillage (Locke and Harper, 1991a, b). On the contrary, the formation of alachlor metabolites was faster but their degradation was slower under conventional tillage than under conservation tillage (Locke et al., 1996). Similarly, the accumulation of a fluometuron metabolite, trifluoromethylphenylurea (TFMPU), occurred mainly under conventional tillage (Zablotowicz et al., 2000), and another of its metabolites, dimethylfluometuron,

seemed to be formed mainly when the air flow conditions, due notably to tillage operations, were favourable (Locke et al., 2005).

Pesticide formulation also appeared to have important consequences on their persistence. For example, whatever the tillage system, starch encapsulation of atrazine and alachlor tended to increase their field persistence (Gish et al., 1994).

Spatial variability of local conditions seems to be an important source of pesticide degradation variation. In their study, Gaynor et al. (1987) observed that the ridge tops retained more herbicide than the furrows, resulting in differences in pesticide persistence. Recently, Alletto et al. (2008) have examined the degradation of the diketonitrile metabolite of isoxaflutole under two tillage systems. Under conventional tillage, the herbicide persistence was found to be highly variable vertically and laterally according to soil sample location within the tilled horizon. The main source of variation in this case was the tillage operation with the mouldboard plough. Under mulch tillage with disk harrowing, MT_{Dk} (Tab. I), the effect of tillage was minimised but variations in diketonitrile persistence were associated with the vertical distribution of organic carbon. Both of these studies highlight the importance of the soil sampling strategy in order to provide accurate assessment of environmental impacts of agricultural practices. Last, the mulch can reduce water content and temperature variations (Bragagnolo and Mielniczuk, 1990; Unger, 1987), leading to a lower variability of degradation under conservation tillage compared with conventional tillage (Mills and Witt, 1991).

3.3. Summary and recommendations

Results about tillage effects on pesticide degradation are highly contrasted. Studies of the effects of management practices on soil properties and microorganism activity have to deal with complex interactions between soil physics, physicochemistry and microbiology which control the microbial activities involved in pesticide breakdown. Particularly the strong coupling between retention and degradation processes highly depends on soil physical and hydrodynamic properties, ensuring the access of microbes to water, substrates and the movement of solutes such as pesticides to sorption and degradation sites. Degradation studies should therefore pay great attention to the soil conditions (dynamics of temperature, water content, pH, N content, etc.), which have been shown to be largely modified by the presence of the mulch at the soil surface. In the same way, studies focusing on the understanding and prediction of pesticide degradation in soil should use dynamic experimental systems where soil solution is allowed to move and be renewed (e.g. soil columns) rather than static (batch) experimental systems. Improving knowledge and quantitative prediction of pesticide degradation in soils also implies an accurate estimation of the spatial distribution and temporal dynamics of active degradative populations. Research efforts should thus be oriented at (a) improving field study for a better understanding of the in situ dynamics of these microorganisms and their location ('hot-spots'), and (b) establishing a comprehensive link between this dynamics and agricultural

operations. Particular attention should be paid to the soil sampling strategy in this regard.

4. TRANSFER OF PESTICIDES

Depending on their mobility and their persistence, pesticides can migrate within and outside the soil and contaminate other compartments of the environment, such as water and air. The three main transfer processes are volatilisation, leaching and runoff. The relative importance of each of these processes depends on the application conditions, the pesticide properties, the climatic conditions and the soil properties partly governed by agricultural practices.

4.1. Volatilisation

Volatilisation is an important pathway for the loss of pesticide that is controlled by the pesticide properties (such as saturated vapour pressure, Henry constant, K_{OC}), the soil properties (temperature, water content, organic carbon content), the farming operations (mode of application, soil roughness, presence of a mulch) and the climatic conditions (wind, solar radiation, temperature) (Bedos et al., 2002). Although volatilisation is often mentioned in the literature to explain differences in the pesticide persistence due to tillage operations (Banks and Robinson, 1982; Curran et al., 1992; Gaynor et al., 2000), this phenomenon remains poorly studied. Whang et al. (1993) highlighted a higher volatilisation in conservation tillage than in conventional tillage that was attributed to the presence of a mulch. In their study, four days after treatment, transfer by volatilisation accounted for 48 and 18% of applied fonofos, 23 and 7% of applied chlorpyrifos and 0.9 and 0.7% of applied atrazine under no-tillage and conventional tillage, respectively. Wienhold and Gish (1994) also observed a larger volatilisation of alachlor and atrazine under conservation tillage, but only until the first rainfall occurred (5 days after treatment). Following this rainfall, volatilisation in conservation tillage was strongly slowed down, which could be explained by the migration of the pesticides from the mulch to the soil. After 35 days, the cumulative loss accounted for 9 and 14% of applied alachlor and 4 and 9% of applied atrazine under conservation tillage and conventional tillage, respectively. In addition, this study showed a significant effect of pesticide formulation on volatilisation. Starch-encapsulated alachlor was less volatilised under conservation tillage than the commercial formulation, and a similar effect was measured for atrazine under both conservation tillage and conventional tillage systems. Volatilisation directly depends on environmental conditions and is favoured first by high temperatures (Glotfelty, 1987; Weber et al., 2002) and also by wet soils that keep the molecules available in water solution. Weber et al. (2006) observed during a year of monitoring that volatilisation reached 22 and 32% of applied metolachlor in conservation tillage and conventional tillage, respectively. Due to the mulch, soil surface water content was higher under conservation tillage, which slowed down the soil warming and thus reduced the amount of herbicide lost.

4.2. Leaching

Pesticide properties play a decisive role in the modifications of the leaching risk by tillage. First, retention properties determine the mobility of the molecules (Singh N. et al., 2002) and directly influence their transfer to groundwater. Masse et al. (1998) reported that leaching of atrazine and deethylatrazine was more significant under conservation tillage, whereas metolachlor leaching, that has a stronger sorption capacity, was not affected by tillage. Water solubility of pesticides also influences their leaching. As an example, alachlor and cyanazine leaching was related to the interaction between tillage and water solubility (Sadeghi and Isensee, 1997). Under no-tillage, the migration depth of alachlor in soil was lower than that of cyanazine, whereas the opposite was observed under conventional tillage. The highest solubility of alachlor compared with that of cyanazine allowed a migration through the soil matrix, thus increasing the possibilities of adsorption, while cyanazine rather circulated via macropores under conservation tillage. Water solubility of the molecules may be modified by soil pH, which could increase leaching risk (Li et al., 2003). For sulfentrazone, solubility in water was, for example, multiplied by a factor of 16 when soil pH rose from 6 to 7.5 (Reddy and Locke, 1998). Other studies have shown a positive correlation between the half-life of several molecules (alachlor, atrazine, cvanazine, metolachlor, metribuzin and simazine) and their concentrations measured in drains without any tillage effect on these transfers (Logan et al., 1994; Ritter et al., 1996).

Formulation, by modifying solubility in water, persistence and retention in soils of pesticides, also influences their transport. Micro-encapsulation of alachlor increased its solubility in water and its transport through the soil matrix (Sadeghi et al., 1998). Starch-encapsulation of atrazine limited its losses by leaching in both no-tillage and conventional tillage (Gish et al., 1994, 1995). Hall et al. (1998) also observed a good efficiency of starch-encapsulation of atrazine, allowing a reduction of the leaching losses by a factor of 2 to 4 compared with the commercial formulation, depending on the year and on the application rate. Likewise, coating of metolachlor with a polyurea polymer helped reduce losses by a factor of 2 compared with the commercial formulation (Hall et al., 1998). According to these results, an efficient control of leaching seems achievable via adequate pesticide formulations.

Pesticide leaching depends on soil physical properties, such as the hydraulic conductivity, which is directly influenced by the soil structure created by tillage. Many studies on pesticide leaching have been conducted under field conditions, on plots equipped with ceramic cups, lysimeters or drains (Tab. VI), and under laboratory conditions, on undisturbed soil columns with simulated rainfalls (Tab. VII). Although most of these works showed greater losses under conservation tillage (Isensee and Sadeghi, 1997; Isensee et al., 1990; Masse et al., 1998; Singh et al., 2002; Weber et al., 2006), results are contrasted, some of them indicating no effect of tillage (Clay et al., 1998; Fomsgaard et al., 2003; Gaynor et al., 2000; Granovsky et al., 1993; Weed et al., 1995), or even greater losses under conventional tillage (Düring and Hummel, 1993; Gish et al., 1995; Levanon et al., 1993).

On one hand, mulch on the soil surface absorbs rainfall energy, thus avoiding the formation of soil crust (Baumhardt and Lascano, 1996; Blevins and Frye, 1993) and pore sealing (Ela et al., 1992). On the other hand, the continued deposition of crop residues on the soil surface appears to contribute to macropore development by stimulating earthworm activity (Bouché, 1972; Edwards et al., 1988; Rovira et al., 1987; Satchell, 1983). Under conservation tillage, and more particularly under no-tillage, the macropore network formed by earthworm burrows, root channels and cracks is not disrupted by tillage and thus may allow downward flows of water and solutes at a higher rate than if movements occurred only through the soil matrix. Because of this bypass of the soil matrix, this type of water and solute movement is called 'preferential flow through macroporosity'. It occurs mainly during saturated conditions. In the soil matrix, solutes move by convection-dispersion and, due to significant contact between the liquid and solid phases, opportunities for pesticide retention are greater than in macropores (Shipitalo and Edwards, 1996). Ogden et al. (1999) showed that soil tillage destroyed the connectivity of the macropores and thus increased fluxes through the soil matrix.

Proportionally to their quantity, crop residues can limit losses due to leaching. By reducing the infiltration rate at the soil surface, i.e. at the crop residues/soil interface, the mulch promotes pesticide fluxes within the soil matrix, thus avoiding preferential flows through macropores (Sigua et al., 1993). On the other hand, at the residues/soil interface or when residues and soil are mixed, the presence of these two compartments that have different water-holding capacities and hydraulic conductivities can also generate preferential flow (Kasteel et al., 2007; Ma and Selim, 2005).

There is no direct relationship between the number of macropores and the intensity of preferential flow, since some macropores do not participate in the conduction of water (Shipitalo et al., 1990). Preferential flow activation seems to depend, in particular, on the initial soil water content, but in contrasted ways according to studies (Granovsky et al., 1993; Shipitalo and Edwards, 1996). Although preferential flows through macropores occur in both conservation and conventional tillage (Andreini and Steenhuis, 1990; Essington et al., 1995; Gish et al., 1991; Granovsky et al., 1993), this dependency on initial water content was found to be greater under conservation tillage (Flury et al., 1995; Sigua et al., 1995). At low initial water content, the hydrophobicity of organic materials at the soil surface under conservation tillage could limit the entry of water into the soil matrix, creating locally and temporarily saturation conditions, thus favouring macropore fluxes (Edwards et al., 1989, 1992a; Phillips et al., 1989; Shipitalo et al., 1990). Despite earthworm burrows being found to have high pesticide sorption capacities due to high amounts of organic compounds in burrow linings (Edwards et al., 1992b; Stehouwer et al., 1993), water and pesticide transfer rates through macropores are generally higher than for the soil matrix (Shipitalo and Edwards, 1996). Other studies showed a higher leaching of atrazine (Kitchen et al., 1998;

L. Alletto et al.

Table VI.	Field studies	of pesticide	leaching (4	% of applied	dose) under	different tillage practices.
-----------	---------------	--------------	-------------	--------------	-------------	------------------------------

				Water		Soil			Reference
Pesticide	Dose	Tillage ¹	Depth	sampling system	3	properties		Leaching	
				sjotem	Clay	Organic	pН		
	1					carbon		Of any line data as	
	kg a.i. na	NIT	cm		g	kg ·	5770	% applied dose	
		IN I DT		-	-	17.5-21.4	5.7-7.0	0.0002-0.102	
Alachlor	2.2	KI MT	0-120	-	-	196 21 2	5760.1		(Weed et al., 1995)
		CT		-	-	10.7.20.4	5.7-0.8		
		NT		-	-	19.7-20.4	0.5-0.0	0.00 0.08 0.15 0.05 ³	
Atrozina		CT			-	-	-	0.04, 0.02, 0.12, 0.02	
Desethylatrazine	$2.2, 1.5, 1.8, 1.9^3$	NT	0-100	D^4	-	-	-	0.04, 0.02, 0.12, 0.02	(Masse et al., 1996)
Desettiylatiazine		CT			_	-	_	$0.07 \ 0.03 \ 0.15 \ 0.04$	
		NT		_	-	17 3-21 4	57-70	0.07, 0.05, 0.15, 0.04	
		RT		-	-	17.6-20.8	59-67		
Atrazine	2.8	MTch	0-120	-	-	18 6-21 3	57-68	0.02-0.35	(Weed et al., 1995)
		CT		-	-	19.7-20.4	6.3-6.6		
		MT _{Ch}	-	-	120-300	15.4-24.2	5.4-4.9	0.07-0.11, 0.08-0.22	
Atrazine	1.3	CT	-	-	120-300	15.7-23.5	5.3-5.8	0.03-0.07, 0.11-0.12	(Fortin et al., 2002)
		NT	0.120	T 4	224 421	0.0.10.0		0.15-0.86, 0.21-9.60	
Atrazine	1.7	CT	0-120	Lª	324-421	2.0-12.0	5.5-6.7	< 0.01 - 0.19, 0.75 - 0.85	(Hall et al., 1989)
Atrazine,					-			20.51	(Watts and Hall
Cyanazine	1.7, 2.2, 1.7	MT mulch	0-120	L	-	-	-	3.0-5.1	(watts and 11an, 1006)
Simazine		CT				-	-	0.69-0.93	1770)
Cyanazine	2.2	NT	0-120	L	324-421	2.0-12.0	5.5-6.7	0.03-0.23, <0.10-4.73	(Hall et al. 1989)
Cyunuzine		CT						< 0.01 - 0.15, 0.32 - 0.56	(11411 et 41., 1909)
Dicamba	0.56	NT	-	-	-	2.0-12.0	5.5-6.7	1.99, 0.39, 5.56, 1.05, 2.45	(Hall and Mumma,
		CT	-	-	-			< 0.01, 0.20, 0.20, 0.0, 0.58	1994)
Fluometuron	1.6	NT	0-90	L	130	1.3-17.7	5.61	29.9-37.4, 37.9-50.8,0-10.3, 0.8-54.7	(Essington et al.,
		CT				1.1-10.2	5.85	11.9-69.3, 30.0-79.5, 6.1-28.7, 3.4-73.3	1995)
Glyphosate		NT (20) ³			134-227	1.1-19.2	5.9-7.6	0.022	
AMPA	0.8	NT	0-110	L				0.018	(Fomsgaard et al.,
Glyphosate		CT			83-268	0.5-11.3	6.4-8.7	0.0305	2003)
AMPA		CT						0.0205	0.6
Metolachlor	2.6, 2.6, 2.6, 2.4	NT	0-100	D	-	-	-	0.00, 0.02, 0.02, 0.04	(Masse et al., 1996)
					-	-	-	0.00, 0.00, 0.02, 0.01	
Metolachlor	2.2	CT	0-120	L	-	-	-	2.46	(watts and Hall,
		MT _{ct}		_	120-300	15 4-24 2	54-49	0.02-0.04_0.06-0.12	1770)
Metolachlor	2.5	CT		_	120-300	15.7-23.5	5.3-5.8	0.01-0.03, 0.07-0.10	(Fortin et al., 2002)
	4.40	NT (10)	0.07		(0.000	20.64	4461	1.4. 6.7	
Metolachlor	4.48	CT	0-97	L	60-290	3.0-6.4	4.4-6.1	0.7, 4.4	(Weber et al., 2006)
Matalaahlar	2.2	NT	0.120	T	324 421	20120	5567	0.01-0.47, <0.10-4.19	(Holl at al. 1080)
Wetoraciiioi	2.2	CT	0-120	L	324-421	2.0-12.0	5.5-0.7	$< 0.01 - 0.10, \ 0.25 - 0.61$	(Hall et al., 1989)
		NT		-	-	17.3-21.4	5.7-7.0		
Metribuzine	0.45	RT	0-120	-	-	17.6-20.8	5.9-6.7	0 14-0 87	(Weed et al. 1005)
WIEUTOUZIIIE	0.10	MT_{Ch}	0 120	-	-	18.6-21.3	5.7-6.8	0.11 0.07	(weed et al., 1993)
		СТ		-	-	19.7-20.4	6.3-6.6		
Simazine	1.7	NT CT	0-120	L	324-421	2.0-12.0	5.5-6.7	0.06-1.76, 0.18-8.36 0.01-0.18, 1.50-1.63	(Hall et al., 1989)

¹ Codes of tillage practices are reported in Table I.
² Hyphens are used to indicate the range of variation of values.
³ Commas are used to separate different years of study.
⁴ D: water sampling in drains; L: water sampling in lysimeters.
⁵ Numbers in parenthesis indicate the age in years of the tillage system.

Pesticide	Dose	Tillage ¹	Depth		Soil properties		Rainfall intensity	Leaching	Reference
		.1		Clay	Organic carbon	pН	1 –1	<i>(</i> / 1' 1 1	
	kg a.i. ha	NT (1()?	cm		g kg		mm h '	% applied dose	(W/11
Alachlor	2.2	NI (16) ² CT	0-30	-	-	-	50	1.6 0.4	(weed et al., 1998)
Alashlar	2.2	NT (8)	0.10	-	-	-	50	29.7	(Clay et al.,
Alachior	5.5	CT	0-10	-	-	-	30	14.4	1991)
		NT			$10.2 - 15.3^3$	6.4-5.9	-	14.7	
A +	20	CT	0.19	170	11.8-12.6	6.4-6.7	-	19.3	(Levanon et al.,
Atrazine	2.0	NT	0-18	170	10.2-15.3	6.4-5.9	-	12.9	1993)
		CT			11.8-12.6	6.4-6.7	-	20.4	
		NT		-		-	9	58.4	
		CT		-		-	9	42.7	
		NT		-	NT. 75 044	-	9	47.2	
A 4	1.2	CT	0.10	-	IN I: 7.3 - 24.4	-	9	33.2	(Sigua et al.,
Atrazine	1.3	NT	0-10	-	CT. 7.0	-	9	31.6	1995)
		CT		-	C1: 7.0	-	9	29.4	
		NT		-		-	9	26.7	
		CT		-		-	9	20.1	
		NT		-		-	9	50.7	
		NT		-		-	9	35.7	
		NT		-		-	9	34.9	
		NT	0.40	_	NT: 24.4 – 7.5	-	9	35.9	(Sigua et al.,
Atrazine	1.3	СТ	0-10	-		-	9	40.3	1995)
		СТ		-	CT: 7.0	-	9	40.1	,
		СТ		_		-	9	38.2	
		СТ		_		-	9	37.0	
		NT			10.2-15.3	6.4-5.9	-	4.5	
		СТ			11.8-12.6	6.4-6.7	-	18.7	(Levanon et al.
Carbofuran	1.9	NT	0-18	170	10.2-15.3	6.4-5.9	-	4.2	(1993)
		CT			11.8-12.6	64-67	-	77	1)))))
		NT			10.2-15.3	64-59	-	0.25	
		CT			11.8-12.6	64-67	_	0.35	(Levanon et al
Diazinon	2.5	NT	0-18	170	10.2-15.3	64-59	_	0.0	(Levanon et al., 1993)
		CT			11.8-12.6	64-67	_	0.0	1775)
		NT (20)		112-210	9.0-18.0	0.4 0.7	0 16 0 38 0 46	0.03 3.0 10.1	(Düring and
Metamitron	4	CT	0-30/40	151-217	9.0-1.0		0.10, 0.50, 0.40	0.3 5 0 12 2	Hummel 1999)
		NT		151 217	10.2-15.3	64-59		8.2	Hummer, 1999)
		CT			11.8-12.6	64-67		10.2	(Levanon et al
Metolachlor	2.25	NT	0-18	170	10.2.15.3	6450	-	6.5	(Levanon et al., 1003)
		CT			11.8.12.6	6467	-	0.3	1995)
		NT (20)		112 210	0.0.18.0	0.4-0.7	-	1857	(Düring and
Metolachlor	10	NT (20)	0-30/40	151 217	9.0-18.0	-	0.25, 0.42	1.0, J.7 2.6, 12.4	(During and
		NT		131-217	9.0-1.0	-		3.0, 12.4	(Singh M at al
Metolachlor	20		0-15	280	19.0	7.3	saturation	30 27	(3 mgn N, et al., 2002)
		NT (20)		112 210	0.0.18.0			25.07	(Dijining and
Terbuthylazine	5	NT (20) CT	0-30/40	112-210	9.0-18.0 9.0-1.0	-	0.25, 0.42	5.2, 12.2	(During and Hummel, 1999)
		NT			2.0 1.0			11	(Singh N et al
Terbuthylazine	10	СТ	0-15	280	19.0	7.3	saturation	6	2002)

Table VII. Laboratory studies on pesticide leaching (% of applied dose) under different tillage practices.

¹ Codes of tillage practices are reported in Table I.
 ² Numbers in parenthesis indicate the age in years the tillage system.
 ³ Hyphens are used to indicate the range of variation of values.

Seyfried and Rao, 1987; Sigua et al., 1995) and fluometuron (Essington et al., 1995) under wet soil initial conditions. A better understanding of the effect of the initial water content on the dynamics of macropore flow is thus needed to improve the control of pesticide leaching and to complement modelling efforts. An illustration is given by Sigua et al. (1995), who observed under laboratory conditions that a decrease in soil matric potential from -1 kPa (near saturation) to -33 kPa led to a decrease of 15% in atrazine leaching.

In addition, the temporal dynamics of preferential flow through macropores needs to be taken into account. During wetting, the number of macropores hydraulically active tends to increase, thus increasing the possibilities of preferential flow (Jaynes et al., 2001; Kung et al., 2000; Malone et al., 2001). However, under unsaturated conditions, macropores are inactive but still the observed movements of solutes through the soil matrix may be faster than those estimated by the convection-dispersion equation.

To describe solute movements in the soil matrix better, the Mobile-Immobile water Model (MIM) has been developed (Coats and Smith, 1964). This model considers that the waterfilled pore space is partitioned into two domains: a mobile domain where water can move and solute transport is due to convection and dispersion, and an immobile domain where water is stagnant and solutes move only by diffusion (Coats and Smith, 1964; van Genuchten and Wierenga, 1976). In this model, it is possible to distinguish different degradation and retention kinetics between the two domains (van Genuchten and Wagenet, 1989), allowing a better description of preferential fluxes under unsaturated conditions (Gaston and Locke, 1996; Gaston and Locke, 2000; Pot et al., 2005). This type of preferential flow has been identified for pesticides under both conventional and conservation tillage (Gaston and Locke, 1996; Gaston and Locke, 2000; Singh N. et al., 2002), but laboratory studies on undisturbed soil columns showed that immobile water fractions were higher under conservation tillage than under conventional tillage, with 56 vs. 49% (Singh and Kanwar, 1991) and 56 vs. 35% (Singh N. et al., 2002), respectively. As for macropore fluxes, preferential flows within the soil matrix also seem to increase under wet soil conditions (Shipitalo and Edwards, 1996). Despite its strong impact on solute transfers, little data about the impact of tillage on MIMtype preferential flow has been published and thus further experimental studies are needed.

Intensity and timing of rainfall after treatment are major factors affecting pesticide leaching (Granovsky et al., 1993; Heatwole et al., 1997; Isensee and Sadeghi, 1994; Masse et al., 1996). Their effects may be greater than those generated by tillage (Gaynor et al., 1995; Granovsky et al., 1993; Otto et al., 1997). During small, low-intensity rainfalls, pesticides intercepted by the mulch can be washed off and then can penetrate slowly into the soil matrix where sorption processes can reduce their leaching (Shipitalo et al., 1990). On the contrary, during high-intensity rainfalls, wash-off is significant and the hydrophobicity of organic residues can slow down infiltration and activate preferential flow in macropores (Isensee and Sadeghi, 1994). By removing the mulch under conservation tillage, Sadeghi and Isensee (1997) observed that leaching was reduced and finally, lower losses under conservation tillage than under conventional tillage were measured. Furthermore, high-intensity rainfall generally leads to greater water and solute fluxes than low-intensity rainfall (Quisenberry et al., 1994; Trojan and Linden, 1992), with most pesticide leaching occurring during the first 2 or 3 rains (Fortin et al., 2002). For atrazine, Sigua et al. (1993) showed on undisturbed soil columns collected under no-tillage plots that the intensity of rainfall determines the percentage of losses: 33% of the applied dose for an intensity of 3 mm h^{-1} and 52% at 9 mm h^{-1} . In addition, if a small and low-intensity rainfall (1.5 mm for Sadeghi and Isensee, 1997, or 5 mm for Shipitalo et al., 1990) preceded a leaching event, then pesticide transport could be reduced by 50% compared with treatments that did not receive this preliminary rain. Moreover, these studies also showed that allowing time for sorption and diffusion in the soil matrix to occur can reduce pesticide movements in soils.

Some questions remain about how rapidly soil hydraulic properties change and preferential flow paths develop when tillage is modified. Very few data have been published on this aspect. In their study, Isensee and Sadeghi (1996) reversed the tillage of 7-year-old no-tillage and conventional tillage plots on which preferential flows had been previously observed (Isensee and Sadeghi, 1994; Isensee et al., 1990; Sadeghi and Isensee, 1992). They found that, immediately after ploughing, preferential flows under the new conventional tillage had disappeared and that sufficient macropore pathways were developed under the new no-tillage in only one year without tillage, leading to significant leaching differences. However, it seemed that two years were required before preferential transport systems became well established. Moreover, in the new conventional tillage, the macropore network inherited from 7 years of no-tillage and situated below the newly ploughed horizon appeared to be functional and preferential flow could still occur (Isensee and Sadeghi, 1997). Soil structural stability, which depends on soil texture, also greatly influences the dynamics of preferential flow (Singh et al., 2002). For example, in four soils of various textures under no-tillage, clay content enhanced macropore stability, leading to a leaching of atrazine 40% higher than for a sandy soil (Sadeghi et al., 2000).

Most of the studies about the effects of tillage on leaching compare contrasted techniques such as ploughing and notillage. Other conservation techniques such as mulch tillage or ridge tillage are still poorly studied in terms of pesticide leaching. In some studies of mulch tillage systems, greater leaching of atrazine, cyanazine, simazine and metolachlor was measured compared with conventional tillage (Watts and Hall, 1996). To the contrary, other studies did not find any differences in atrazine and metolachlor leaching between mulch tillage and conventional tillage, concluding that the differentiation between these systems was too weak (Fortin et al., 2002). Between mulch tillage and ridge tillage systems, differences in the migration of atrazine were also poorly marked (Kitchen et al., 1998). However, localisation of the herbicide in seed rows under ridge tillage allowed a decrease by a factor of 3 in the applied doses and could significantly reduce losses by leaching (Lamb et al., 1998; Lowery et al., 1998). In their study comparing three different tillage systems, Mazzoncini et al.

(1998) observed a deeper migration of diclofop-methyl after a heavy rainfall under no-tillage, while the pesticide was found mainly in the seedbed layer (0–5-cm depth) under mulch and conventional tillage.

4.3. Runoff

Agricultural runoff is the primary mechanism contributing to pesticide contamination of surface waters. The main objective of conservation tillage is to reduce runoff and soil erosion (Gebhardt et al., 1985).

Runoff is also affected by pesticide properties. Sorption properties directly act on the dominant mode of transport. For pesticides that have a high sorption capacity on organomineral particles, such as glyphosate, trifluralin, paraquat or organochlorine pesticides, surface transport is associated with erosion and soil particle transport (Wauchope, 1978). Potter et al. (2004) indicated that 55% of pendimethalin (K_{OC} = 5000 L kg⁻¹) losses were bound to sediment. However, for most pesticides, transport to surface water is realised in solution in water runoff (Wauchope, 1978). Losses in solution represented from 88 to 97% of total losses of alachlor and cyanazine (Hansen et al., 2001), 99.8% of atrazine losses (Basta et al., 1997) and, according to tillage practices, from 89 to 98% of fluometuron losses (Potter et al., 2004). For herbicides transported only in solution, such as chlorimuron and nicosulfuron, the interest of conservation techniques, especially no-tillage used to control erosion, is very limited (Afyuni et al., 1997). For some authors, however, pesticide transport in solution may be overestimated due to desorption from suspended sediments during runoff or sample storage and thus the importance of the solid phase as a source of pesticides in runoff would be underestimated (Hansen et al., 2001).

Logan et al. (1994) compared the effects of conventional tillage and no-tillage on the transport of atrazine, alachlor, metolachlor and metribuzin by runoff. No difference between tillage systems was found but runoff losses were positively correlated with the half-lives of the molecules. In other studies, water solubility directly determines the loss by runoff and explains differences in behaviour between alachlor and cyanazine (Hansen et al., 2001) or between atrazine and metribuzin (Gaynor et al., 2001). Isoxaflutole has a low solubility in water but its hydrolysis forms a diketonitrile metabolite (active ingredient) which is 50 times more soluble and thus transported by runoff (Rector et al., 2003). However, mainly because of analytical costs, metabolite monitoring in runoff is not systematic even in research studies. For fluometuron, Potter et al. (2004) indicated that 50% of the herbicide losses occurred as desmethylfluometuron metabolite. Results of runoff losses for some molecules, such as atrazine or fluometuron, are often contrasted under conservation tillage despite similar water solubility and retention by organic compounds. Potter et al. (2003) suggested a relationship to predict the mode of transport of pesticides (in solution or adsorbed) based on their K_{OC} and their solubility.

Pesticide solubility may vary according to formulation. For example, micro-encapsulation of alachlor increases its solubility, allowing a faster migration within the soil matrix and a decrease in its runoff compared with the commercial formulation (Isensee and Sadeghi, 1993). Metolachlor solubility is higher than that of atrazine but it may be reduced by its formulation, which may then limit its runoff compared with atrazine (Sadeghi and Isensee, 2001).

Concerning soil properties, increasing soil organic matter content in topsoil improves soil aggregate stability and cohesion, leading to a significant decrease in soil loss (Rhoton et al., 2002). It appears, however, that tillage effects on pesticide transfer by runoff or erosion are, as for leaching, mixed (Tab. VIII). Many works highlighted a decrease in pesticide losses by runoff under conservation tillage due to a decrease in water runoff volume as the degree of tillage is reduced (Seta et al., 1993; Tebrügge and During, 1999; Watts and Hall, 1996; Webster and Shaw, 1996). However, in some studies, pesticide concentrations in runoff from conservation tillage plots, especially no-tillage plots, were higher than in runoff from conventional tillage plots. As a consequence, conservation tillage plots may generate greater pesticide losses despite a lower water runoff volume (Heatwole et al., 1997; Kenimer et al., 1987; Shipitalo and Owens, 2006; Webster and Shaw, 1996). In other studies, water runoff volume was higher under conservation tillage (Gaynor et al., 1995; Myers et al., 1995). A review of tillage effects by Fawcett et al. (1994) concluded that, under natural rainfall, conservation tillage was efficient in controlling runoff, erosion and pesticide losses. Under simulated rainfall, the amount and intensities of water applied shortly after treatment resulted in very mixed results in conservation tillage. For Fawcett et al. (1994), intensive rainfall was more an indicator of the limits of the efficiency of conservation tillage to control runoff rather than an accurate representation of their effects under real field conditions. Modelling was used to identify and classify the most suitable practices to control atrazine runoff in a watershed (Harman et al., 2004). In this study, conservation tillage was found to be marginally effective. The most effective practices were: sediment retention ponds, grass filter strips, atrazine application at planting time in bands and wetlands construction. According to these contradictory results, it seems necessary to clarify the main conditions and mechanisms involved in pesticide transfer by runoff that may lead to an efficient or inefficient control of losses by conservation tillage.

Rainfall is the most important parameter controlling water runoff, soil erosion and pesticide loss. Depending on its arrival time after treatment, its intensity and the interval between two rainfall events, rainfall can lead to very contradictory results for the same study site, the same molecule or even the same practice (Baker and Johnson, 1979). Pesticide transport in soils directly depends on the interval between treatment and the first rainfall (Baker and Johnson, 1979; Potter et al., 2004; Shipitalo and Owens, 2003). Several studies indicated that conservation tillage was inefficient in controlling runoff when heavy rainfall occurs quickly after pesticide application (Rector et al., 2003; Shipitalo and Owens, 2003; Shipitalo and Owens, 2006). Moreover, most pesticide loss by

Pesticide	Dose	Tillage ¹		Soil properties		Slope	Rainfalls		F	Runoff	Reference
	kg a.i. ha ⁻¹	l	Clay	Organic carbon (OC) g kg ⁻¹	Hd	%	шш	Type	water % applied dose	pesticide % applied dose	
Alachlor	2.24	NT RT CT		5.8-17.4 ²		12-18 ²	657, 487, 463, 425 ³	Z 4	$13, 4, 12, 8^3 \\17, 4, 8, 8 \\17, 8, 18, 9$	$\begin{array}{c} 7.1,0.45,1.8,0.1^3\\ 8.0,0.8,0.45,<\!0.1\\ 14.3,0.7,4.1,0.2 \end{array}$	(Baker and Johnson, 1979)
Alachlor	2.80	NT MT _{Ch} CT	220	17.4		9	136	z	34, 41 28, 54 35, 54 44, 48	0.32, 4.8 0.25, 2.1 0.36, 6.0 0.29, 3.0	(Sauer and Daniel, 1987)
Alachlor	1.79	NT CT		7.4-23.7 6.6-6.9		3-5			4-6, 3-4 ¹ 5-6, 3-5	0.22-0.24, 0.15-0.24 0.48-0.57, 0.15-0.28	(Isensee and Sadeghi, 1993)
Alachlor	3.36	NT C ⁵ NT U/D ⁵ CT C CT U/D		12.8		7-11	67 66 56	S^4	19 34 27 58	0.06 0.08 0.1 0.2	(Felsot et al., 1990)
Alachlor	3.36	NT C ⁵ NT U/D ⁵ ST C ST U/D RT C RT U/D MT _{Ch} C MT _{Ch} U/D CT C		12.8		8 8 9 9 9 8 8 10 10 10 10 10 10 10 10 10 10 10 10 10	63 64 65 65 64 64 64 64 64 64 64 64 64 64 64 64 64	N	6 23 10 22 33 33 43	0.09 2.1 0.06 0.75 0.75 1.3 1.2 2.3 6.5	(Felsot et al., 1990)
Alachlor	2.24-3.36	NT 1 NT 2 MT $_{Ch}$ 1 MT $_{Ch}$ 2				11 10 7	629-1012 686-1025 609-981 663-1007	z	1-10 2-14 < 0.01 - 1 1-15	Tr-0.23 0.01-0.19 Tr-<0.01 Tr-0.22	(Shipitalo et al., 1997)
Alachlor	2.2 2.2 2.2 0.7 Loc ⁶ 0.7 Loc 0.7 Loc	MT _{ch} RT CT MT _{ch} CT CT		14.0 17.4 13.4 14.0 17.4 13.4	7.1	8-10	240, 480	Z		0.4, 0.2 0.1, 0.1 1.0, 0.6 0.1, 0.06 0.1, 0.06 0.2, 0.06	(Hansen et al., 2001)
Alachlor ⁷	1.68 3.36	$\underset{DT}{\operatorname{MT}}_{Ch}$		1 1 1		10-11 7-13 6-9	976 953 960	Z	8 7 11	0.28 0.11 0.12	(Shipitalo and Owens, 2006)
Atrazine	1.7	NT MT mulch CT				3-5			1 1 1	0.06, - -, 0.28 0.18, 0.77	(Watts and Hall, 1996)
Atrazine	1.68	NT CT	67 59	9.7 4.4	5.64 5.91	2				0.7 1.5	(Heatwole et al., 1997)

Table VIII. Field studies of water and pesticide runoff under simulated (S) or natural (N) rainfall under different tillage practices.

	¢	1000	r c	5	: : :			8	c F
Pesticide	Dose	- 111age	Class Oscill properties	Stope	Kaintalls		and our	Kunon	Kererence
	kg a.i. ha ⁻¹		ciay Organic carbon (OC) g kg ⁻¹	% III	mm	Type	% applied dose	% applied dose	
	2.24	NT 1	י נ נ	- 13		5 '	4-7	0.027-0.82	
	2.24	NT 2		- 7	ı		6-15	0.049-4.95	
	2.24	MT_{Ci} . 1		-	ı	,	0.1-2	Tr-0.011	
Atrazine	2.24	$MT_{Ch} 2$		- 10		,	9-18	0.0045-2.62	(Shipitalo and
(+ DEA + DIA)	1.12	MT_{Dk} 3		- 6		,	7-17	0.034-0.64	Owens, 2003)
	1.12	MT_{Dk} 4		- 7		,	4-8	0.0010-0.59	
	1.12	MT_{Dk} 5	1	- 9		ı	10-22	0.0043-6.37	
		NT					13, 4	7.1, 0.75	(Deltan and
Atrazine	2.24	RT	5.8-17.4	12-18	657, 487	z	17, 4	19.2, 1.8	(Daker and Tetress 1070)
		CT	ı	I			18, 8	20.1, 0.09	JOUNSON, 1979)
		NT					34, 41	0.32, 7.3	
· · · · · · · · · · · · · · · · · · ·	7 80	MT_{Ch}	V L1 UCC	y	136	,	28, 54	0.21, 3.0	(Sauer and Daniel,
Autazine	000.7	RT	1.11 077	0	0.01	,	35, 54	0.34, 8.7	1987)
		CT				,	44, 48	0.25, 4.3	
Attoring	2 7 4	NT	1	ı 1	941 1158 824 926 1029 860		5, 6, 7, 18, -, 1	2.5, 1.9, 1.0, 5.1, -, 2.0	(Ghidey et al.,
Auazino		MT			/11) 1100, 011, /TO, 101/, 000		1, 4, 7, 18, 18, 1	1.1, 0.3, 0.6, 1.5, 4.0, 2.1	2005)
Atrazina	1 34	NT	- 7.4-23.7	3-5	1	,	4-6, 3-4	1.39 - 1.51, 0.56 - 0.90	(Isensee and
Auazine		CT	- 6.6-6.9	<i>נ</i>		,	5-6, 3-5	0.81-0.97, 0.21-0.47	Sadeghi, 1993)
Attoring	1 7	NT	- 12.0	67 3-5	т		-, 0.1, <0.01, 0.02	-, 0.10, <0.01, <0.01	(Holl at al 1001)
Auazine		CT	0.71	3-4	ı	,	0.3, 0.3, 0.03, 0.03, 0.04	0.36, 0.33, 0.02, < 0.01	(Hall Cl al., 1991)
		NT	1				9	0.7	
Atrazine	2.24	MT	1	- 9	132	S	22	1.3	(Seta et al., 1993)
		CT		ı			34	1.8	
Atrazine	2.24	TN	232 21.5	_ 10.3	100	S	0.2	0.28	(Kenimer et al.,
		CT		10.6			3	2.85	1987)
		NT (1) ⁸						7.9	
Atrazine	0.56	MT_{Ch}	- 9.5	4.9 4	80	S		10.5	(Basta et al., 1997)
		MT_{Dk}						7.5	
		NT 1	1	- 11	629-1012		1-10	< 0.01 - 0.08	
Attoring	2 74	NT 2		- 10	686-1025	z	2-14	< 0.01 - 1.10	(Shipitalo et al.,
Auazine		MT_{Ch} 1	1	- 13	609-981	-	< 0.01 - 1	< 0.01	1997)
		$MT_{Ch} 2$	1	- 7	663-1007		1-15	< 0.01 - 0.91	
		NT $(2) - M.^{9}$					31-50, 35-38	2.2-9.4, 2.1-4.7	(Muarc at al
Atrazine	1.6	NT – Sil. ⁸	210 9.1	5.9 2-6	31.7-50.8	S	31-56, 35-48	1.7-8.1, 1.9-2.5	1005)
		CT					16-24, 34-47	0.5-1.6, 1.7-3.0	(0001
	0.9	NT (4)					-	4.99 / 10.51	
Atrazine	$0.9 \mathrm{Inc.}^{10}$	CT	- 13.4	6.0 1-2		N/N+S		1.26/2.73	(Rector et al.,
	0.9	CT						2.48 / 7.72	2003)
	1.8	NT (4)						2.64 / 7.52	
	0.55	$MT_{Ch} + Cov.C^{1}$	1				8, 7	0.1, 1.8	
Atrazine		MT_{Ch}	- 14.5	- <1	995, 688	z	14, 9	0.4, 1.5	(Gaynor et al.,
		CT + Cov.C					8, 6	0.4, 0.7	2001)
		۲. بیب		11 01	100	14	12,1	0.0, 1.0	
A turning	۲ <i>с</i> с	IN		- 10-11 7 13	9/6	Z	7 X	1.29	(Shipitalo and
Atrazine	7.24	MI_{Ch}		- /-15	5CY			0.00	Owens, 2006)
		MT_{Dk}		- 6-9	960		11	0.92	

Table VIII. Continued.

	I	-		-			3			8	- -
Pesticide	Dose	Tillage	ξ	Soil properties	E.	Slope	Rainfalls			Kunott	Reference
	kg a.i. ha ⁻¹		Ulay	Organic carbon (OC) g kg ⁻¹	цц	%	mm	Type	water % applied dose	pesuctue % applied dose	
		NTC					20 20		19	0.2	ť
Carbofuran	1.12			12.8		7-11	00	S	54 77	0.0	(Felsot et al.,
		CT U/D					56		<i>21</i> 58	1.21	(0661
Chlorimuron	0.014	LN	80	5.3	5.9	0-1	,		1	2.0	(Afyuni et al.,
		NT							34 41	0.04 0.10	(1661
-3:	1 2/	MT_{Ch}		V 11		4	136		28, 54	0.05, 0.29	(Sauer and Daniel,
Chlorpyntos	+C.1	RT	077	1/.4		0	0.01	ı	35, 54	0.06, 0.28	1987)
		CT							44, 48	0.08, 0.20	
	, c c	. La		1 21 0 3		01 01	307 637	N	12, 8 0 0	4.5, 0.2	(Baker and
Cyanazine	+7.7	CT		4./ I-0.C		01-71	403, 423	2	o, o 18, 9	2.0, 0.1 8.0, 0.3	Johnson, 1979)
Cvanazina	1 34	NT		7.4-23.7		2-5			4-6, 3-4	1.30-1.91, 0.50-0.68	(Isensee and
Cyanazure		CT	'	6.6-6.9					5-6, 3-5	0.52-0.84, 0.15-0.27	Sadeghi, 1993)
Cyanazine	2.2	NT CT	ı	12.0	6.7	3-5 3-4	I	ı	-, 0.1, < 0.01, 0.02 0.3, 0.3, 0.03, 0.04	-, 0.06, <0.01, <0.01 0.13, 0.30, 0.01, <0.01	(Hall et al., 1991)
	2.1	MT_{Ch}		14.0					I.	2.0, 0.9	
	2.1	RT	ı	17.4					ı	0.38, 0.8	
Cvanazine	2.1	CT	ı	13.4	7.1	8-10	240.480		·	4.4, 1.0	(Hansen et al.,
c) unumit	$0.7 \mathrm{Loc}^{6}$	MT_{Ch}	ı	14.0					ı	0.8, 0.2	2001)
	0.7 Loc	RT	·	17.4					ı	0.6, 0.2	
	0.4 Loc	CT	ı.	13.4						0.8, 0.2	
		Lu							ı	0.02, -	(Watts and Hall.
Cyanazıne	2.2	MT mulch		1		C-5	ı		1	-, 0.17 0 11 0 30	1996)
		NT				25			01 /001 000	012 -001 -001	(Holl and
Dicamba	0.56	CT	ı	12.0	6.7	2-0 3-4	-		-, 0.1, < 0.01, 0.02 0.3, 0.3, 0.03, 0.04	-, 0.12, <0.01, <0.01 0.46, 0.81, <0.01, <0.01	(rian anu Mumma, 1994)
Dimethipin	0.35	ST (1)	32	5.1	6.5	3-4		s	23	5.0	(Potter et al.,
7		CT							23	1.6	2003)
ī	I.	LN						;	- 9	8.0, 5.3	(Baughman et al.,
Fluometuron	1./	CT CT		18.0	0.4	0	3/0,480	2+2	0 0	8.2, 3.2 9.9, 4.4	2001)
		ST					63.8	$S-I_{Var}^{12}$	20	1.4	
	00	ST	37	5 1	59	2.4	64.3	$S-I_{cons}^{12}$	23	1.9	(Potter et al.,
Fluomeuron	0.7	CT	4 C	7.1	<u></u>	+	62.7	S- I _{Var}	45	0.8	2006)
		CT					60.0	S- I _{cons}	50	0.6	
, ,		LN							4, 12, 8	0.18, 1.3, 0.05	(Baker and
Fonotos	1.12	RT CT		5.8-I7.4		12-18 4	87, 463, 425		4, 8, 8 8 18 0	0.09, 0.25, nd 0.36 1 0.007	Johnson, 1979)
		SS (3)								1 22 3 39	(Seifert et al
Imazaquin	0.14	CT	660	16.9	6.4	NA	1	·		1.53, 3.63	2001b)
	0.11	NT (4)								2.82 / 9.24	
Isoxaflutole/ Aiteronitrile	0.11 Inc. ¹⁰	CT	,	13.4	6.0	1-2	I	N/N+S	1	0.93 / 3.35 1 63 / 6 75	(Rector et al.,
AIRMANI	0.05	NT (4)							1 1	2.35 / 6.84	10007

Table VIII. Continued.

de	Dose	Tillage ¹		Soil properties	Slope	Rainfalls			Runoff	Reference
1		1	Clay C	rganic carbon (OC)	, Hq			water	pesticide	I
k	g a.i. ha ⁻¹			g kg ⁻¹	%	mm	Type	% applied dose	% applied dose	
		I TN			- 11	629-1012		1-10	< 0.01 - 0.01	
-	12	NT 2			- 10	686-1025	z	2-14	< 0.01 - 1.04	(Shipitalo et al.,
-	11.	MT_{Ch} 1			- 13	609-981	-	< 0.01 - 1	< 0.01	1997)
		MT_{Ch} 2			- 7	663-1007		1-15	< 0.01 - 0.46	
L		NT			- 10-11	976		8	0.70	(PL
-	.12	MT_{Ch}			- 7-13	953	z	7	0.29	Outputato and
		MT_{Dk}			- 6-9	096		11	0.21	Uwens, 2000)
1		NT - S/W ¹³						8, 7, 9	0.4, 0.2, 0.6	
¢		$NT - Soy^{13}$	-		, ,	101 101	0.14	-, 14, 18	-, 1.4, 0.3	(Webster and
•)	4.	CT - S/W		0.0	0.4 3	213, 191, 188	2+2	9.9.15	0.8.1.1.0.6	Shaw. 1996)
		CT - Sov						9, 5, 10	0.4, 0.5, 0.8	
1		NT (2) – $M.^9$						31-50, 35-38	1.2-9.0, 0.9-4.3	,
	6	$NT - Sil^9$	210 9	-	5.9 2-6	31.7-50.8	s.	31-56, 35-48	1.4-8.0.0.8-1.8	(Myers et al.,
	2	CL		1	1		2	16-24, 34-47	0.4-1.5, 1.1-2.7	1995)
1		NT	-		2-5			0.1. <0.01. 0.02	0.06. <0.01. <0.01	
1	7	CT	-	0.7	0.7 3-4	-	ı	0.3, 0.3, 0.03, 0.04	0.16, 0.25, 0.01, <0.01	(Hall et al., 1991)
1		NT							0.03, -	(Watts and Hall
C 1	.2	MT mulch			- 3-5	585, 748, 486, 562	z		-, 0.22	Walls and Hall,
		CT							0.29, 0.55	1996)
<u>`</u>	10	NT	6 29	L	5.64 ,				0.5	(Heatwole et al.,
4	+7.	CT	59 4	4.	5.91 ²	T	ı		1.2	1997)
	10 1 01	NT				011 1150 831 036 1030 860	N	5, 6, 7, 18, -, 1	5.5, 1.0, 0.9, 2.1, -, 0.3	(Ghidey et al.,
-	.12-1.0/	MT				941, 11.30, 024, 920, 1029, 000	2	1, 4, 7, 18, 18, 1	2.0, 0.4, 1.2, 1.6, 5.4, 0.4	2005)
1		$MT_{Ch} + Cov.C^{1}$	-					8,7	0.06, 0.8	
<	0.1	MT_{Ch}	-	2 1	~	002 600	N	14, 9	0.2, 1.0	(Gaynor et al.,
ر ا	10.	CT + Cov.C		0.4	- /	777, 000		8, 6	0.3, 0.5	2001)
		CT						12, 7	0.4, 1.4	
C	Ľ	NT (4)							4.22 / 8.28	
0	7 Inc. ¹⁰	CT		7 4	60 1-2	ľ	STN/N		1.01 / 1.97	(Rector et al.,
0	7.7	CT		+	7-1 0.0	·			2.22 / 5.98	2003)
-	4.	NT (4)							2.44 / 6.01	
		NT 1			- 11	629-1012		1-10	< 0.01 - 0.79	
0	38	NT 2			- 10	686-1025	Z	2-14	0.08-0.22	(Shipitalo et al.,
		MT_{Ch} 1			- 13	609-981		< 0.01 - 1	< 0.01 - 0.02	1997)
		$MT_{Ch} 2$			- 7	663-1007		1-15	< 0.01 - 0.01	
1		NT	· ·		- 10-11	926		8	0.83	F 1-+;;-13/
0	.38	MT_{Ch}			- 7-13	953	z	7	0.04	(Surprise and
		MT_{Dk}			- 6-9	096		11	0.55	Uwens, 2000)
1		$MT_{Ch} + Cov.C^{1}$	T					8,7	0.04, 1.1	
0	u C	MT_{Ch}	-	L.			F	14, 9	0.3, 1.4	(Gavnor et al
ر	C7.	CT + Cov.C	-	C. 1	- -	1 000	7	8.6	0.3, 0.6	2001)
		CT						12, 7	0.3, 1.7	
		NT - S/W ¹³						8, 7, 9	0.7, 0.5, 0.7	
0	Ţ	NT - Soy ¹³	-	90	61 2	213 101 188	S IN	-, 14, 18	-, 1.9, 0.6	(Webster and
ر	ţ.	CT - S/W		0.0	C + 0	213, 131, 100	0+1	9, 9, 15	1.7, 1.6, 0.9	Shaw, 1996)
		CT - Soy						9, 5, 10	0.9, 0.8, 1.4	
	016	ΤN	1 1001	0.0	10	Puc	Z	0.2	0.01	(Malone et al.,
ر	018.	CT	1 120 1	5.0	- 10	774	2	15	0.64	1996)

Table VIII. Continued.

Tillage management effects on pesticide fate in soils. A review

Pesticide	Dose	Tillage ¹		Soil properties		Slope	Rainfalls			Runoff	Reference
	ke a.i. ha ⁻¹	Ĭ	Clay	y Organic carbon (OC) _{φ kφ⁻¹}	Ηd	%	a	Tvne	water % annlied dose	pesticide % annlied dose	I
Nicosulfuron	0.014	NT CT	80	5.3	5.9	0-1	-			2.3 1.1	(Afyuni et al., 1997)
Norflurazon	1.7	NT RT CT		18.6	6.4	3	376, 480	N+S	e 2 e	4.2, 4.2 4.2, 2.5 4.8, 3.6	(Baughman et al., 2001)
Pendimethalin	1.0	ST ST CT CT	32	5.1	6.5	3-4	63.8 64.3 62.7 60.0	$S - \frac{12}{Var}$ $S - \frac{12}{12}$ $S - \frac{12}{Var}$ $S - \frac{1}{Var}$ $S - \frac{1}{Var}$	20 23 50	0.4 0.4 5.0 4.1	(Potter et al., 2006)
Simazine	1.7	NT CT		12.0	6.7	3-5 _ 3-4 _		1	-, 0.1, <0.01, 0.02 0.3, 0.3, 0.03, 0.04	-, 0.18, < 0.01, < 0.01 0.62, 0.51, 0.03, < 0.01	(Hall et al., 1991)
Simazine	1.7	NT MT mulch CT		1		3-5	I.		1 1 1	0.02, - -, 0.13 0.11, 0.35	(Watts and Hall, 1996)
		NT C NT U/D ST C ST U/D				8 8 6 6	63 64 63		6 23 10	0.03 0.3 0.2	
Terbufos + Terbufos metabolites	1.12	RT C, RT U/D MT _{C/} C MT _{C/} U/D CT C CT U/D		12.8	1	0 1 1 0 1 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	5 5 5 5 5 5 5 5 5 5 5 5	ŝ	24 29 33 33 33 43	0.4 2.3 1.4 1.6 7.5	(Felsot et al., 1990)
Thidiazuron	0.05	ST (1) ⁸ CT	32	5.1	6.5	3-4		S	23 23	13.7 12.2	(Potter et al., 2003)
Tribufos	0.30	ST (1) CT	32	5.1	6.5	3-4		S	23 23	12.8 14.5	(Potter et al., 2003)
2,4-D	0.56	NT CT	232	21.5		10 11	100	S	0.2 3	0.02 0.27	(Kenimer et al., 1987)
¹ Codes of tilla ² Hyphens are 1 ³ Commas are 1 ⁴ N: natural rain 5 C: contoured 1 5 Loc: seed row 6 Loc: seed row 7 Average resul 8 Numbers in p. 9 M.: maize = r ¹⁰ Inc.: preplant ¹¹ Cov.C: cover	ge practices ar ised to indicat ised to separat islope: U/D: ur islope: U/D: ur islope: U/D: ur islope: U/D: ur islope: U/D: ur islope: ur islo	e reported in Tal e the range of va e different years ated rainfall. and-down slop of the pesticide tr of study. cate the age in y ted; Sil.: silage - ation of herbicid. ity; L _{cons} : constan	ble I. uriation e. ceatrnel ears of e = no cr e.	o f values. dy. nt. f the soil tillage. cop residues. fall intensity.							

L. Alletto et al.

runoff occurred during the first few runoff events after application (Seifert et al., 2001a; Shipitalo and Owens, 2006). In a 9-year-study comparing three types of conservation tillage (NT, MT_{Ch} and MT_{Dk}), Shipitalo and Owens (2006) indicated that 60 to 99% of herbicide (alachlor, atrazine, linuron, and metribuzin) losses were due to the five largest transport events. After a rainfall representing 3% of annual rainfall and occurring 2 days after treatment, atrazine losses reached 4.7% of the applied dose (Shipitalo and Owens, 2003). Triplett et al. (1978) proposed a linear relationship to predict atrazine concentrations in runoff ([atrazine]_{runoff}) based on the number of days after treatment (ln ([atrazine]_{runoff} = a + b. ln (number of days after treatment)). This relationship has been tested and used for several pesticides such as alachlor (Shipitalo et al., 1997), atrazine and its metabolites (deethylatrazine and deisopropylatrazine) (Gaynor et al., 1995; Shipitalo and Owens, 2003; Shipitalo et al., 1997), cyanazine (Franti et al., 1998), linuron (Shipitalo et al., 1997), dimethipin (Potter et al., 2003), metolachlor (Gaynor et al., 1995; Ghidey et al., 2005; Webster and Shaw, 1996), metribuzin (Shipitalo et al., 1997), thidiazuron and tribufos (Potter et al., 2003). However, Hansen et al. (2001) indicated that the relationship between ln [pesticide]_{runoff} and ln (number of days after treatment) was sensitive to annual variability in weather and may not be a good variable for simple estimates of herbicide concentration in runoff. They recommended the use of cumulative rainfall as a variable for estimating concentration in runoff: In $[\text{pesticide}]_{runoff} = a + b.ln$ (cumulative rainfall). The intensity of the first rainfall also influences the fate of the applied molecules. A small, low-intensity rainfall occurring a few days after treatment allows the incorporation of pesticide within the topsoil horizon, thus reducing losses during subsequent runoff events (Afyuni et al., 1997; Gaynor et al., 1995; Olson et al., 1998).

As for leaching, the initial soil water content is also a key factor controlling runoff. Runoff due to exceedance of infiltration capacity was found to occur earlier under no-tillage than under conventional tillage (Isensee and Sadeghi, 1993; Rector et al., 2003; Sadeghi and Isensee, 2001). In hydromorphic soils, saturation occurred more rapidly under no-tillage and runoff was more significant than under conventional tillage (Ghidey et al., 2005). Soil tillage, even superficial, increases soil infiltration capacity and drainage, which can limit runoff (Olson et al., 1998). The time interval between two rainfalls may significantly affect runoff volumes under no-tillage. Isensee and Sadeghi (1993) indicated that, for a time interval lower than 7 days between two rainfalls, the high soil water content remaining under no-tillage, favoured by the mulch, led to a significantly higher runoff than under conventional tillage.

The mulch increases soil surface roughness, thus reducing runoff (Isensee and Sadeghi, 1993; Selim et al., 2003). However, the mulch effectiveness in controlling runoff and erosion depends on the nature and quantity of plant residues. For example, Olson et al. (1998) observed that after a soybean crop, the abundance of residues under no-tillage was not sufficient to limit (water and atrazine) runoff and higher runoff volumes than under mulch tillage were measured. For quantities of residues increasing from 0 to 1.5 t ha⁻¹ either remaining on the soil surface in no-tillage or mixed in the ploughed horizon in conventional tillage, runoff was reduced by 96 and 40%, respectively (Kenimer et al., 1987).

Strip tillage is a conservation tillage well-developed in the US which is starting to be used for spring productions by farmers in Europe. On one hand, strip tillage was found to reduce runoff compared with conventional tillage but, on the other hand, to increase lateral subsurface flow (Bosch et al., 2005). Potter et al. (2003) indicated that, after only one year of strip tillage, no effect on runoff volumes was measured compared with conventional tillage but erosion rates were lower. On the same study site, a few years later, runoff volumes were decreased by a factor of 4 compared with conventional tillage (Potter et al., 2004). Concerning pesticide losses, fluometuron losses were 2 to 3 times higher and pendimethalin losses were 12 times lower under strip tillage than under conventional tillage (Potter et al., 2006). Under strip tillage, herbicide treatments are applied only on the seed row representing only 1/3 of the field surface. In inter-row positions, the soil surface is most often covered by a cover crop, which reduces the risk of runoff (Hansen et al., 2001). Ridge tillage can also be used to reduce runoff but, in some cases, pesticides could migrate from the ridge tops to the furrows and then be transferred by runoff (Gaynor et al., 1987; Olson et al., 1998). Very few studies reported results on the effects of sub-soiling on runoff. Seifert et al. (2001a) did not observe any effect of this tillage practice on runoff and pesticide loss compared with conventional tillage. Pesticide incorporation into soil through superficial tillage was found to significantly reduce losses by runoff (Franti et al., 1998; Olson et al., 1998; Rector et al., 2003) but this technique is not suitable for no-tillage systems and may lead to an increase in losses by leaching. Finally, whatever the tillage system, soil tillage along isotopographic lines rather than up-and-down slope was efficient in reducing transfers by runoff (Felsot et al., 1990).

4.4. Summary and recommendations

Concerning pesticide transfer, it clearly appears that initial soil conditions (water content and temperature) and climatic conditions (rainfall intensity, interval between treatment and the first rainfall) play a large role in the dynamics of water and solutes and may explain the contrasted effects of tillage on pesticide transfers. Improving the understanding of pesticide transport related to tillage operations implies clarifying the temporal and spatial dynamics of solute flows, especially preferential flows, which are in this case particularly relevant. Most of the studies mentioned in this review highlighted the transient and local nature of transport mechanisms, making them difficult to characterise. In the same order of ideas, the lack of efficient and robust ways to quantify soil structure and its dynamics is an impediment to the prediction of pesticide fate and transport, especially in topsoil layers.

The significant effect of rainfall characteristics on pesticide transfers suggests that tillage effects are highly dependent on the type of climate, and its eventual modifications to come. Regional studies should be of great help in developing generic models for more robust predictions. Tillage is certainly a relevant means of controlling agricultural pesticide impacts on the environment. Pesticide fate in soil implies numerous processes in complex interactions. If one is to improve tillage systems to mitigate these impacts, one needs first to understand how the various tillage practices modify the functional characteristics of soil (water retention, hydraulic conductivity, solute transport, etc.) through detailed, comprehensive studies. Such studies should consider the diversity of tillage systems, which are currently insufficiently explored and documented. Such research efforts would imply long-term studies (> 10 years) to allow significant and well-established differentiation between the cropping systems being compared and would thus involve stability in funding research programmes.

5. CONCLUSION

This review article outlines the four main following points.

- 1. Pesticide interception is enhanced under conservation tillage practices. The intensity of this process depends on (a) the amount and nature of crop residues which have sorption capacities 10 to 60 times higher than soil, and (b) the climatic conditions, such as the timing between pesticide treatment and the first rainfall and its intensity.
- Pesticide retention, which is generally positively correlated with organic carbon content, is increased in the topsoil layer under conservation tillage. Desorption of pesticides is also affected by tillage and greater proportions of pesticides remained sorbed on soil particles and on mulch under conservation tillage.
- 3. As a consequence of points 1 and 2, a lower fraction of pesticide remains available for biological degradation under conservation tillage and, in several cases, pesticide persistence in soils increases. In addition to this lower bioavailability due to retention processes, the presence of crop residues, a lower temperature and a higher acidity under conservation tillage can disrupt or slow down microorganisms' activity. Moreover, a greater availability of carbon under conservation tillage can defer the use of pesticides as a source of carbon.
- 4. Transport of pesticides is affected by tillage management and by its interactions with climatic conditions – more particularly by the intensity of rainfall, its arrival time after treatment, the interval between two rainfall events – and pesticide properties – e.g. water solubility, retention, halflife and formulation. In a general way, conservation tillage is more efficient in reducing runoff than leaching. Indeed, a higher soil surface roughness due to the presence of crop residues and a greater aggregate stability lead to a significant decrease in pesticide loss in water runoff or bound to eroded sediment, while reducing tillage intensity maintains a well-connected macropore network through which leaching of pesticides is enhanced.

REFERENCES

- Afyuni M.M., Wagger M.G., Leidy R.B. (1997) Runoff of two sulfonylurea herbicides in relation to tillage system and rainfall intensity, J. Environ. Qual. 26, 1318–1326.
- Agreste (2004) Enquête sur les pratiques culturales en 2001, Chiffres et Données Série Agriculture n° 159.
- Agreste (2008) Enquête sur les pratiques culturales en 2006, Chiffres et Données Série Agriculture n° 200.
- Alletto L., Benoit P., Bergheaud V., Coquet Y. (2008) Temperature and water pressure head effects on the degradation of the diketonitrile metabolite of isoxaflutole in a loamy soil under two tillage systems, Environ. Pollut. 156, 678–688.
- Andreini M.S., Steenhuis T.S. (1990) Preferential paths of flow under conventional and conservation tillage, Geoderma 46, 85–102.
- Arshad M.A., Franzluebbers A.J., Azooz R.H. (1999) Components of surface soil structure under conventional and no-tillage in northwestern Canada, Soil Till. Res. 53, 41–47.
- Aubertot J.N., Barbier J.M., Carpentier A., Gril J.J., Guichard L., Lucas P., Savary S., Savini I., Voltz M. (2005) Pesticides, agriculture et environnement. Réduire l'utilisation des pesticides et en limiter les impacts environnementaux, Rapport d'Expertise scientifique collective, INRA et Cemagref (France).
- Baker J.L., Johnson H.P. (1979) The effect of tillage systems on pesticides runoff from small watersheds, T. ASAE 22, 554–559.
- Balesdent J., Mariotti A., Boisgontier D. (1990) Effect of tillage on soil organic carbon mineralization estimated from ¹³C abundance in maize fields, J. Soil Sci. 41, 587–596.
- Banks P.A., Robinson E.L. (1982) The influence of straw mulch on the soil reception and persistence of metribuzin, Weed Sci. 30, 164– 168.
- Banks P.A., Robinson E.L. (1986) Soil reception and activity of acetochlor, alachlor, and metolachlor as affected by wheat (Triticum aestivum) straw and irrigation, Weed Sci. 34, 607–611.
- Barnes C.J., Goetz A.J., Lavy T.L. (1989) Effects of imazaquin residues on cotton (*Gossypium hirsutum*), Weed Sci. 37, 820–824.
- Barr D.P., Aust S.D. (1994) Mechanisms white rot fungi use to degrade pollutants, Environ. Sci. Technol. 28, 78A.
- Barriuso E., Calvet R. (1992) Soil type and herbicide adsorption, Int. J. Environ. An. Ch. 46, 117–128.
- Barriuso E., Feller C., Calvet R., Cerri C. (1992) Sorption of atrazine, terbutryn and 2,4-D herbicides in two Brazilian Oxisols, Geoderma 53, 155–167.
- Basta N.T., Huhnke J.H., Stiegler J.H. (1997) Atrazine runoff from conservation tillage systems: a simulated rainfall study, J. Soil Water Conserv. 52, 44–48.
- Baughman T.A., Shaw D.R., Rhodes G.N., Mueller T.C. (1996) Effects of tillage on chlorimuron persistence, Weed Sci. 44, 162–165.
- Baughman T.A., Shaw D.R., Webster E.P., Boyette M. (2001) Effect of cotton (Gossypium hirsutum) tillage systems on off-site movement of fluometuron, norflurazon, and sediment in runoff, Weed Technol. 15, 184–189.
- Baumhardt R.L., Lascano R.J. (1996) Rain infiltration as affected by wheat residue amount and distribution in ridged tillage, Soil Sci. Soc. Am. J. 60, 1908–1913.
- Bedos C., Cellier P., Calvet R., Barriuso E., Gabrielle B. (2002) Mass transfer of pesticides into the atmosphere by volatilization from soils and plants: overview, Agronomie 22, 21–33.
- Berglof T., Koskinen W.C., Brucher J., Kylin H. (2000) Linuron sorptiondesorption in field-moist soils, J. Agr. Food Chem. 48, 3718–3721.

- Bernoux M., Cerri C.C., Cerri C.E.P., Neto M.S., Metay A., Perrin A.S., Scopel E., Razafimbelo T., Blavet D., Piccolo M.D., Pavei M., Milne E. (2006) Cropping systems, carbon sequestration and erosion in Brazil, a review, Agron. Sustain. Dev. 26, 1–8.
- Biederbeck V.O., Campbell C.A., Hunter J.H. (1997) Tillage effects on soil microbial and biochemical characteristics in a fallow-wheat rotation in a dark brow soil, Can. J. Soil Sci. 77, 309–316.
- Blevins R.L., Frye W.W. (1993) Conservation tillage: an ecological approach to soil management, Adv. Agron. 51, 33–78.
- Bosch D.D., Potter T.L., Truman C.C., Bednarz C.W., Strickland T.C. (2005) Surface runoff and lateral subsurface flow as a response to conservation tillage and soil-water conditions, T. ASAE 48, 2137–2144.
- Bouché M.B. (1972) Lombriciens de France. Écologie et systématique, Annales de zoologie-écologie animale, numéro hors-série, Ed. INRA, 671 p.
- Boyd S.A., Xiangcan J., Lee J.F. (1990) Sorption of nonionic organic compounds by corn residues from a no-tillage field, J. Environ. Qual. 19, 734–738.
- Bragagnolo N., Mielniczuk J. (1990) Mulch from eight crop sequences and its effects on soil temperature, moisture, emergence and initial growth of corn, Rev. Bras. Cienc. Solo 14, 91–98.
- Brown B.A., Hayes R.M., Tyler D.D., Mueller T.C. (1994) Effect of tillage and cover crop on fluometuron adsorption and degradation under controlled conditions, Weed Sci. 42, 629–634.
- Brown B.A., Hayes R.M., Tyler D.D., Mueller T.C. (1996) Effect of longterm Vetch (*Vicia villosa*) cover crop and tillage system on fluometuron dissipation from surface soil, Weed Sci. 44, 171–175.
- Chapman R.A., Cole C.M. (1982) Observations on the influence of water and soil pH on the persistance of insecticides, J. Environ. Sci. Health 17, 487–504.
- Chin Y.P., Weber W.J., Eadie B.L. (1990) Estimating the effects of dispersed organic polymers on the sorption of contaminants by natural solids. 2. Sorption in the presence of humic and other natural macromolecules, Environ. Sci. Technol. 24, 837–842.
- Clay S.A., Koskinen W.C. (1990) Adsorption and desorption of atrazine, hydroxyatrazine, and s-glutathione atrazine on two soils, Weed Sci. 38, 262–266.
- Clay S.A., Koskinen W.C., Carlson P. (1991) Alachlor movement through intact soil columns taken from two tillage systems, Weed Technol. 5, 485–489.
- Clay S.A., Clay D.E., Koskinen W.C., Berg R.K. (1998) Application method: impacts on atrazine and alachlor movement, weed control, and corn yield in three tillage systems, Soil Till. Res. 48, 215–224.
- Coats K.H., Smith B.D. (1964) Dead-end pore volume and dispersion in porous media, SPE J. 4, 73–84.
- Comia R.A., Stenberg M., Nelson P., Rydberg T., Håkansson I. (1994) Soil and crop responses to different tillage systems, Soil Till. Res. 29, 335–355.
- CTIC (2006) National crop residues management survey, Conservation tillage information center, Ames, IA.
- Curran W.S., Liebl R.A., Simmons F.W. (1992) Effects of tillage and application method on clomazone, imazaquin, and imazethapyr persistence, Weed Sci. 40, 482–489.
- Dalal R.C., Henderson P.A., Glasby J.M. (1991) Organic matter and microbial biomass in a vertisol after 20 yr of zero-tillage, Soil Biol. Biochem. 23, 435–441.
- Dao T.H. (1991) Field decay of wheat straw and its effects on metribuzin and S-ethyl metribuzin sorption and elution from crop residues, J. Environ. Qual. 20, 203–208.

- Dao T.H., Lavy T.L. (1978) Atrazine adsorption on soil as influence by temperature, moisture content and electrolyte concentration, Weed Sci. 26, 303–309.
- Ding G., Novak J.M., Amarasiriwardena D., Hunt P.G., Xing B. (2002a) Soil organic matter characteristics as affected by tillage management, Soil Sci. Soc. Am. J. 66, 421–429.
- Ding G.W., Novak J.M., Herbert S., Xing B.S. (2002b) Long-term tillage effects on soil metolachlor sorption and desorption behavior, Chemosphere 48, 897–904.
- Ding G.W., Liu X.B., Herbert S., Novak J., Amarasiriwardena D., Xing B.H. (2006) Effect of cover crop management on soil organic matter, Geoderma 130, 229–239.
- Doran J.W. (1980) Soil microbial and biochemical changes associated with reduced tillage, Soil Sci. Soc. Am. J. 44, 765–771.
- Drijber R.A., Doran J.W., Parkhurst A.M., Lyon D.J. (2000) Changes in soil microbial community structure with tillage under long-term wheat-fallow management, Soil Biol. Biochem. 32, 1419–1430.
- Drury C.F., Tan C.S., Welacky T.W., Oloya T.O., Hamill A.S., Weaver S.E. (1999) Red clover and tillage influence on soil temperature, water content, and corn emergence, Agron. J. 91, 101–108.
- Düring R.A., Hummel H.E. (1993) Soil tillage as a parameter influencing the fate of three selected soil herbicides, Med. Fac. Landbouww. Univ. Gent 58/3a, 827–835.
- Düring R.A., Hummel H.E. (1999) Herbicide and metabolite movement in different soils as studied by computer assisted microlysimeters, Chemosphere 39, 641–654.
- Düring R.A., Hoss T., Gath S. (2002) Depth distribution and bioavailability of pollutants in long-term differently tilled soils, Soil Till. Res. 66, 183–195.
- Edwards W.M., Norton L.D., Redmond C.E. (1988) Characterizing macropores that affect infiltration in nontilled soil, Soil Sci. Soc. Am. J. 52, 483–487.
- Edwards W.M., Shipitalo M.J., Owens L.B., Norton L.D. (1989) Water and nitrate movement in eathworm burrows within long-term notill cornfields, J. Soil Water Conserv. 44, 240–243.
- Edwards W.M., Shipitalo M.J., Dick W.A., Owens L.B. (1992a) Rainfall intensity affects transport of water and chemicals through macropores in no-till soil, Soil Sci. Soc. Am. J. 56, 52–58.
- Edwards W.M., Shipitalo M.J., Traina S.J., Edwards C.A., Owens L.B. (1992b) Role of *Lumbricus terrestris* (L.) burrows on quality of infiltrating water, Soil Biol. Biochem. 24, 1555–1561.
- Ela S.D., Gupta S.C., Rawles W.J. (1992) Macropore and surface seal interactions affecting water infiltration into soil, Soil Sci. Soc. Am. J. 56, 714–721.
- Erbach D.C., Lovely W.G. (1975) Effect of plant residue on herbicide performance in no-tillage corn, Weed Sci. 23, 512–515.
- Essington M.E., Tyler D.D., Wilson G.V. (1995) Fluometuron behavior in long-term tillage plots, Soil Sci. 160, 405–414.
- Fawcett R.S., Christensen B.R., Tierney D.P. (1994) The impact of conservation tillage on pesticide runoff into surface water: a review and analysis, J. Soil Water Conserv. 49, 126–135.
- Felsot A.S., Mitchell J.K., Kenimer A.L. (1990) Assessment of management practices for reducing pesticide runoff from sloping cropland in Illinois, J. Environ. Qual. 19, 539–545.
- Ferri M.V.W., Vidal R.A., Gomes J., Dick D.P., de Souza R.F. (2002) Activity of the herbicide acetochlor in soil under no-till and conventional tillage system, Pesqui. Agropecu. Bras. 37, 1697–1703.
- Flury M., Leuenberger J., Studer B., Flühler H. (1995) Transport of anions and herbicides in a loamy and a sandy field soil, Water Resour. Res. 30, 1945–1954.

- Fomsgaard I.S., Spliid N.H., Felding G. (2003) Leaching of pesticides through normal-tillage and low-tillage soil - A lysimeter study. II, Glyphosate J. Environ. Sci. Health B. 38, 19–35.
- Fortin J., Gagnon-Bertrand E., Vezina L., Rompre M. (2002) Preferential bromide and pesticide movement to tile drains under different cropping practices, J. Environ. Qual. 31, 1940–1952.
- Franti T.G., Peter C.J., Tierney D.P., Fawcett R.S., Myers J.L. (1998) Reducing herbicide losses from tile-outlet terraces, J. Soil Water Conserv. 53, 25–31.
- Gan J., Becker R.L., Koskinen W.C., Buhler D.D. (1996) Degradation of atrazine in two soils as a function of concentration, J. Environ. Qual. 25, 1064–1072.
- Gao J.P., Maguhn J., Spitzauer P., Kettrup A. (1998) Sorption of pesticides in the sediment of the Teufelsweiher pond (Southern Germany). II: Competitive adsorption, desorption of aged residues and effect of dissolved organic carbon, Water Res. 32, 2089–2094.
- Gaston L.A., Locke M.A. (1996) Bentazon mobility through intact, unsaturated columns of conventional and no-till Dundee soil, J. Environ. Qual. 25, 1350–1356.
- Gaston L.A., Locke M.A. (2000) Acifluorfen sorption, degradation, and mobility in a Mississippi delta soil, Soil Sci. Soc. Am. J. 64, 112– 121.
- Gaston L.A., Locke M.A., Zablotowicz R.M. (1996) Sorption and degradation of bentazon in conventional- and no-till Dundee soil, J. Environ. Qual. 25, 120–126.
- Gaston L.A., Boquet D.J., Bosch M.A. (2001) Fluometuron wash-off from cover crop residues and fate in a loessial soil, Soil Sci. 166, 681–690.
- Gaston L.A., Boquet D.J., Bosch M.A. (2003) Fluometuron sorption and degradation in cores of silt loam soil from different tillage and cover crop systems, Soil Sci. Soc. Am. J. 67, 747–755.
- Gaynor J.D., Stone J.A., Vyn T.J. (1987) Tillage systems and atrazine and alachlor residues on a poorly drained soil, Can. J. Soil Sci. 67, 959–963.
- Gaynor J.D., MacTavish D.C., Findlay W.I. (1995) Atrazine and metolachlor loss in surface and subsurface runoff from three tillage treatments in corn, J. Environ. Qual. 24, 246–256.
- Gaynor J.D., MacTavish D.C., Labaj A.B. (1998) Atrazine and metolachlor residues in Brookston CL following conventional and conservation tillage culture, Chemosphere 36, 3199–3210.
- Gaynor J.D., Tan C.S., Ng H.Y.F., Drury C.F., Welacky T.W., vanWesenbeeck I.J. (2000) Tillage and controlled drainage-subirrigated management effects on soil persistence of atrazine, metolachlor, and metribuzin in corn, J. Environ. Qual. 29, 936–947.
- Gaynor J.D., Tan C.S., Drury C.F., Ng H.Y.F., Welacky T.W., van Wesenbeeck I.J. (2001) Tillage, intercrop, and controlled drainagesubirrigation influence atrazine, metribuzin, and metolachlor loss, J. Environ. Qual. 30, 561–572.
- Gebhardt M.R., Daniel T.C., Schweizer E.E., Allmaras R.R. (1985) Conservation tillage, Science 230, 625–630.
- Ghadiri H., Shea P.J., Wicks G.A., Haderlie L.C. (1984) Atrazine dissipation in conventional-till and no-till sorghum, J. Environ. Qual. 13, 549–552.
- Ghidey F., Blanchard P.E., Lerch R.N., Kitchen N.R., Alberts E.E., Sadler E.J. (2005) Measurement and simulation of herbicide transport from the corn phase of three cropping systems, J. Soil Water Conserv. 60, 260–273.
- Gish T.J., Helling C.S., Mojasevic M. (1991) Preferential movement of atrazine and cyanazine under field conditions, T. ASAE 34, 1699– 1705.

- Gish T.J., Shirmohammadi A., Wienhold B.J. (1994) Field-scale mobility and persistence of commercial and starch-encapsulated atrazine and alachlor, J. Environ. Qual. 23, 355–359.
- Gish T.J., Shirmohammadi A., Vyravipillai R., Wienhold B.J. (1995) Herbicide leaching under tilled and no-tillage fields, Soil Sci. Soc. Am. J. 59, 895–901.
- Glotfelty D.E. (1987) The effects of conservation tillage practices on pesticide volatilization and degradation, in: Logan T.J. et al. (Eds.), Effects of conservation tillage on groundwater quality: Nitrates and pesticides. Lewis, Chelsea, MI, pp. 56–62.
- Granovsky A.V., McCoy E.L., Dick W.A., Shipitalo M.J., Edwards W.M. (1993) Water and chemical transport through long-term no-till and plowed soils, Soil Sci. Soc. Am. J. 57, 1560–1567.
- Grant R.F., Izaurralde R.C., Chanasyk D.S. (1990) Soil temperature under conventional and minimum tillage: simulation and experimental verification, Can. J. Soil Sci. 70, 289–304.
- Grey T.L., Walker R.H., Wehtje G.R.H., Hancock H.G. (1997) Sulfentrazone adsorption and mobility as affected by soil and pH, Weed Sci. 45, 733–738.
- Gupta S.C., Schneider E.C., Swan J.B. (1988) Planting depth and tillage interactions on corn emergence, Soil Sci. Soc. Am. J. 52, 1122– 1127.
- Hall J.K., Mumma R.O. (1994) Dicamba mobility in conventionally tilled and non-tilled soil, Soil Till. Res. 30, 3–17.
- Hall J.K., Murray M.R., Hartwig N.L. (1989) Herbicide leaching and distribution in tilled and untilled soil, J. Environ. Qual. 18, 439–445.
- Hall J.K., Mumma R.O., Watts D.W. (1991) Leaching and runoff losses of herbicides in a tilled and untilled field, Agr. Ecosyst. Environ. 37, 303–314.
- Hall J.K., Jones G.A., Hickman M.V., Amistadi M.K., Bogus E.R., Mumma R.O., Hartwig N.L., Hoffman L.D. (1998) Formulation and adjuvant effects on leaching of atrazine and metolachlor, J. Environ. Qual. 27, 1334–1347.
- Hansen N.C., Moncrief J.F., Gupta S.C., Capel P.D., Olness A.E. (2001) Herbicide banding and tillage system interactions on runoff losses of alachlor and cyanazine, J. Environ. Qual. 30, 2120–2126.
- Hanson J. (2006) History of tillage. Improving Soil Enhancing Profit. Manitoba-North Dakota Zero Tillage Farmers Association Workshop, Bismarck, ND, 9–10 February 2006.
- Harman W.L., Wang E., Williams J.R. (2004) Reducing atrazine losses: Water quality implications of alternative runoff control practices, J. Environ. Qual. 33, 7–12.
- Heatwole C.D., Zacharias S., Mostaghimi S., Dillaha T.A. (1997) Movement of field-applied atrazine, metolachlor, and bromide in a sandy loam soil, T. ASAE 40, 1267–1276.
- Isensee A.R., Sadeghi A.M. (1993) Impact of tillage practice on runoff and pesticide transport, J. Soil Water Conserv. 48, 523–527.
- Isensee A.R., Sadeghi A.M. (1994) Effects of tillage and rainfall on atrazine residue levels in soil, Weed Sci. 42, 462–467.
- Isensee A.R., Sadeghi A.M. (1996) Effect of tillage reversal on herbicide leaching to groundwater, Soil Sci. 161, 382–389.
- Isensee A.R., Sadeghi A.M. (1997) Interactions of tillage and rainfall on atrazine leaching under field and laboratory conditions, Chemosphere 34, 2715–2723.
- Isensee A.R., Nash R.G., Helling C.S. (1990) Effect of conventional vs. no-tillage on pesticide leaching to shallow groundwater, J. Environ. Qual. 19, 434–440.

- Jaynes D.B., Ahmed S.I., Kung K.J.S., Kanwar R.S. (2001) Temporal dynamics of preferential flow to a subsurface drain, Soil Sci. Soc. Am. J. 65, 1368–1376.
- Johnson M.D., Wyse D.L., Lueschen W.E. (1989) The influence of herbicide formulation on weed control in four tillage systems, Weed Sci. 37, 239–249.
- Kabir Z. (2005) Tillage or no-tillage: Impact on mycorrhizae, Can. J. Plant Sci. 85, 23–29.
- Kasteel R., Garnier P., Vachier P., Coquet Y. (2007) Dye tracer infiltration in the plough layer after straw incorporation, Geoderma 137, 360– 369.
- Kenimer A.L., Mostaghimi S., Young R.W., Dillaha T.A., Shanholtz V.O. (1987) Effects of residue cover on pesticide losses from conventional and no-tillage systems, T. ASAE 30, 953–959.
- Kitchen N.R., Hughes D.E., Donald W.W., Alberts E.E. (1998) Agrichemical movement in the root-zone of claypan soils: ridgeand mulch-tillage systems compared, Soil Till. Res. 48, 179–193.
- Kung K.J.S., Steenhuis T.S., Kladivko E.J., Gish T.J., Bubenzer G., Helling C.S. (2000) Impact of preferential flow on the transport of adsorbing and non-adsorbing tracers, Soil Sci. Soc. Am. J. 64, 1290–1296.
- Lafrance P., Marineau L., Perreault L., Villeneuve J.P. (1994) Effect of natural dissolved organic matter found in groundwater on soil adsorption and transport of pentachlorophenol, Environ. Sci. Technol. 28, 2314–2320.
- Lal R., Mahboubi A.A., Fausey N.R. (1994) Long-term tillage and rotation effects on properties of a Central Ohio Soil, Soil Sci. Soc. Am. J. 58, 517–522.
- Lal R., Reicosky D.L., Hanson J.D. (2007) Evolution of the plow over 10,000 years and the rationale for no-till farming, Soil Till. Res. 93, 1–12.
- Lamb J.A., Dowdy R.H., Anderson J.L., Allmaras R.R. (1998) Water quality in an irrigated sandy soil: ridge tillage in rotated corn and soybean compared with full-width tillage in continuous corn, Soil Till. Res. 48, 167–177.
- Lavorenti A., Rocha A.A., Prata F., Regitano J.B., Tornisielo V.L., Pinto O.B. (2003) Reactions of diclosulam in a Rhodic Hapludox soil under no-till and conventional tillage systems, Rev. Bras. Cienc. Solo 27, 183–190.
- Levanon D., Codling E.E., Meisinger J.J., Starr J.L. (1993) Mobility of agrochemicals through soil from two tillage systems, J. Environ. Qual. 22, 155–161.
- Levanon D., Meisinger J.J., Codling E.E., Starr J.L. (1994) Impact of tillage on microbial activity and the fate of pesticides in the upper soil, Water Air Soil Poll. 72, 179–189.
- Li K., Liu W.P., Xu D.M., Lee S.J. (2003) Influence of organic matter and pH on bentazone sorption in soils, J. Agr. Food Chem. 51, 5362– 5366.
- Li K., Xing B.S., Torello W.A. (2005) Effect of organic fertilizers derived dissolved organic matter on pesticide sorption and leaching, Environ. Pollut. 134, 187–194.
- Locke M.A. (1992) Sorption-desorption kinetics of alachlor in surface soil from two soybean tillage systems, J. Environ. Qual. 21, 558– 566.
- Locke M.A., Zablotowicz R.M., Gaston L.A. (1995) Fluometuron herbicide interactions in crop residue-managed soils, in: Buehring N.W., Kingery W.A. (Eds.), Proc. Southern Conservation Tillage Conf. for Sustainable Agriculture: A focus on water quality. Jackson, MS. MAFES, Mississippi State University, Mississippi State, Starkville, MS, pp. 55–58.

- Locke M.A., Gaston L.A., Zablotowicz R.M. (1996) Alachlor biotransformation and sorption in soil from two soybean tillage systems, J. Agr. Food Chem. 44, 1128–1134.
- Locke M.A., Harper S.S. (1991a) Metribuzin degradation in soil: I-Effects of residue amendment, metribuzin level, and soil depth, Pestic. Sci. 31, 221–237.
- Locke M.A., Harper S.S. (1991b) Metribuzin degradation in soil: II-Effects of tillage, Pestic. Sci. 31, 239–247.
- Locke M.A., Bryson C.T. (1997) Herbicide-soil interactions in reduced tillage and plant residue management systems, Weed Sci. 45, 307– 320.
- Locke M.A., Zablotowicz R.M., Bauer P.J., Steinriede R.W., Gaston L.A. (2005) Conservation cotton production in the southern United States: herbicide dissipation in soil and cover crops, Weed Sci. 53, 717–727.
- Logan T.J., Eckert D.J., Beak D.G. (1994) Tillage, crop and climatic effects on runoff and tile drainage losses of nitrate and four herbicides, Soil Till. Res. 30, 75–103.
- Loux M.M., Liebl R.A., Slife F.W. (1989) Availability and persistence of imazaquin, imazethapyr, and clomazone in soil, Weed Sci. 37, 259–267.
- Lowery B., Hartwig R.C., Stoltenberg D.E., Fermanich K.J., McSweeney K. (1998) Groundwater quality and crop-yield responses to tillage management on a Sparta sand, Soil Till. Res. 48, 225–237.
- Ma L.W., Selim H.M. (2005) Predicting pesticide transport in mulchamended soils: A two-compartment model, Soil Sci. Soc. Am. J. 69, 318–327.
- Malone R.W., Warner R.C., Byers M.E. (1996) Runoff losses of surfaceapplied metribuzin as influenced by yard waste compost amendments, no-tillage, and conventional-tillage, B. Environ. Contam. Tox. 57, 536–543.
- Malone R.W., Shipitalo M.J., Ma L., Ahuja L.R., Rojas K.W. (2001) Macropore component assessment of the root zone water quality model (RZWQM) using no-till soil blocks, T. ASAE 44, 843–852.
- Martin C.D., Baker J.L., Erbach D.C., Johnson H.P. (1978) Washoff of herbicides applied to corn residue, T. ASAE 21, 1164–1168.
- Masse L., Patni N.K., Jui P.Y., Clegg B.S. (1996) Tile effluent quality and chemical losses under conventional and no tillage - Part 2: atrazine and metolachlor, T. ASAE 39, 1673–1679.
- Masse L., Patni N.K., Jui P.Y., Clegg B.S. (1998) Groundwater quality under conventional and no tillage: II. Atrazine, deethylatrazine, and metolachlor, J. Environ. Qual. 27, 877–883.
- Masutti C. (2004) Le Dust Bowl, la politique de conservation des ressources et les écologues aux États-Unis dans les années 1930, Thèse de Doctorat de l'Université Louis Pasteur.
- Mazzoncini M., Lorenzi R., Risaliti R., Sorce C., Ginanni M., Curadi M., Pini R. (1998) Diclofop-methyl dissipation in clay soil under different tillage systems in central Italy, Soil Till. Res. 46, 241–250.
- Mills J.A., Witt W.W. (1991) Dissipation of imazaquin and imazethapyr under conventional and no-tillage soybean (Glycine max), Weed Technol. 5, 586–591.
- Mills J.A., Witt W.W., Barrett M. (1989) Effects of tillage on the efficacy and persistence of clomazone in soybean (Glycine max), Weed Sci. 37, 233–238.
- Monks C.D., Banks P.A. (1993) Effect of straw, ash, and tillage on dissipation of imazaquin and imazethapyr, Weed Sci. 41, 133–137.
- Moonen A.C., Barberi P. (2004) Size and composition of the weed seedbank after 7 years of different cover-crop-maize management systems, Weed Res. 44, 163–177.

- Myers J.L., Wagger M.G., Leidy R.B. (1995) Chemical movement in relation to tillage system and simulated rainfall intensity, J. Environ. Qual. 24, 1183–1192.
- Nanny M.A., Maza J.P. (2001) Noncovalent interactions between monoaromatic compounds and dissolved humic acids: A deuterium NMR T-1 relaxation study, Environ. Sci. Technol. 35, 379–384.
- Novak J.M., Watts D.W., Hunt P.G. (1996) Long-term tillage effects on atrazine and fluometuron sorption in coastal plain soils, Agr. Ecosyst. Environ. 60, 165–173.
- Ochsner T.E., Stephens B.M., Koskinen W.C., Kookana R.S. (2006) Sorption of a hydrophilic pesticide: Effects of soil water content, Soil Sci. Soc. Am. J. 70, 1991–1997.
- Ogden C.B., van Es H.M., Wagenet R.J., Steenhuis T.S. (1999) Spatialtemporal variability of preferential flow in a clay soil under no-till and plow-till, J. Environ. Qual. 28, 1264–1273.
- Olson B.L.S., Regehr D.L., Janssen K.A., Barnes P.L. (1998) Tillage system effects on atrazine loss in surface water runoff, Weed Technol. 12, 646–651.
- Otto S., Riello L., During R.A., Hummel H.E., Zanin G. (1997) Herbicide dissipation and dynamics modelling in three different tillage systems, Chemosphere 34, 163–178.
- Peigné J., Ball B.C., Roger-Estrade J., David C. (2007) Is conservation tillage suitable for organic farming? A review, Soil Use Manage. 23, 129–144.
- Phillips R.E., Quisenberry V.L., Zeleznik J.M., Dunn G.H. (1989) Mechanism of water entry into simulated macropores, Soil Sci. Soc. Am. J. 53, 1629–1635.
- Pinheiro E.F.M., Pereira M.G., Anjos L.H.C. (2004) Aggregate distribution and soil organic matter under different tillage systems for vegetable crops in a Red Latosol from Brazil, Soil Till. Res. 77, 79–84.
- Pot V., Simunek J., Benoit, P., Coquet Y., Yra A., Martinez-Cordon M. J. (2005) Impact of rainfall intensity on the transport of two herbicides in undisturbed grassed filter strip soil cores, J. Contam. Hydrol. 81, 63–88.
- Potter T.L., Truman C.C., Bosch D.D., Bednarz C.W. (2003) Cotton defoliant runoff as a function of active ingredient and tillage, J. Environ. Qual. 32, 2180–2188.
- Potter T.L., Truman C.C., Bosch D.D., Bednarz C. (2004) Fluometuron and pendimethalin runoff from strip and conventionally tilled cotton in the southern Atlantic Coastal Plain, J. Environ. Qual. 33 2122– 2131.
- Potter T.L., Truman C.C., Strickland T.C., Bosch D.D., Webster T.M., Franklin D.H., Bednarz C.W. (2006) Combined effects of constant versus variable intensity simulated rainfall and reduced tillage management on cotton preemergence herbicide runoff, J. Environ. Qual. 35, 1894–1902.
- Preston C.M., Newman R.H., Rother P. (1994) Using ¹³C CPMAS NMR to assess the effects of cultivation on the organic matter of particle size fractions in a grassland soil, Soil Sci. 157, 26–35.
- Quisenberry V.L., Phillips R.E., Zeleznik J.M. (1994) Spatial distribution of water and chloride macropore flow in a well-structured soil, Soil Sci. Soc. Am. J. 58, 1294–1300.
- Rasmussen K.J. (1991) Reduced soil tillage and Italian ryegrass as catch crop. II. Soil bulk density, root development and soil chemistry, Dan. J. Plant Soil Sci. 95, 139–154.
- Réal B., Malaval C., Bonin L., Labreuche J., Barriuso E., Benoit P., Bedos C., Köller R., Heddadj D., Quéré L., Cariolle M., Caboulet D., Alletto L. (2007) Impacts des TCSL sur les transferts de produits phytosanitaires, Évaluation des impacts environnementaux des techniques culturales sans labour (TCSL) en France, Edition ADEME, pp. 250–268.

- Rector R.J., Regehr D.L., Barnes P.L., Loughin T.M. (2003) Atrazine, Smetolachlor, and isoxaflutole loss in runoff as affected by rainfall and management, Weed Sci. 51, 810–816.
- Reddy K.N., Locke M.A. (1996) Imazaquin spray retention, foliar washoff, and runoff losses under simulated rainfall, Pestic. Sci. 48, 179–187.
- Reddy K.N., Locke M.A. (1998) Sulfentrazone sorption, desorption, and mineralization in soils from two tillage systems, Weed Sci. 46, 494– 500.
- Reddy K.N., Zablotowicz R.M., Locke M.A. (1995a) Chlorimuron adsorption, desorption, and degradation in soils from conventional tillage and no-tillage systems, J. Environ. Qual. 24, 760–767.
- Reddy K.N., Locke M.A., Wagner S.C., Zablotowicz R.M., Gaston L.A., Smeda R.J. (1995b) Chlorimuron ethyl sorption and desorption kinetics in soils and herbicide-dessicated cover crop residues, J. Agr. Food Chem. 43, 2752–2757.
- Reddy K.N., Locke M.A., Gaston L.A. (1997a) Tillage and cover crop effects on cyanazine adsorption and desorption kinetics, Soil Sci. 162, 501–509.
- Reddy K.N., Locke M.A., Zablotowicz R.M. (1997b) Soil type and tillage effects on sorption of cyanazine and degradation products, Weed Sci. 45, 727–732.
- Renner K.A., Schabenberger O., Kells J.J. (1998) Effect of tillage and application method on corn (*Zea mays*) response to imidazolinone residues in soil, Weed Technol. 12, 281–285.
- Rhoton F.E., Shipitalo M.J., Lindbo D.L. (2002) Runoff and soil loss from midwestern and southeastern US silt loam soils as affected by tillage practice and soil organic matter content, Soil Till. Res. 66, 1–11.
- Ritter W.F., Chirnside A.E.M., Scarborough R.W. (1996) Movement and degradation of triazines, alachlor, and metolachlor in sandy soils, J. Environ. Sci. Health 31, 2699–2721.
- Rovira A.D., Smettem K.R.J., Lee K.E. (1987) Effect of rotation and conservation tillage on earthworms in a Red-brown earth under wheat, Aust. J. Soil Res. 38, 829–834.
- Sadeghi A.M., Isensee A.R. (1992) Effect of tillage systems and rainfall patterns on atrazine distribution in soil, J. Environ. Qual. 21, 464– 469.
- Sadeghi A.M., Isensee A.R. (1996) Impact of reversing tillage practices on movement and dissipation of atrazine in soil, Soil Sci. 161, 390– 397.
- Sadeghi A.M., Isensee A.R. (1997) Alachlor and cyanazine persistence in soil under different tillage and rainfall regimes, Soil Sci. 162, 430–438.
- Sadeghi A.M., Isensee A.R. (2001) Impact of hairy vetch cover crop on herbicide transport under field and laboratory conditions, Chemosphere 44, 109–118.
- Sadeghi A.M., Isensee A.R., Shelton D.R. (1998) Effect of tillage age on herbicide dissipation: A side-by-side comparison using microplots, Soil Sci. 163, 883–890.
- Sadeghi A.M., Isensee A.R., Shirmohammadi A. (2000) Influence of soil texture and tillage on herbicide transport, Chemosphere 41, 1327– 1332.
- Salloum M.J., Dudas M.J., McGill W.B. (2001) Variation of 1-naphthol sorption with organic matter fractionation: the role of physical conformation, Org. Geochem. 32, 709–719.
- Satchell J.E. (1983) Earthworm Ecology from Darwin to vermiculture. Ed. Chapman and Hall, London, 495 p.
- Sauer T.J., Daniel T.C. (1987) Effect of tillage system on runoff losses of surface-applied pesticides, Soil Sci. Soc. Am. J. 51, 410–415.

- Seifert S., Shaw D.R., Kingery W.L., Snipes C.E., Wesley R.A. (2001a) Imazaquin mobility and persistence in a Sharkey clay soil as influenced by tillage systems, Weed Sci. 49, 571–577.
- Seifert S., Shaw D.R., Zablotowicz R.M., Wesley R.A., Kingery W.L. (2001b) Effect of tillage on microbial characteristics and herbicide degradation in a Sharkey clay soil, Weed Sci. 49, 685–693.
- Selim H.M., Zhou L., Zhu H. (2003) Herbicide retention in soil as affected by sugarcane mulch residue, J. Environ. Qual. 32, 1445– 1454.
- Senesi N., Testini C. (1982) Physico-chemical investigations of interaction mechanisms between s-triazine herbicides and soil humic acids, Geoderma 28, 129–146.
- Seta A.K., Blevins R.L., Frye W.W., Barfield B.J. (1993) Reducing soil erosion and agricultural chemical losses with conservation tillage, J. Environ. Qual. 22, 661–665.
- Seyfried M.S., Rao P.S.C. (1987) Solute transport in undisturbed columns of an aggregated tropical soil: preferential flow effects, Soil Sci. Soc. Am. J. 51, 1434–1444.
- Shang C., Arshad M.A. (1998) Sorption of clopyralid, dicamba and MCPA by two soils with conventional and no-till management, Can. J. Soil Sci. 78, 181–186.
- Shelton D.R., Sadeghi A.M., Isensee A.R. (1998) Effect of tillage on atrazine bioavailability, Soil Sci. 163, 891–896.
- Shipitalo M.J., Edwards W.M. (1996) Effects of initial water content on macropore/matrix flow and transport of surface-applied chemicals, J. Environ. Qual. 25, 662–670.
- Shipitalo M.J., Owens L.B. (2003) Atrazine, deethylatrazine, and deisopropylatrazine in surface runoff from conservation tilled watersheds, Environ. Sci. Technol. 37, 944–950.
- Shipitalo M.J., Owens L.B. (2006) Tillage system, application rate, and extreme event effects on herbicide losses in surface runoff, J. Environ. Qual. 35, 2186–2194.
- Shipitalo M.J., Edwards W.M., Dick W.A., Owens L.B. (1990) Initial storm effects on macropore transport of surface-applied chemicals in no-till soil, Soil Sci. Soc. Am. J. 54, 1530–1536.
- Shipitalo M.J., Edwards W.M., Owens L.B. (1997) Herbicide losses in runoff from conservation-tilled watersheds in a corn-soybean rotation, Soil Sci. Soc. Am. J. 61, 267–272.
- Sigua G.C., Isensee A.R., Sadeghi A.M. (1993) Influence of rainfall intensity and crop residue on leaching of atrazine through intact notill soil cores, Soil Sci. 156, 225–232.
- Sigua G.C., Isensee A.R., Sadeghi A.M. (1995) Influence of tillage, antecedent moisture, and rainfall timing on atrazine transport, Weed Sci. 43, 134–139.
- Singh N., Kloeppel H., Klein W. (2002) Movement of metolachlor and terbuthylazine in core and packed soil columns, Chemosphere 47, 409–415.
- Singh P., Kanwar R.S. (1991) Preferential solute transport through macropores in large undisturbed saturated soil columns, J. Environ. Qual. 20, 295–300.
- Six J., Elliott E.T., Paustian K. (1999) Aggregate and soil organic matter dynamics under conventional and no-tillage systems, Soil Sci. Soc. Am. J. 63, 1350–1358.
- Sleutel S., Kader M.A., Leinweber P., D'Haene K., De Neve S. (2007) Tillage management alters surface soil organic matter composition: A pyrolysis mass spectroscopy study, Soil Sci. Soc. Am. J. 71, 1620–1628.

- Sorenson B.A., Shea P.J., Roeth F.W. (1991) Effects of tillage, application time and rate on metribuzin dissipation, Weed Res. 31, 333–345.
- Stearman G.K., Lewis R.J., Tortorelli L.J., Tyler D.D. (1989) Herbicide reactivity of soil organic matter fractions in no-tilled and tilled cotton, Soil Sci. Soc. Am. J. 53, 1690–1694.
- Stehouwer R.C., Dick W.A., Traina S.J. (1993) Characteristics of earthworm burrow lining affecting atrazine sorption, J. Environ. Qual. 22, 181–185.
- Suba J.D., Essington M.E. (1999) Adsorption of fluometuron and norflurazon: Effect of tillage and dissolved organic carbon, Soil Sci. 164, 145–155.
- Szajdak L., Jezierski A., Cabrera M.L. (2003) Impact of conventional and no-tillage management on soil amino acids, stable and transient radicals and properties of humic and fulvic acids, Org. Geochem. 34, 693–700.
- Tatzber M., Stemmer M., Spiegel H., Katzlberger C., Haberhauer G., Gerzabek M. H. (2008) Impact of different tillage practices on molecular characteristics of humic acids in a long-term field experiment – An application of three different spectroscopic methods, Sci. Total Environ. 406, 256–268.
- Teasdale J.R., Shelton D.R., Sadeghi A.M., Isensee A.R. (2003) Influence of hairy vetch residue on atrazine and metolachlor soil solution concentration and weed emergence, Weed Sci. 51, 628–634.
- Tebrügge F., During R.A. (1999) Reducing tillage intensity a review of results from a long-term study in Germany, Soil Till. Res. 53, 15–28.
- Torresen K.S., Skuterud R., Tandsaether H.J., Hagemo M.B. (2003) Long-term experiments with reduced tillage in spring cereals. I. Effects on weed flora, weed seedbank and grain yield, Crop Prot. 22, 185–200.
- Triplett G.B., Conner B.J., Edwards W.M. (1978) Transport of atrazine and simazine in runoff from conventional and no-tillage corn, J. Environ. Qual. 7, 77–84.
- Trojan M.D., Linden D.R. (1992) Microrelief and rainfall effects on water and solute movement in earthworm burrows, Soil Sci. Soc. Am. J. 56, 727–733.
- Ulbrich A.V., Souza J.R.P., Shaner D. (2005) Persistence and carryover effect of imazapic and imazapyr in Brazilian cropping systems, Weed Technol. 19, 986–991.
- Unger P.W. (1987) Straw mulch effects on soil temperature and sorghum germination and growth, Agron. J. 70, 858–864.
- van Genuchten M.T., Wierenga P.J. (1976) Mass transfer studies in sorbing porous media. I. Analytical solutions, Soil Sci. Soc. Am. J. 40, 473–480.
- van Genuchten M.T., Wagenet R.J. (1989) Two-site/two-region for pesticide transport and degradation: theoretical development and analytical solutions, Soil Sci. Soc. Am. J. 53, 1303–1310.
- Vasileiadis V.P., Froud-Williams R.J., Eleftherohorinos I.G. (2007) Vertical distribution, size and composition of the weed seedbank under various tillage and herbicide treatments in a sequence of industrial crops, Weed Res. 47, 222–230.
- Wagner S.C., Zablotowicz R.M., Gaston L.A., Locke M.A., Kinsella J. (1996) Bentazon degradation in soil: Influence of tillage and history of bentazon application, J. Agric. Food Chem. 44, 1593–1598.
- Walker A. (1971) Effects of soil moisture content on the availability of soil-applied herbicides to plants, Pestic. Sci. 2, 56–59.
- Watts D.W., Hall J.K. (1996) Tillage and application effects on herbicide leaching and runoff, Soil Till. Res. 39, 241–257.

L. Alletto et al.

- Wauchope R.D. (1978) The pesticide content of surface water draining from agrigultural fields - a review, J. Environ. Qual. 7, 459–472.
- Weber J.B., Hardy D.H., Leidy R.B. (2002) Laboratory, green-house, and field lysimeter studies of ¹⁴C-atrazine volatilization, in: Phelps W. et al. (Eds.), Pesticide environmental fate. ACS Symp. Ser. 813. Am. Chem. Soc., Washington, DC, pp. 125–142.
- Weber J.B., Taylor K.A., Wilkerson G.G. (2006) Soil cover and tillage influenced metolachlor mobility and dissipation in field lysimeters, Agron. J. 98, 19–25.
- Webster E.P., Shaw D.R. (1996) Off-site runoff losses of metolachlor and metribuzin applied to differing soybean (Glycine max) production systems, Weed Technol. 10, 556–564.
- Weed D.A.J., Kanwar R.S., Stoltenberg D.E., Pfeiffer R.L. (1995) Dissipation and distribution of herbicides in the soil profile, J. Environ. Qual. 24, 68–79.
- Weed D.A.J., Kanwar R.S., Cambardella C., Moorman T.B. (1998) Alachlor dissipation in shallow cropland soil, J. Environ. Qual. 27, 767–776.
- Whang J.M., Schomburg C.J., Glotfelty D.E., Taylor A.W. (1993) Volatilization of fonofos, chlorpyrifos, and atrazine from conventional and no-till surface soils in the field, J. Environ. Qual. 22, 173–180.

- Wienhold B.J., Gish T.J. (1994) Effect of formulation and tillage practice on volatilization of atrazine and alachlor, J. Environ. Qual. 23, 292– 298.
- Worsham A.D. (1991) Role of cover crops in weed management and water quality, in: Hargrove W.L. (Ed.), Cover Crops for Clean Water, Jackson, TN, USA, pp. 141–145.
- Xing B. (2001) Sorption of naphthalene and phenanthrene by soil humic acids, Environ. Pollut. 111, 303–309.
- Xing B., Pignatello J.J. (1997) Dual-mode sorption of low-polarity compounds in glassy poly(vinylchloride) and soil organic matter, Environ. Sci. Technol. 31, 792–799.
- Xue S.L., Chen S., Selim H.M. (1997) Modeling alachlor transport in saturated soils from no-till and conventional tillage systems, J. Environ. Qual. 26, 1300–1307.
- Yuan G.S., Xing B.S. (1999) Site-energy distribution analysis of organic chemical sorption by soil organic matter, Soil Sci. 164, 503–509.
- Zablotowicz R.M., Locke M.A., Smeda R.J. (1998) Degradation of 2,4-D and fluometuron in cover crop residues, Chemosphere 37, 87–101.
- Zablotowicz R.M., Locke M.A., Gaston L.A., Bryson C.T. (2000) Interactions of tillage and soil depth on fluometuron degradation in a Dundee silt loam soil, Soil Till. Res. 57, 61–68.