



INFLUENCE OF DIFFERENT FOOD RESTRICTION METHODS APPLIED DURING THE LAYING PERIOD ON LAYERS PERFORMANCES

M. Bougon, Mireille Levitoux, M. Le Menec, Y. Heuze

► To cite this version:

M. Bougon, Mireille Levitoux, M. Le Menec, Y. Heuze. INFLUENCE OF DIFFERENT FOOD RESTRICTION METHODS APPLIED DURING THE LAYING PERIOD ON LAYERS PERFORMANCES. *Annales de zootechnie*, 1975, 24 (1), pp.162-162. hal-00887440

HAL Id: hal-00887440

<https://hal.science/hal-00887440>

Submitted on 11 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

INFLUENCE OF DIFFERENT FOOD RESTRICTION METHODS APPLIED DURING THE LAYING PERIOD ON LAYERS PERFORMANCES

M. BOUGON

avec la collaboration technique de Mireille LEVITOUX, M. LE MENEC
et Y. HEUZE

*Station expérimentale d'Aviculture,
22440 Ploufragan*

2304 W. S. S. L. layers randomised in 6 groups received a food with a metabolisable energy content of 2.750 kcal/kg from 21 to 66 weeks of age.

The food was distributed *ad libitum* to group A and in limited quantities to groups B-C-D from 30 weeks so that each bird received an average of 116.4 g per day (320 kcal).

The restriction was modified in group C so that 111 g (305 kcal) per day were fed for the last 21 weeks of the period.

Group D was restricted as group B, except that the food was distributed at 3 p.m. instead of 8 a.m. as in group B.

Groups E and F had free access to food from 21 to 25 weeks of age. Group E was then restricted by allowing access to food for 2 × 2 hour periods (8-10 h and 15-17 h). From 25 to 37 weeks of age, group F received food for 4 consecutive hours (8 to 12 a.m.) followed by 6 hours of feeding (8 a.m. to 2 p.m.) from 37 to 66 weeks of age.

While groups B and D receiving measured amounts of feed showed a 9.1 p. 100 restriction in practice, group C ate 11.1 p. 100 less than the controls. Limiting access time resulted in a similar level of restriction, group E consuming 11.4 p. 100 less, while group F was subjected to the severest restriction at 13.8 p. 100.

In all restricted treatments, the food efficiency was improved and the cost of production of 1 kg of egg decreased. The best result in economic terms, was achieved by treatment E.

The study suggests that it is better to distribute the food at 3 p.m. instead of 8 a.m. and it is not advisable to increase the level of restriction towards the end of the laying period.

Access to food for 4 consecutive hours leads to an insufficient food intake and poor production. It is necessary to increase the time of access to 6 hours or split it to two hours in the morning and two in the afternoon.

IV. — Pathology

UTILIZATION OF ANTIFUNGAL ANTIBIOTICS IN GUINEA-FOWL EXPERIMENTALLY INFECTED WITH *CANDIDA ALBICANS*

G. JOLIVET et M. MORMEDE*

*Laboratoire de Parasitologie,
École nationale vétérinaire d'Alfort, I. N. R. A.,*

* *Laboratoire Squibb, Sofcf,
Paris*