



Colibacillar enterotoxic syndrome of the pig in France

L. Renault, Cl. Maire, J. Vaissaire

► To cite this version:

| L. Renault, Cl. Maire, J. Vaissaire. Colibacillar enterotoxic syndrome of the pig in France. Annales de zootechnie, 1976, 25 (3), pp.450-450. hal-00887638

HAL Id: hal-00887638

<https://hal.science/hal-00887638v1>

Submitted on 11 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Study of blood platelets in the pig

M. SAVEY, B. BONEU, J. ESPINASSE

*École nationale vétérinaire de Toulouse,
Chemin des Capelles,
31076 Toulouse*

As part of a systematic study of hemostasis in the pig, we studied the properties of platelets in that species. After briefly describing their basic role in the hemostatic mechanism, we examined the volume and density distribution of platelets in the pig, comparing our results with those obtained in man.

Colibacillar enterotoxic syndrome of the pig in France

L. RENAULT, CI. MAIRE, J. VAISSAIRE

*Établissements Sanders,
17, quai de l'Industrie,
91260 Juvisy-sur-Orge*

A serological grouping of 147 *Escherichia coli* strains, isolated from 52 newborn piglets and from 105 piglets at weaning revealed that 3/4 of the strains were enteropathogenic. In the new born piglet, the strains isolated were mainly 0149 : K 91, K 88 ac and in the piglet at weaning, by order of frequency : 0141 : K 85 ab, K 88 ac, 0149 : K 91, K 88 ac and 08 : K 87, K 88 ac.

Colistine excepted, all strains showed a considerable lowering of sensitivity towards all currently used antibiotics.

Vaccinations made peroral or from specific fractions of enterotoxines seem to be promissing.