



HAL
open science

Amino acid requirements of preruminant lambs

P. Patureau-Mirand, M. Theriez

► **To cite this version:**

P. Patureau-Mirand, M. Theriez. Amino acid requirements of preruminant lambs. *Annales de zootechnie*, 1977, 26 (2), pp.287-287. <hal-00887695>

HAL Id: hal-00887695

<https://hal.science/hal-00887695v1>

Submitted on 11 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

was partially replaced by dry peas or urea, the latter not exceeding 1.5 % of the concentrate in any case;

— The problem of protein supplementation of farm cereal mixtures is not completely solved. Even if there is free access to cereals and protein feed, over-consumption of the latter is almost always the rule.

Amino acid requirements of preruminant lambs

P. PATUREAU-MIRAND (*) and M. THERIEZ (**)

(*) *Laboratoire d'étude du métabolisme azoté,*
 (**) *Station de Recherches sur l'Élevage des Ruminants,*
Centre de Recherches de Clermont-Ferrand, I.N.R.A.,
Theix 63110 Beaumont (France).

The amino acid requirements of preruminant lambs were tentatively determined by different methods:

— ewe milk amino acids are assumed to satisfy the amino acid requirement of lambs and are compared to those of cows milk (Table I);

— the amounts of essential amino acids ingested by lambs in which the protein requirement is satisfied may be used as rough estimations of the amino acid requirements of preruminant lambs (Table 2). This method of evaluation tends to underestimate the requirements for the limiting amino acids of the experimental protein, and to overestimate the requirements for amino acids in excess of that protein.

The requirements for some essential amino acids were directly determined by measuring the nitrogen balance and/or the blood level of free amino acids of lambs fed increasing amounts of those amino acids. Thus, the methionine requirement of preruminant lambs was about 2.0 g/d (Fig. I), i.e. 2.6 g/d of sulphur amino acids.

The amino acid requirements of fast-growing lambs (250 to 300 g/d), evaluated by the different methods, are summarized in Table 5. It may be inferred that amino acid requirements vary according to age and feeding level. The amino acid requirements of 8-day old preruminant lambs fed *ad libitum* seemed to be satisfied by 67 g/d of cow milk protein supplemented by 0.3 g/d of DL methionine and 0.9 g/d of L lysine. When the lambs were 21 day old, they needed 72 g/d of cow milk protein supplemented by 0.1 g/d of DL methionine and 0.5 g/d of L lysine, if fed *ad libitum*: 76 g/d of cow milk protein supplemented by 0.3 g/d of DL methionine if the feeding level was only 80 % of *ad libitum* feed consumption.

Effect of dehydrated pelleted hay on calcium and phosphate metabolism in lambs

J. P. BARLET

Station de Physiopathologie de la Nutrition,
Centre de Recherches de Clermont-Ferrand, I.N.R.A.,
Theix, 63110 Beaumont (France).

In 3-month-old ram lambs fed dehydrated pelleted lucerne hay, the urinary and faecal excretion of calcium was significantly higher than in control lambs fed normal lucerne hay (Table II). The causes for these differences are still unknown. This may be due to an increase in the speed