



The digestion of fresh perennial ryegrass fertilized at 2 levels of nitrogen in lactating dairy cows

Jl Peyraud, L Astigarraga, Philippe Faverdin

► To cite this version:

Jl Peyraud, L Astigarraga, Philippe Faverdin. The digestion of fresh perennial ryegrass fertilized at 2 levels of nitrogen in lactating dairy cows. *Annales de zootechnie*, 1993, 42 (2), pp.138-138. hal-00888883

HAL Id: hal-00888883

<https://hal.science/hal-00888883>

Submitted on 11 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The digestion of fresh perennial ryegrass fertilized at 2 levels of nitrogen in lactating dairy cows

JL Peyraud, L Astigarraga, P Faverdin

INRA, station de recherches sur la Vache laitière, 35590 Saint-Gilles, France

The digestion of protein in intensively fertilized grasses is characterized by high losses of nitrogen (N) in the urine. This experiment aimed at studying the effect of lowering N fertilization on N metabolism in the dairy cow.

The effect of 2 levels of N fertilization (0 or 80 Kg N/ha before and after the first cut in spring 1991) of fresh perennial ryegrass on digestion was tested with a second cut (30-d-old) in a cross-over design on 4 fistulated dairy cows (20 kg milk). Grass was given *ad libitum* in 3 meals per day. The flow of duodenal digesta was calculated from PEG and Yb recovered in faeces and their respective concentration in digesta. Feeding behaviour was also recorded.

The absence of N fertilizer resulted in a reduction in N content of grass (17.4 vs 24.2 g/kg DM) and N intake (NI) (table I) whereas organic matter (OM) intake was not affected (14 kg/d). OM and NDF digestibilities were reduced (0.79 vs 0.81 and 0.72 vs 0.78, respectively for OM and NDF, $P < 0.05$). Due to a higher DM content in unfertilized grass (197 vs 145 g/kg) time spent eating was reduced (495 vs 607 min/d). The non-ammonia nitrogen flow into the duodenum (NAN) expressed in g/d or as a fraction of digestible OM intake (32.5 g/kg DOMI), was similar in the 2 treatments ($P > 0.1$). NAN was equal to NI in the fertilized grass but exceeded NI by 95 g/d in the unfertilized grass. This net recycling matched the quantities of N excreted in urine (NU) which were indeed reduced by 95 g/d (in fact 86 g/d urea-N) with the unfertilized grass and could be related to the

lower rumen concentration of ammonia (NH_3). Reduced N fertilization led both to a slight decrease in feed N degradability estimated *in sacco* (0.74 vs 0.77) and in the absolute amount of feed N escaping the rumen degradation – from 96 to 76 g/d if predicted from the *in sacco* degradability (Vérité and Peyraud, 1989). The efficiency of microbial synthesis was calculated to be 25 g N/kg DOMI and was unlikely to be modified according to the treatment.

In conclusion, these results suggest that reduced N fertilization greatly affects N metabolism in the dairy cow, resulting in lower N excretion in urine.

Vérité R, Peyraud JL (1989) *Ruminant Nutrition* (Jarrige R, ed) John Libbey and Co, London, 33-47

Table I. Effect of N fertilization of N metabolism in dairy cows.

Fertilization	NI (g/d)	NAN (g/d)	NU (g/d)	NH_3 (mg/l)
0	263 ^a	358	69 ^a	15 ^a
80	367 ^b	367	164 ^b	99 ^b
rsd	22	4	5	19

a,b: means with different superscripts in the same column are significantly different ($P < 0.05$); rsd: from the model: $Y_{ijk} = \text{mean} + (\text{fertilization})_i + (\text{period})_j + (\text{cow})_k + e_{ijk}$.