



**HAL**  
open science

## Effect of grinding of hay on rumen particle kinetics in sheep

Laurence Bernard, Jp Chaise, C Poncet

► **To cite this version:**

Laurence Bernard, Jp Chaise, C Poncet. Effect of grinding of hay on rumen particle kinetics in sheep. Annales de zootechnie, 1993, 42 (2), pp.155-155. hal-00888896

**HAL Id: hal-00888896**

**<https://hal.science/hal-00888896>**

Submitted on 11 May 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## Effect of grinding of hay on rumen particle kinetics in sheep

L Bernard, JP Chaise, C Poncet

INRA de Theix, laboratoire de la Dynamique de la digestion, 63122 Saint-Genès-Champanelle, France

Grinding of hay has been found to have variable effects on rumen mean retention time (MRT). The aim of the present work was to clarify this point by studying components of MRT and underlying mechanisms.

Four mature Texel wethers fitted with rumen and duodenal cannulae were fed a mixture of a chopped and a ground (8-mm screen) and pelleted orchard-grass hay (2.4% N, 59.4% NDF) in the ratios (90:10, 50:50, 30:70 or 10:90) according to a 4 x 4 latin-square design. Food (1 200 g/d) was delivered in 8 equal meals. The amount of DM and indigestible acid detergent lignin (IADL) contents and size distribution of rumen particles were determined by emptying and wet-sieving. Duodenal digesta were analyzed in the same manner and their flows were estimated using the double-marker technique ( $^{169}\text{Yb}$  acetate and  $^{51}\text{Cr}$ -EDTA).

Particle MRT in the rumen estimated with IADL decreased and then plateaued when the proportion of ground hay in the diet was increased. The comminution rate of large particles (LP) was always higher than the small particle (SP) outflow rate, indicating that comminution is not the limiting step of passage. These results were the consequence of the curvilinear and opposite evolutions of the particulate DM pool in and outflow from the rumen (table I). Passage rate variations could not be explained by aperture number of the reticulo-omasal orifice.

*In conclusion*, the curvilinear effect of grinding grass hay at the same level of intake on components of MRT (pools and flow) may explain the discrepancy between published results in this field.

**Table I.** Particle kinetics in the rumen as affected by grinding a grass hay.

Chopped / Ground hay		90:10	50:50	30:70	10:90
Rumen DM pool (g)	Total	1123 <sup>a</sup>	851 <sup>b</sup>	840 <sup>b</sup>	858 <sup>b</sup>
	LP	335 <sup>a</sup>	174 <sup>b</sup>	146 <sup>b</sup>	137 <sup>b</sup>
	SP	721 <sup>a</sup>	602 <sup>b</sup>	626 <sup>b</sup>	665 <sup>ab</sup>
Duodenal DM flow (g/d)	Total	808	888	833	826
	SP	621 <sup>b</sup>	715 <sup>a</sup>	662 <sup>ab</sup>	648 <sup>b</sup>
IADL MRT (h)		42.2 <sup>a</sup>	28.0 <sup>b</sup>	31.5 <sup>b</sup>	30.8 <sup>b</sup>
LP comminution rate (%·h <sup>-1</sup> )		6.4	7.1	7.2	5.5
SP Passage rate (%·h <sup>-1</sup> )		3.7 <sup>d</sup>	5.0 <sup>a</sup>	4.4 <sup>b</sup>	4.1 <sup>c</sup>

a, b, c  $P < 0.05$ .