



An Improved technique for cecal cannulation in the rabbit. Effect of sampling time after feeding on cecal VFA pattern

Thierry Gidenne, R Bellier, T Bouyssou

► To cite this version:

Thierry Gidenne, R Bellier, T Bouyssou. An Improved technique for cecal cannulation in the rabbit. Effect of sampling time after feeding on cecal VFA pattern. *Annales de zootechnie*, 1993, 42 (2), pp.163-163. hal-00888902

HAL Id: hal-00888902

<https://hal.science/hal-00888902>

Submitted on 11 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

An Improved technique for cecal cannulation in the rabbit . Effect of sampling time after feeding on cecal VFA pattern

T Gidenne ¹, R Bellier ¹, T Bouyssou ²

¹ INRA, station de Recherches cunicoles, BP 27, 31326 Castanet;

² École nationale vétérinaire de Toulouse, 23 Ch des Capelles, 31076 Toulouse, France

The *in vivo* study of cecal fermentation in the rabbit required repetitive digesta sampling. Instead of a traditional plastic cannula (Carman and Waynforth, 1984), a glass cannula was used to reduce accidental damage by the rabbit and facilitate cecal digesta outflow during sampling.

A glass cannula (diameter 10 mm, length 45 mm) was inserted between the 4th and the 5th caecum whorl and secured to the cecal wall with a purse-string suture. Ten d after surgery, rabbits had returned to their pre-surgery feed intake and live weight. Cecal digesta collection (10–20 g each) was carried out at 7.30, 13.00, 15.00, and 17.00 h for 4 consecutive d (1 collection/day), on 4 adult rabbits (3 010 ± 200 g) receiving a standard diet (CP 17%, NDF 30%) given in 2 meals (8–11 h and 17–19 h).

Five h after feeding (13.00 h) the volatile fatty acid level was 4 times greater than at

7.30 h (starvation state), whereas ammonia and pH decreased sharply (table I). VFA molar proportions were close to the classic rabbit VFA profile (butyrate > propionate) except before feeding, when the propionate level was higher than that of butyrate. The glass cannula remained functional and allowed repetitive digesta sampling over a period of 4–6 months without significant disturbance to the animal.

In conclusion, this technique may be useful for the elaboration of an *in vivo* experimental model to study cecal fermentation under different nutritional status.

Carman RJ, Waynforth HB (1984) *Lab Anim* 18, 258–260

Table I. Post-feeding changes of caecal fermentation in the adult rabbit.

Time (h)	pH	Ammonia	VFA (mM/l)	Acetate (%)	Propionate (%)	Butyrate (%)
7.30	6.56 ^a	18.0 ^a	22.8 ^c	84.5	8.4 ^a	4.7 ^a
13.00	6.02 ^a	5.7 ^b	96.4 ^a	84.2	4.1 ^b	10.0 ^b
15.00	6.27 ^{ab}	5.0 ^b	70.1 ^b	83.5	4.4 ^b	10.1 ^b
17.00	6.47 ^a	9.6 ^b	55.2 ^b	83.2	4.8 ^b	10.1 ^b
Root MSE	0.20	1.8	8.1	1.3	1.0	0.4
Time (<i>P</i> level)	0.09	0.004	0.001	0.07	0.02	0.001
Indiv (<i>P</i> level)	0.28	0.17	0.97	0.003	0.08	0.001

In the same column, means with a different letter differ at the level *P* < 0.05 according to time effect.