



HAL
open science

Autocatalytic models describing ruminal in situ digestion

Jaap J. van Milgen

► **To cite this version:**

Jaap J. van Milgen. Autocatalytic models describing ruminal in situ digestion. *Annales de zootechnie*, 1994, 43 (3), pp.275-275. hal-00889014

HAL Id: hal-00889014

<https://hal.science/hal-00889014v1>

Submitted on 11 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Autocatalytic models describing ruminal *in situ* digestion

J Van Milgen

INRA, Station de Recherche sur la Nutrition des Herbivores, Theix,
 63122 Saint-Genès-Champanelle, France

Digestion of particulate matter in the rumen is a microbial enzymatic process and probably depends on both substrate and enzyme availability. Although microbes can be considered the cause of digestion, they are also the result of digestion, being one of the digestion end-products. Thus, digestion partially is an autocatalytic process.

In the classical digestion model (model 1), rate of digestion (dS/dt) is proportional only to the concentration of potentially digestible substrate (S). The effect of other factors affecting digestion is represented by the fractional digestion rate, which is assumed to be constant (k_d). To account for time-varying effects of microbes (or enzymes) on digestion of substrate, 2 models are proposed which are based on a non-constant fractional digestion rate (Z), such that $dS/dt = -ZS$:

1. $dZ/dt = 0$; $Z = 0$ for $t \leq t_i$ and $Z = k_d$ for $t > t_i$
2. $dZ/dt = (Z_i - Z)Z$ with $Z_0 \neq 0$; $Z = Z_i Z_0 / (Z_0 + (Z_i - Z_0) \exp(-Z_i t))$
3. $dZ/dt = (Z_i - Z)k_z$ with $Z_0 = 0$; $Z = Z_i (1 - \exp(-k_z t))$

which yield for Y , the total quantity of substrate remaining (Stf_i):

1. $Y = f_d + f_i$ for $t \leq t_i$; $S = f_d \exp(-k_d(t - t_i)) + f_i$ for $t > t_i$
2. $Y = f_d Z / (Z_0 \exp(Z_i t) + Z_i - Z_0) + f_i$
3. $Y = f_d \exp(Z_i(1 - k_z t - \exp(-k_z t)/k_z)) + f_i$

where t is the incubation time (h), f_d the potentially digestible fraction (g/g), f_i the indigestible fraction (g/g), k_d the fractional digestion rate constant (1/h), t_i the discrete lag time (h), Z_0 the initial value of Z (1/h), Z_i the asymptotic value of Z (1/h), and k_z the fractional rate of change of Z (1/h). The models were evaluated by fitting the equations to *in situ* data of 4 roughages.

Microbes involved in digestion of substrate are either synthesized within the system (*in situ* bag) or originate from the environment (rumen fluid surrounding *in situ* bags). Model 2 (Y is the logistic function) is based on the conversion of substrate into microbes (or enzymes), but does not account for exchange of microbes between the system and its environment. Model 3 assumes that the effect of microbes on digestion is time-dependent. Although it does not explicitly account for the origin of microbial mass, it has properties that cannot be included in model 1 (eg, $Z_0 = 0$).

The residual standard deviation (RSD) and estimates for f_d and f_i were generally similar between models. The Z_i (representing the fractional digestion rate after infinite incubation) for model 3 was of a similar magnitude to k_d in model 1. However, Z_i for model 2 was 35 to 95% higher than k_d . The t_i , Z_0 and k_z are difficult to compare between models because they are based on different concepts. Parameter estimates for orchard grass hay are given in table I as an example. Both models are suitable alternatives to the classical model.

Table I. Digestion kinetic parameters for orchard grass hay.

	Model 1	Model 2	Model 3
RSD (g/g)	0.023	0.024	0.021
f_d	0.597	0.615	0.604
f_i	0.227	0.237	0.230
k_d or Z_i	0.034	0.047	0.036
t_i	4.61	—	—
Z_0	—	0.023	—
k_z	—	—	0.198