



HAL
open science

Comparative study of the cellulolytic activity of caecum microbes in ponies and donkeys

F Faurie, JI Tisserand

► **To cite this version:**

F Faurie, JI Tisserand. Comparative study of the cellulolytic activity of caecum microbes in ponies and donkeys. *Annales de zootechnie*, 1994, 43 (3), pp.281-281. hal-00889020

HAL Id: hal-00889020

<https://hal.science/hal-00889020v1>

Submitted on 11 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Comparative study of the cellulolytic activity of caecum microbes in ponies and donkeys

F Faurie, JL Tisserand

Laboratoire Associé de Recherches Zootechniques INRA-ENSSAA, 26, bd Docteur-Petitjean, 21000 Dijon, France

In the context of comparative research on the digestive physiology of donkeys and ponies we have compared the *in sacco* degradation of fodder in the caeca of donkeys and ponies.

Three adult ponies and 3 adult donkeys with permanent caecum cannulas (225 and 248 kg live weight, respectively) were given a pelleted feed made up of wheat straw, maize grain and soya cake (crude protein CP in dry matter DM = 12.1%; NDF = 64.2%; ADF = 33.4%) at the maintenance level. After a 15 d adaptation period, the straw degradation in the caecum (CP = 3.5% DM; NDF = 77.1%; ADF = 46.5%; CF = 42.4%) was assessed by the nylon-bag technique (Miraglia *et al*, 1988) as follows: 1) degradation after 24 h; 2) degradation after 48 h; 3) degradation after 24 h in one species and 24 h in the other; with 3 repetitions per animal, *ie* 9 results per treatment.

Determination of the cell-wall content was according to Van Soest (1963).

Figure 1 shows that after 24 and 48 h the straw NDF and ADF DM degradation is higher in the donkeys than in the ponies. These results are statistically significant for a 48 h degradation period. The increase in DM, NDF and ADF degradation is higher in donkeys after a period of 24 h in ponies than in ponies after a 24 h period in donkeys.

These results show that the capacity of the caecum to degrade cell walls is higher in donkeys than in ponies.

Miraglia N, Martin-Rosset W, Tisserand JL (1988) *Ann Zootech* 37, 13-20

Van Soest PJ (1963) *J Am Off Agric Chim* 46, 829-835




Fig 1. *In sacco* degradation in the caecum of ponies (▨) and donkeys (□, 48 h top, 24 h bottom).