



HAL
open science

In vitro study of the dose effect of *Saccharomyces cerevisiae* on rumen digestion of a mixed diet

Jp Jouany, B Lassalas, G Bertin

► **To cite this version:**

Jp Jouany, B Lassalas, G Bertin. In vitro study of the dose effect of *Saccharomyces cerevisiae* on rumen digestion of a mixed diet. *Annales de zootechnie*, 1995, 44 (Suppl1), pp.155-155. <hal-00889317>

HAL Id: hal-00889317

<https://hal.science/hal-00889317v1>

Submitted on 11 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

In vitro study of the dose effect of *Saccharomyces cerevisiae* on rumen digestion of a mixed diet

JP Jouany ¹, B Lassalas ¹, G Bertin ²

¹INRA, SRNH, Theix, 63122 St-Genès-Champanelle ; ²CSI, Direction R and D, 94, rue Edouard Vaillant, 92532 Levallois-Perret, France

Live yeasts, used as feed additives, have been suggested as a mean of stabilizing rumen microbial digestion in animals fed diets rich in starch (RJ Wallace and J Newbold, 1992, Probiotics. The scientific basis, Chapman and Hall, 317-353). Three doses of *Saccharomyces cerevisiae* (I-1077)* (SC) were tested in Rusitec to study their effects on rumen digestive parameters. The four fermentors were first inoculated with mixed rumen digesta sampled from three sheep fed on grass hay (700 g/d), barley (400 g/d) and soybean meal (100 g/d). Each fermentor (1 l) was fed daily 10 g grass hay + 6 g barley + 2 g fishmeal + 1 g sucrose in a nylon bag 50 µm in mesh size, which was changed every 48 h. A Mc Dougall-type saliva with trace elements, urea (200 mg/d) and Na₂SO₄ (90.4 mg/d), was infused continuously at the rate of 700 ml/d. Nylon bags were vertically agitated at a rate of 40 cm/min. Yeast were given in dry form, once per day, when the fermentors were fed.

The addition of SC significantly improved the digestibility of OM and cell wall components

whatever the dose ; the increase was greatest at the dose of 10⁵ CFU. VFA production only increased with the 10⁵ CFU ; the composition of VFA mixture was not influenced by SC. As a consequence, fermented organic matter (FOM) calculated from DI Demeyer and CJ Van Nevel (1975, Digestion and Metabolism in the Ruminants, University of New England Publishing Co, 366-382) was highest with the 10⁵ CFU. Gas production and composition, NH₃-N concentration and pH in fermentors were not altered by SC, nor the efficiency of microbial protein synthesis (EMPS). Administered at the concentration of 10⁵ CFU/ml, SC had the highest effect on plant cell wall digestion and VFA production in Rusitec. Compared to results of *in vivo* studies, the response of microbial cellulolytic activities to SC in Rusitec was amplified. This suggests that Rusitec could not be suitable for probiotic studies.

(This work was supported by CSI Levallois-Perret, France. The authors thank M Fabre and P Journaix for their skilled technical assistance).

	Control	Yeast supply (CFU/ml)		
		10 ⁴	10 ⁵	10 ⁶
NDF digestibility (%)	36.4 ^a	42.8 ^b	47.5 ^c	46.1 ^c
VFA production (mMol/d)	63.6 ^a	58.4 ^a	76.8 ^b	69.9 ^{ab}
Acetate/propionate	2.4 ^a	2.5 ^a	2.5 ^a	2.3 ^a
FOM (g/d)	5.6 ^a	5.2 ^a	6.8 ^b	6.2 ^b
Gas production (l/d)	1.9 ^a	1.7 ^a	2.0 ^a	2.0 ^a
CO ₂ /CH ₄	4.5 ^a	3.9 ^a	3.8 ^a	4.1 ^a
EMPS (g N/kg FOM)	22.9 ^a	22.7 ^a	19.3 ^a	23.3 ^a

* registered at Institut Pasteur, Paris.