

Daily intake and the selection of feeding sites by horses in heterogeneous wet grasslands

Géraldine Fleurance, Patrick Duncan, Bruno Mallevaud

▶ To cite this version:

Géraldine Fleurance, Patrick Duncan, Bruno Mallevaud. Daily intake and the selection of feeding sites by horses in heterogeneous wet grasslands. Animal Research, 2001, 50 (2), pp.149-156. 10.1051/animres:2001123 . hal-00889847

HAL Id: hal-00889847 https://hal.science/hal-00889847

Submitted on 11 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Daily intake and the selection of feeding sites by horses in heterogeneous wet grasslands

Géraldine FLEURANCE^{a,b*}, Patrick DUNCAN^a, Bruno MALLEVAUD^c

 ^a Centre d'Études Biologiques de Chizé, CNRS-UPR 1934, 79360 Beauvoir-sur-Niort, France
 ^b Present address: Les Haras Nationaux, Direction du Développement, Station Expérimentale des Haras, 19370 Chamberet, France
 ^c Parc Interrégional du Marais Poitevin, place de l'Église, 79510 Coulon, France

(Received 10 January 2000; accepted 16 March 2001)

Abstract — In spite of the importance of grazed forage in horse nutrition, little information is available on their daily intake at pasture. We determined the intake of 4 non-breeding mares of a heavy breed (average body weight = 674 kg), grazing during the summer in heterogeneous natural grasslands of the Marais Poitevin (France), an internationally important wetland where grazing is an essential process which maintains biodiversity. The mares ate large quantities of forage (21.9 ± 2.4 kg of organic matter per day, i.e. 166.2 ± 20.8 g of organic matter per kg LW^{0.75} per day) in comparison with previous published values and with the estimated requirements of these horses. The use of the vegetation was very selective, the mares spent about 70% of their feeding time on short grass lawns (sward surface ≤ 4 cm, biomass < 100 g·m⁻²), that represented only 10% of the area. This behaviour maintained the plants at young growing stages which are of better quality than ungrazed plants. These results are discussed in relation to the dynamics of the plant communities.

horse / intake /grassland / heterogeneity / wetland

Résumé — **Quantités ingérées et sélection alimentaire réalisées par des chevaux en prairies humides hétérogènes.** Malgré l'importance de l'herbe dans l'alimentation du cheval, peu de données sont disponibles concernant les quantités ingérées au pâturage. En juillet-août 1997, nous avons déterminé les niveaux d'ingestion de 4 juments de race lourde à l'entretien (poids moyen : 674 kg) pâturant au sein de prairies naturelles hétérogènes du Marais Poitevin (France), un marais d'importance internationale où le pâturage est un processus essentiel au maintien de la biodiversité. Les juments ingéraient d'importantes quantités d'herbe (21,9 ± 2,4 kg de matière organique par jour, i.e. $166,2 \pm 20,8$ g de matière organique par kg PV^{0,75} par jour) en comparaison des résultats publiés antérieurement et des besoins estimés pour ces animaux. Les juments utilisaient la végétation de manière très sélective, passant environ 70 % de leur temps d'alimentation sur des zones d'herbes rases (hauteur ≤ 4 cm, biomasse < 100 g·m⁻²) qui couvraient 10 % de la surface totale. Ce comportement

^{*} Correspondence and reprints

Tel.: 33 (0)5 55 98 31 46; fax: 33 (0)5 55 73 47 70; e-mail: geraldine.fleurance@haras-nationaux.fr

maintenait les plantes dans de jeunes stades de croissance, de meilleure qualité que celle d'une végétation non pâturée. Les résultats sont discutés en rapport aux dynamiques des groupements végétaux présents sur le site.

cheval / ingestion / prairie / hétérogénéité / zone humide

1. INTRODUCTION

Intake rates by horses of green or dried forages distributed ad libitum are well documented [2, 4, 5, 18]. In spite of the importance of grazed forages in horse nutrition (these can represent more than 70% of the annual food, [17]), little information is available on intake at pasture [17], principally because in the field it is difficult to estimate faecal output and diet digestibility accurately. Rittenhouse et al. [24], using neutral detergent fiber (NDF) as an external marker for non productive and lactating mares in a natural grassland, measured intakes of 113 and 138 g DM·LW^{-0.75}·d⁻¹ respectively. Nonetheless, the reliability of this method, and of others based on external markers such as chromic oxid (e.g. in yearling horses, $80 \text{ g DM} \cdot LW^{-0.75} \cdot d^{-1}$ [14] is debatable [13, 15, 26]. Mesochina et al. [18], in a recent study, demonstrated that fecal crude protein is a good predictor of the digestibility of forages by horses (dry matter digestibility: rsd: 0.038; $r^2 = 0.65$; organic matter digestibility: rsd: 0.036, $r^2 = 0.74$). Duncan [8] used this approach to determine the intake of lactating mares of the Camargue breed (intake: $170 \text{ g } \text{DM} \cdot \text{LW}^{-0.75} \cdot \text{d}^{-1}$).

This study was part of a long term programme of research on the effects of different herbivores on the dynamics of the vegetation in a wetland of international importance, the Marais Poitevin. The site was in natural wet grasslands of the Communal des Magnils-Reigniers, which is of importance for the conservation of wetland plants (e.g. *Ranunculus ophioglossus*), birds (e.g. breeding *Tringa totanus*) and mammals (e.g. otters, *Lutra lutra*), and has traditionally been grazed by cattle. The experimental

set-up consisted of 10 plots of 1-2 ha which were grazed by cattle, horses or both species. Under cattle at densities comprised between 341 and 1246 kg·ha⁻¹, the plant community dynamics depended on the biomass density of the animals [1]: with high densities there are many rosette plants (hemicryptophytes), while at low animal densities these plants are replaced by competitive species (sensu Grime [12], mainly perennial graminoids). Horse grazing lead heterogeneous plant communities, as the two dynamics were observed whatever the density of the animals (345 to 900 kg·ha⁻¹). Tall grasses appear in patches of lightly grazed plants, where the animals deposit faeces. This leads to a loss of plant species diversity in the tall grass patches, but an increase in structural diversity. Elsewhere, in heavily grazed lawns, halophytes and hemicryptophytes appear.

The aim of this work was first to provide data on the rates of daily food intake in the horses on these pastures and then to contribute to this programme by describing the use of these grasslands by horses. We measured the time-budgets of the horses and their selective use of structural types (lawns and patches of tall grasses) and of the plant communities.

2. MATERIALS AND METHODS

2.1. Study site

The Magnils-Reigniers common is part of an area of brackish grasslands where the topography leads to a specific pattern of plant communities [1]. The shallow natural drainage channels, which are flooded from late autumn to early spring, are characterized by hygrophile vegetation with subaquatic species (Tab. I). The top of the catena is occupied by graminoids, forming the mesophile community. The slopes, whose soils are saline, are favourable for sub-halophile species, forming the mesohygrophile community.

We defined two structural vegetation types, which were represented in all communities, lawns (sward surface ≤ 4 cm) and taller grasses (mostly at the reproductive stage during the study) where the horses deposited all their faeces.

2.2. The animals

We studied 4 non-reproductive mares (*Mulassier Poitevin* breed), 2 to 7 years old. They were distributed in 2 sets, homogeneous for weight, each grazing a 2 ha pasture (E2D or E2) from April to October 1997. The mean weight of the animals was 674 kg during July and August.

2.3. Measurements

2.3.1. Daily food intake

The daily food intake was measured as OMI = F/(1-OMD) where OMI is the organic matter intake, F is the weight of faeces produced over 24 h (g) and OMD is the organic matter digestibility estimated by the crude protein (CP) content of the faeces [18]. The results are expressed in organic matter (OM) in order to avoid inaccuracies due to contamination by soil. Faecal output was measured by collecting the total amount of faeces produced 3 times a day over 4 successive days (29.VII.97 to 01.VIII.97) [15]. Faecal dry and organic matter content were measured in one sample per plot per day, dried at 80 °C to constant weight and by mineralization at 550 °C. Nitrogen was determined by the Kjeldahl method. The results were corrected for the mean weight of the animals, expressed as live or metabolic weight.

2.3.2. Spatio-temporal organization of the feeding behaviour

The mares were observed for a total of 48 h, between 21.VII-08.VIII.97. Individual behaviours were recorded using scan sampling during periods of 3 h by day and by night in order to cover the 24 h in 4 days. The 4 mares, previously accustomed to the presence of the observer in the pasture and to the use of a lamp, were checked successively by the same person at 5 minute intervals.

An animal was recorded in feeding activity if one of these actions was observed: searching (sorting or gripping food with the upper-lip), cutting, chewing or swallowing. The structural type and the community grazed were noted. In each plot, 100 measures of the sward surface were made (50 on each diagonal, with a measure every 4 m, on 21.VII.97, 28.VII.97, 06.VIII.97) using a 50×50 cm sward stick. At each measure, the structural type and the community were recorded. The mean of the 3 measurements enabled us to determine the area covered by each community and structural type in each plot, and to calculate for each plot a selection index S for the 2 mares of the pasture (the ratio of the percentage of the feeding time passed on the particular structural type or community, and the percentage of the area covered by that structural type or community). The null hypothesis was that the animals used the vegetation types evenly, i.e. S = 1.0.

2.3.3. Above-ground plant biomass, and forage quality

In each plot, samples (Tab. I) were clipped with shears in a rectangle $(50 \times 25 \text{ cm})$ placed at random in each structual type of each community (mesophile, meso-hygrophile, hygrophile). The samples were dried at 60 °C to constant weight to calculate the plant biomass (dry weight/area clipped). Some samples were analyzed for nitrogen (Kjeldahl method), cellulose (Weende

		Top of the catena	Slopes	Drainage channels				
		Mesophile community	Meso-hygrophile community	Hygrophile community				
		Hordeum secalinum,	Carex divisa,	Agrostis stolonifera,				
		Agrostis stolonifera,	Hordeum marinum,	Glyceria fluitans,				
		Elymus repens,	Juncus gerardii,	Alopecurus geniculatus,	р	Lawns	Tall grasses	р
		Cynosurus cristatus,	Hordeum secalinum,	Eleocharis palustris,				
		Bromus commutatus,	Alopecurus bulbosus	Oenanthe fistulosa				
		Lolium perenne	Agrostis stolonifera					
a) Biomass	E2D	199.4 ± 141.6 (<i>n</i> = 4)	$25.6 \pm 1.6 (n = 2)$	459.8 ± 271.8 (<i>n</i> = 4)	0.073	91.1 ± 71.4 (<i>n</i> = 6)	535.4 ± 196.7 (<i>n</i> = 4)	0.011
$(g DM \cdot m^{-2})$	E2	$293.0 \pm 116.7 \ (n = 4)$	$97.1 \pm 65.6 \ (n = 3)$	$319.2 \pm 272.2 \ (n = 4)$	0.176	$86.1 \pm 36.3 \ (n = 5)$	$403.2 \pm 200.2 \ (n = 5)$	0.018
	mean	$246.2 \pm 137.9 \ (n = 8)$	$68.5 \pm 61.7 \ (n = 5)$	$389.5 \pm 281.0 \ (n=8)$	0.023	$88.8 \pm 58.2 \ (n = 11)$	$461.9 \pm 209.2 \; (n=9)$	0.000
					M-MH: 0.028			
					MH-H: 0.013			
					M-H: 0.401			
b) CP (% DM)		$10.1 \pm 1.7 \ (n = 4)$	12.2 ± 1.9 (<i>n</i> = 2)	$10.7 \pm 2.1 \ (n = 10^1)$	0.596	$12.8 \pm 1.3 \ (n = 6)$	9.3 ± 1.2 (<i>n</i> = 9)	0.009
c) Cellulose (% DM)		$32.1 \pm 1.7 \ (n = 4)$	$27.1 \pm 2.3 (n = 2)$	$28.2 \pm 2.0 \ (n = 10^1)$	0.153	$26.7 \pm 2.2 \ (n = 6)$	30.8 ± 1.6 (<i>n</i> = 9)	0.056
NDF (% DM)		$70.5 \pm 2.1 \ (n = 4)$	$62.1 \pm 3.2 \ (n=2)$	$62.7 \pm 2.6 \ (n = 10^1)$	0.106	$62.1 \pm 2.8 \ (n = 6)$	$67.1 \pm 2.9 (n = 9)$	0.059

Table I. (a) Comparison of the mean aerial biomasses between communities and between lawns and patches of tall grasses; (b) and (c) comparison of crude protein (CP), cellulose and Neutral Detergent Fiber (NDF) contents between communities and between lawns and patches of tall grass, across both plots.

 $\frac{1}{18}$ samples from biomass measurements + 2 at random; DM: dry matter.

method) and cell walls: Neutral Detergent Fiber (NDF) [28].

2.4. Statistics

The significance of selection for vegetation types was determined by chi-square tests, by comparing the numbers of feeding observations with the values expected if the animals used the vegetation types evenly. Kruskal-Wallis and Mann-Whitney tests were used to compare plant biomass and grass quality respectively, between plant communities (across structural types) and between structural types (across plant communities).

Statistical tests were bilateral, for p < 0.05 using Statistica [25].

3. RESULTS

3.1. Daily food intake at pasture

The mares in both pastures selected diet of good quality (estimated organic matter digestibility: $61.8 \pm 0.8\%$) with little variability. Intake was on average 21.9 kg OM·kg LW^{-0.75}·d⁻¹, i.e. 3.3% of liveweight, i.e.

166.2 \pm 20.8 g OM·kg LW^{-0.75}·d⁻¹. The mean value in terms of dry matter was 172.5 \pm 23.1g DM·kg LW^{-0.75}·d⁻¹. The digestible organic matter intake (nutrient assimilation) was 102.8 \pm 13.8 g DOM·kg LW^{-0.75}·d⁻¹.

3.2. Spatio-temporal organization of the feeding behaviour

The mares fed on average $60 \pm 1\%$ of the 24 h. This activity occupied about ${}^{3/_{4}}$ of the night $(74 \pm 4\%)$ and half of the daylight $(54 \pm 1\%)$. The use of vegetation communities was not homogeneous (chi-square test, p < 0.01). The mares selected the meso-hygrophile community, using this 2.5 times as much as expected from the hypothesis of even use and they avoided the mesophile community (Tab. II).

At the time of the study, the structural heterogeneity of the cover was not very pronounced in the pasture E2, with small patches of lawns beginning to appear. The use of the structural types by the mares was measured only in the E2D pasture. Here the mares fed on both the structural types, but the use of the lawns was 7 times as much as expected (chi-square test, p < 0.01), their

 Table II. Use of the plant communities, lawns and patches of tall grasses and the values of the selection index.

Community	Pasture	% of the feeding time	% of the total area	Selection index
Top of the catena Mesophile community	E2 E2D mean	$8.1 \pm 1.2 (n = 2) 17.8 \pm 0.3 (n = 2)$	39.1 59.8	0.21 0.30 0.26
Slopes Meso-hygrophile community	E2 E2D mean	$30.6 \pm 0.8 \ (n = 2) \\ 8.6 \pm 0.8 \ (n = 2)$	10.1 5.4	3.03 1.59 2.53
Drainage channels Hygrophile community	E2 E2D mean	$\begin{array}{l} 61.3 \pm 0.0 \; (n=2) \\ 73.6 \pm 0.7 \; (n=2) \end{array}$	50.8 34.8	1.21 2.11 1.58
Structural type Lawns Tall grass	E2D E2D	$69.5 \pm 2.8 \ (n = 2)$ $30.5 \pm 2.8 \ (n = 2)$	9.8 90.2	7.09 0.34

selectivity index being 20 times that on the taller grass. The preference for the lawns is verified in each community (M, MH, H), where the percentage of the area covered by the lawns were respectively: 9.4%, 37.5%, 10.7% and the selection index were 6.24, 2.58, 7.15 for the lawns and 0.36, 0.05, 0.36 for the tall grasses (chi-square test, p < 0.01).

3.3. Aerial biomass and vegetation quality

The meso-hygrophile community, which was selected by the mares, had the least biomass (p < 0.05) (Tab. I), but the difference was not significant within pastures because the variances are high. The mean biomass of the patches of tall grasses was 5 times higher than the biomass of the lawns (p < 0.01, Tab. I).

There were no significant difference between the plant communities for crude protein, cellulose and NDF contents (Tab. I). Nonetheless, the lawns which were strongly selected by the mares were characterized by a crude protein content 38% higher than the tall grasses. The cellulose and NDF contents tended to be lower for lawns (p < 0.06) (Tab. I).

4. DISCUSSION

4.1. Daily intake

The daily food intake found here $(172.5 \pm 23.1 \text{ g DM} \cdot \text{kg LW}^{-0.75} \cdot \text{d}^{-1}$ or $166.2 \pm 20.8 \text{ g}$ OM $\cdot \text{kg LW}^{-0.75} \cdot \text{d}^{-1}$) is much higher than the few values available for non-productive horses at pasture (e.g. 113 g DM \cdot kg LW $^{-0.75} \cdot \text{d}^{-1}$ from Rittenhouse et al. [24]) and higher than those of Mesochina [19] and Ellis et Lawrence [10] for growing saddle horses (92 g OM \cdot kg LW $^{-0.75} \cdot \text{d}^{-1}$ and 80 g DM \cdot kg LW $^{-0.75} \cdot \text{d}^{-1}$). The values obtained in the Marais Poitevin are comparable to the values for breeding and growing wild equid species [2, 11, 21] and for lactating mares (e.g. in the Camargue, France [8]). Nonetheless, even if the needs for production seem to be the main determinant of food intake [2], Meyer [20] has shown that horses can have intakes higher than their needs, and maintain these for a long time. These intakes are also much higher than those of comparable beef cattle at pasture (3.3% of liveweight in horses compared with 2–3% in cattle (Dulphy et al. [7]).

In view of the fact that there are considerable inter-individual variations in intake in some herbivores [6, 23], and the small number of animals used in this study, the measures should be repeated. Nonetheless the slight variations between days on food intake (coefficient of variation: 12%) show that data from four days allow estimation of daily intake for horses on these pastures with reasonable accuracy. An obvious improvement would be to individualise the faeces with colored plastic marks.

4.2. Duration of feeding activity

The time spent feeding, $60 \pm 1\%$ (or 14 h 30) is consistent with previous published values (14 h–17 h, [8]) and the animals in the experimental plots appeared to have normal feeding behaviour. Nocturnal feeding comprised 38% of the total time spent feeding, which is typical for horses (20 to 50%, [16]), which is why it was necessary to conduct 24 h observations in this study of their feeding strategies.

4.3. Selection of vegetation types

The horses showed more marked selection for lawns than for any of the vegetation communities, as has been observed previously in dry grasslands [3, 9, 22]. The mares maintained lawns where the quality of the grass was high in comparison with the remainder of the pasture (38% higher crude protein, lower fibre). In this pasture, the horses choose their food plants principally on the basis of their structure (growth stage) rather than their botanical species.

In spite of the short period covered by this study, the feeding strategies of the mares described here clarify causes of the differences in the vegetation dynamics under cattle or horse grazing described by Amiaud [1]. The intense use of the meso-hygrophile community by horses explains the rejuvenation of sub-halophyte communities and the development of hemicryptophytes on the slopes. In the same way, the opening of the hygrophile community leads to the emergence of annual species (Galium debile, Ranunculus sardous) [1]. The development of highly competitive species at the top of the catena seems to be the result of low grazing pressure and perhaps favourable conditions of mineral and nitrogen availability brought by the faeces.

Finally, the high intake levels compared with cattle and strong selection for short grass lawns explain why the horses create and maintain structural diversity in such grasslands, whatever their stocking level.

The current hypothesis to explain the heterogeneous use of grasslands by horses proposes that the animals minimize their feeding time on patches of tall grasses where the faeces are deposited in order to minimize the intake of gastro-intestinal parasites [27]. The results of this study suggest another hypothesis: if the feeding strategy of these horses is to maximise daily nutrient assimilation, the maintenance of lawns of good quality could be a nutritional as well as an antiparasite tactic.

ACKNOWLEDGEMENTS

This work was funded by the Parc Interrégional du Marais Poitevin. We wish to thank D. Naudon for his support of the project and for his help in the collection of the data. We are also grateful to J.B. Bouzillé (CNRS-University of Rennes I UMR 5558) for including us in this study.

REFERENCES

- Amiaud B., Dynamique végétale d'un écosystème prairial soumis à différentes modalités de pâturage, exemple des communaux du marais Poitevin, thèse, Université de Rennes I, France, 1998, 317 p. + annexes.
- [2] Boulot S., L'ingestion chez la jument. Étude au cours du cycle gestation-lactation : implications nutritionnelles et métaboliques, thèse, ENSA Rennes, France, 1987, 147 p.
- [3] Carson K., Wood–Gush D.G.M., Equine behaviour: II. A review of the litterature on feeding, eliminative and resting behaviour, Appl. Anim. Ethol. 10 (1983) 179–183.
- [4] Chenost M., Martin-Rosset W., Comparaison entre espèces (mouton, cheval, bovin) de la digestibilité et des quantités ingérées des fourrages verts, Ann. Zootech. 34 (1985) 291–312.
- [5] Cymbaluk N.F., Comparison of forage digestion by cattle and horses, Can. J. Anim. Sci. 70 (1990) 601–610.
- [6] Doreau M., Boulot S., Martin-Rosset W., Effect of parity and physiological state on intake, milk production and blood parameters in lactating mares differing in body size, Anim. Prod. 53 (1991) 111–118.
- [7] Dulphy J.P., Jouany J., Martin-Rosset W., Thériez M., Aptitudes comparées de différentes espèces d'herbivores domestiques à ingérer et digérer des fourrages distribués à l'auge, Ann. Zootech. 43 (1994) 11–32.
- [8] Duncan P., Horses and Grasses. The nutritional ecology of equids and their impact on the Camargue, Springer-Verlag, NY, 1992, 287 p.
- [9] Edwards P.J., Hollis S., The distribution of excreta on New Forest grassland used by cattle, ponies and deer, J. Appl. Ecol. 19 (1982) 953–964.
- [10] Ellis R.N.W., Lawrence L.J., Energy under nutrition in the weanling filly foal. III. Effects on heart rate and subsequent voluntary food intake, Brit. Vet. J. 134 (1978) 333–341.
- [11] Gakahu C.G., Feeding strategies of the Plains zebra, *Equus quagga bohmi*, in the Amboseli ecosystem, Ph.D. thesis, University of Nairobi, Kenya, 1982.
- [12] Grime J.P., Plant strategies and vegetation processes, Wiley and Sons, Chichester, 1979, 222 p.
- [13] Haenlein G.F.W., Smith R.C., Yoon M., Determination of the fecal excretion rate of horses with chromic oxide, J. Anim. Sci. 25 (1966) 1091–1095.
- [14] Lawrence L.M., Moore K.J., Hintz H.F., Jaster E.H., Wischover L., Acceptability of alfafa hay treated with an organic acid preservative for horses, Can. J. Anim. Sci. 67 (1987) 217–220.
- [15] Le Du Y.L.P., Penning P.D., Animal based techniques for estimating herbage intake, in: Leaver J.D. (Ed.), Herbage intake handbook, Brit. Grassld. Soc., 1982, pp. 37–75.

- [16] Martin-Rosset W., Doreau M., Consommation d'aliments et d'eau par le cheval, in: Jarrige R., Martin-Rosset W. (Ed.), Le Cheval: Reproduction, Sélection, Alimentation, Exploitation, INRA Publications, Paris, France, 1984, pp. 333–354.
- [17] Martin-Rosset W., Trillaud-Geyl C., Jussiaux M., Agabriel J., Loiseau P., Béranger C., Exploitation du pâturage par le cheval en croissance ou à l'engrais, in: Jarrige R., Martin-Rosset W. (Ed.), Le Cheval: Reproduction, Sélection, Alimentation, Exploitation, INRA Publications, Paris, France, 1984, pp. 583–599.
- [18] Mésochina P., Martin–Rosset W., Peyraud J.L., Duncan P., Micol D., Boulot S., Prediction of the digestibility of the diet of horses: evaluation of faecal indices, Grass Forage Sci. 53 (1998) 189–196.
- [19] Mésochina P., Niveau d'ingestion du cheval en croissance au pâturage : mise au point méthodologique et étude de quelques facteurs de variation, thèse, Institut National Agronomique Paris Grignon, France, 2000, 158 p.
- [20] Meyer H., Ein Beitrag zur Regulation der Futteraufnahme beim Pferd, Dtsch. Tierarztl.Wschr. 87 (1980) 404–408.
- [21] Ngethe J.C., Preference and daily intake of five East African grasses by zebra, J. Range Manage. 29 (1976) 510–511.

- [22] Odberg F.O., Francis-Smith K., Studies on the formation of ungrazed eliminative areas in fields used by horses, Appl. Anim. Ethol. 3 (1977) 27–34.
- [23] Ralston S.L., van den Broek G., Baile C.A., Feed intake patterns and associated blood glucose, free fatty acid and insuline changes in ponies, J. Anim. Sci. 49 (1979) 838–845.
- [24] Rittenhouse L.R., Johnson D.E., Borman M.M., A study of food consumption rates and nutrition of horses and cattle, Bureau of Land Management, US, Department of Interior, Washington, 1982, 116 p.
- [25] Statsoft, 2nd ed., Statistica for Windows: General conventions and statistics I, Vol. 1, StatSoft Inc., Tulsa, 1995.
- [26] Sutton E.I., Bowland O.P., Mac Carthy O.F., Studies with horses comparing 4N-HCL insoluble ash as an index material with total faecal collection in the determination of apparent digestibilities, Can. J. Anim Sci. 57 (1977) 543–549.
- [27] Taylor E.L., Grazing behaviour and helmintic disease, Brit. J. Anim. Behav. 2 (1954) 61–62.
- [28] Van Soest P.J., Wine R.H., Use of detergents in the analysis of fibrous feeds. IV. Determination of plant cell wall constituents, J. Assoc. Off. Agric. Chem. 50 (1967) 50–55.

To access this journal online: www.edpsciences.org

156