



# **Numbers of spermatozoa in queens and drones indicate multiple mating of queens in *Apis andreniformis* and *Apis dorsata***

G Koeniger, N Koeniger, M Mardan, Rwk Punchihewa, Gw Otis

## **► To cite this version:**

G Koeniger, N Koeniger, M Mardan, Rwk Punchihewa, Gw Otis. Numbers of spermatozoa in queens and drones indicate multiple mating of queens in *Apis andreniformis* and *Apis dorsata*. *Apidologie*, 1990, 21 (4), pp.281-286. <hal-00890836>

**HAL Id: hal-00890836**

**<https://hal.science/hal-00890836v1>**

Submitted on 11 May 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

## Numbers of spermatozoa in queens and drones indicate multiple mating of queens in *Apis andreniformis* and *Apis dorsata*

G Koeniger <sup>1\*</sup>, N Koeniger <sup>1</sup>, M Mardan <sup>2</sup>,  
RWK Punchihewa <sup>3</sup>, GW Otis <sup>2</sup>

<sup>1</sup> Institut für Bienenkunde, Polytechnische Gesellschaft; FB Biologie  
an der JW Goethe-Universität, Frankfurt, K-v-Frisch-Weg 2, D-6370 Oberursel, FRG;

<sup>2</sup> Universiti Pertanian Malaysia, Department of Plant Protection,  
43400 Serdang, Selangor, Malaysia;

<sup>3</sup> Sri Lanka Department of Agriculture, Agriculture Research Station,  
Makandura, Gonawila (NWP), Sri Lanka

(Received 24 November 1989; accepted 3 May 1990)

**Summary** — Drones of *Apis dorsata* had an average of  $2.46 \cdot 10^6$  spermatozoa in their vesiculae seminales. Two queens had  $3.67 \cdot 10^6$  spermatozoa in their spermathecae. In *A. andreniformis*, drones had an average of  $0.13 \cdot 10^6$  and the spermathecae of 2 queens contained 0.98 and  $1.09 \cdot 10^6$  spermatozoa. In both *A. dorsata* and *A. andreniformis* the spermathecae of queens contained more spermatozoa than the vesiculae seminales of a single drone of either species. Therefore, we conclude that multiple mating occurs in both species as is the case for *A. mellifera*, *A. cerana* and *A. florea*.

multiple mating / spermatozoon / *Apis andreniformis* / *Apis dorsata*

### INTRODUCTION

In the past, sperm counts in drones and queens have provided valuable information on mating behavior. The high numbers of spermatozoa in the oviducts of freshly mated queens relative to the numbers in mature drones indicate that queens of *A. mellifera* and *A. cerana* are mated by an average of 8–10 drones on 1 mating flight (Tryasko, 1956; Woyke, 1960, 1975).

The numbers of spermatozoa in males and in spermathecae of queens have

barely been studied in other species. Recently, in *A. florea*, sperm counts of freshly mated queens revealed that, in this species, sperm is not injected into the oviducts but directly into the spermatheca. Drones only have about 1/3 as many spermatozoa in the vesiculae seminales as are contained in the spermatheca of the queen, suggesting that queens mate with 2–3 males (Koeniger *et al.*, 1989).

In this paper we report numbers of spermatozoa of queens and drones of *A. andreniformis* and *A. dorsata*. Furthermore, the size of the spermathecae was

\* Correspondence and reprints

measured and the number of ovarioles was counted.

## MATERIAL AND METHODS

Samples of *A andreniformis* were collected in the Johore state, Malaysia. *A dorsata* drones were collected in the Selangor state, Malaysia and queens in the Anuradhapura district, Sri Lanka.

Drones were kept for some hours without food until they were too weak to empty the vesiculae seminales when killed by ether. Afterwards the drones were dissected and the vesiculae seminales were isolated. Drones were considered to be sexually mature when the testis had become degenerate and when the cornua had an orange color.

Queens were dissected and the spermatheca isolated. After removal of the tracheal net, the diameter of the globular spermatheca was measured under a microscope.

In each case, the organs were transferred into 1 ml 0.9% NaCl in order to count the spermatozoa. The organ was ruptured with fine forceps and the emerging spermatozoa were dispersed evenly in the fluid by heavily blowing air through it with a syringe. If the sperm concentration was high (opaque fluid), the sample was further diluted with up to 9 ml of distilled water.

A haemocytometer (Fuchs-Rosenthal or the "improved Neubauer") was used to count the sperm.

The fresh ovaries were isolated and divided into 4 portions; the number of ovarioles counted ranged between 10 to 30 per portion. Then 70% ethanol was poured over it; thus, the ovarioles became white and were easy to count.

## RESULTS

### *Vesiculae seminales*

In drones of *A andreniformis* ( $n = 5$ ), the average number of spermatozoa in the vesiculae seminales was  $0.13 \pm 0.01 \cdot 10^6$ . In

drones of *A dorsata* ( $n = 5$ ) it was  $2.46 \pm 0.15 \cdot 10^6$  spermatozoa (table I).

### *Spermatheca*

All spermathecae were surrounded by a dense tracheal net and had a spermathecal gland similar in shape to that of *A mellifera*. The diameter of the spermathecae in both queens of *A andreniformis* was 0.8 mm. Because the spermathecal wall, as in all *Apis* species, is very thin, the volume of the spermatheca can be calculated directly from the diameter. It is  $0.27 \text{ mm}^3$  in *A andreniformis*. The numbers of spermatozoa were  $0.98$  and  $1.09 \cdot 10^6$ . The diameter of the spermathecae of 2 *A dorsata* queens was 1.1 mm (volume:  $0.7 \text{ mm}^3$ ). The sperm number was  $3.69$  and  $3.64 \cdot 10^6$  (table II).

### *Ovarioles*

Two queens of *A andreniformis* had 100 and 104 ovarioles (50 and 52 per ovary respectively), the 2 *A dorsata* queens had 230 and 236 ovarioles.

**Table I.** Sperm numbers of vesiculae seminales ( $\times 10^6$ ).

\* (Koeniger *et al*, 1989)

<i>A andreniformis</i>	<i>A florea</i> *	<i>A dorsata</i>
0.12	0.45	2.24
0.13	0.46	2.56
0.14	0.40	2.46
0.12	0.39	2.44
0.13	0.48	2.64
	0.45	
	0.41	
$\bar{x} = 0.13$	$\bar{x} = 0.43$	$\bar{x} = 2.46$
$s = \pm 0.01$	$s = \pm 0.03$	$s = \pm 0.15$

**Table II.** Data of spermathecae of *A andreniformis*, *A florea*, *A dorsata*, *A mellifera* and *A cerana*.  
 \* (Koeniger *et al*, 1989), \*\* (Woyke, 1966), \*\*\* (Woyke, 1975).

	Diameter (mm)	Volume (mm <sup>3</sup> )	No sperm (10 <sup>6</sup> )	Concentr (10 <sup>6</sup> /mm <sup>3</sup> )
<i>A a</i>	0.8	0.27	1.09	4.03
	0.8	0.27	0.98	3.63
<i>A d</i>	1.1	0.70	3.64	5.20
	1.1	0.70	3.69	5.27
<i>A f</i> *	0.8	0.27	1.35	5.00
	0.8	0.27	1.20	4.44
	0.8	0.27	1.20	4.44
	0.8	0.27	1.17	4.33
	0.75	0.22	0.85	3.86
<i>A m</i> **	1.14	0.78	4.73	6.12
<i>A c</i> ***	0.98	0.48	1.35	2.79

## DISCUSSION

In the species *A dorsata* and *A andreniformis*, the numbers of spermatozoa in the spermatheca exceeded those of the seminal vesicles of the respective drones. This verifies multiple mating for both species. Now multiple mating has been found in all known *Apis* species except *A koschevnikovi* where no data are yet available. Thus, polyandry appears to be a general feature of the *Apis* genus.

In *A mellifera* and *A cerana*, the number of matings cannot be deduced from the relation of the number of spermatozoa in the vesiculæ seminales and in the spermatheca, because, in these species, drones inject the sperm into the oviducts while finally in the spermatheca less than 5% of the spermatozoa received during mating is stored.

In contrast to these results, no spermatozoa were found in the oviducts of freshly mated *A florea* queens; they were

found only in the spermatheca (Koeniger *et al*, 1989). The endophallus does not end in a bulb as in *A mellifera* and *A cerana* but in a thin tip (Simpson, 1960; Ruttner, 1988), which seems to be an adaptation for direct sperm transfer into the spermatheca.

In *A andreniformis* the endophallus has a similar shape as in *A florea* and also ends in a thin tip (Wongsiri *et al*, 1990). Supposing that, in *A andreniformis*, the endophallus also enables direct sperm transfer, the number of matings would be 7–8. If the spermatozoa first reach the oviducts and the queen discards a lot of it, as in *A cerana* and *A mellifera*, a much higher number of matings can be assumed.

In *Apis dorsata* the size of the bulb is reduced and the cervix elongated compared to *A mellifera* and *A cerana* (Simpson, 1970; McEvoy and Underwood, 1988). There is no evidence to suggest that sperm is first injected into the oviducts or if a direct sperm transfer into the spermathe-

ca exists. In the case of direct sperm transfer, 2 drones are required to fill the spermatheca. If sperm transfer occurs as in *A cerana*, the number of matings would probably be much higher.

Sperm numbers in drones of *A florea* and *A andreniformis* are significantly different (table I; *t*-test,  $P > 0.001$ ). This result also supports the view that these should be considered as different species.

The high number of ovarioles in queens of *A dorsata* suggests that she produces more eggs than the other 2 species (230 ovarioles compared with about 100). In queens from the Philippines, even more ovarioles per bee (260) were reported (Velthuis *et al*, 1971). Also the number of  $3.7 \cdot 10^6$  spermatozoa, which is 3 times higher than in *A florea* and *A andreniformis*, supports this idea.

The concentrations of spermatozoa within the spermatheca appear to be similar in all *Apis* species, on average between 3 and  $6 \cdot 10^6/\text{mm}^3$  (table II). The lowest value was found in *A cerana* (Woyke, 1975) with an average of  $2.79 \cdot 10^6$  spermatozoa per  $1 \text{ mm}^3$  (13 queens) and highest in *A mellifera* (Woyke, 1966) with  $6.12 \cdot 10^6$  spermatozoa per  $1 \text{ mm}^3$  on the average (126 queens).

The anatomy of the spermathecae is similar in all *Apis* species. They all are surrounded by a dense tracheal net and have a spermathecal gland as described for *A mellifera* by Snodgrass (1956). This may indicate that the mechanism of sperm storage and utilisation is similar in all species.

## ACKNOWLEDGMENTS

We thank NA Zainal and RC Hindurangala for their valuable assistance in locating and catching the wild colonies.

GW Otis received a Research Leave Grant from the University of Guelph, and funds from

the International Development Research Centre of Canada.

**Résumé — Le nombre de spermatozoïdes chez les reines et les mâles, preuve d'accouplement multiple des reines chez *Apis andreniformis* et *Apis dorsata*.** Chez *Apis dorsata*, nous avons trouvé une moyenne de  $2,46 \cdot 10^6$  spermatozoïdes dans les vésicules séminales des mâles et  $3,67 \cdot 10^6$  de spermatozoïdes dans la spermathèque de 2 reines; chez les mâles d'*Apis andreniformis*,  $0,13 \cdot 10^6$  de spermatozoïdes et chez les 2 reines examinées 0,98 et  $1,09 \cdot 10^6$  (tableaux I et II). Le nombre de spermatozoïdes contenu dans la spermathèque des reines des 2 espèces est donc supérieur à celui produit par un seul mâle de chaque espèce. Ceci n'est possible que si les reines s'accouplent avec plusieurs mâles. Les reines d'*A dorsata* doivent s'accoupler avec au moins 2 mâles. Et si, lors de l'accouplement, comme c'est le cas chez *A mellifica* et *A cerana*, le sperme passe dans les oviductes et que seule une petite partie atteint ensuite la spermathèque, le nombre d'accouplement doit être sensiblement plus élevé.

Chez *A florea*, les spermatozoïdes passent directement dans la spermathèque. L'endophallus se termine par une pointe, et non par un bulbe, ce qui est considéré comme une adaptation au transfert direct (Koeniger *et al*, 1989). L'endophallus d'*A andreniformis* se termine également par une pointe (Wongsiri *et al*, 1990). C'est pourquoi nous admettons un transfert direct dans la spermathèque chez *A andreniformis* également. Dans ce cas les reines doivent s'accoupler avec 7–8 mâles. Nous avons donc une preuve de l'accouplement multiple des reines chez toutes les espèces récentes d'*Apis*, à l'exception d'*A koschevnikovi*, chez laquelle aucun comptage de spermatozoïdes n'a été effectué.

Les reines d'*A dorsata* renferment dans leur spermathèque presque 4 fois plus de spermatozoïdes que celles d'*A andreniformis*. Les concentrations par mm<sup>3</sup> sont par contre semblables ( $5,2 \cdot 10^6$  et  $3,8 \cdot 10^6$ , tableau II). La structure de la spermathèque est semblable chez toutes les espèces; c'est la taille qui diffère. Les spermathèques sont toutes entourées d'un réseau dense de trachées et possèdent toutes une glande spermathéciale bifide (Snodgrass, 1956). Il en résulte vraisemblablement un stockage et une utilisation des spermatozoïdes pour la fécondation des œufs qui sont semblables chez toutes les espèces d'*Apis*.

#### **Accouplement multiple / spermatozoïde / *Apis andreniformis* / *Apis dorsata***

**Zusammenfassung — Die Anzahl Spermatozoen in Königinnen und Drohnen beweist eine Mehrfachpaarung der Königinnen bei *Apis andreniformis* und *Apis dorsata*.** Bei Drohnen von *Apis dorsata* fanden wir im Durchschnitt  $2,46 \cdot 10^6$  Spermatozoen in den Vesiculae seminales; zwei Königinnen hatten je  $3,67 \cdot 10^6$  Spermatozoen in ihren Spermatheken. Bei Drohnen von *Apis andreniformis* wurden  $0,13 \cdot 10^6$  Spermatozoen gezählt, bei den beiden Königinnen 0,98 und  $1,09 \cdot 10^6$  (Tabelle I, und II). Die Spermatheken der beiden Honigbienenarten enthielten demnach mehr Spermatozoen, als ein einzelner Drohn der jeweiligen Art produziert. Das ist nur möglich, wenn sich die Königinnen mit mehreren Drohnen paaren. Die *A dorsata*-Königinnen müssen sich mit mindestens 2 Drohnen gepaart haben. Sollten jedoch die Spermien bei der Paarung wie bei *A mellifera* und *A cerana* zunächst in die Ovidukte übertragen werden, von denen nur ein kleiner Teil in die Spermatheka gelangt, muß die Anzahl der Paarungen wesentlich höher sein.

Bei *A florea* werden die Spermatozoen direkt in die Spermatheka übertragen. Der spitz auslaufende Bulbus des Endophallus wird als eine Anpassung an diese Funktion gedeutet (Koeniger *et al*, 1989). Der Endophallus von *A adreniformis* endet ebenfalls mit einer Spitze (Wongsiri *et al*, 1990). Deshalb nehmen wir auch bei *A andreniformis* eine direkte Spermaübertragung in die Spermatheka an. In diesem Fall sind die Königinnen von 7–8 Drohnen begattet worden.

Damit ist für alle rezenten *Apis* Arten – mit Ausnahme von *A koschevnikovi*, über die bisher keine Spermazählungen vorliegen – eine Mehrfachpaarung der Königinnen nachgewiesen. Die *A dorsata*-Königinnen enthalten fast viermal so viel Spermatozoen in den Spermatheken wie die Königinnen von *A andreniformis*. Die Konzentrationen pro mm<sup>3</sup> dagegen sind ähnlich ( $5,2 \cdot 10^6$  bzw  $3,8 \cdot 10^6$ , Tabelle II). Der Bau der Spermatheka differiert vor allem in der Größe, sonst sind sie ähnlich wie in *A mellifera* von einem dichten Tracheennetz umhüllt und haben eine zweiarmlige Spermathekaldrüse (Snodgrass, 1956). Wahrscheinlich erfolgt die Speicherung der Spermatozoen und deren Aktivierung zur Eibefruchtung in allen *Apis* Arten nach einem ähnlichen Prinzip.

#### **Mehrfachpaarung / Spermatozoon / *Apis andreniformis* / *Apis dorsata***

#### **REFERENCES**

- Koeniger N, Koeniger G, Wongsiri S (1989) Mating and sperm transfer in *Apis florea*. *Apidologie* 20, 413-418
- McEvoy M, Underwood BA (1988) The drone and species status of the Himalayan honey bee, *Apis laboriosa* (Hymenoptera, Apidae). *J Kans Entomol Soc* 61, 246-249
- Ruttner F (1988) *Biogeography and Taxonomy of Honeybees*. Springer Verlag Berlin, 85-86

- Simpson J (1960) Male genitalia of *Apis* species. *Nature* 185, 56
- Simpson J (1970) The male genitalia of *Apis dorsata* (F) (Hymenoptera, Apidae). *Proc R Entomol Soc Lond A* 45, 169-171
- Snodgrass RE (1956) *Anatomy of the honey bee*. Cornell University Press Ltd, Ithaca and London, 305 p
- Tryasko VV (1956) Multiple mating of queen bees. *Pchelovodstvo* 33, 50-54 (in Russian)
- Velthuis HHW, Clement J, Morse RA, Laigo FM (1971) The ovaries of *Apis dorsata* workers and queens from the Philippines. *J Apic Res* 10, 63-66
- Wongsiri S, Limbipichai K, Tangkanasing P, Mardan M, Rinderer T, Sylvester HA, Koeniger G, Otis G (1990) Evidence of reproductive isolation confirms that *Apis andreniformis* (Smith 1858) is a separate species from sympatric *Apis florea* (Fabricius 1787). *Apidologie* 21, 47-52
- Woyke J (1960) Natural and artificial insemination of queen honeybees. *Pszczel Zesz Nauk* 4, 183-275 (English summary in *Bee World*, 1962, 43, 21-25)
- Woyke J (1966) Wovon hängt die Zahl der Spermien in der Samenblase der auf natürlichem Wege begatteten Königin ab? *Z Bienenforsch* 8, 236-247
- Woyke J (1975) Natural and artificial insemination of *Apis cerana* in India. *J Apic Res* 14, 153-159