



HAL
open science

A scientific note on the ITS-1 region of *Apis mellifera* subspecies

Pilar de La Rúa, Stefan Fuchs, José Serrano

► **To cite this version:**

Pilar de La Rúa, Stefan Fuchs, José Serrano. A scientific note on the ITS-1 region of *Apis mellifera* subspecies. *Apidologie*, 2007, 38 (4), pp.378-379. hal-00892264

HAL Id: hal-00892264

<https://hal.science/hal-00892264>

Submitted on 11 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A scientific note on the ITS-1 region of *Apis mellifera* subspecies*

Pilar De La RÚA^a, Stefan FUCHS^b, José SERRANO^a

^a Área de Biología Animal, Facultad de Veterinaria, Apdo. 4021, Universidad de Murcia, 30100 Murcia, Spain

^b Institut fuer Bienenkunde, J.W. Goethe-Universitaet Frankfurt-am-Main, Karl von Frisch-Weg 2, 61440 Oberursel, Germany

Received 9 March 2006 – Revised 9 February 2007 – Accepted 5 March 2007

Apis mellifera / honeybee subspecies / ITS-1 region / DNA sequence / molecular markers

Several molecular markers have been widely used in insect molecular systematics in the last 20 years; among them nuclear DNA markers as the internal transcribed spacers (ITS1 and ITS2) of the nuclear rDNA (revision in Caterino et al., 2000). The sequence of these regions, which lie between functional 18S, 5.8S and 28S ribosomal genes, has proven suitable for phylogenetic studies at the species and intraspecific levels due to the fact that they are relatively free from selective constraints and correspondingly accumulate sequence differences rapidly. The sequence variation of this marker has been characterized in several genera within the order Hymenoptera as in *Ageniapis* (Álvarez and Hoy, 2002) and in *Tetranychus* (Navajas et al., 1998) and also it has been applied to phylogenetic analysis in *Melipona* species (Fernandes-Salomão et al., 2005). Although the size of the ITS-1 region was determined by Sheppard and McPherson (1991) to 132 bp in a sample of *Apis mellifera* from Argentina, nothing is published regarding sequence characteristics. Therefore, we characterized the sequence of the internal transcribed region (ITS-1) and evaluated its potential usefulness for phylogenetic and population studies among subspecies of *A. mellifera*.

The ITS-1 region was PCR-amplified (Ji et al., 2003) in ten subspecies of *Apis mellifera* (GenBank accession numbers DQ195225–DQ195236) (see material online at <http://www.apidologie.org>). It had a size of 132 bp, one of the smallest ITS-1 regions known so far in bees. In other Apidae this region varies from 1387–1417 bp in *Melipona*

subspecies (Fernandes-Salomão et al., 2005) to 289 bp in *Bombus lapidarius* (Ji et al., 2003). The G+C content (21.6%) was similar to that found in *Drosophila melanogaster* (i.e. 27%, Tautz et al., 1988) but in contrast, it was different to that observed in *Melipona* in which it varies from 49 to 54%. This A-T richness of the ITS sequence can be explained by processes as biased occurrence or fixation of point mutations. The presence of microsatellite loci located within the ITS-1 region is well documented in many organisms (see Harris and Crandall, 2000 for rationale) and has also been observed in *Melipona* subspecies, in the form of repeated elements of one to four nucleotides (Fernandes-Salomão et al., 2005). Despite the variation found in these regions, the ITS-1 region had the same sequence in eight of the ten *mellifera* subspecies studied and differed only in two, *A. m. scutellata* and *A. m. intermissa*, with exhibit a southern and northern African distribution, respectively.

The homogeneity observed in the ITS-1 region of the honeybee presumably results from the process of concerted evolution leading to a homogenization of the rDNA copies within a species. Vogler and DeSalle (1994) hypothesized that the homogenization proceeds more efficiently within a single chromosome than among different chromosomes, which is in congruence with the observations of Beye and Moritz (1993) concerning the location of the rDNA copies in only two chromosomes of the chromosomal set of the honeybee ($n = 16$). In contrast, a considerable amount of sequence divergence in the mitochondrial DNA molecule exists within *Apis mellifera*, which has been widely used to establish the phylogenetic relationships

Corresponding author: P. De La Rúa,
pdelarua@um.es

* Manuscript editor: Walter S. Sheppard

among the *A. mellifera* subspecies (Garnery et al., 1992; Arias and Sheppard, 1996; De la Rúa et al., 2005 and references therein). These results suggest that suitable molecular markers may vary notably within the order Hymenoptera. Each potential marker has to be carefully evaluated to test its suitability for phylogenetic or population use within particular taxa. In the case of *Apis mellifera* we conclude that the ITS-1 region is unlikely to be useful at the subspecific level. Detailed analyses of the ITS-2 region will provide a more complete resolution of the sequence variation of the intergenic transcribed spacers of honeybee rDNA.

ACKNOWLEDGEMENTS

The authors wish to thank the financial support of the projects BOS2003-09765 and API06-010COORD-2 and the Program Ramón y Cajal of the Spanish Ministry of Education and Science.

Note scientifique sur la région ITS-1 chez les sous-espèces d'*Apis mellifera*.

Zusammenfassung – Eine wissenschaftliche Notiz über die ITS-1 Region der Unterarten von *Apis mellifera*.

REFERENCES

- Álvarez J.M., Hoy M.A. (2002) Evaluation of the ribosomal ITS2 DNA sequences in separating closely related populations of the parasitoid *Ageniopsis* (Hymenoptera: Encyrtidae), *Ann. Entomol. Soc. Am.* 95, 250–256.
- Arias M.C., Sheppard W.S. (1996) Molecular phylogenetics of honeybee subspecies (*Apis mellifera* L.) inferred from mitochondrial DNA sequence, *Mol. Phylogenet. Evol.* 5, 557–566.
- Beye M., Moritz R.F.A. (1993) In situ hybridization of rDNA on chromosomes of the honeybee, *Apis mellifera* L., *Experientia* 49, 337–338.
- Caterino M.S., Cho S., Sperling F.A.H. (2000) The current state of insect molecular systematics: a thriving tower of Babel, *Annu. Rev. Entomol.* 45, 1–54.
- De La Rúa P., Hernández-García R., Jiménez Y., Galián J., Serrano J. (2005) Biodiversity of *Apis mellifera iberica* (Hymenoptera: Apidae) from north-eastern Spain assessed by mitochondrial analysis, *Insect Syst. Evol.* 36, 21–28.
- Fernandes-Salomão T.M., Rocha R.B., Campos L.A.O., Araújo E.F. (2005) The first internal transcribed spacer (ITS-1) of *Melipona* species (Hymenoptera, Apidae, Meliponini): characterization and phylogenetic analysis, *Insect. Soc.* 52, 11–18.
- Garnery L., Cornuet J.M., Solignac M. (1992) Evolutionary history of the honey bee *Apis mellifera* inferred from mitochondrial DNA analysis, *Mol. Ecol.* 1, 145–154.
- Harris D.J., Crandall K.A. (2000) Intra-genomic variation within ITS-1 and ITS2 of freshwater crayfishes (Decapoda: Cambaridae): implications for phylogenetic and microsatellite studies, *Mol. Biol. Evol.* 17, 284–291.
- Ji Y.J., Zhang D.X., He L.J. (2003) Evolutionary conservation and versatility of a new set of primers for amplifying the ribosomal internal transcribed spacer (ITS) regions in insects and other invertebrates, *Mol. Ecol. Notes* 3, 581–585.
- Navajas M., Lagnel J., Gutierrez J., Boursot P. (1998) Species-wide homogeneity of nuclear ribosomal ITS2 sequences in the spider mite *Tetranychus urticae* contrast with extensive mitochondrial *COI* polymorphism, *Heredity* 80, 742–752.
- Sheppard W.S., McPheron B.A. (1991) Ribosomal DNA diversity in Apidae, in: D. Smith R. (Ed.), *Diversity in the genus Apis*, Westview Press, Oxford, pp. 89–102.
- Tautz D., Hancock M., Webb D.A. (1988) Complete sequence of the rRNA genes of *Drosophila melanogaster*, *Mol. Biol. Evol.* 5, 366–376.
- Vogler A.P., DeSalle R. (1994) Evolution and phylogenetic information content of the ITS-1 region in the tiger beetle *Cicindela dorsalis*, *Mol. Biol. Evol.* 11, 393–405.

Figure 1. Sequence alignment of the partial 18S, complete ITS-1 and partial 5.8S sequence of the ten *A. mellifera* subspecies. Gaps are indicated by hyphens and identical nucleotides by dots. An asterisk indicates the location of the repeated unit.

	[18S→				
<i>A. m. mellifera</i> 1	AGTGAGGTCT	TCGGACTGGT	GCGCGGCCAA	TGTCTCGGCA	40
<i>A. m. mellifera</i> 2	
<i>A. m. iberiensis</i> 1	
<i>A. m. iberiensis</i> 2	
<i>A. m. caucasica</i>	
<i>A. m. carnica</i>	
<i>A. m. ligustica</i>	
<i>A. m. scutellata</i>	
<i>A. m. sahariensis</i>	
<i>A. m. intermissa</i>	
<i>A. m. jemenitica</i>	
<i>A. m. meda</i>	
<i>A. m. mellifera</i> 1	TTGCCGATGT	TACCGGGAAG	ATGACCAAAC	TTGATCATT	80
<i>A. m. mellifera</i> 2	
<i>A. m. iberiensis</i> 1	
<i>A. m. iberiensis</i> 2	
<i>A. m. caucasica</i>	
<i>A. m. carnica</i>	
<i>A. m. ligustica</i>	
<i>A. m. scutellata</i>	
<i>A. m. sahariensis</i>	
<i>A. m. intermissa</i>	
<i>A. m. jemenitica</i>	
<i>A. m. meda</i>	
<i>A. m. mellifera</i> 1	AGAGGAAGTA	AAAGTCGTAA	CAAGGTTTCC	GTAGGTGAAC	120
<i>A. m. mellifera</i> 2	
<i>A. m. iberiensis</i> 1	
<i>A. m. iberiensis</i> 2	
<i>A. m. caucasica</i>	
<i>A. m. carnica</i>	
<i>A. m. ligustica</i>	
<i>A. m. scutellata</i>	
<i>A. m. sahariensis</i>	
<i>A. m. intermissa</i>	
<i>A. m. jemenitica</i>	
<i>A. m. meda</i>	
	[ITS-1→				
<i>A. m. mellifera</i> 1	CTGCGGAAGG	ATCATTAACG	AATGAAACAT	ACAAAAAGAG	160
<i>A. m. mellifera</i> 2	
<i>A. m. iberiensis</i> 1	
<i>A. m. iberiensis</i> 2	
<i>A. m. caucasica</i>	
<i>A. m. carnica</i>	
<i>A. m. ligustica</i>	
<i>A. m. scutellata</i>	
<i>A. m. sahariensis</i>	
<i>A. m. intermissa</i>	
<i>A. m. jemenitica</i>	
<i>A. m. meda</i>	

Figure 1. Continued.

<i>A. m. mellifera</i> 1	AAAATGCGTA	ACAAAACAAA	CAGAGAATTG	AAAGAAGGGG	200
<i>A. m. mellifera</i> 2	
<i>A. m. iberiensis</i> 1	
<i>A. m. iberiensis</i> 2	
<i>A. m. caucasica</i>	
<i>A. m. carnica</i>	
<i>A. m. ligustica</i>	
<i>A. m. scutellata</i>A..-A..	
<i>A. m. sahariensis</i>	
<i>A. m. intermissa</i>R..R..	
<i>A. m. jemenitica</i>	
<i>A. m. meda</i>	
				*	
<i>A. m. mellifera</i> 1	ATGATAAATA	TATATATTTT	ATATATATAT	ATATTTATAA	240
<i>A. m. mellifera</i> 2	-----	
<i>A. m. iberiensis</i> 1	
<i>A. m. iberiensis</i> 2	-----	
<i>A. m. caucasica</i>	-----	
<i>A. m. carnica</i>	-----	
<i>A. m. ligustica</i>	-----	
<i>A. m. scutellata</i>	-----	
<i>A. m. sahariensis</i>	-----	
<i>A. m. intermissa</i>	-----	
<i>A. m. jemenitica</i>	-----	
<i>A. m. meda</i>	-----	
				[5.8S→	
<i>A. m. mellifera</i> 1	AATGGATTTT	TTGAATCCAT	TATCAAATAC	CAAACCTTTG	280
<i>A. m. mellifera</i> 2	
<i>A. m. iberiensis</i> 1	
<i>A. m. iberiensis</i> 2	
<i>A. m. caucasica</i>	
<i>A. m. carnica</i>	
<i>A. m. ligustica</i>	
<i>A. m. scutellata</i>	
<i>A. m. sahariensis</i>	
<i>A. m. intermissa</i>	
<i>A. m. jemenitica</i>	
<i>A. m. meda</i>	
<i>A. m. mellifera</i> 1	AACATCGACA	TTTCGAACCG	CACATA		306
<i>A. m. mellifera</i> 2		302
<i>A. m. iberiensis</i> 1		306
<i>A. m. iberiensis</i> 2		302
<i>A. m. caucasica</i>	-----	-----	-----		269
<i>A. m. carnica</i>		302
<i>A. m. ligustica</i>		302
<i>A. m. scutellata</i>		301
<i>A. m. sahariensis</i>	-----		292
<i>A. m. intermissa</i>		300
<i>A. m. jemenitica</i>		302
<i>A. m. meda</i>	-----	-----	-----		269