



# THE ROLE OF CHROMOSOME DEFECTS AND VARIANTS IN CATTLE BREEDING

B. Mayr, W. Schleger

## ► To cite this version:

B. Mayr, W. Schleger. THE ROLE OF CHROMOSOME DEFECTS AND VARIANTS IN CATTLE BREEDING. Annales de génétique et de sélection animale, 1978, 10 (4), pp.593-593. <hal-00893067>

**HAL Id: hal-00893067**

**<https://hal.science/hal-00893067v1>**

Submitted on 11 May 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

A perfect knowledge of the effects of chromosomal abnormalities in domestic animal requires an exact determination of the chromosome involved in each type of translocation as well as of the segregation pattern and the resulting proportion of unbalanced gametes.

From a practical point of view, the accurate identification by means of new banding methods of the chromosomes involved in each type of abnormality will lead to a better understanding of the meiotic behaviour, the segregation pattern and the consequences on animal production characters.

#### THE ROLE OF CHROMOSOME DEFECTS AND VARIANTS IN CATTLE BREEDING

B. MAYR and W. SCHLEGER

*Institut f. Tierzucht und Haustiergenetik  
der Vet.-med. Universität, Linke Bahngasse 11, 1030 Wien Austria*

Chromosomal polymorphism, in the form of a Robertsonian Translocation 1/29, was detected in 3 out of 280 individuals tested in Austria. The report deals with meiotic findings explaining the tendencies, in cattle, to centric fusions of this nature.

G-band testing of 50 bulls brought no results as to a new type of polymorphism. Reference is made to 4 bulls with XY/XX complement. No unbalanced complements have so far been observed in this country.

#### ROUTINE CHROMOSOMAL EXAMINATION OF AI BULLS IN HUNGARY,

M. PAPP and A. KOVACS

*Central Hungarian Station for Artificial  
Insemination 1440 Budapest 70, P.O.B. 19, Hungary*

In Hungary the routine chromosomal examinations started at the beginning of 1975. Till now, a total of 324 bulls were karyotypically evaluated in five A. I. stations. 313 bulls were found to be normal and 11 showed chromosomal abnormalities. The bulls belonged to eight pure breeds and three crossed ones. The number of bulls in the breeds examined were as follows: 76 *Hungarian Simmental*, 21 *Austrian Simmental*, 4 *German Simmental*, 174 *Holstein-Friesian*, 3 *Dutch Friesian*, 1 *Swedish Friesian*, 7 *Limousine*, 5 *Hereford*, 1 *Kostroma* and 32 cross-bred.

Chromosomal abnormalities were found in the *Hungarian Simmental*, *German Simmental* and *Holstein-Friesian* breeds. In *Hungarian Simmental* three cases of translocation (two 1/29 and one 14/21) and two chimaerisms, in *German Simmental* one 1/29 translocation, in *Holstein-Friesian* breed one case of mosaic 13/21 translocation (?) and four chimaerisms were disclosed.

#### PRELIMINARY STUDIES ON 6 TO 7 DAYS OLD BLASTOCYSTS NON-SURGICALLY RECOVERED FROM CATTLE, III. — METHOD FOR CHROMOSOME PREPARATION

W. A. KING\*, T. LINARÈS\*\*, J. GUSTAVSSON \* and A. BANE\*\*

*\* Dep. of Animal Breeding and Genetics  
The Swedish University of Agricultural Sciences,  
S-750 07 Uppsala 7, Sweden*

*\*\*Dep. of Obstetrics and Gynaecology, same University*

A simple technique for making chromosome preparation from 6 to 7 day old bovine blastocysts is described. It is noted that the morphological features of blastocysts and total number of cells greatly influence the quality of the preparation.