



THYROXINE AND TRIIODOTHYRONINE DEGRADATION IN LINES OF PIGS SELECTED FOR RATE OF GAIN AND THICKNESS OF BACKFAT

N. Standal, B. Tveit, A. Eggum, P. M. Dahl

► To cite this version:

N. Standal, B. Tveit, A. Eggum, P. M. Dahl. THYROXINE AND TRIIODOTHYRONINE DEGRADATION IN LINES OF PIGS SELECTED FOR RATE OF GAIN AND THICKNESS OF BACKFAT. *Annales de génétique et de sélection animale*, 1980, 12 (1), pp.126-126. <hal-00893175>

HAL Id: hal-00893175

<https://hal.science/hal-00893175v1>

Submitted on 11 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

THYROXINE AND TRIIODOTHYRONINE DEGRADATION
IN LINES OF PIGS SELECTED FOR RATE OF GAIN AND THICKNESS OF BACKFAT

N. STANDAL, B. TVEIT, A. EGGUM, P. M. DAHL

Dept of Animal Genetics and Breeding, Agricultural University of Norway, Ås - N.L.H.

A selection experiment in two directions has produced lines for pigs that differ considerably in fat/lean ratio and also in growth rate. It has been hypothesized that the selection for thin backfat may have altered the metabolic rate of the pigs.

In order to investigate this hypothesis, thyroid activity parameters were estimated in 12 pigs, from each of the two selection lines, at a live weight of about 40 kg.

There was a slight difference between the two selection lines in triiodothyronine degradation constant, and T_3 serum level. For the other tested parameters, T_3 and T_4 distribution space, T_3 and T_4 degradation/day, T_4 degradation constant and T_4 serum level there was no significant difference between the two selection lines. The line difference in percent dissected fat in the half carcass amounted to about 13 percentage units, and was very highly significant.

It was concluded that the thyroid activity measurements did not give clear evidence of a difference in metabolic rate between the two selection line.

DIFFICULTIES WITH THE GENETIC IMPROVEMENT OF FEED EFFICIENCY
IN BROILER LINES USING GROUP CAGE INFORMATION

D. K. FLOCK, F. MARAHRENS

Lohmann Tierzucht GmbH, Cuxhaven, W. Germany

Progeny or sib testing in group cages may become a useful tool for broiler breeders who wish to improve feed efficiency directly. Problem areas today are : (1) the investment and operating cost for measuring family feed consumption on a large enough scale to have an impact on the commercial broiler; (2) mortality, which tends to be higher than under floor management and contributes to the inaccuracies of measuring feed intake; (3) leg problems, which limit the possible selection intensity for growth rate and feed efficiency; and (4) the expected decrease in rate of progress for growth rate.

SELECTION ON FEED EFFICIENCY IN LAY HENS : DIRECT AND CORRELATED RESPONSE

G. HEIL*, F. PIRCHNER

Lehrstuhl für Tierzucht, Weiheustephan, Bayern

* Institut f. Kleintierzucht der Bundesforschungsanstalt f. Landwirtschaft, Braunschweig-Volkenræde
D-3101 Celle

Two *Leghorn*-lines were selected for feed efficiency over three generations with a reciprocal recurrent selection program. Selection response was estimated by using a repeated mating scheme. Furthermore a comparison could be made between generation 3 of this experiment and lines of the same origin selected for egg production. The following results could be observed. Body weight feed consumption and egg weight decreased due to selection for feed efficiency. A slight increase in the egg production rate could be seen. Feed efficiency changed from 2.46 in generation 0 to 2.22 in generation 3 in terms of kg feed used to produce one kg eggmass. The lines selected for egg production laid more and heavier eggs, had a higher body weight and a greater feed intake than the lines selected directly for feed efficiency. For the production of one kg eggmass they used 2.34 kg feed.